

Intracerebroventricular injection of L-proline and D-proline induces sedative and hypnotic effects by different mechanisms under an acute stressful condition in chicks

K. Hamasu · K. Shigemi · Y. Tsuneyoshi ·
H. Yamane · H. Sato · D. M. Denbow · M. Furuse

Received: 3 September 2008 / Accepted: 27 October 2008 / Published online: 21 November 2008
© Springer-Verlag 2008

Abstract The central effects of L-proline, D-proline and *trans*-4-hydroxy-L-proline were investigated by using the acute stressful model with neonatal chicks in Experiment 1. Sedative and hypnotic effects were induced by all compounds, while plasma corticosterone release under isolation stress was only attenuated by L-proline. To clarify the mechanism by which L-proline and D-proline induce sedative and hypnotic effects, the contribution of the strychnine-sensitive glycine receptor (glycine receptor) and *N*-methyl-D-aspartate glutamate receptor (NMDA receptor) were further investigated. In Experiments 2–3, the glycine receptor antagonist strychnine was co-injected intracerebroventricular (i.c.v.) with L-proline or D-proline. The suppression of isolation-induced stress behavior by D-proline was attenuated by strychnine. However, the suppression of stress behavior by L-proline was not attenuated. In Experiment 4, the NMDA receptor antagonist (+)-MK-801 was co-injected i.c.v. with L-proline. The suppression of stress behavior by L-proline was attenuated by (+)-MK-801. These results indicate that L-proline and D-proline

differentially induce sedative and hypnotic effects through NMDA and glycine receptors, respectively.

Keywords CNS · Stress · Strychnine · (+)-MK-801 · Strychnine-sensitive glycine receptor · NMDA receptor

Introduction

There are many studies suggesting that L-proline may be a neuronal modulator or transmitter candidate in the central nervous system (CNS) (Snyder et al. 1973; Yoneda and Roberts 1982; Fremeau et al. 1992; Renick et al. 1999; Gogos et al. 1999). We have previously demonstrated that the amount of L-proline in the brain was reduced under stressful conditions, and intracerebroventricular (i.c.v.) injection of L-proline had sedative and hypnotic effects under an acute stressful condition in neonatal chicks (Hamasu et al. 2009). From these facts, it appears that L-proline might function through several neurotransmitter systems in the CNS. However, the mechanism by which L-proline attenuates stress behavior is presently unknown.

The behavioral model is based on the response of neonatal chicks to acute stress (Panksepp et al. 1980; Sahley et al. 1981; Feltenstein et al. 2003a, b). In brief, chicks grown in crowds feel acute stress when they are isolated. This isolation-induced stress increases vocalization and plasma corticosterone concentration. This isolation-induced stress paradigm has been used as an objective model for anxiolytic drug screening, with vocalization and plasma corticosterone concentration as the index of the behaviors induced by the stressor.

L-Proline is an amino acid. However, it differs from other amino acids in that it has imino group and its side chain is bound to both nitrogen and α -carbon atoms.

K. Hamasu · K. Shigemi · Y. Tsuneyoshi · H. Yamane ·
M. Furuse (✉)
Laboratory of Advanced Animal and Marine Bioresources,
Graduate School of Bioresource and Bioenvironmental Sciences,
Kyushu University, Fukuoka 812-8581, Japan
e-mail: furuse@brs.kyushu-u.ac.jp

H. Sato
Ajinomoto Co., Inc., Kawasaki, Kawasaki 210-8681, Japan

D. M. Denbow
Department of Animal and Poultry Sciences,
Virginia Polytechnic Institute and State University,
Blacksburg, VA 24061-0306, USA

L-Pipecolic acid has the molecular structure that looks like L-proline very well. Takagi et al. (2001) demonstrated that i.c.v. injection of L-pipecolic acid induced sleep-like behavior in neonatal chicks. From these findings, it is suggested that these molecular structures are important in inducing sedative and hypnotic effects.

The purpose of the present study was (1) to confirm the effect of L-proline and proline-related compounds such as D-proline and *trans*-4-hydroxy-L-proline under the acute stressful model in neonatal chicks (Experiment 1), and (2) to determine which receptors mediate the actions of these compounds (Experiments 2–4).

The strychnine-sensitive glycine receptor (glycine receptor) is a pentameric ligand-gated anion channel composed of two different subunits, α and β . Also the *N*-methyl-D-aspartate glutamate receptor (NMDA receptor) is an excitatory amino acid ligand-gated ion channel that is selectively activated by NMDA and regulated at several pharmacologically distinct sites. This receptor is present in both the hypothalamus (Halpain et al. 1984) and pituitary (Lindstrom and Ohlsson 1992), suggesting its role in neuroendocrine regulation (Mahesh et al. 1999). Past studies showed that although L- and D-proline can activate the glycine receptor, the NMDA receptor is activated by L-proline alone (Henzi et al. 1992; Ortiz et al. 1996). In addition, glycine (Asechi et al. 2006) and L-glutamate (Yamane et al., unpublished data) caused a sedative and hypnotic effect similar to that observed with L-proline. So, we hypothesized that these receptors are involved in the action of proline to induce sedative and hypnotic effects which was investigated using the glycine receptor antagonist strychnine and NMDA receptor antagonist (+)-MK-801.

Materials and methods

Animals and food

One-day-old male layer chicks (Julia) purchased from a local hatchery (Murata Hatchery, Fukuoka, Japan) were maintained in a windowless room at a constant temperature of $30 \pm 1^\circ\text{C}$. Lighting was provided continuously for 24 h. Chicks were given free access to a commercial starter diet (Toyohashi Feed and Mills Co. Ltd, Aichi, Japan) and water. On the experimental day, chicks (4- or 5-day-old) were distributed into groups based on their body weight, so that the average body weight was as uniform as possible within the same experiment. Experimental procedures followed the guidance for Animal Experiments in the Faculty of Agriculture and in the Graduate Course of Kyushu University and the Law (No. 105) and Notification (No. 6) of the Japanese Government. The number of animals used and their discomfort were minimized.

Preparation of drugs

L-Proline and *trans*-4-hydroxy-L-proline were a gift from Kyowa Hakko Kogyo (Tokyo, Japan). The D-proline and glycine receptor antagonist strychnine and the NMDA receptor antagonist (+)-MK-801 maleate were purchased from Sigma (St Louis, MO) and Tocris Bioscience (Ellisville, USA), respectively. In Experiments 1 and 4, L-proline, D-proline, *trans*-4-hydroxy-L-proline, and (+)-MK-801 were dissolved in 0.85% saline containing 0.1% Evans Blue solution. In Experiments 2 and 3, strychnine was dissolved in 0.85% saline containing 5% dimethyl sulfoxide (DMSO) and 0.1% Evans Blue solution. L-Proline and D-proline were treated similarly.

Intracerebroventricular (i.c.v.) injection and behavioral observation

In all experiments, drugs were injected into the left lateral ventricle of the chicks using a microsyringe according to the method of Davis et al. (1979) and Koutoku et al. (2005). The stress and discomfort by this method is minimal as described elsewhere (Koutoku et al. 2005).

After the injection, chicks were placed in an acrylic glass monitoring cage (40 cm \times 30 cm \times 20 cm) with paper on the floor, and behavioral observations were made for 10 min at a constant temperature of $30 \pm 1^\circ\text{C}$. The monitoring systems were set in a separate room to avoid disturbing the animals. The number of distress vocalizations, which are shrill and intense calls, was simultaneously recorded and counted, using a computer with Gretchen software (Excla Inc., Japan). Chicks were recorded by three video cameras positioned in different directions. Based on the method by van Luijtelaaar et al. (1987), the behaviors were classified into four categories: (1) active wakefulness; (2) standing/sitting motionless with eyes opened; (3) standing motionless with eyes closed; and (4) sitting motionless with head drooped (sleeping posture) by watching the videotapes. They demonstrated the correlation between sleeping posture and electrophysiological sleep with EEG measurement (van Luijtelaaar et al. 1987). During the monitoring period, chicks were not given food or water. These behavioral experiments were conducted according to the methods of Takagi et al. (2001), Koutoku et al. (2005) and Asechi et al. (2006).

Experiment 1: Effects of L-proline, D-proline and *trans*-4-hydroxy-L-proline on the stress response under isolation-induced stress

Chicks were given i.c.v. injections of 0.50 μmol of L-proline, D-proline, *trans*-4-hydroxy-L-proline, or saline

as the control. This dosage of L-proline was based on a previous experiment (Hamasu et al. 2009). The dosage of D-proline and *trans*-4-hydroxy-L-proline was same as L-proline. After the behavioral observations, blood was collected in heparinized microtubes from the jugular vein. In addition, the intact group (with neither i.c.v. injection nor isolation stress) was also sampled. Blood was centrifuged at 4°C and 10,000×*g* for 4 min, and the plasma collected and stored at −30°C until analysis. Plasma corticosterone concentration was determined using a corticosterone enzyme immunoassay kit (Assay Designs Inc., MI, USA).

Experiment 2: Effects of strychnine on sedative and hypnotic behaviors induced by L-proline

Chicks were divided into four groups that received either saline, 2.99 nmol of strychnine, 0.50 μmol of L-proline, or strychnine plus L-proline. This dosage of strychnine was based on a previous experiment (Sigemi et al. 2008). After the injection, behavioral observations were conducted.

Experiment 3: Effects of strychnine on sedative and hypnotic behaviors induced by D-proline

This experiment was similar to Experiment 2 except that L-proline was replaced with D-proline (0.50 μmol).

Experiment 4: Effects of (+)-MK-801 on sedative and hypnotic behaviors induced by L-proline

Chicks were divided into four groups that received either saline, 0.50 nmol of (+)-MK-801, 0.50 μmol of L-proline, or (+)-MK-801 plus L-proline. This dosage of (+)-MK-801 was determined in a pilot study. After injection, behavioral observations were conducted.

Statistical analysis

The chicks were decapitated after an overdose of sodium pentobarbital. The brains were removed and location of the Evans Blue dye was confirmed. Data of chicks without dye in the lateral ventricle were deleted. In Experiment 1, data were statistically analyzed by one-way analysis of variance (ANOVA). In Experiments 2–4, data were statistically analyzed by two-way ANOVA. Tukey–Kramer test was used as a post hoc test. Statistical analysis was conducted using a commercially available package StatView (version 5, SAS Institute, Cary, USA, 1998).

Results

Experiment 1

Figure 1 shows the effect of i.c.v. injection of saline, L-proline, D-proline, *trans*-4-hydroxy-L-proline on distress vocalizations during the 10 min isolation-induced stress. Significant effects on total distress vocalizations ($F(3, 16) = 25.347$, $P < 0.001$) were detected. All drugs significantly decreased distress vocalizations compared to the control.

Figure 2 shows the effect of i.c.v. injection of saline, L-proline, D-proline, *trans*-4-hydroxy-L-proline on plasma corticosterone concentration immediately after 10 min of isolation. The effect of the drugs on plasma corticosterone concentration was significant ($F(4,19) = 5.046$, $P < 0.01$). Only L-proline significantly lowered the corticosterone level compared to the control group, and these values were as low as that of the intact group. D-Proline and *trans*-4-hydroxy-L-proline had no effect on plasma corticosterone levels.

Table 1 shows the effect of i.c.v. injection of L-proline, D-proline, and *trans*-4-hydroxy-L-proline on various behavioral categories of chicks during the 10 min behavior observation period under isolation stress. Significant effects were observed in active wakefulness ($F(3,16) = 40.296$, $P < 0.0001$) and sleeping posture ($F(3,16) = 17.850$, $P < 0.0001$). L-Proline, D-proline, and *trans*-4-hydroxy-L-proline induced sedative and hypnotic effects compared with the saline control.




Fig. 1 Effect of i.c.v. injection of L-proline, D-proline, and *trans*-4-hydroxy-L-proline (Hydroxyl-proline) on distress vocalizations during 10 min isolation in 4- or 5-day-old layer chicks. Results are expressed as means ± SEM. The number of chicks used in each group was five. Groups with different superscripts are significantly different ($P < 0.05$)




Fig. 2 Effect of i.c.v. injection of L-proline, D-proline, and *trans*-4-hydroxy-L-proline (hydroxyl-proline) on plasma corticosterone concentration after isolation stress for 10 min in 4- or 5-day-old layer chicks. Results are expressed as means \pm SEM. The number of chicks used in each group was as follows: control, five; L-proline, five; D-proline, five; *trans*-4-hydroxy-L-proline, five; intact, four. Groups with different superscripts are significantly different ($P < 0.05$)

Experiment 2

Figure 3 shows the effect of i.c.v. injection of L-proline with or without strychnine on distress vocalizations during the 10 min isolation-induced stress. A significant effect of L-proline ($F(1,19) = 31.049$, $P < 0.0001$) was detected, but no significant effect was observed with strychnine ($F(1, 19) = 0.046$, $P > 0.05$). The lack of a significant ($F(1, 19) = 1.338$, $P > 0.05$) interaction between L-proline and strychnine implies that the effect of L-proline could not be attenuated by strychnine.

The effect of i.c.v. injection of strychnine on behaviors induced by L-proline is shown in Table 2. Interactions between L-proline and strychnine were not significant for any category indicating that strychnine did not alter behaviors induced by L-proline.




Fig. 3 Effect of i.c.v. injection of either 2.99 nmol of strychnine, 0.50 μ mol of L-proline, or strychnine plus L-proline on distress vocalizations during 10 min isolation in 4- or 5-day-old layer chicks. Results are expressed as means \pm SEM. The number of chicks used in each group was as follows: control, five; strychnine, six; L-proline, six; strychnine + L-proline, six

Experiment 3

Figure 4 shows the effect of i.c.v. injection of D-proline with or without strychnine on distress vocalizations during the 10 min isolation-induced stress. Significant effects of D-proline ($F(1,22) = 7.256$, $P < 0.05$) and strychnine ($F(1, 22) = 4.627$, $P < 0.05$) were detected. An interaction between D-proline and strychnine was nearly significant ($F(1,22) = 3.482$, $P = 0.075$) suggesting that the suppressive effect of D-proline on distress-induced vocalizations was moderately attenuated by co-injected of strychnine.

The effect of i.c.v. injection of strychnine on behaviors induced by D-proline is shown in Table 3. Interactions between D-proline and strychnine were significant in the following categories: active wakefulness ($F(1,22) = 6.913$, $P < 0.05$) and sitting motionless with head drooped (sleeping posture) ($F(1,22) = 19.330$, $P < 0.001$). D-Proline increased sitting motionless with head drooped (sleeping posture) and decreased active wakefulness, and strychnine attenuated these effects. Since strychnine did not affect the behavioral categories under the control conditions, these

Table 1 Influence of i.c.v. injections of saline, L-proline, D-proline, or *trans*-4-hydroxy-L-proline on various behavioral categories of chicks 10 min post-injection

	Control	L-Proline	D-Proline	<i>Trans</i> -4-hydroxy-L-proline
Active wakefulness	474 \pm 39 ^a	64 \pm 41 ^b	7 \pm 7 ^b	45 \pm 39 ^b
Standing/sitting motionless with eyes open	126 \pm 39	152 \pm 25	155 \pm 52	158 \pm 23
Standing motionless with eyes closed	0 \pm 0	0 \pm 0	25 \pm 25	0 \pm 0
Sitting motionless with head drooped (sleeping posture)	0 \pm 0 ^a	384 \pm 37 ^b	413 \pm 78 ^b	397 \pm 38 ^b
Total (s)	600	600	600	600

Values are means \pm SEM in seconds. The number of chicks used in each group was five. Groups with different letters are significantly different ($P < 0.05$)

Table 2 Influence of i.c.v. injections of strychnine and L-proline on various behavioral categories of chicks 10 min post-injection

Strychnine	Saline		L-Proline		<i>P</i>	L-Proline	Strychnine	L-Proline × Strychnine
	-	+	-	+				
Active wakefulness	508 ± 42 ^a	521 ± 24 ^a	185 ± 63 ^b	240 ± 83 ^b	<0.001	NS	NS	NS
Standing/sitting motionless with eyes open	89 ± 39	74 ± 21	170 ± 55	130 ± 23	NS	NS	NS	NS
Standing motionless with eyes closed	3 ± 3	5 ± 5	132 ± 60	109 ± 83	<0.05	NS	NS	NS
Sitting motionless with head drooped (sleeping posture)	0 ± 0	0 ± 0	113 ± 67	121 ± 68	<0.05	NS	NS	NS
Total (s)	600	600	600	600				

Values are means ± SEM in seconds. The numbers of chicks used were: saline, five; strychnine, six; L-proline, six; L-proline + strychnine, six. NS, not significant. Groups with different letters are significantly different ($P < 0.05$)




Fig. 4 Effect of i.c.v. injection of either 2.99 nmol of strychnine, 0.50 μmol of D-proline, or strychnine plus D-proline on distress vocalizations during 10 min isolation in 4- or 5-day-old layer chicks. Results are expressed as means ± SEM. The number of chicks used in each group was as follows: control, seven; strychnine, six; D-proline, five; strychnine + D-proline, eight

results indicate that strychnine altered behaviors induced by D-proline. Sleeping posture induced by D-proline was attenuated by co-injection with strychnine and consequently, the behavior of chicks tended toward wakefulness.

Experiment 4

Figure 5 shows the effect of i.c.v. injection of L-proline with or without (+)-MK801 on distress vocalizations during the 10 min isolation-induced stress. Significant effects of L-proline ($F(1, 26) = 8.530$, $P < 0.01$) were detected, but the effect of (+)-MK801 was not significant ($F(1,26) = 1.390$, $P > 0.05$). A significant ($F(1, 26) = 11.708$, $P < 0.01$) interaction between L-proline and (+)-MK801 suggested that the suppressive effect of L-proline on distress-induced vocalizations was attenuated by co-injected of (+)-MK801.

The effect of i.c.v. injection of (+)-MK801 on behaviors induced by L-proline is shown in Table 4. The interactions between L-proline and (+)-MK801 were significant in the

following categories: active wakefulness ($F(1, 26) = 19.217$, $P < 0.001$), standing/sitting motionless with eyes opened ($F(1,26) = 6.876$, $P < 0.05$) and sitting motionless with head drooped (sleeping posture) ($F(1, 26) = 11.725$, $P < 0.01$). These results indicate that (+)-MK801 altered behaviors induced by L-proline. Sleeping posture induced by L-proline was attenuated by co-injection with (+)-MK801 and consequently, the behavior of chicks tended toward wakefulness.

Discussion

In order to examine whether D-proline and *trans*-4-hydroxy-L-proline have sedative and hypnotic effects similar to L-proline, all drugs were centrally administered and then the behavior and plasma corticosterone concentration of neonatal chicks under isolation-induced stress were determined in Experiment 1. D-Proline and *trans*-4-hydroxy-L-proline as well as L-proline attenuated isolation stress-induced behaviors during the 10 min post-injection period (Fig. 1; Table 1). However, L-proline, but not D-proline or *trans*-4-hydroxy-L-proline, significantly lowered plasma corticosterone level compared to the control group (Fig. 2). Therefore, L-proline might have the ability to directly suppress the secretion of corticosterone from the adrenal glands through the hypothalamic-pituitary-adrenal (HPA) axis in response to stress. On the other hand, D-proline and *trans*-4-hydroxy-L-proline appear not to have such an effect. As a result, the mechanism of sedative and hypnotic effects of L-proline in chicks appears different from D-proline and *trans*-4-hydroxy-L-proline. Therefore, the mechanism of L-proline and D-proline was examined in Experiments 2–4.

L-Proline has long been recognized as a neuronal modulator or transmitter candidate in the CNS (Snyder et al. 1973). In addition, Henzi et al. (1992) revealed that L- and D-proline can activate the glycine receptor, while L-proline alone can also activate the NMDA receptor. This

Table 3 Influence of i.c.v. injections of strychnine and D-proline on various behavioral categories of chicks 10 min post-injection

Strychnine	Saline		D-Proline		<i>P</i>	D-Proline	Strychnine	D-Proline × Strychnine
	-	+	-	+				
Active wakefulness	434 ± 48 ^a	443 ± 86 ^a	71 ± 45 ^b	397 ± 51 ^a	<0.01	<0.05	<0.05	<0.05
Standing/sitting motionless with eyes open	127 ± 29	76 ± 36	166 ± 39	102 ± 32	NS	NS	NS	NS
Standing motionless with eyes closed	39 ± 22	43 ± 28	10 ± 10	64 ± 34	NS	NS	NS	NS
Sitting motionless with head drooped (sleeping posture)	0 ± 0 ^a	38 ± 24 ^a	353 ± 90 ^b	37 ± 28 ^a	<0.001	<0.01	<0.01	<0.001
Total (s)	600	600	600	600				

Values are means ± SEM in seconds. The numbers of chicks used were: saline, seven; strychnine, six; D-proline, five; D-proline + strychnine, eight. NS, not significant. Groups with different letters are significantly different ($P < 0.05$)




Fig. 5 Effect of i.c.v. injection of either 0.50 nmol of (+)-MK-801, 0.50 μmol of L-proline, or strychnine plus L-proline on distress vocalizations during 10 min isolation in 4- or 5-day-old layer chicks. Results are expressed as means ± SEM. The number of chicks used in each group was as follows: control, seven; (+)-MK-801, eight; L-proline, seven; (+)-MK-801 + L-proline, eight. Groups with different superscripts are significantly different ($P < 0.05$)

suggests that these receptors might be involved in the mechanisms whereby L- and D-proline induce sedative and hypnotic effects. To examine these possibilities, the effect of co-injection of L- or D-proline with the glycine receptor antagonist strychnine on behavior of neonatal chicks under isolation-induced stress was investigated. Sedative and hypnotic effects induced by L-proline were not attenuated by co-injection of strychnine (Fig. 3; Table 2). However, effects induced by D-proline were partially attenuated by strychnine (Fig. 4; Table 3). The i.c.v. injection of 2.99 nmol of strychnine alone dose not influence isolation stress-induced behaviors (Shigemi et al., unpublished data). These results indicate that the glycine receptor is involved in sedative and hypnotic effects induced by D-proline, but not L-proline. In vitro, L-proline can activate the glycine receptor (Henzi et al. 1992; Ortiz et al. 1996). However,

this receptor is not involved in the action of L-proline to induce sedative and hypnotic effects in vivo.

Previous studies reported that this receptor is playing important roles in controlling motor functions, sensory signaling in vision and audition (Lopez-Corcuera et al. 2001) and inflammatory pain sensitization (Harvey et al. 2004). However, few studies have examined its role in the stress response. Our results suggest that the glycine receptor might be involved in part of the stress response.

Secondly, the effect of co-injection of L-proline with the NMDA receptor antagonist (+)-MK-801 on behavior of neonatal chicks under isolation-induced stress was investigated. The sedative and hypnotic effects induced by L-proline were attenuated by co-injection of (+)-MK-801 (Fig. 5; Table 4). These results indicate that the NMDA receptor is involved in sedative and hypnotic effects induced by L-proline.

The NMDA receptor is well established as playing important roles in neuroendocrine regulation (Mahesh et al. 1999). In fact, NMDA stimulates corticotrophin-releasing hormone release from rat hypothalamic slices, and this effect is inhibited by an NMDA receptor antagonist (Joanny et al. 1997). Several studies examined the role of the NMDA receptor in vivo. Systemic administration of NMDA enhanced adrenocorticotropin (ACTH) and serum corticosterone secretion (Farah et al. 1991), and the secretion of ACTH in response to immobilization stress was inhibited by a NMDA centrally acting receptor antagonist (Jezova et al. 1995). Thus, this receptor is involved in the regulation of the HPA axis. In our study, the increase in plasma corticosterone induced by isolation stress was suppressed with i.c.v. injections of L-proline. Presumably, this effect was mediated via the NMDA receptor activity which was modulated by L-proline.

Previous studies indicated that activation of dopaminergic and serotonergic pathways are related to behavioral and emotional changes (Abercrombie et al. 1989; Gruss et al. 1996). For example, isolation-induced stress resulted in a significant increase of 5-hydroxyindoleacetic acid and

Table 4 Influence of i.c.v. injections of MK801 and L-proline on various behavioral categories of chicks 10 min post-injection

MK801	Saline		L-Proline		<i>P</i>	L-Proline	MK801	L-Proline × MK801
	–	+	–	+				
Active wakefulness	512 ± 46 ^a	343 ± 66 ^a	110 ± 38 ^b	407 ± 52 ^a	<0.01	NS		<0.001
Standing/sitting motionless with eyes open	43 ± 15 ^a	193 ± 41 ^b	189 ± 31 ^b	157 ± 39 ^{ab}	NS	NS		<0.05
Standing motionless with eyes closed	40 ± 27	34 ± 24	77 ± 25	36 ± 18	NS	NS		NS
Sitting motionless with head drooped (sleeping posture)	5 ± 5 ^a	30 ± 17 ^a	224 ± 75 ^b	0 ± 0 ^a	<0.05		<0.05	<0.01
Total (s)	600	600	600	600				

Values are means ± SEM in seconds. The numbers of chicks used were: saline, seven; MK801, eight; L-proline, seven; L-proline + MK801, eight. NS, not significant. Groups with different letters are significantly different ($P < 0.05$)

only a slight increase of homovanillic acid in the forebrain of domestic chicks (Gruss et al. 1996). Additionally, some experiments suggested a modulatory interaction between the glutamatergic and monoaminergic pathway, which is mediated via NMDA receptor activation (Gruss et al. 1999). Arco and Mora (2001) showed that local activation of prefrontal NMDA receptors reduced the stress-induced dopamine release in the prefrontal cortex of the rat. From these findings, we assumed that L-proline modulates activation of NMDA receptors and alters activation of dopaminergic and serotonergic pathways. Through this mechanism, i.c.v. injection of L-proline might induce sedative and hypnotic effects under an acute stressful condition in neonatal chicks. Indeed, i.c.v. injection of L-proline suppressed the increase in serotonin turnover rate and slightly suppressed the increase in dopamine turnover rate induced by isolation stress in the telencephalon (Hamasu et al., unpublished data). However, the effect of L-proline on the stress-induced monoamines release is still obscure. Further study is necessary to clarify this potential.

In conclusion, the i.c.v. injection of L-proline and D-proline induces sedative and hypnotic effects under an acute stressful condition in neonatal chicks. However, L-proline, but not D-proline significantly lowered plasma corticosterone level. Probably, it is attributed to difference of action mechanism. The mechanism of attenuation of the acute stress response by L-proline is mediated by the modulation of NMDA receptor in the CNS. On the contrary, D-proline is mediated by the modulation of glycine receptor.

Acknowledgments This work was supported by a Grant-in-Aid for Scientific Research from Japan Society for the Promotion of Science (No. 18208023) and the SKYLARK Food Science Institute.

References

- Abercrombie ED, Keefe KA, Difrischia DS, Zigmond MJ (1989) Differential effect of stress on in vivo dopamine release in striatum, nucleus accumbens, and medial frontal cortex. *J Neurochem* 52:1655–1658. doi:10.1111/j.1471-4159.1989.tb09224.x
- Arco AD, Mora F (2001) Dopamine release in the prefrontal cortex during stress is reduced by the local activation of glutamate receptors. *Brain Res Bull* 56:125–130. doi:10.1016/S0361-9230(01)00616-5
- Asechi M, Tomonaga S, Tachibana T, Han L, Hayamizu K, Denbow DM, Furuse M (2006) Intracerebroventricular injection of L-serine analogs and derivatives induces sedative and hypnotic effects under an acute stressful condition in neonatal chicks. *Behav Brain Res* 170:71–77. doi:10.1016/j.bbr.2006.02.005
- Bhattacharjee A, Bansal M (2005) Collagen structure the Madras triple helix and the current scenario. *IUBUB Life* 57:161–172. doi:10.1080/15216540500090710
- Davis JL, Masuoka DT, Gerbrandt LK, Cherkin A (1979) Autoradiographic distribution of L-proline in chicks after intracerebral injection. *Physiol Behav* 22:693–695. doi:10.1016/0031-9384(79)90233-6
- Delport M, Maas S, Van der merwe SW, Laurens JB (2004) Quantitation of hydroxyproline in bone by gas chromatography–mass spectrometry. *J Chromatogr A* 804:345–351
- Farah JM Jr, Rao TS, Mick SJ, Coyne KE, Iyengar S (1991) N-methyl-D-aspartate treatment increases circulating adrenocorticotropic and luteinizing hormone in the rat. *Endocrinology* 128:1875–1880
- Feltenstein MW, Lambdin LC, Ganzera M, Ranjith H, Dharmaratne W, Nanayakkara NP (2003a) Anxiolytic properties of Piper methysticum extract samples and fractions in the chick social-separation-stress procedure. *Phytother Res* 17:210–216. doi:10.1002/ptr.1107
- Feltenstein MW, Lambdin LC, Webb HE, Warnick JE, Khan SI, Khan IA, Acevedo ED, Sufka KJ (2003b) Corticosterone response in the chick separation-stress paradigm. *Physiol Behav* 78:489–493. doi:10.1016/S0031-9384(03)00030-1
- Freneau RT Jr, Caron MG, Blakely RD (1992) Molecular cloning and expression of a high affinity L-proline transporter expressed in putative glutamatergic pathways of rat brain. *Neuron* 8:915–926. doi:10.1016/0896-6273(92)90206-S
- Gogos JA, Sautha M, Takacs Z, Beck KD, Luine V, Lucas LR, Nadler JV, Karayiorgou M (1999) The gene encoding proline dehydrogenase modulates sensorimotor gating in mice. *Nat Genet* 21:434–439. doi:10.1038/7777
- Gruss M, Braun K (1996) Distinct activation of monoaminergic pathways in chick brain in relation to auditory imprinting and stressful situations: a microdialysis study. *Neuroscience* 76:891–899. doi:10.1016/S0306-4522(96)00385-5

- Gruss M, Bredenkotter M, Braun K (1999) *N*-methyl-D-aspartate receptor-mediated modulation of monoaminergic metabolites and amino acids in the chicks forebrain: an in vivo microdialysis and electrophysiology study. *J Neurobiol* 40:116–135. doi:10.1002/(SICI)1097-4695(199907)40:1<116::AID-NEU10>3.0.CO;2-M
- Halpain S, Wiczorek CM, Rainbow TC (1984) Localization of L-glutamate receptors in rat brain by quantitative autoradiography. *J Neurosci* 4:2247–2258
- Hamasu K, Haraguchi T, Kabuki Y, Adachi N, Tomonaga S, Sato H, Denbow DM, Furuse M (2009) L-Proline is a sedative regulator of acute stress in the brain of neonatal chicks. *Amino Acids* (in press)
- Harvey R, Depner U, Wassle H, Ahmadi S, Heindl C, Reinold H, Smart T, Harvey K, Schutz B, Akbari O, Zimmer A, Poisbeau P, Welzl H, Wolfer H, Betz H, Zeilhofer U, Muller U (2004) Glycine receptor $\alpha 3$: an essential target for spinal PGE₂-mediated inflammatory pain sensitization. *Science* 304:884–888. doi:10.1126/science.1094925
- Henzi V, Reichling DB, Helm SW, Macdermott AB (1992) L-Proline activates glutamate and glycine receptors in cultured rat dorsal horn neurons. *Mol Pharmacol* 41:793–801
- Jazova D, Tokarev D, Rusnak M (1995) Endogenous excitatory amino acids are involved in stress-induced adrenocorticotropin and catecholamine release. *Neuroendocrinology* 62:326–332. doi:10.1159/000127021
- Joanny P, Steinberg J, Oliver C, Grino M (1997) Glutamate and *N*-methyl-D-aspartate stimulate rat hypothalamic corticotropin-releasing factor secretion in vitro. *J Neuroendocrinol* 9:93–97. doi:10.1046/j.1365-2826.1997.00548.x
- Koutoku T, Takahashi H, Tomonaga S, Oikawa D, Saito S, Tachibana T, Han L, Hayamizu K, Denbow DM, Furuse M (2005) Central administration of phosphatidylserine attenuates isolation stress-induced behavior in chicks. *Neurochem Int* 47:183–189. doi:10.1016/j.neuint.2005.03.006
- Lindstrom P, Ohlsson L (1992) Effect of *N*-methyl-D, L-aspartate on isolated rat somatotrophs. *Endocrinology* 131:1903–1907. doi:10.1210/en.131.4.1903
- Lopez-Corcuera B, Geerlings A, Aragon C (2001) Glycine neurotransmitter transporters: an update. *Mol Membr Biol* 18:13–20
- Mahesh VB, Zamorano P, Sevilla LD, Lewis D, Brann DV (1999) Characterization of ionotropic glutamate receptors in rat hypothalamus, pituitary and immortalized gonadotropin-releasing hormone (GnRH) neurons (GT1–7 cells). *Neuroendocrinology* 69:397–407. doi:10.1159/000054442
- Messia MC, Falco TD, Panfili G, Marconi E (2008) Rapid determination of collagen in meat-based foods by microwave hydrolysis of proteins and HPAEC–PAD analysis of 4-hydroxyproline. *Meat Sci* (in press)
- Ortiz JG, Cordero ML, Rosado A (1996) Proline-glutamate interactions in the CNS. *Prog Neuro-Psychoph* 21:141–152. doi:10.1016/S0278-5846(96)00164-9
- Panksepp J, Bean NJ, Bishop P, Vilberg T, Sahley TL (1980) Opioid blockade and social comfort in chicks. *Pharmacol Biochem Behav* 13:673–683. doi:10.1016/0091-3057(80)90011-8
- Powell J (2006) Skin physiology. *Women Health Med* 3:130–133. doi:10.1383/wohm.2006.3.3.130
- Renick SE, Kleven DT, Chan J, Stenius K, Milner TA, Pickel VM, Freneau RT Jr (1999) The mammalian brain high-affinity L-proline transporter is enriched preferentially in synaptic vesicles in a subpopulation of excitatory nerve terminals in rat forebrain. *J Neurosci* 19:21–33
- Sahley TL, Panksepp J, Zolovick AJ (1981) Cholinergic modulation of separation distress in the domestic chick. *Eur J Pharmacol* 72:261–264. doi:10.1016/0014-2999(81)90283-1
- Sigemi K, Tsuneyoshi Y, Hamasu K, Han L, Hayamizu K, Denbow PM, Furuse M (2008) L-Serine induces sedative and hypnotic effects acting at GABA (A) receptors in neonatal chicks. *Eur J Pharmacol* 599:86–90
- Snyder SH, Young AB, Bennett JP, Mulder AH (1973) Synaptic biochemistry of amino acids. *Fed Proc* 32:2039–2047
- Takagi T, Ando R, Ohgushi A, Yamashita T, Dobashi E, Hussain-Yusuf H, Onodera R, Bungo T, Sato H, Furuse M (2001) Intracerebroventricular injection of pipercolic acid inhibits food intake and induces sleeping-like behaviors in the neonatal chick. *Neurosci Lett* 310:97–100. doi:10.1016/S0304-3940(01)02059-6
- Tanida M, Katsuyama M, Sakatani K (2007) Relation between mental stress-induced prefrontal cortex activity and skin conditions: a near-infrared spectroscopy study. *Brain Res* 1184:210–216. doi:10.1016/j.brainres.2007.09.058
- Uitto J (1971) Collagen biosynthesis in human skin. *Ann Clin Res* 3:250–258
- Van Luijckelaar ELJM, van der Grinten CPM, Blokhuis HJ, Coenen AML (1987) Sleep in the domestic hen (*Gallus domesticus*). *Physiol Behav* 41:409–414. doi:10.1016/0031-9384(87)90074-6
- Yoneda S, Roberts E (1982) A new synaptosomal biosynthetic pathway of proline from ornithine and its negative feedback inhibition by proline. *Brain Res* 239:479–488. doi:10.1016/0006-8993(82)90523-6