

Acute hyperammonemia activates branched-chain amino acid catabolism and decreases their extracellular concentrations: different sensitivity of red and white muscle

Milan Holecek · Roman Kandar · Ludek Sispera ·
Miroslav Kovarik

Received: 10 November 2009 / Accepted: 25 June 2010 / Published online: 8 July 2010
© Springer-Verlag 2010

Abstract Hyperammonemia is considered to be the main cause of decreased levels of the branched-chain amino acids (BCAA), valine, leucine, and isoleucine, in liver cirrhosis. In this study we investigated whether the decrease in BCAA is caused by the direct effect of ammonia on BCAA metabolism and the effect of ammonia on BCAA and protein metabolism in different types of skeletal muscle. M. soleus (SOL, slow-twitch, red muscle) and m. extensor digitorum longus (EDL, fast-twitch, white muscle) of white rat were isolated and incubated in a medium with or without 500 μ M ammonia. We measured the exchange of amino acids between the muscle and the medium, amino acid concentrations in the muscle, release of branched-chain keto acids (BCKA), leucine oxidation, total and myofibrillar proteolysis, and protein synthesis. Hyperammonemia inhibited the BCAA release (81% in SOL and 60% in EDL vs. controls), increased the release of BCKA (133% in SOL and 161% in EDL vs. controls) and glutamine (138% in SOL and 145% in EDL vs. controls), and increased the leucine oxidation in EDL (174% of controls). Ammonia also induced a significant increase in glutamine concentration in skeletal muscle. The effect of ammonia on intracellular BCAA concentration, protein synthesis and on total and myofibrillar proteolysis was insignificant. The data indicates that hyperammonemia

directly affects the BCAA metabolism in skeletal muscle which results in decreased levels of BCAA in the extracellular fluid. The effect is associated with activated synthesis of glutamine, increased BCAA oxidation, decreased release of BCAA, and enhanced release of BCKA. These metabolic changes are not directly associated with marked changes in protein turnover. The effect of ammonia is more pronounced in muscles with high content of white fibres.

Keywords Ammonia · Branched-chain amino acids · Leucine metabolism · Protein metabolism · Encephalopathy

Abbreviations

BCAA	Branched-chain amino acids
BCKA	Branched-chain keto acids
BCA-CoA	Branched-chain acyl-CoA
α -KG	Alpha-ketoglutarate
Glu	Glutamate
Gln	Glutamine
Ala	Alanine
ECF	Extracellular fluid
SA _{Leu}	Leucine-specific activity

M. Holecek (✉) · L. Sispera · M. Kovarik
Department of Physiology, Medical Faculty in Hradec Kralove,
Charles University in Prague, Simkova 870,
500 38 Hradec Kralove, Czech Republic
e-mail: holecek@lfhk.cuni.cz

R. Kandar
Department of Biological and Biochemical Sciences,
Faculty of Chemical Technology, University of Pardubice,
Pardubice, Czech Republic

Introduction

The branched-chain amino acids (BCAA), valine, leucine, and isoleucine, are essential amino acids which exert a specific regulatory effect on the rates of protein synthesis and degradation (Tischler et al. 1982; Nair and Short 2005). The decrease in plasma levels of these BCAA is a characteristic abnormality of patients with liver cirrhosis,

and the decreased BCAA levels are considered an important pathogenic factor in hepatic encephalopathy and protein-energy malnutrition development (Iber et al. 1957; Fischer and Baldessarini 1971; Blonde-Cynober et al. 1994; Holecek et al. 1996a, b). These alterations in BCAA concentrations and their metabolic properties serve as the rationale for the recommendation to use the BCAA-supplemented diets in treatment of different forms of liver disease (Fischer et al. 1976; Marchesini et al. 2003; Nakaya et al. 2007; Urata et al. 2007).

The pathogenesis of BCAA deficiency in patients with chronic liver disease is not completely clear. Several factors have been proposed as the cause of decreased plasma BCAA levels, including hyperinsulinemia, hyperglucagonemia, catecholamines, and starvation (Soeters and de Boer 1984; Marchesini et al. 1979; Eriksson et al. 1982; Hayashi et al. 1981; Yamato et al. 1995). Now it appears that ammonia has a crucial role. Several studies have shown an inverse relationship between the plasma ammonia and BCAA concentrations and/or that ammonia infusion decreases BCAA levels both in blood and in tissues (Hayashi et al. 1981; Leweling et al. 1996; Yamato et al. 1995; Holecek et al. 2000b). It was suggested that skeletal muscle takes up ammonia from the blood and detoxifies it via the synthesis of glutamine from glutamate. BCAA deficiency may result from the intensified synthesis of glutamate from α -ketoglutarate and BCAA (Leweling et al. 1996). However, the direct evidence of the supposed effect of ammonia on BCAA metabolism in skeletal muscle is absent.

Considering the physiological functions and pharmacological properties of BCAA, the relationship between ammonia and the development of protein-energy malnutrition, characterized by loss of energy stores and by wasting of skeletal muscle, also seems important. The prevalence of cachexia in patients with liver cirrhosis is about 50%, its pathogenesis is not fully understood, and effective therapeutic possibilities are not known (Peng et al. 2007).

The main objective of this study was to test the hypothesis whether the reduction of BCAA levels in the extracellular fluid of cirrhotic patients is caused by a direct effect of hyperammonemia on the BCAA metabolism in muscle, excluding the effect of neurohumoral changes associated with hyperammonemia, such as alterations in glucose and insulin concentrations and changes in pH (Fernandez et al. 1988). Therefore, we studied the effect of ammonia on BCAA metabolism using the *in vitro* technique. We examined the effect of ammonia addition into the incubation medium on amino acid concentrations in skeletal muscle, on amino acid exchange between the muscle and the extracellular fluid, on BCAA oxidation, and on release of branched-chain keto acids (BCKA) from the

muscle. To solve the question of possible effects of hyperammonemia on protein metabolism, the changes both in protein synthesis and proteolysis were examined. As demonstrated in several studies, muscles mostly composed of white (fast-twitch) fibres have different metabolic features compared to muscles mostly composed of red (slow-twitch) fibres (Hasselgren et al. 1986; Muthny et al. 2008). Concerning possible differences in BCAA metabolism between red and white skeletal muscle regarding ammonia, the higher BCAA aminotransferase activity may be of importance in the red muscle (Yang and Birkhahn 1997) and higher glutamine synthetase activity in the white one (Hickson et al. 1996). Therefore, our studies were performed in both types of skeletal muscle, in *m. soleus* (SOL, slow-twitch, red muscle) and in *m. extensor digitorum longus* (EDL, fast-twitch, red muscle).

Materials and methods

Animals

Male Wistar rats (body weight 40–60 g) obtained from BioTest, Konarovice, CZ, were used in this study. The rats were housed under controlled conditions (12-h light–dark cycle, 22°C, 55–65% relative humidity) with free access to standard laboratory chow and water. All procedures involving animal manipulation were performed in accordance with guidelines set by the Institutional Animal Use and Care Committee of Charles University, Prague, CZ.

Materials

L-[1-¹⁴C]leucine was purchased from GE Healthcare Life Sciences (Buckinghamshire, UK); amino acids, cycloheximide, Folin–Ciocalteu's phenol reagent, *N*-succinyl-Leu-Leu-Val-Tyr-7-amido-4-methylcoumarin (Suc-LLVY-MCA), and albumin were purchased from Sigma Chemical (St Louis, MO, US); Aminoplasmal 15 from B. Braun Medicals (Melsungen, DE). Hyamine hydroxide was obtained from Packard Instrument (Meriden, CT). The remaining chemicals were obtained from Sigma Chemicals (St Louis, MO, US), Waters (Milford, MA, US), and Lachema (Brno, CZ).

Experimental design

The animals were killed in pentobarbital narcosis (6 mg/100 g body weight, intraperitoneally) by exsanguination via abdominal aorta. Soleus (SOL) and extensor digitorum longus (EDL) muscles of both legs were dissected according to Maizels et al. (1977) and fixed via the tendons to stainless steel clips to provide slight tension (at

approximately resting length), and immediately transferred into 2.5 ml of modified Krebs–Heinseleit bicarbonate buffer with 6 mM glucose and 2 mU/ml insulin (pH 7.4, 37°C). The medium was saturated with O₂/CO₂ (19:1).

The muscles were preincubated for 30 min in a thermostatically controlled bath (37°C) with a shaking device (70 cycles/min) to ensure stable intramuscular concentrations of components present in the medium. After the preincubation, the muscles were quickly rinsed in 0.9% NaCl, blotted and transferred to a second set of vials containing fresh incubation media enriched with an ammonium acetate/bicarbonate buffer to reach 500 μM ammonia in the medium (ammonia group) or sodium acetate/bicarbonate buffer. Muscles of the left leg were used for determination of the effect of ammonia, and muscles of the right leg served as controls. Other components present in the medium were dependent on the parameter measured as described below. The viability of the incubated muscles was previously confirmed in our laboratory (Safranek et al. 2003) as well as by other authors (Fang et al. 2005). Three separate studies were performed.

Study 1: the effect of hyperammonemia on amino acid exchange between the muscle and the medium, the release of BCKA, and amino acid concentration in muscle (18 animals)

The muscles were incubated for 2 h in the medium enriched with amino acids in approximately physiological concentrations. At the end of incubation, the muscles were quickly removed, rinsed and homogenized in 0.6 ml of 2% (v/v) perchloric acid and then centrifuged for 5 min at 12,000g. The supernatant was used for determination of free amino acid concentration in muscle. Results were expressed in μmol/g wet muscle.

The amino acid exchange and the BCKA release were calculated by the formula:

$$\text{Amino acid exchange (nmol/g wet muscle/hour)} = \frac{(C_{t120} - C_{t0})(\text{nmol/l}) \times V \text{ (l)}}{\text{Muscle weight (g)} \times t \text{ (hours)}}$$

where C_{t120} and C_{t0} are amino acid concentrations in the medium at the end and at the beginning of the incubation, V is the volume of incubation medium, and t is the duration of incubation.

Amino acid concentrations were determined with liquid chromatograph (Waters, Milford, MA) after precolumn derivatization with 6-aminoquinolyl-*N*-hydroxysuccinimidyl carbamate, BCKA with liquid chromatograph (Shimadzu, Kyoto, Japan) after precolumn derivatization with *o*-phenylenediamine (Kandar et al. 2009).

Study 2: the effect of hyperammonemia on proteolysis and chymotrypsin-like activity (15 animals)

The total and myofibrillar protein breakdown was estimated after a 2-h incubation of the muscle in a medium enriched with cycloheximide (0.5 mM), which prevented the reincorporation of the released amino acids into the proteins. Since tyrosine is neither synthesized nor degraded in skeletal muscle (Fulks et al. 1975), its release into the medium reflects total proteolysis. The release of 3-methylhistidine, a characteristic product of the myofibrillar breakdown, served for the myofibrillar proteolysis calculation. The calculation of the rates of amino acid release was based on their respective concentrations in the medium, the weight of the muscle, and time of the incubation. Tyrosine and 3-methylhistidine were determined via liquid chromatography after a precolumn derivatization with 6-aminoquinolyl-*N*-hydroxysuccinimidyl carbamate (Cohen and Michaud 1993) or fluorecamine (Wassner et al. 1980).

The chymotrypsin-like activity of proteasome was determined using the fluorogenic substrate Suc-LLVY-MCA (Gomes-Marcondes and Tisdale 2002) as follows. The muscles were homogenized in 0.4 ml of ice-cold 20 mM Tris buffer, pH 7.5, containing 2 mM ATP, 5 mM MgCl₂ and 1 mM dithiothreitol. The homogenates were centrifuged for 10 min at 18,000g at 4°C. The cellular supernatant (0.1 ml) was incubated with 0.1 ml of substrate Suc-LLVY-MCA (0.1 mM), with and without inhibitor MG132 (0.02 mM), for 1 h on ice. Adding 0.4 ml of 100 mM sodium acetate buffer of pH 4.3, stopped the reaction. The fluorescence of the samples was immediately determined at the excitation wavelength of 340 nm and the emission wavelength of 440 nm (Tecan Infinite™ 200). The standard curve was established for 7-amino-4-methylcoumarin (AMC), which permitted the expression of CTLA as nmol of AMC/g protein per hour. The activity was adjusted for the protein concentration of the supernatant (Lowry et al. 1951). The differences after the subtraction of inhibited from non-inhibited activities were used for the calculations.

Study 3: the effect of hyperammonemia on leucine oxidation and protein synthesis (9 animals)

Leucine oxidation and protein synthesis rates were measured after 1-h incubation of the muscle in 2.5 ml of medium enriched with amino acids in approximately physiological concentrations and L-[1-¹⁴C]leucine (0.6 μCi/ml) (Carbó et al. 2000; Safranek et al. 2003).

Protein synthesis

The muscles were removed from the incubation flasks, quickly rinsed in cold 6% (v/v) perchloric acid, blotted, and homogenized in 0.6 ml of 6% (v/v) perchloric acid. The homogenate was centrifuged for 5 min at 12,000g. The supernatant was used for measurement of L-[1-¹⁴C]leucine radioactivity and leucine concentration to calculate muscle free leucine specific activity (SA_{Leu}). The precipitated pellet was washed three times by 2% (v/v) perchloric acid and then hydrolyzed in 1 M NaOH. Aliquots were taken for measurement of protein content (Lowry et al. 1951) and L-[1-¹⁴C]leucine radioactivity in muscle protein. SA_{Leu} and protein synthesis rates were calculated by the following formulas:

$$SA_{Leu} \text{ (dpm/nmol)} = \frac{\text{Leu radioactivity (dpm/ml)}}{\text{Leu concentration (nmol/ml)}}$$

$$\text{Protein synthesis (nmol Leu/g protein per hour)} = \frac{\text{Leu radioactivity in protein (dpm/g protein per hour)}}{SA_{Leu} \text{ (dpm/nmol)}}$$

Leucine oxidation

At the end of the incubation period, 0.4 ml of hyamine hydroxide was added into the well hanging above the incubation medium, the reaction was stopped by the addition of 35% (v/v) perchloric acid solution (0.2 ml) into the incubation medium, and the flasks were shaken for 1 h to ensure complete absorption of ¹⁴CO₂ into the hyamine hydroxide. The hanging wells containing hyamine were placed into counting vials containing 10 ml of scintillation mixture and counted in the liquid scintillation radioactivity counter LS 6000 (Beckman Instruments, CA, USA). The efficiency of ¹⁴CO₂ recovery estimated by adding sodium [¹⁴C]bicarbonate to the medium was 96%. Therefore the recovery factor (FR) 0.96 was used for calculation of leucine oxidation by the formula:

$$\text{Leucine oxidation (nmol/g wet muscle/h)} = \frac{\text{radioactivity of released } ^{14}\text{CO}_2 \text{ (dpm/h)}}{SA_{Leu} \text{ (dpm/nmol)} \times \text{muscle weight (g)} \times \text{FR}}$$

Statistical analysis

The results are expressed as the mean ± SEM. *F* test followed by paired *t* test (to estimate the effect of ammonia on specific type of the muscle) and unpaired *t* test (to estimate the difference in sensitivity to ammonia between SOL and EDL) have been used for the analysis of the data. Differences were considered significant at *P* < 0.05. Statistical software NCSS 2001 (Kaysville, UT, US) was applied.

Results

Exchange of amino acids between the muscle and the medium

The enhanced release of glutamine and decreased release of BCAA, glutamate and alanine was observed from the muscles incubated under hyperammonemic conditions. The effect of ammonia on BCAA release was more pronounced in EDL (60% of controls) than in SOL (81% of controls) (Table 1).

Release of BCKA

Addition of ammonia into the incubation medium significantly enhanced the release of all three BCKA. The effect was more pronounced in EDL (161% of controls) than in SOL (133% of controls) (Table 2).

Leucine oxidation

Ammonia significantly stimulated the leucine oxidation in EDL (174% of controls). The effect in SOL was insignificant (Fig. 1).

Protein synthesis and proteolysis

Insignificant changes in leucine incorporation in muscle protein, in tyrosine and 3-MH release, and in CHTLA activity indicate that the incubation of skeletal muscle in the medium with high concentration of ammonia had no effect on protein synthesis and proteolysis. The higher values of SA_{Leu} in EDL are related to lower BCAA concentrations in EDL than in SOL. The effect of ammonia on SA_{Leu} was insignificant (Table 3).

Amino acid concentrations in muscle

The most impressive effect of incubation of isolated muscles in the medium with ammonia was a significant increase in glutamine concentration. Changes in the BCAA concentration were insignificant (Table 4).

Discussion

There are differences in relative distribution of BCAA aminotransferase and BCKA dehydrogenase, and in their activities in human and rat tissues (Khatra et al. 1977; Sweatt et al. 2004; Suryawan et al. 1998), and therefore the results should be interpreted cautiously. In a noteworthy study, Suryawan et al. (1998) demonstrated that some BCAA aminotransferase activity is present in human liver while

Table 1 Effect of ammonia on amino acid exchange between the incubated muscle and incubation medium

	SOL (<i>n</i> = 16)		EDL (<i>n</i> = 16)	
	Control	Ammonia	Control	Ammonia
Glu	1,009 ± 118	757 ± 73*	683 ± 41	514 ± 137*
% of control		80 ± 6		80 ± 6
Gln	3,612 ± 275	4,696 ± 249*	2,220 ± 153	3,186 ± 220*
% of control		138 ± 10		145 ± 5
Ala	2,619 ± 216	2,090 ± 146*	1,263 ± 113	860 ± 123*
% of control		84 ± 6		69 ± 7
Val	802 ± 51	646 ± 46*	472 ± 23	295 ± 35*
% of control		83 ± 6		65 ± 8
Ile	548 ± 50	400 ± 37*	328 ± 19	206 ± 17*
% of control		81 ± 8		64 ± 4
Leu	651 ± 75	500 ± 53*	318 ± 34	139 ± 17*
% of control		82 ± 7		49 ± 7 [#]
BCAA	2,001 ± 148	1,546 ± 115*	1,118 ± 69	641 ± 40*
% of control		81 ± 6		60 ± 5 [#]

Values are means ± SE in nmol/g per hour. Positive values indicate net release; negative values (not found) indicate net uptake. Effect of ammonia in each type of the muscle was estimated by paired *t* test

BCAA branched-chain amino acids

* *P* < 0.05 versus control. The differences between relative changes in SOL and EDL induced by ammonia were estimated by unpaired *t* test

[#] *P* < 0.05

Table 2 Effect of ammonia on BCKA release from incubated muscle

	SOL (<i>n</i> = 18)		EDL (<i>n</i> = 18)	
	Control	Ammonia	Control	Ammonia
KIV	47.2 ± 2.8	55.8 ± 2.6*	42.2 ± 2.4	62.0 ± 2.9*
% of control		127 ± 10		156 ± 12
KIC	99.8 ± 6.0	127.9 ± 10.6*	84.3 ± 5.0	126.3 ± 5.1*
% of control		138 ± 14		161 ± 13
KMV	48.5 ± 2.6	59.2 ± 2.9*	40.6 ± 2.6	63.6 ± 2.9*
% of control		129 ± 9		170 ± 15
Total BCKA	195.5 ± 11.2	242.8 ± 14.3*	167.0 ± 9.7	251.9 ± 10.8*
% of control		133 ± 11		161 ± 13

Values are means ± SE in nmol/g per hour. Effect of ammonia in each type of the muscle was estimated by paired *t* test

BCKA branched-chain keto acids, KIV α -ketoisovalerate, KIC α -ketoisocaproate, KMV α -keto- β -methylvalerate

* *P* < 0.05 versus control. The differences between relative changes in SOL and EDL induced by ammonia were estimated by unpaired *t* test

[#] *P* < 0.05

almost absent in rat liver, and that human muscle contains approximately 54% of total BCKA dehydrogenase activity with only 13% in the liver, while in rats, 3% is found in the muscle and 83% in the liver. Aside from skeletal muscle and the liver, other organs, particularly the brain and kidneys, contribute significantly to whole-body BCAA oxidation.

The results obtained in this study clearly demonstrate that hyperammonemia induces marked alterations in the

BCAA metabolism in skeletal muscle, resulting in the decrease in BCAA concentration in the extracellular fluid. The results strongly support the hypothesis that the mechanism by which hyperammonemia decreases the BCAA level is based on a simple biochemical regulation principle of enzymatic reactions, by alterations in reactant concentrations. The main changes are summarized in Fig. 2 and discussed below.




Fig. 1 Effect of ammonia on leucine oxidation. Values are means \pm SE; $n = 9$ in each group. Paired test, $*P < 0.05$. The effect of ammonia was significant only in EDL (174% vs. control)

Hyperammonemia enhances the conversion of BCAA to BCKA in muscle

The initial step in the BCAA catabolism, which occurs mostly in skeletal muscle due to a high activity of BCAA aminotransferase, involves the transfer of the amino group with α -ketoglutarate to form glutamate and BCKA. Glutamate can be converted to glutamine (in the presence of ammonia) or it can pass its amino group to pyruvate to form alanine. Since these reactions are reversible and near equilibrium, the principal regulatory mechanism of the BCAA aminotransferase are changes in the concentration of substrates and products of these reactions (Harper et al. 1984).

When hyperammonemia develops, the skeletal muscle takes up ammonia from the blood and detoxifies it by the

synthesis of glutamine from glutamate in the reaction catalysed by glutamine synthetase. As BCAA are the main source of nitrogen for synthesis of glutamate from α -ketoglutarate, the consequence of activated synthesis of glutamine is the enhanced consumption of BCAA and the production of BCKA in BCAA aminotransferase reaction. The increase in BCAA aminotransferase activity in rat muscle by hyperammonemia was demonstrated by Hod et al. (1982).

As BCAA are known also as essential donors of nitrogen for synthesis of alanine from pyruvate, the increased demands for glutamate caused by hyperammonemia should decrease the alanine synthesis and release from the muscle. Therefore, the finding of inverse alteration in the concentrations of glutamine (increase) and alanine (decrease) under hyperammonemic conditions indicates, together with decreased release of BCAA and increased release of BCKA, the direct relationship among hyperammonemia, glutamine synthesis, the rate of BCAA aminotransferase reaction, and the BCAA consumption.

The higher increase in glutamine concentration in muscles incubated in medium with ammonia (around 2,000 nmol/g per hour) than can be accounted for BCAA losses (around 500 nmol/g per hour) demonstrates that other sources of glutamate participate in synthesis of glutamine. An important source is undoubtedly extracellular fluid, as indicated in the present study by decreased net release of glutamate in both muscles. We also assume the involvement of other sources of amino nitrogen in synthesis of glutamate and/or the shift from glutamate utilization in other reactions, such as synthesis of glutathione, to glutamine synthesis. Glutathione depletion has been

Table 3 Effect of ammonia on protein metabolism in isolated skeletal muscle

	<i>n</i>	SOL		EDL	
		Control	Ammonia	Control	Ammonia
Protein synthesis (nmol Leu/g per hour)	9	1,417 \pm 168	1,317 \pm 143	1,138 \pm 130	1,071 \pm 104
% of control			101 \pm 14		102 \pm 12
SA _{Leu} (dpm/nmol)	9	5,179 \pm 481	5,863 \pm 647	7,825 \pm 532	7,681 \pm 100
% of control			120 \pm 13		100 \pm 7
Tyrosine release (nmol Tyr/g per hour)	13	278 \pm 14	279 \pm 13	144 \pm 6	152 \pm 31
% of control			102 \pm 6		106 \pm 4
3-MH release (nmol 3-MH/g per hour)	15	1.70 \pm 0.10	1.71 \pm 0.07	1.06 \pm 0.07	1.06 \pm 0.07
% of control			103 \pm 4		101 \pm 2
CHTLA (nmol AMC/mg per hour)	14	447 \pm 38	516 \pm 34	445 \pm 46	452 \pm 51
% of Control			129 \pm 14		110 \pm 12

Values are means \pm SE. Effect of ammonia in each type of the muscle was estimated by paired *t* test

SA_{Leu} specific activity of free leucine in muscle

* $P < 0.05$ versus control. The differences between relative changes in SOL and EDL induced by ammonia were estimated by unpaired *t* test

$P < 0.05$

Table 4 Effect of ammonia on amino acid concentrations in incubated muscle

	SOL (<i>n</i> = 16)		EDL (<i>n</i> = 16)	
	Control	Ammonia	Control	Ammonia
Glu	5,386 ± 413	5,024 ± 175	4,711 ± 248	3,183 ± 81*
% of control		99 ± 8		70 ± 5 [#]
Gln	9,572 ± 713	13,354 ± 608*	12,269 ± 465	15,342 ± 791*
% of control		145 ± 8		126 ± 6
Ala	3,306 ± 177	3,257 ± 167	3,380 ± 176	2,608 ± 125*
% of control		99 ± 3		79 ± 5 [#]
Val	393 ± 30	387 ± 22	217 ± 9	212 ± 10
% of control		103 ± 9		98 ± 4
Ile	272 ± 26	253 ± 13	131 ± 6	127 ± 8
% of control		101 ± 11		98 ± 7
Leu	491 ± 32	503 ± 29	257 ± 13	241 ± 14
% of control		107 ± 10		95 ± 5
BCAA	1,156 ± 81	1,143 ± 57	604 ± 27	580 ± 32
% of control		103 ± 9		97 ± 5

Values are means ± SE in nmol/g. Effect of ammonia in each type of the muscle was estimated by paired *t* test

BCAA branched-chain amino acids

* *P* < 0.05 versus control. The differences between relative changes in SOL and EDL induced by ammonia were estimated by unpaired *t* test

[#] *P* < 0.05




Fig. 2 Hypothetical scheme of the effect of hyperammonemia on BCAA metabolism in skeletal muscle resulting in the decrease of BCAA concentration in extracellular fluid. Ammonia is in skeletal muscle detoxified by synthesis of glutamine from glutamate. The preferential source of nitrogen for synthesis of glutamine from α -ketoglutarate are the BCAA. The BCKA formed from BCAA in aminotransferase reaction are oxidized in skeletal muscle and/or released to the blood. The suggested mechanism of decreased net release of BCAA (or increased uptake from ECF) is their enhanced exchange with glutamine via L-transporter system. ECF the extracellular fluid, BCAA branched-chain amino acids, BCKA branched-chain keto acids, BCA-CoA branched-chain acyl-CoA, α -KG alpha-ketoglutarate, Glu glutamate, Gln glutamine, Ala alanine. 1 branched-chain aminotransferase, 2 branched-chain α -ketoacid dehydrogenase, 3 glutamine synthetase

observed both in hyperammonemia (Essa and Subramanian 2006) and other conditions of activated glutamine synthesis (Hammarqvist et al. 1997).

Hyperammonemia enhances both the BCKA release to the extracellular fluid and BCAA oxidation in muscle

The enhanced flux of BCAA through the BCAA aminotransferase reaction results in an increased production of BCKA, which can be oxidized in skeletal muscle or released to the extracellular fluid. In our study, both these pathways were found to have been activated, as demonstrated by the enhanced release of BCKA into the incubation medium, and by the stimulation of leucine oxidation.

The main cause of release of BCKA from skeletal muscle to the blood is the low activity of the BCKA dehydrogenase in muscle. These BCKA are then taken up and oxidized by tissues with high BCKA dehydrogenase activity, such as adipose tissue, liver and the heart (Harper et al. 1984), or reaminated to BCAA, particularly in the liver (Walser et al. 1973). The latter may occur in a condition of enhanced supply of glutamate or BCKA and/or in case of a lack of BCAA or alpha-ketoglutarate. The enhanced production of glutamate and activated resynthesis of BCAA may be associated with glutaminase reaction in the liver to provide ammonia for urea synthesis. This suggestion can be supported by finding a higher synthesis

of BCAA from BCKA in the liver perfused by a medium with glutamine than by a glutamine-deficient medium (Holecek et al. 2003). We suppose that in case of severe hepatic injury, the conversion of BCKA to BCAA induced by the enhanced availability of glutamine, may be impaired and may be the additional mechanism decreasing the BCAA levels in cirrhotic patients.

The way in which hyperammonemia activates the BCAA oxidation might be explained by changes in the BCKA dehydrogenase reaction, which is the first irreversible and the rate-limiting step in the BCAA oxidation in skeletal muscle, and therefore commits the BCAA to degradation (Harper et al. 1984). The BCKAD is a multi-enzyme complex located on the inner surface of the inner mitochondrial membrane which is highly regulated by a phosphorylation (inhibition)–dephosphorylation (activation) mechanism. The BCKA dehydrogenase kinase, which is considered to be the key regulator of BCKA dehydrogenase activity (Harris et al. 2004), is allosterically inhibited by BCKA, particularly by α -ketoisocaproate (Schauder 1988; Brosnan and Brosnan 2006). Therefore the activation of BCKA dehydrogenase by the enhanced supply of BCKA (produced by activated aminotransferase reaction as described above), seems to be the main mechanism by which hyperammonemia enhances the BCAA oxidation.

Our measurements show that the ammonia-induced fall of BCAA release from muscle (around 500 nmol/g per hour) was higher than the increase in their oxidation and the release of BCKA from muscle. Considering insignificant changes in intracellular BCAA levels, the possible cause should be the increase in BCKA concentration in intracellular space and/or enhanced BCAA utilization in protein synthesis, although the increase in protein synthesis observed both in red and white muscle was insignificant.

Hyperammonemia affects transport of BCAA across the cell membrane

The decrease in BCAA levels in incubation medium and the enhanced release of BCKA and leucine oxidation in muscles incubated in hyperammonemic conditions were not associated with the decrease in BCAA levels in skeletal muscle. Considering unaffected rates of protein synthesis and proteolysis, the explanation should be sought in terms of alterations in amino acid transport across the cell membrane.

There are several transport systems with overlapping substrate specificity which may be involved. An example may be a System L, that serves for neutral amino acids. System L operates as an obligatory amino acid exchanger which can couple the uptake of BCAA with the efflux of cytoplasmic amino acids such as glutamine (Meier et al. 2002). Therefore, the enhanced efflux of glutamine could

prevent intracellular depletion of BCAA via increase in the influx of BCAA from the extracellular fluid, in spite of their enhanced catabolism induced by hyperammonemia.

Effect of hyperammonemia on protein synthesis and proteolysis in muscle

In our experimental conditions we did not succeed in demonstrating that hyperammonemia affects protein synthesis, proteolysis, or BCAA concentration in muscle. We assume that this finding indicates that the fall of BCAA in extracellular fluid and enhanced concentrations of glutamine both in intracellular and extracellular space preceded the fall in intracellular BCAA levels. This suggestion is supported by observations of Leweling et al. (1996), who infused ammonium salts in rats for 2 or 6 h. Administration of ammonium brought about a decline in BCAA plasma concentrations after 2 h and in muscle after 6 h; although the increase of glutamine associated with the fall of glutamate and alanine was observed both in plasma and muscle already after 2 h.

Therefore, alterations in protein metabolism may develop later, and may be activated by humoral changes induced by ammonia, especially by alterations in leucine, BCKA, and glutamine concentrations. Leucine has a potent stimulatory effect on protein synthesis, probably via activation of translation initiation factors (Nair and Short 2005), BCKA have an inhibitory effect on proteolysis (Tischler et al. 1982), and glutamine probably has both a stimulatory effect on protein synthesis (Iresjo et al. 2005) and an inhibitory effect on proteolysis (Holecek et al. 2000a). These findings indicate that changes induced by hyperammonemia may in fact result in a decrease in muscle protein turnover rather than in an increase, and that the cause of any potential muscle wasting induced by ammonia should be a greater decrease in protein synthesis compared to the respective decrease in proteolysis.

Effect of hyperammonemia on the BCAA metabolism in muscle depends on the type of fibre

The results indicate that EDL (muscle composed of white, fast-twitch fibres, which derives energy mostly from anaerobic glycolysis and is adapted to short duration contraction movements) is more susceptible to the effect of hyperammonemia than SOL (muscle composed of red, slow-twitch fibres, with a greater capacity for aerobic metabolism and adapted to maintaining a relatively sustained contraction). This is clearly demonstrated not only by lower release of BCAA into medium, but also by higher release of BCKA, and higher increase in leucine oxidation in EDL than in SOL. We assume that the higher increase in BCAA utilization in white muscle may be due to higher

glutamine synthetase activity, which, under hyperammonemic conditions enables more rapid consumption of glutamate and thus activates the flux of amino nitrogen through BCAA aminotransferase and BCKA dehydrogenase reactions. The physiological importance of this phenomenon may be related to higher capacity of fast-twitch muscle to produce ammonia during contraction (Terjung et al. 1985). When considering that red fibres are the major components of respiratory muscles (Polla et al. 2004), the finding that hyperammonemia affects groups of muscles with a higher proportion of white fibres may be also of clinical importance.

Conclusion

In conclusion, our study provides clear evidence that hyperammonemia activates the BCAA catabolism in skeletal muscle, which is associated with enhanced release of BCKA to the extracellular fluid, with increased BCAA oxidation in muscle, and with decreased BCAA concentration in extracellular fluid. These metabolic changes are not associated with marked changes in protein turnover. The effect of ammonia is more pronounced in muscles with high amount of white fibres.

Acknowledgments This study was supported by Research Project MSM 0021620820. We are grateful for the technical assistance of H. Buzkova and R. Rysava. Many thanks to Nishant Sunil Joshi from London, UK, for his assistance with revision of English.

References

- Blonde-Cynober F, Plassart F, Rey C, Coudray-Lucas C et al (1994) Assessment of the carbon tetrachloride-induced cirrhosis model for studies of nitrogen metabolism in chronic liver disease. *Ann Nutr Metab* 38:238–248
- Brosnan JT, Brosnan ME (2006) Branched-chain amino acids: enzyme and substrate regulation. *J Nutr* 136:207S–211S
- Carbó N, Ribas V, Busquets S et al (2000) Short-term effects of leptin on skeletal muscle protein metabolism in the rat. *J Nutr Biochem* 11:431–435
- Cohen SA, Michaud DP (1993) Synthesis of a fluorescent derivatizing reagent, 6-aminoquinolyl-*N*-hydroxysuccinimidyl carbamate, and its application for the analysis of hydrolysate amino acids via high-performance liquid chromatography. *Anal Biochem* 211:279–287
- Eriksson LS, Hagenfeldt L, Wahren J (1982) Intravenous infusion of alpha-oxoisocaproate: influence on amino acid and nitrogen metabolism in patients with liver cirrhosis. *Clin Sci (Lond)* 62:285–293
- Essa MM, Subramanian P (2006) *Pongamia pinnata* modulates the oxidant–antioxidant imbalance in ammonium chloride-induced hyperammonemic rats. *Fundam Clin Pharmacol*. 20:299–303
- Fang CH, Li BG, James JH et al (2005) Protein breakdown in muscle from burned rats is blocked by insulin-like growth factor and glycogen synthase kinase-3beta inhibitors. *Endocrinology* 146:3141–3149
- Fernandez JM, Croom WJ Jr, Johnson AD et al (1988) Subclinical ammonia toxicity in steers: effects on blood metabolite and regulatory hormone concentrations. *J Anim Sci* 66:3259–3266
- Fischer JE, Baldessarini RJ (1971) False neurotransmitters and hepatic failure. *Lancet* 2:75–80
- Fischer JE, Rosen HM, Ebeid AM et al (1976) The effect of normalization of plasma amino acids on hepatic encephalopathy in man. *Surgery* 80:77–91
- Fulks RM, Li JB, Goldberg AL (1975) Effects of insulin, glucose, and amino acids on protein turnover in rat diaphragm. *J Biol Chem* 250:290–298
- Gomes-Marcondes MCC, Tisdale MJ (2002) Induction of protein catabolism and the ubiquitin-proteasome pathway by mild oxidative stress. *Cancer Lett* 180:69–74
- Hammarqvist F, Luo JL, Cotgreave IA et al (1997) Skeletal muscle glutathione is depleted in critically ill patients. *Crit Care Med* 25:78–84
- Harper AE, Miller RH, Block KP (1984) Branched-chain amino acid metabolism. *Annu Rev Nutr* 4:409–454
- Harris RA, Joshi M, Jeoung NH (2004) Mechanisms responsible for regulation of branched-chain amino acid catabolism. *Biochem Biophys Res Commun* 313:391–396
- Hasselgren PO, Talamini M, James JH et al (1986) Protein metabolism in different types of skeletal muscle during early and late sepsis in rats. *Arch Surg* 121:918–923
- Hayashi M, Ohnishi H, Kawade Y et al (1981) Augmented utilization of branched-chain amino acids by skeletal muscle in decompensated liver cirrhosis in special relation to ammonia detoxication. *Gastroenterol Jap* 16:64–70
- Hickson RC, Wegrzyn LE, Osborne DF, Karl IE (1996) Glutamine interferes with glucocorticoid-induced expression of glutamine synthetase in skeletal muscle. *Am J Physiol* 270:E912–E917
- Hod G, Chaouat M, Haskel Y et al (1982) Ammonia uptake by skeletal muscle in the hyperammonaemic rat. *Eur J Clin Invest* 12:445–450
- Holecek M, Mraz J, Tilser I (1996a) Plasma amino acids in four models of experimental liver injury in rats. *Amino Acids* 10:229–241
- Holecek M, Tilser I, Skopec F et al (1996b) Leucine metabolism in cirrhotic rats. *J Hepatol* 24:209–216
- Holecek M, Skopec F, Skalska H et al (2000a) Effect of alanyl-glutamine on leucine and protein metabolism in endotoxemic rats. *JPEN J Parenter Enteral Nutr* 24:215–222
- Holecek M, Sprongl L, Tichy M et al (2000b) Effect of hyperammonemia on leucine and protein metabolism in rats. *Metabolism* 49:1330–1334
- Holecek M, Rysava R, Safranek R et al (2003) Acute effects of decreased glutamine supply on protein and amino acid metabolism in hepatic tissue. A study using isolated perfused rat liver. *Metabolism* 52:1062–1067
- Iber FL, Rosen H, Levenson SM et al (1957) The plasma amino acids in patients with liver failure. *J Lab Clin Med* 50:417–425
- Iresjo BM, Svanberg E, Lundholm K (2005) Reevaluation of amino acid stimulation of protein synthesis in murine- and human-derived skeletal muscle cells assessed by independent techniques. *Am J Physiol Endocrinol Metab* 288:E1028–E1037
- Kandar R, Zakova P, Jirosova J et al (2009) Determination of branched chain amino acids, methionine, phenylalanine, tyrosine and α -keto acids in plasma and dried blood samples using HPLC with fluorescence detection. *Clin Chem Lab Med* 47:565–572
- Khatra BS, Rajender KC, Sewell CW, Rudman D (1977) Distribution of branched-chain α -keto acid dehydrogenases in primate tissues. *J Clin Invest* 59:558–564
- Leweling H, Breitkreutz R, Behne F et al (1996) Hyperammonemia-induced depletion of glutamate and branched-chain amino acids in muscle and plasma. *J Hepatol* 25:756–762

- Lowry OH, Rosebrough NJ, Farr AL et al (1951) Protein measurement with the Folin phenol reagent. *J Biol Chem* 193:265–275
- Maizels EZ, Ruderman NB, Goodman MN et al (1977) Effect of acetoacetate on glucose metabolism in the soleus and extensor digitorum longus muscles of the rat. *Biochem J* 162:557–568
- Marchesini G, Forlani G, Zoli M et al (1979) Insulin and glucagon levels in liver cirrhosis. Relationship with plasma amino acid imbalance of chronic hepatic encephalopathy. *Dig Dis Sci* 24:594–601
- Marchesini G, Bianchi G, Merli M et al (2003) Nutritional supplementation with branched-chain amino acids in advanced cirrhosis: a double-blind, randomized trial. *Gastroenterology* 124:1792–1801
- Meier C, Ristic Z, Klausner S, Verrey F (2002) Activation of system L heterodimeric amino acid exchangers by intracellular substrates. *EMBO J* 21:580–589
- Muthy T, Kovarik M, Sispera L et al (2008) Protein metabolism in slow- and fast-twitch muscle during turpentine-induced inflammation. *Int J Exp Pathol* 89:64–71
- Nair KS, Short KR (2005) Hormonal and signaling role of branched-chain amino acids. *J Nutr* 135:1547S–1552S
- Nakaya Y, Okita K, Suzuki K, Hepatic Nutritional Therapy (HNT) Study Group et al (2007) BCAA-enriched snack improves nutritional state of cirrhosis. *Nutrition* 23:113–120
- Peng S, Plank LD, McCall JL et al (2007) Body composition, muscle function, and energy expenditure in patients with liver cirrhosis: a comprehensive study. *Am J Clin Nutr* 85:1257–1266
- Polla B, D'Antona G, Bottinelli R, Reggiani C (2004) Respiratory muscle fibres: specialisation and plasticity. *Thorax* 59:808–817
- Safranek R, Holecek M, Kadlcikova J et al (2003) Effect of acute acidosis on protein and amino acid metabolism in rats. *Clin Nutr* 22:437–443
- Schauder P (1988) Do branched chain keto amino acids regulate their own metabolism? *Hepatology* 8:1716–1718
- Soeters PB, de Boer J (1984) Why are plasma branched chain amino acid levels diminished in patients with liver cirrhosis. In: Adibi SA, Fekl W, Langenbeck U, Schauder P (eds) *Branched chain amino and keto acids in health and disease*. Karger, Basel, pp 483–496
- Suryawan A, Hawes JW, Harris RA et al (1998) A molecular model of human branched-chain amino acid metabolism. *Am J Clin Nutr* 68:72–81
- Sweatt AJ, Wood M, Suryawan A et al (2004) Branched-chain amino acid catabolism: unique segregation of pathway enzymes in organ systems and peripheral nerves. *Am J Physiol Endocrinol Metab* 286:E64–E76
- Terjung RL, Dudley GA, Meyer RA (1985) Metabolic and circulatory limitations to muscular performance at the organ level. *J Exp Biol* 115:307–318
- Tischler ME, Desautels M, Goldberg AL (1982) Does leucine, leucyl-tRNA, or some metabolite of leucine regulate protein synthesis and degradation in skeletal and cardiac muscle? *J Biol Chem* 257:1613–1621
- Urata Y, Okita K, Korenaga K et al (2007) The effect of supplementation with branched-chain amino acids in patients with liver cirrhosis. *Hepatol Res* 37:510–516
- Walser M, Lund P, Ruderman NB, Coulter AW (1973) Synthesis of essential amino acids from their α -keto analogue by perfused rat liver and muscle. *J Clin Invest* 52:2865–2877
- Wassner SJ, Schlitzer L, Li JB (1980) A rapid, sensitive method for the determination of 3-methylhistidine levels in urine and plasma using high-pressure liquid chromatography. *Anal Biochem* 104:284–289
- Yamato M, Muto Y, Yoshida T (1995) Clearance rate of plasma branched-chain amino acids correlates significantly with blood ammonia level in patients with liver cirrhosis. *Int Hepatol Commun* 3:91–96
- Yang Q, Birkhahn RH (1997) Branched-chain transaminase and keto acid dehydrogenase activities in burned rats: evidence for a differential adaptation according to sex. *Nutrition* 13:640–645