

Use of platelet-rich fibrin in regenerative dentistry: a systematic review

Richard J. Miron^{1,2} · Giovanni Zucchelli³ · Michael A. Pikos⁴ · Maurice Salama^{1,5,6} · Samuel Lee⁷ · Vincent Guillemette⁸ · Masako Fujioka-Kobayashi^{1,9,10} · Mark Bishara¹¹ · Yufeng Zhang¹² · Hom-Lay Wang² · Fatiha Chandad⁸ · Cleopatra Nacopoulos¹³ · Alain Simonpieri^{14,15,16} · Alexandre Amir Aalam¹⁷ · Pietro Felice³ · Gilberto Sammartino¹⁸ · Shahram Ghanaati¹⁹ · Maria A Hernandez¹ · Joseph Choukroun²⁰

Received: 3 December 2016 / Accepted: 15 May 2017 / Published online: 27 May 2017
© Springer-Verlag Berlin Heidelberg 2017

Abstract

Objectives Research across many fields of medicine now points towards the clinical advantages of combining regenerative procedures with platelet-rich fibrin (PRF). This systematic review aimed to gather the extensive number of articles published to date on PRF in the dental field to better understand the clinical procedures where PRF may be utilized to enhance tissue/bone formation.

Materials and methods Manuscripts were searched systematically until May 2016 and separated into the following categories: intrabony and furcation defect regeneration, extraction

socket management, sinus lifting procedures, gingival recession treatment, and guided bone regeneration (GBR) including horizontal/vertical bone augmentation procedures. Only human randomized clinical trials were included for assessment.

Results In total, 35 articles were selected and divided accordingly ($\kappa = 0.94$). Overall, the use of PRF has been most investigated in periodontology for the treatment of periodontal intrabony defects and gingival recessions where the majority of studies have demonstrated favorable results in soft tissue management and repair. Little to no randomized clinical trials

Electronic supplementary material The online version of this article (doi:10.1007/s00784-017-2133-z) contains supplementary material, which is available to authorized users.

✉ Richard J. Miron
richard.miron@zmk.unibe.ch

- ¹ College of Dental Medicine, Department of Periodontology, Nova Southeastern University, Fort Lauderdale, FL, USA
- ² Department of Periodontics and Oral Medicine, University of Michigan, Ann Arbor, MI, USA
- ³ Department of Biomedical and Neuromotor Sciences, University of Bologna, Bologna, Italy
- ⁴ Pikos Institute, Tampa Bay, FL, USA
- ⁵ Department of Periodontology, Georgia University, Athens, GA, USA
- ⁶ Goldstein Garber & Salama, Atlanta, GA, USA
- ⁷ International Academy of Dental Implantology, San Diego, CA, USA
- ⁸ Department of Periodontology, Laval University, Quebec City, Canada
- ⁹ Department of Cranio-Maxillofacial Surgery, Inselspital, Bern University Hospital, University of Bern, Bern, Switzerland

- ¹⁰ Department of Oral Surgery, Clinical Dentistry, Institute of Biomedical Sciences, Tokushima University Graduate School, Tokushima, Japan
- ¹¹ West Bowmanville Dental, Bowmanville, Ontario, Canada
- ¹² Department of Oral Implantology, University of Wuhan, Wuhan, China
- ¹³ Laboratory for Research of the Musculoskeletal System, KAT Hospital, School of Medicine, National and Kapodistrian, University of Athens, Athens, Greece
- ¹⁴ Oral Surgery Department, University Federico II Naples, Naples, Italy
- ¹⁵ Periodontology and Implantology, Beausoleil, France
- ¹⁶ Periodontology and Implantology, Marseille, France
- ¹⁷ Department of Advanced Periodontics, USC School of Dentistry, Los Angeles, CA, USA
- ¹⁸ Department of Neuroscience, Reproductive Science and Odontostomatology, University of Naples Federico II, Naples, Italy
- ¹⁹ FORM, Frankfurt Oral Regenerative Medicine, Clinic for Maxillofacial and Plastic Surgery, Johann Wolfgang Goethe University, Frankfurt am Main, Germany
- ²⁰ Pain Clinic, Nice, France

were found for extraction socket management although PRF has been shown to significantly decrease by tenfold dry sockets of third molars. Very little to no data was available directly investigating the effects of PRF on new bone formation in GBR, horizontal/vertical bone augmentation procedures, treatment of peri-implantitis, and sinus lifting procedures.

Conclusions Much investigation now supports the use of PRF for periodontal and soft tissue repair. Despite this, there remains a lack of well-conducted studies demonstrating convincingly the role of PRF during hard tissue bone regeneration. Future human randomized clinical studies evaluating the use of PRF on bone formation thus remain necessary.

Clinical relevance PRF was shown to improve soft tissue generation and limit dimensional changes post-extraction, with little available data to date supporting its use in GBR.

Keywords Platelet-rich fibrin · Tissue regeneration · Bone augmentation · Soft tissue regeneration

Introduction

Regenerative therapy in dentistry involves the replacement and/or regeneration of oral tissues altered as a result of disease or injury. One of the reported aspects complicating this endeavor has been the complex nature of the tissues found in the oral cavity. These include both mineralized tissues such as the cementum, alveolar bone, and dentin, as well as soft tissues connected by ligaments (periodontal ligament), each comprising distinct cell populations from various tissue origins (ectodermal and mesodermal). These cell populations reside in specialized extracellular matrices organized in complex fashions [1, 2]. In the past, a variety of regenerative procedures utilizing highly sophisticated biomaterials were introduced to attempt their regeneration. These included ambitious attempts with barrier membranes to perform guided tissue/bone regeneration and the use of a variety of bone grafting materials from human, animal and synthetic sources, as well as bioactive growth factors such as bone morphogenetic proteins (BMPs) and enamel matrix derivative (EMD). Other investigators proposed that the use of three-dimensional scaffolds fabricated from the patient's own peripheral blood could be utilized [2]. This new approach is based on the concepts that were introduced over a decade ago consisting of a platelet concentrate without the use of anticoagulants. Platelet-rich fibrin (PRF) was therefore developed as an improved formulation of the previously utilized platelet-rich plasma (PRP) [3].

Unlike PRP, which requires the addition of anticoagulants such as bovine thrombin during initial blood collection, PRF is obtained simply by centrifugation without anticoagulants and is therefore strictly autologous. This fibrin matrix contains platelets and leukocytes as well as a variety of growth factors and cytokines including transforming growth factor-beta 1

(TGF- β 1), platelet-derived growth factor (PDGF), vascular endothelial growth factor (VEGF), interleukin (IL)-1 β , IL-4, and IL-6 [4]. Furthermore, fibrin that forms during the final stages of the coagulation cascade, combined with cytokines secreted by platelets, makes PRF a highly biocompatible matrix especially in damaged sites where the fibrin network acts also as a reservoir of tissue growth factors [5]. These factors act directly on promoting the proliferation and differentiation of osteoblasts, endothelial cells, chondrocytes, and various sources of fibroblasts [6, 7]. Despite this, many questions remain about the actual clinical performance of PRF. Therefore, the purpose of the present systematic review is to report the current state of knowledge and clinical potential of PRF in regenerative dental therapy when compared to both standardized controls and well-established standard regenerative biomaterials from human clinical randomized trials.

Brief history of platelet concentrates

The original concept leading towards the preparation of platelet concentrates was that concentrated platelets and autologous growth factors could be collected in plasma solutions that could then be utilized in a surgical site to promote local healing [8, 9]. It was given the popular working name "platelet-rich plasma" (PRP), introduced in the late 1990s [10–12]. PRP is composed of over 95% platelets, a cell type that actively secretes growth factors for initiating wound healing and secreting factors responsible for enhancing cell adhesion, proliferation, and migration of various cell types [10, 13]. Around the same time period, Anitua et al. formulated a second platelet concentrate also utilizing anticoagulants termed platelet-rich growth factor (PRGF) [14, 15].

Despite this, several factors have been shown to limit the use of PRP and PRGF. Their preparation requires the additional use of bovine thrombin or CaCl₂ in addition to coagulation factors. Furthermore, the preparation must be centrifuged in two separate stages in order to increase platelet concentration without incorporation of leukocytes (sometimes requiring 1 h). It has further been reported that the liquid nature of PRP also complicates its handling and reduces its potential application since it must be utilized in combination with other biomaterials. Lastly, the clinical potential for bone regeneration with PRP is limited having a very short release of growth factor profile [16–18]. All these limitations have led to the emergence of a second-generation platelet concentrate termed PRF fabricated from 100% autologous sources [19].

Advantages of PRF over PRP

PRF differs from its predecessor (PRP/PRGF) by several parameters which can be summarized as follows: the simplicity

of its preparation and its implementation. The time of preparation and cost of preparation are both significantly lower as PRF does not necessitate the direct activation with additional factors such as bovine thrombin or extrinsic anticoagulants [3]. Because of its fibrous structure, PRF retains a larger number of cytokines and growth factors in a supportive three-dimensional fibrin scaffold for cell migration [20]. In tissue, PRF dissolves more slowly than PRP, forming a solid fibrin matrix slowly remodeled in the style of a natural blood clot. Platelets and cytokines are then effectively retained and released gradually over time [18]. The PRF scaffold allows a continuous slow release of growth factors and cytokines over a period of 10 days, in contrast to PRP which has been shown to release the majority of its growth factors within the first day [18]. Therefore, migrating cells in near proximity to PRF scaffolds are in an environment with fibrin and growth factors throughout their entire growth cycle [21].

Once blood is collected (in the absence of anticoagulants), samples must be centrifuged immediately to avoid the activation of the coagulation cascade. During centrifugation, fibrinogen is concentrated to the top of the collection tube until the circulating thrombin transforms it into a fibrin network. This results in a fibrin clot rich in platelets, trapped between an acellular plasma layer and erythrocytes. The solid fibrin clot is found between the supernatant and the reddish background formed by red blood cells. The clot may then be removed immediately and condensed in a metal box so as to obtain a solid covering membrane or a filling cylinder (Fig. 1). The resulting exudate may be cut and used to hydrate graft materials if required (Fig. 1) [19, 20].

Implications of PRF in wound healing

Although leukocyte and platelet cytokines play an important role in the PRF healing capacity, it has often been suggested that it is the fibrin matrix supporting these elements which is actually responsible for its therapeutic potential [20]. The keys to tissue regeneration lie in their angiogenic potential, their immune system control, their potential to recruit circulating stem cells, and their ability to ensure undisturbed wound closure/healing by epithelial tissues [20]. The angiogenic properties of PRF may therefore be explained by the three-dimensional structure of the fibrin matrix which holds a number of growth factors and cytokines simultaneously embedded in the matrix including PDGF, TGF- β 1, IGF, and VEGF. The regenerative potential of these cytokines has been abundantly studied in tissue wound healing and regeneration [4, 12, 22–33]. Furthermore, the fibrin matrix stimulates the expression of integrin α v β 3 which allows cells to bind to fibrin, fibronectin, and vitronectin [33]. This cascade of events is of utmost importance to initiate the process of angiogenesis and thus tissue wound healing [33].

Moreover, the fibrin degradation products directly stimulate neutrophil migration and facilitate transmigration into the vascular endothelium. This neutrophil activation causes secretion of proteases that facilitate their penetration in the basement membrane of blood vessels, in addition to their contribution to degrade the fibrin clot. Neutrophils trapped within the fibrin clot act to eliminate incoming bacteria and pathogens in the wound site by phagocytosis and the production of toxic free radicals and digestive enzymes. This contributes towards the prevention of bacterial contamination within the surgical site [34]. PRF also contains macrophages that are involved in the healing and repair process by playing a key role in the transition between inflammation and wound repair during osteogenesis [33, 34].

Methods

Development of a protocol

This review was conducted and reported according to the PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) [35]. A protocol including all aspects of a systematic review methodology was developed prior to initiation of this review. This included definition of the focused question; a PICO (patient, intervention, comparison, outcome) question; a defined search strategy; study inclusion criteria; determination of outcome measures; screening methods, data extraction, and analysis; and data synthesis.

PICO question

- P: Do patients in need of clinical bone, cementum, soft tissue, and/or PDL gain
- I: Undergoing to dental treatments (i.e., guided bone/tissue repair/regeneration or pulp repair/regeneration) using defined non/surgical approaches combined with the use of PRF as sole/combined biomaterial
- C: Defined regenerative/reparative approaches without the use of PRF
- O: Soft and/or hard tissue reconstruction of the periodontium, alveolar bone, peri-implant tissues, or tooth structure

Defining the focused question

The following focused question was defined: “What indications has platelet rich fibrin (PRF) been shown effective for tissue repair/regeneration of either soft or hard tissues in dentistry?”




Fig. 1 Fabrication of various PRF membranes from 10 mL autologous blood. Thereafter, two PRF clots were mixed with a bone grafting material accordingly (clinical images were kindly provided by Dr. Michael A. Pikos)

Search strategy

Electronic and manual literature searches were conducted independently by two authors (RJM and MFK) in several databases, including MEDLINE (OVID), EMBASE (OVID), Cochrane Central Register of Controlled Trials (Cochrane Library), Cochrane Oral Health Group Trials Register (Cochrane Library), Web of Science (Thomson Reuters), and SciVerse (Elsevier). The electronic literature was searched for articles published up to and including May 14, 2016: combinations of several search terms and search strategies were applied to identify appropriate studies (Supplemental Tables 1–4). These include search strategies to identify the effects of PRF on (1) intrabony defect regeneration, (2) furcation defect regeneration, (3) management of gingival recessions, (4) guided bone regeneration and extraction socket healing, and (5) sinus floor elevation procedures. Reference lists of review articles and of the included articles in the present review were screened. Finally, a hand search of the *Journal of Clinical Periodontology*, *Journal of Dental Research*, *Journal of Periodontal Research*, *Journal of Periodontology*, *Clinical Oral Implants Research*, *Clinical Implant Dentistry and Related Research*, *Clinical Oral Investigations*, and *The International Journal of Periodontics and Restorative Dentistry* was performed from January 2000 to May 2016.

Criteria for study selection and inclusion

Study selection considered only articles published in English, describing the human clinical evaluation of PRF for the above-indicated search strategies. Only human studies evaluating the comparative effects of PRF to an appropriate control or to another regenerative modality in human studies were included. All human studies evaluating PRF in a case report

or case series were excluded if controls were not present. All animal and in vitro studies were also excluded.

Outcome measure determination

The primary outcome of interest was to determine the regenerative/repairative potential of PRF in a variety of clinical settings utilized in dentistry. For each of the investigated clinical indications, different primary outcomes were considered. For studies dealing with intrabony defect regeneration, probing pocket depth (PPD) and clinical attachment levels (CAL) were measured. For studies dealing with gingival recessions, root coverage was calculated as percentage. Studies investigating the use of PRF for furcation defect regeneration quantified CAL gains as a primary outcome measure. For studies dealing with bone regeneration, dimensional change/density of hard tissues was compared. Similarly, sinus floor elevation procedures quantified new bone formation and/or implant success rates following sinus lifting procedures with/without PRF. Outcomes were summarized in Supplemental Tables 1–4 for the various clinical studies accordingly.

Screening method

Titles and abstracts of the selected studies were independently screened by three reviewers (RJM, MFK, and VG). The screening was based on the question: “Does platelet rich fibrin (PRF) have the ability to affect primary outcomes measured across a variety of procedures commonly performed in dentistry?” Full-text articles were obtained if the response to the screening question was “yes” or “uncertain.” The level of agreement between reviewers was determined by kappa scores. Disagreement regarding inclusion was resolved by discussion between authors. For necessary missing data, the authors of the studies were contacted.

Data extraction and analysis

The following data were extracted: general characteristics (authors and year of publication), defect type, number of patients, healing period, treatment groups, primary outcome measurements, and significant value. Due to the size of the study and the number of treatment procedures compared using PRF, no meta-analysis was performed. Instead, the data is reported in a systematic fashion with an overview of all studies fitting the search descriptions. Thereafter, data was extracted from the collection of articles and summarized in separate tables and discussed accordingly.

Results

Search outcomes

In total, the initial search strategies generated 152 articles that were separated accordingly into intrabony, furcation, gingival recession, guided bone regeneration (GBR)/extraction socket, and sinus lift accordingly (Fig. 2). Of the initial searches, 45 abstracts were retained for further investigation. In total, 10 articles were excluded primarily based on their lack of controls or appropriate endpoints matching our search criteria. In total, 35 articles were kept for further evaluation. This section aims to present viable treatment options utilizing PRF and evaluate its performance according to published studies.

Intrabony defect regeneration with PRF

One of the main uses for PRF has been for the repair/regeneration of periodontal intrabony defects [36–45]. To date, 11 randomized clinical trials (RCTs) have reported the use of PRF for intrabony defect regeneration, most often comparing PRF to open flap debridement alone (Table 1). Clinical improvements were reported via PPD reductions as well as CAL gains following regenerative periodontal therapy. All seven studies found that the additional use of PRF increased PPD reductions and CAL gains when compared to open flap debridement (OFD) alone (Table 1). One study comparing the regeneration of intrabony defects utilizing either PRF or a bone grafting material (demineralized freeze-dried bone allograft (DFDBA)) found no significant differences between treatment groups [42]. Two studies reported the effectiveness of PRF in combination with a bone grafting material when compared to bone grafting material alone [36, 38]. In both these studies, the additional use of PRF enhanced the filling of intraosseous defects. Most recently, Panda et al. most recently found that the supplemental use of PRF for intrabony defect regeneration in combination with a barrier membrane also led to statistically better results [39]. In summary, the collected RCTs have demonstrated that the use of PRF leads

to statistically superior periodontal repair of intrabony defects when compared to OFD alone and may further be combined with regenerative biomaterials such as bone grafts or collagen barrier membranes to further enhance periodontal regeneration of intrabony defects. Despite the widespread use of PRF demonstrating the reduction of PPD and CAL gains, it remains of interest to note that no histological findings have yet been utilized to demonstrate true histological periodontal regeneration in human subjects. Therefore, future research to characterize intrabony defect regeneration versus repair utilizing PRF as a biomaterial remains necessary.

Furcation defect regeneration with PRF

Similarly, PRF has also been utilized in three studies investigating periodontal regeneration of class II furcation defects (Supplemental Table 5) [46–48]. In all studies, PRF was compared to OFD alone, thereby fully characterizing its regenerative potential utilizing appropriate well-designed controls in all human clinical studies. In all three studies conducted by Sharma et al. 2011, Bajaj et al. 2013, and Pradeep et al. 2016, the use of PRF led to a significant improvement in CAL gains when compared to controls [46–48]. These findings report a gain in vertical CAL of 2.33, 2.87, and 4.17 mm in test PRF groups when compared to 1.28, 1.37, and 1.82 mm, respectively, in OFD controls [46–48]. These results demonstrate the potential for tissue repair utilizing PRF for furcation defects. One remaining issue to address is that the results have not confirmed the regenerative potential of PRF via histological evaluation and therefore the process can solely be defined as tissue “repair.” Furthermore, to date, no study has compared the use of PRF to other effective regenerative materials such as bone grafting materials or other regenerative bioactive growth factors. In the future, its clinical performance could be better assessed if compared to other leading regenerative agents.

Root coverage of gingival recessions with PRF

PRF has also been widely utilized as a bioactive matrix in numerous studies for root coverage of gingival recessions (Table 2, Fig. 3) [49–61]. Of the 13 listed studies, six studies compared the use of coronally advanced flap (CAF) to CAF + PRF. Of these studies, three found that PRF induced a significant increase in root coverage [49, 51, 58] whereas the other three found no significant differences [52, 54, 60]. Of the remaining studies, one study compared PRF to EMD and found no differences in the reported root coverage [56]. Four studies compared CAF + PRF to CAF + connective tissue graft (CTG) and also found no difference in average root coverage [50, 53, 55, 61]. One study compared CAF + CTG with CAF + CTG and PRF and found a significant increase in root coverage for the combination approach utilizing both CTG with PRF [57]. Rajaram et al. utilized a double lateral sliding




Fig. 2 Flowchart of the screened relevant publications

bridge flap with and without PRF for the treatment of gingival Miller class II defects and found no significant differences between control and test groups [59]. These results seem to hint at the fact that the use of PRF favors a slight gain in root coverage when compared to CAF alone but does not lead to better results when compared to EMD or to CTG. Furthermore, several reports show that CAF in combination with CTG leads to more width in keratinized tissue when compared to PRF.

Interestingly, two reports commented on the added advantage of PRF in pain management [55, 56]. Jankovic et al. found in two separate studies comparing PRF to either EMD or CTG that PRF led to lower morbidity and faster wound healing [55, 56]. In a similar study investigating the wound healing properties of PRF, the palatal donor site of the epithelialized CTG was treated with PRF or a gelatin sponge on the healing of palatal donor sites [62]. It was reported that

the PRF-enriched palatal bandage significantly accelerated palatal wound healing and reduced the patient's morbidity [62].

In summary, the use of PRF for the treatment of gingival recessions is limited. Evidence from another systematic review from 2016 concluded that the additional use of PRF for the treatment of gingival recessions did not lead to any additional benefit in root coverage or CAL ($P = 0.57$ and $P = 0.50$, respectively) [63]. Furthermore, the reported keratinized mucosa width (KMW) gain was significantly greater in the subgroup treated with CTG when compared to PRF for studies greater or equal to 6 months in duration [63]. The results of that meta-analysis suggest that the use of PRF does not improve the root coverage, KMW, or CAL of Miller class I and II gingival recessions compared with other treatment modalities including EMD or CTG but can be obtained easily at low cost when compared to other regenerative modalities [63].

Table 1 Effects of PRF on intrabony defect regeneration

Author	Defect no.	Healing time (months)	Groups	Δ PPD (mm)	CAL gain (mm)	<i>P</i> value
Thorat 2011	32	9	OFD	3.56	2.13	Δ PPD <0.01
			OFD + PRF	4.56	3.69	CAL <0.01
Sharma 2011	56	9	OFD	3.21	2.77	Δ PPD 0.006
			OFD + PRF	4.55	3.31	CAL n.s.
Pradeep 2012	90	9	OFD	2.97	2.67	Δ PPD 0.002
			OFD + PRF	3.90	3.03	GAL n.s.
Pradeep 2012	90	9	OFD	2.97	2.83	Δ PPD 0.018
			OFD + PRF	3.77	3.17	GAL n.s.
Shah 2015	40	6	OFD + DFDBA	3.70	2.97	n.s.
			OFD + PRF	3.67	2.97	
Pradeep 2015	120	9	OFD	3.01	2.96	Δ PPD <0.001
			OFD + PRF	4.01	4.03	CAL <0.001 both treatment groups
			ORF + 1% MF	3.93	3.93	
			OFD + 1% MF + PRF	4.9	4.9	
Ajwani 2015	40	9	OFD	1.60	1.3	PPD <0.001
			OFD + PRF	1.90	1.8	CAL <0.001
Elgandhy 2015	40	6	OFD + HA	3.42	3.55	PPD <0.02
			OFD + HA + PRF	3.82	3.9	CAL <0.027
Agawal 2016	60	12	OFD + DFDBA	3.60	2.61	PPD <0.05
			OFD + DFDBA + PRF	4.15	3.73	CAL <0.05
Panda 2016	32	9	Barrier membrane	3.19	3.38	PPD 0.002
			Membrane + PRF	3.88	4.44	CAL = 0.001

PPD probing periodontal depth, CAL clinical attachment level, OFD open flap debridement, PRF platelet-rich fibrin, DFDBA demineralized freeze-dried bone allograft, MF metformin, HA hydroxyapatite

Guided bone regeneration and extraction socket management with PRF

One area of research that has gained tremendous popularity in recent years is the management of dimensional changes of the alveolar bone directly following tooth extraction [64–66]. These changes have been reported to occur within 8 weeks following extraction [67] as a consequence of decreased blood supply following tooth removal (periodontal ligament absence). Several advantages have been reported when filling extraction sockets with PRF (Supplemental Table 6) [68–71]. Hauser et al. found in a study of 23 patients that PRF reduced dimensional changes prior to implant placement when compared to natural socket healing [71]. Furthermore, it was reported that raising a peri-mucosteal flap reduced the effectiveness of PRF [71]. Girish Rao et al. found that following third molar extractions, the filling of sockets with PRF led to a non-significant increase in bone volume [68]. Hoaglin et al. reported that filling third molar extraction sockets with

PRF led to a nearly tenfold decrease in osteomyelitis infections when compared to natural healing. This study was conducted bilaterally in 200 patients, thus providing some of the highest scientific evidence for the reduced rate of infection following use of PRF [70]. Lastly, Suttapreyasri et al. found that PRF reduced dimensional changes in premolar extraction sites when compared to blank controls [69].

Despite the limited number of studies, the use of PRF acts as an ideal material post-extraction by improving bone healing/regeneration, preserving the quality and density of the residual ridge, reducing infection, and decreasing the time of surgery when compared to the use of a covering membrane. These benefits are increasingly associated with a low cost of operation and a minimal risk of infection. PRF may further be utilized around immediate implant placement to pack gaps or additionally to speed soft tissue wound healing (Fig. 4). There remains however a great necessity to further evaluate dimensional changes utilizing PRF in various clinical situations using appropriately designed studies. Future clinical research

Table 2 Effects of PRF on root coverage of gingival recessions

Author	Study type	Patient no.	Healing period (months)	Treatment groups	Root coverage (%)	P value
Aroca 2009	Split-mouth; Miller class I or II	20	6	CAF	91.5	<0.004
				CAF + PRF	80.7	
Jankovic 2010	Split-mouth; Miller class I or II	20	12	CAF + EMD	70.5	n.s.
				CAF + PRF	72.1	
Aleksic 2010	Split-mouth; Miller class I or II	19	12	CAF + CTG	88.6	n.s.
				CAF + PRF	79.9	
Jankovic 2012	Split-mouth; Miller class I or II	15	6	CAF + CTG	88.7	n.s.
				CAF + PRF	92	
Padma 2013	Split-mouth; Miller class I or II	15	1, 3, and 6	CAF	68.4	<0.0001
				CAF + PRF	100	
Eren 2014	Split-mouth; Miller class I or II	22	6	CAF + CTG	94.2	n.s.
				CAF + PRF	92.7	
Tunaliota 2015	Split-mouth; Miller class I or II	22	12	CAF + CTG	77.4	n.s.
				CAF + PRF	76.6	
Thamaraiselvan 2015	Split-mouth; Miller class I or II	20	3 and 6	CAF	65	n.s.
				CAF + PRF	74.2	
Gupta 2015	Split-mouth; Miller class I or II	26	3 and 6	CAF	86.6	n.s.
				CAF + PRF	91	
Keceli 2015	Split-mouth; Miller class I or II	40	3 and 6	CAF + CTG	79.9	<0.05
				CAF + CTG + PRF	89.6	
Dogan 2015	Split-mouth; Miller class I or II	20	6	CAF	82.1	n.s.
				CAF + PRF	86.7	
Rajaram 2015	Split-mouth; Miller class II	20	12 and 24	DLSBF	80	n.s.
				DLSBF + PRF	78.8	
Agarwal 2016	Split-mouth; Miller class I or II	30	3 and 6	CAF	33	<0.05
				CAF + AM	36	
				CAF + PRF	56	

CAF coronally advanced flap, PRF platelet-rich fibrin, EMD enamel matrix derivative, CTG connective tissue graft, DLSBF double lateral sliding bridge flap, AM amniotic membrane

is therefore necessary. Furthermore, it remains unknown what effect PRF may play in combination with GBR techniques. While a collagen barrier membrane is routinely used during such procedures, additional use or replacement with PRF may provide further regenerative advantages when compared to collagen barrier membranes alone. Future studies are thus necessary to validate these potential advantages.


Sinus elevation procedures with PRF

The use of PRF for sinus elevation is relatively new with little comparative studies or standardized protocols available. Although the success rate of surgeries utilizing the addition of PRF is very high, it is difficult to compare the results between various treatment methods. Three authors using PRF alone as a graft material for sinus lift concluded that PRF significantly promoted bone healing with bone gains of 7.52 mm [72], 10.1 mm [73], and 10.4 mm [74] between the sinus floor and the top of the alveolar ridge. No controls were utilized in these studies. Nevertheless, no implants were lost at 6 months, 1 year, and 6 years in their respective studies

[72–74]. Some authors further claim that the use of PRF alone may be a valid treatment protocol for the majority of sinus lift cases although the lack of controls and limited number of studies have thus far limited its use [73].

Other studies compared the use of a bone grafting material with and without the additional use of PRF [75–78]. The results show that despite the large bone gains observed with PRF, no statistically significant differences were reported. One reported plausible advantage of combining PRF with a bone grafting material seems to result in a decrease in the overall healing time and better graft material handling [75–78]. A recent systematic review on the topic published in 2016 by Ali et al. found that of 290 initial publications searched, only eight met the inclusion criteria with approximately half not utilizing controls [79]. It was reported that the identified studies showed great heterogeneity regarding surgical technique, grafting material, implant placement time, surgical protocols, outcome measures, healing time for biopsy, implant placement, and follow-up periods [79]. In summary, although the results do not seem to confirm that PRF is better than other biomaterials, its ease of use, combined with its

Fig. 3 a Multiple gingival recessions from the canine to the molar in the upper jaw. *A* frontal view. *B–D* Lateral views (case performed by Dr. Giovanni Zucchelli). **b** Surgical technique: *A* A flap for multiple gingival recessions has been elevated with a split-full-split thickness approach. *B* A-PRF prepared. *C* A-PRF has been applied to cover all teeth affected by gingival recessions. Multiple layers have been applied. *D, E* Lateral view showing the thickness of A-PRF material applied to the root exposures. *F, G* Lateral view showing the flap coronally advanced and covering completely the A-PRF material. *H* Frontal view showing the flap covering in excess all gingival recessions (case performed by Dr. Giovanni Zucchelli). **c** Six months follow-up. *A* Complete root coverage with increase in keratinized tissue height has been achieved in all treated gingival recessions. *B–D* Lateral view showing the increase in gingival thickness at all teeth previously affected by gingival recessions (case performed by Dr. Giovanni Zucchelli)



minimal costs and high success rates, seems to illustrate that high success rates with minimal costs can be obtained by using PRF during sinus lifting procedures. Nevertheless, much further research is needed to support the beneficial effect of PRF.


Discussion

In this systematic review, all randomized clinical studies using PRF in dentistry were selected without discrepancies as compared to controls or commonly utilized surgical methods. The aim was to evaluate the current literature with respect to the clinical indications where PRF has been investigated for wound healing and/or tissue regeneration/repair. As was

observed in the analysis of its clinical applications, the performance of PRF was often compared either to conventional treatments such as OFD during intrabony/furcation defects or to naturally healing “blank” defects during extraction socket management. The analysis of the generated publications revealed a great heterogeneity of results with a general lack of conclusive evidence in large part due to the lack of study number with appropriate controls. Therefore, only guidelines can be drawn from the sum of these general conclusions with an obvious need for further research.

One factor that has been frequently reported in this systematic literature search was the ability for PRF to stimulate regeneration over a wide range of tissues. PRF has been shown to quickly stimulate tissue healing by significantly increasing the recruitment and proliferation of a variety of cells including

Fig. 4 Immediate implant placement in combination with the use of PRF. **a, b** Maxillary right cuspid fracture, extraction. **c** PRF placed into fresh extraction socket. **d** Dental implant placement with PRF fragments visible following placement (arrow in **e**). **f** Three-week follow-up, excellent soft tissue wound healing. **g** Three-year follow-up (case performed by Dr. Michael A. Pikos)



endothelial cells, gingival fibroblast, chondrocytes, and osteoblasts, thereby heavily promoting tissue repair and angiogenesis at the site of injury [80, 81]. These processes are regulated by local concentrations of cytokines and growth factors trapped within the fibrous scaffolding, most notably derived from autologous sources. In comparison, the growing use of two of the main growth factors approved by the FDA are PDGF and BMP derived from recombinant sources [82–85]. While these products sell for hundreds of dollars and are fabricated in mammalian cells or bacteria, the use of PRF are growth factors harvested purely from autologous sources via low-cost methods. It is therefore of interest to determine the benefit of using high supra-physiological concentrations of growth factors in recombinant form (i.e., BMP and PDGF) versus lower concentrations from autologous form (PRF). Although recombinant proteins have a regenerative potential well documented in the literature [86–88], many biological limitations to their use (swelling and edema), coupled with their low stability in vivo [89, 90], remain a limiting factor. Therefore, future research should target the comparison of the half-life and bioactivity of the growth factors found in PRF in comparison to commercially available recombinant growth factors.

Another interesting aspect that requires further study is to determine the regenerative/reparative potential of PRF on soft versus hard tissue formation. Thus far, this systematic review seems to point to the fact that the reparative potential of PRF favors soft tissue formation/ligament regeneration. Periodontitis is known to be one of the most common diseases with breakdown at the periodontium, causing destruction of

the cementum and periodontal ligament and intrabony defects [91]. The use of PRF specifically for intrabony defect repair showed significantly higher PPD reductions and CAL gains when compared to control OFD in all seven studies. Furthermore, a bone grafting material (DFDBA) could be combined with PRF to further generate statistically better CAL gains and PPD reductions [36]. Therefore, these findings demonstrate that PRF is able to support periodontal ligament repair as effective or potentially more effectively than commonly utilized biomaterials. Despite these positive findings, it remains of interest to determine if the reparative potential of PRF leads to true periodontal regeneration in humans. Therefore, future human histological studies are needed.

With respect to treatment and management of gingival recessions, PRF has been studied in 13 randomized human clinical studies. In general, it was found that PRF had similar advantages in root coverage of Miller class I and II defects when compared to CTG. Noteworthy, it was however commonly reported that significantly higher keratinized tissue width was found with CTG when compared to PRF. While it is difficult to evaluate significant differences in these treatment procedures due to the high success rates (generally observed over 80% root coverage for all treatment groups), one area of research that remains to be determined is precisely under which clinical situations should one expect similar results between PRF and CTG. Since CTG procedures are associated with high patient morbidity, it may be that in the future, such procedures could be substituted with PRF to prevent high morbidity. Furthermore, the technical ability of the clinician plays a more prominent role during CTG harvesting

when compared to PRF. Future studies are therefore needed to present updated clinical guidelines.

Another reported advantage to the use of PRF is its ability to decrease bacterial infections following surgery such as osteomyelitis commonly reported following third molar extractions [70]. In that study, a 9.5-fold significant decrease in reported cases of osteomyelitis was observed in a clinical trial with 100 patients [70]. Therefore and most likely due to the increase in white blood cells and macrophages capable of fighting infection, the use of PRF offers some antibacterial defense against incoming pathogens. Furthermore, macrophages have been shown to be key implicators in new bone formation both during bone modeling and remodeling, as well as in association with bone biomaterials [92–94]. Despite this, it remains interesting to point out that no human clinical study to date has investigated the effects of PRF in a controlled manner during GBR procedures, and only three studies on extraction socket healing have investigated dimensional changes following tooth extraction utilizing PRF (Supplemental Table 6). A similar trend investigating sinus lift procedures with PRF was also observed whereby a lack of well-designed studies with appropriate controls or endpoints was commonly found. Therefore, the effect of PRF on pure bone regeneration remains questionable and requires more validating studies. Similarly, various reports have now supported the use of PRF for pulp regeneration, cystic bone defect, and papilla augmentation under various clinical indications to improve healing [95–103]. While these reports are rare and anecdotal, future research aimed at better characterizing the regenerative potential of PRF for various other dental procedures remains necessary.

Very recently, a team of researchers has convincingly shown that lower centrifugation speeds and time resulted in higher leukocyte concentrations and release of growth factors [104–108]. Ghanaati et al. demonstrated via histological processing of PRF scaffolds that at higher centrifugation speeds, the majority of leukocytes were found at the bottom of PRF scaffolds [104]. By reducing centrifugation g-force, leukocytes were more evenly distributed throughout the PRF scaffolds [104]. In addition, regenerative growth factors released and gingival fibroblast activity are increased when utilizing slower centrifugation speed and time [108]. While reports from these studies support modifications to centrifugation protocols, the impact these may have on clinical outcomes in the various indications highlighted throughout this review article remains to be investigated. Future clinical study is therefore needed.

In conclusion, this systematic review demonstrates the widespread use of PRF in dentistry in various clinical settings. Although this regenerative modality remains unfamiliar to many clinicians, the evidence supporting its use has accumulated over the years, demonstrating its ability to improve tissue regeneration. The combination of PRF with regenerative

therapy has been shown to be most promising for periodontal repair of intrabony and furcation defects, as well as soft tissue root coverage of gingival recessions. Furthermore, evidence from the literature suggests that PRF is able to decrease infection following tooth extraction and may further limit dimensional changes following tooth loss. It was also concluded that regeneration of bone defects (GBR procedures and sinus elevation) necessitates more study with focused endpoints. Nevertheless, its ease of use, combined with its low cost and autologous source, makes it an ideal biomaterial worth further investigation across a variety of surgical procedures in dentistry.

Compliance with ethical standards

Conflict of interest Joseph Choukroun is the founder of Process of PRF company and the developer and inventor of PRF protocol in Nice, France. All other authors declare no conflict of interest.

Funding This work was fully funded by the Cell Biology Laboratory at Nova Southeastern University, College of Dental Medicine.

Ethical approval No ethical approval was required for this study as human participants or animals were not utilized in this study.

Informed consent Informed consent was not required as no human or animal subjects were utilized in this study.

References

- Dangaria SJ, Ito Y, Walker C, Druzinsky R, Luan X, Diekwisch TG (2009) Extracellular matrix-mediated differentiation of periodontal progenitor cells. *Differentiation* 78:79–90. doi:10.1016/j.diff.2009.03.005
- Hollander A, Macchiarini P, Gordijn B, Birchall M (2009) The first stem cell-based tissue-engineered organ replacement: implications for regenerative medicine and society. *Regen Med* 4:147–148. doi:10.2217/17460751.4.2.147
- Choukroun J, Adda F, Schoeffler C, Vervelle A (2001) Une opportunité en paro-implantologie: le PRF. *Implantodontie* 42:e62
- Dohan DM, Choukroun J, Diss A, Dohan SL, Dohan AJ, Mouhyi J, Gogly B (2006) Platelet-rich fibrin (PRF): a second-generation platelet concentrate. Part II: platelet-related biologic features. *Oral Surg, Oral Med, Oral Pathol, Oral Radiol Endod* 101:e45–e50. doi:10.1016/j.tripleo.2005.07.009
- Kang YH, Jeon SH, Park JY, Chung JH, Choung YH, Choung HW, Kim ES, Choung PH (2011) Platelet-rich fibrin is a bioscaffold and reservoir of growth factors for tissue regeneration. *Tissue Eng A* 17:349–359. doi:10.1089/ten.TEA.2010.0327
- He L, Lin Y, Hu X, Zhang Y, Wu H (2009) A comparative study of platelet-rich fibrin (PRF) and platelet-rich plasma (PRP) on the effect of proliferation and differentiation of rat osteoblasts in vitro. *Oral Surg, Oral Med, Oral Pathol, Oral Radiol Endod* 108:707–713. doi:10.1016/j.tripleo.2009.06.044
- Dohan Ehrenfest DM, Diss A, Odin G, Doglioli P, Hippolyte MP, Charrier JB (2009) In vitro effects of Choukroun's PRF (platelet-rich fibrin) on human gingival fibroblasts, dermal prekeratinocytes, preadipocytes, and maxillofacial osteoblasts in

- primary cultures. *Oral Surg, Oral Med, Oral Pathol, Oral Radiol Endod* 108:341–352. doi:10.1016/j.tripleo.2009.04.020
8. Anfossi G, Trovati M, Mularoni E, Massucco P, Calcamuggi G, Emanuelli G (1989) Influence of propranolol on platelet aggregation and thromboxane B2 production from platelet-rich plasma and whole blood. *Prostaglandins Leukot Essent Fat Acids* 36:1–7
 9. Fijnheer R, Pietersz RN, de Korte D, Gouwerok CW, Dekker WJ, Reesink HW, Roos D (1990) Platelet activation during preparation of platelet concentrates: a comparison of the platelet-rich plasma and the buffy coat methods. *Transfusion* 30:634–638
 10. Jameson C (2007) Autologous platelet concentrate for the production of platelet gel. *Lab Med* 38:39–42
 11. Whitman DH, Berry RL, Green DM (1997) Platelet gel: an autologous alternative to fibrin glue with applications in oral and maxillofacial surgery. *J Oral Maxillofac Surg* 55:1294–1299
 12. Marx RE, Carlson ER, Eichstaedt RM, Schimmele SR, Strauss JE, Georgeff KR (1998) Platelet-rich plasma: growth factor enhancement for bone grafts. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 85:638–646
 13. Marx RE (2004) Platelet-rich plasma: evidence to support its use. *J Oral Maxillofac Surg* 62:489–496
 14. Anitua E (1999) Plasma rich in growth factors: preliminary results of use in the preparation of future sites for implants. *Int J Oral Maxillofac Implants* 14:529–535
 15. Anitua E, Prado R, Troya M, Zalduendo M, de la Fuente M, Pino A, Muruzabal F, Orive G (2016) Implementation of a more physiological plasma rich in growth factor (PRGF) protocol: anticoagulant removal and reduction in activator concentration. *Platelets* 27:459–466. doi:10.3109/09537104.2016.1143921
 16. Lucarelli E, Beretta R, Dozza B, Tazzari PL, O'Connell SM, Ricci F, Pierini M, Squarizoni S, Pagliaro PP, Oprita EI, Donati D (2010) A recently developed bifacial platelet-rich fibrin matrix. *Eur Cell Mater* 20:13–23
 17. Saluja H, Dehane V, Mahindra U (2011) Platelet-rich fibrin: a second generation platelet concentrate and a new friend of oral and maxillofacial surgeons. *Ann Maxillofac Surg* 1:53–57. doi:10.4103/2231-0746.83158
 18. Kobayashi E, Fluckiger L, Fujioka-Kobayashi M, Sawada K, Sculean A, Schaller B, Miron RJ (2016) Comparative release of growth factors from PRP, PRF, and advanced-PRF. *Clin Oral Investig*. doi:10.1007/s00784-016-1719-1
 19. Dohan Ehrenfest DM, Del Corso M, Diss A, Mouhyi J, Charrier JB (2010) Three-dimensional architecture and cell composition of a Choukroun's platelet-rich fibrin clot and membrane. *J Periodontol* 81:546–555. doi:10.1902/jop.2009.090531
 20. Toffler M, Toscano N, Holtzclaw D, Corso M, Dohan D (2009) Introducing Choukroun's platelet rich fibrin (PRF) to the reconstructive surgery milieu. *J Implant Adv Clin Dent* 1:22–31
 21. Tsay RC, Vo J, Burke A, Eisig SB, Lu HH, Landesberg R (2005) Differential growth factor retention by platelet-rich plasma composites. *J Oral Maxillofac Surg* 63:521–528. doi:10.1016/j.joms.2004.09.012
 22. Carlson NE, Roach RB Jr (2002) Platelet-rich plasma: clinical applications in dentistry. *J Am Dent Assoc* 133(13):1383–1386
 23. Andrades JA, Han B, Becerra J, Sorgente N, Hall FL, Nimni ME (1999) A recombinant human TGF-beta1 fusion protein with collagen-binding domain promotes migration, growth, and differentiation of bone marrow mesenchymal cells. *Exp Cell Res* 250:485–498. doi:10.1006/excr.1999.4528
 24. Lind M, Deleuran B, Thestrup-Pedersen K, Soballe K, Eriksen EF, Bunger C (1995) Chemotaxis of human osteoblasts. Effects of osteotropic growth factors. *APMIS* 103:140–146
 25. Dohan DM, Choukroun J, Diss A, Dohan SL, Dohan AJ, Mouhyi J, Gogly B (2006) Platelet-rich fibrin (PRF): a second-generation platelet concentrate. Part III: leucocyte activation: a new feature for platelet concentrates? *Oral Surg, Oral Med Oral Pathol Oral Radiol Endod* 101:e51–e55. doi:10.1016/j.tripleo.2005.07.010
 26. Grando Mattuella L, Poli de Figueiredo JA, Nor JE, de Araujo FB, Medeiros Fossati AC (2007) Vascular endothelial growth factor receptor-2 expression in the pulp of human primary and young permanent teeth. *J Endod* 33:1408–1412. doi:10.1016/j.joen.2007.08.019
 27. Troost E, Hold GL, Smith MG, Chow WH, Rabkin CS, McColl KE, El-Omar EM (2003) The role of interleukin-1beta and other potential genetic markers as indicators of gastric cancer risk. *Can J Gastroenterol* 17(Suppl B):8B–12B
 28. Kishimoto T, Akira S, Narazaki M, Taga T (1995) Interleukin-6 family of cytokines and gp130. *Blood* 86:1243–1254
 29. Kishimoto T (1989) The biology of interleukin-6. *Blood* 74:1–10
 30. Waters JP, Pober JS, Bradley JR (2013) Tumour necrosis factor in infectious disease. *J Pathol* 230:132–147. doi:10.1002/path.4187
 31. Murtaugh MP, Johnson CR, Xiao Z, Scamurra RW, Zhou Y (2009) Species specialization in cytokine biology: is interleukin-4 central to the T(H)1-T(H)2 paradigm in swine? *Dev Comp Immunol* 33:344–352. doi:10.1016/j.dci.2008.06.014
 32. Li Z, Zhang Y, Sun B (2011) Current understanding of Th2 cell differentiation and function. *Protein Cell* 2:604–611. doi:10.1007/s13238-011-1083-5
 33. Choukroun J, Diss A, Simonpieri A, Girard MO, Schoeffler C, Dohan SL, Dohan AJ, Mouhyi J, Dohan DM (2006) Platelet-rich fibrin (PRF): a second-generation platelet concentrate. Part IV: clinical effects on tissue healing. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 101:e56–e60. doi:10.1016/j.tripleo.2005.07.011
 34. Clark RA (2001) Fibrin and wound healing. *Ann N Y Acad Sci* 936:355–367
 35. Swartz MK (2011) The PRISMA statement: a guideline for systematic reviews and meta-analyses. *J Pediatr Health Care* 25:1–2. doi:10.1016/j.pedhc.2010.09.006
 36. Agarwal A, Gupta ND, Jain A (2016) Platelet rich fibrin combined with decalcified freeze-dried bone allograft for the treatment of human intrabony periodontal defects: a randomized split mouth clinical trial. *Acta Odontol Scand* 74:36–43. doi:10.3109/00016357.2015.1035672
 37. Ajwani H, Shetty S, Gopalakrishnan D, Kathariya R, Kulloli A, Dolas RS, Pradeep AR (2015) Comparative evaluation of platelet-rich fibrin biomaterial and open flap debridement in the treatment of two and three wall intrabony defects. *J Int Oral Health* 7:32–37
 38. Elgendy EA, Abo Shady TE (2015) Clinical and radiographic evaluation of nanocrystalline hydroxyapatite with or without platelet-rich fibrin membrane in the treatment of periodontal intrabony defects. *J Indian Soc Periodontol* 19:61–65. doi:10.4103/0972-124x.148639
 39. Panda S, Sankari M, Satpathy A, Jayakumar D, Mozzati M, Mortellaro C, Gallesio G, Taschieri S, Del Fabbro M (2016) Adjunctive effect of autologous platelet-rich fibrin to barrier membrane in the treatment of periodontal intrabony defects. *J Craniofac Surg* 27:691–696. doi:10.1097/scs.0000000000002524
 40. Pradeep AR, Nagpal K, Karvekar S, Patnaik K, Naik SB, Guruprasad CN (2015) Platelet-rich fibrin with 1% metformin for the treatment of intrabony defects in chronic periodontitis: a randomized controlled clinical trial. *J Periodontol* 86:729–737. doi:10.1902/jop.2015.140646
 41. Pradeep AR, Rao NS, Agarwal E, Bajaj P, Kumari M, Naik SB (2012) Comparative evaluation of autologous platelet-rich fibrin and platelet-rich plasma in the treatment of 3-wall intrabony defects in chronic periodontitis: a randomized controlled clinical trial. *J Periodontol* 83:1499–1507. doi:10.1902/jop.2012.110705
 42. Shah M, Patel J, Dave D, Shah S (2015) Comparative evaluation of platelet-rich fibrin with demineralized freeze-dried bone allograft in periodontal infrabony defects: a randomized controlled

- clinical study. *J Indian Soc Periodontol* 19:56–60. doi:10.4103/0972-124x.145803
43. Thorat M, Pradeep AR, Pallavi B (2011) Clinical effect of autologous platelet-rich fibrin in the treatment of intra-bony defects: a controlled clinical trial. *J Clin Periodontol* 38:925–932. doi:10.1111/j.1600-051X.2011.01760.x
 44. Pradeep AR, Bajaj P, Rao NS, Agarwal E, Naik SB (2012) Platelet-rich fibrin combined with a porous hydroxyapatite graft for the treatment of three-wall intrabony defects in chronic periodontitis: a randomized controlled clinical trial. *J Periodontol*. doi:10.1902/jop.2012.110722
 45. Sharma A, Pradeep AR (2011) Treatment of 3-wall intrabony defects in patients with chronic periodontitis with autologous platelet-rich fibrin: a randomized controlled clinical trial. *J Periodontol* 82:1705–1712. doi:10.1902/jop.2011.110075
 46. Sharma A, Pradeep AR (2011) Autologous platelet-rich fibrin in the treatment of mandibular degree II furcation defects: a randomized clinical trial. *J Periodontol* 82:1396–1403. doi:10.1902/jop.2011.100731
 47. Bajaj P, Pradeep AR, Agarwal E, Rao NS, Naik SB, Priyanka N, Kalra N (2013) Comparative evaluation of autologous platelet-rich fibrin and platelet-rich plasma in the treatment of mandibular degree II furcation defects: a randomized controlled clinical trial. *J Periodontol Res*. doi:10.1111/jre.12040
 48. Pradeep AR, Karvekar S, Nagpal K, Patnaik K, Raju A, Singh P (2016) Rosuvastatin 1.2 mg in situ gel combined with 1:1 mixture of autologous platelet-rich fibrin and porous hydroxyapatite bone graft in surgical treatment of mandibular class II furcation defects: a randomized clinical control trial. *J Periodontol* 87:5–13. doi:10.1902/jop.2015.150131
 49. Agarwal SK, Jhingran R, Bains VK, Srivastava R, Madan R, Rizvi I (2016) Patient-centered evaluation of microsurgical management of gingival recession using coronally advanced flap with platelet-rich fibrin or amnion membrane: a comparative analysis. *Eur J Dent* 10:121–133. doi:10.4103/1305-7456.175686
 50. Aleksic Z, Jankovic S, Dimitrijevic B, Divnic-Resnik T, Milinkovic I, Lekovic V (2010) The use of platelet-rich fibrin membrane in gingival recession treatment. *Srp Arh Celok Lek* 138:11–18
 51. Aroca S, Keglevich T, Barbieri B, Gera I, Etienne D (2009) Clinical evaluation of a modified coronally advanced flap alone or in combination with a platelet-rich fibrin membrane for the treatment of adjacent multiple gingival recessions: a 6-month study. *J Periodontol* 80:244–252. doi:10.1902/jop.2009.080253
 52. Dogan SB, Dede FO, Balli U, Atalay EN, Durmuslar MC (2015) Concentrated growth factor in the treatment of adjacent multiple gingival recessions: a split-mouth randomized clinical trial. *J Clin Periodontol* 42:868–875. doi:10.1111/jcpe.12444
 53. Eren G, Atilla G (2014) Platelet-rich fibrin in the treatment of localized gingival recessions: a split-mouth randomized clinical trial. *Clin Oral Investig* 18:1941–1948. doi:10.1007/s00784-013-1170-5
 54. Gupta S, Banthia R, Singh P, Banthia P, Raje S, Aggarwal N (2015) Clinical evaluation and comparison of the efficacy of coronally advanced flap alone and in combination with platelet rich fibrin membrane in the treatment of Miller class I and II gingival recessions. *Contemp Clin Dent* 6:153–160. doi:10.4103/0976-237x.156034
 55. Jankovic S, Aleksic Z, Klokkevold P, Lekovic V, Dimitrijevic B, Kenney EB, Camargo P (2012) Use of platelet-rich fibrin membrane following treatment of gingival recession: a randomized clinical trial. *Int J Periodontics Restorative Dent* 32:e41–e50
 56. Jankovic S, Aleksic Z, Milinkovic I, Dimitrijevic B (2010) The coronally advanced flap in combination with platelet-rich fibrin (PRF) and enamel matrix derivative in the treatment of gingival recession: a comparative study. *Eur J Esthet Dent* 5:260–273
 57. Keceli HG, Kamak G, Erdemir EO, Evginer MS, Dolgun A (2015) The adjunctive effect of platelet-rich fibrin to connective tissue graft in the treatment of buccal recession defects: results of a randomized, parallel-group controlled trial. *J Periodontol* 86:1221–1230. doi:10.1902/jop.2015.150015
 58. Padma R, Shilpa A, Kumar PA, Nagasri M, Kumar C, Sreedhar A (2013) A split mouth randomized controlled study to evaluate the adjunctive effect of platelet-rich fibrin to coronally advanced flap in Miller's class-I and II recession defects. *J Indian Soc Periodontol* 17:631–636. doi:10.4103/0972-124x.119281
 59. Rajaram V, Thyegarajan R, Balachandran A, Aari G, Kanakamedala A (2015) Platelet rich fibrin in double lateral sliding bridge flap procedure for gingival recession coverage: an original study. *J Indian Soc Periodontol* 19:665–670. doi:10.4103/0972-124x.164764
 60. Thamaraiselvan M, Elavarasu S, Thangakumaran S, Gadagi JS, Arthie T (2015) Comparative clinical evaluation of coronally advanced flap with or without platelet rich fibrin membrane in the treatment of isolated gingival recession. *J Indian Soc Periodontol* 19:66–71. doi:10.4103/0972-124x.145790
 61. Tunalioa M, Ozdemir H, Arabaciota T, Gurbuzer B, Pikkoken L, Firatli E (2015) Clinical evaluation of autologous platelet-rich fibrin in the treatment of multiple adjacent gingival recession defects: a 12-month study. *Int J Periodontics Restorative Dent* 35:105–114. doi:10.11607/prd.1826
 62. Femminella B, Iaconi MC, Di Tullio M, Romano L, Sinjari B, D'Arcangelo C, De Ninis P, Paolantonio M (2016) Clinical comparison of platelet-rich fibrin and a gelatin sponge in the management of palatal wounds after epithelialized free gingival graft harvest: a randomized clinical trial. *J Periodontol* 87:103–113. doi:10.1902/jop.2015.150198
 63. Moraschini V, Barboza Edos S (2016) Use of platelet-rich fibrin membrane in the treatment of gingival recession: a systematic review and meta-analysis. *J Periodontol* 87:281–290. doi:10.1902/jop.2015.150420
 64. De Risi V, Clementini M, Vittorini G, Mannocci A, De Sanctis M (2015) Alveolar ridge preservation techniques: a systematic review and meta-analysis of histological and histomorphometrical data. *Clin Oral Implants Res* 26:50–68. doi:10.1111/clr.12288
 65. Jambhekar S, Kernen F, Bidra AS (2015) Clinical and histologic outcomes of socket grafting after flapless tooth extraction: a systematic review of randomized controlled clinical trials. *J Prosthet Dent* 113:371–382. doi:10.1016/j.prosdent.2014.12.009
 66. Moraschini V, Barboza ED (2016) Quality assessment of systematic reviews on alveolar socket preservation. *Int J Oral Maxillofac Surg*. doi:10.1016/j.ijom.2016.03.010
 67. Chappuis V, Engel O, Reyes M, Shahim K, Nolte LP, Buser D (2013) Ridge alterations post-extraction in the esthetic zone: a 3D analysis with CBCT. *J Dental Res* 92:195s–201s. doi:10.1177/0022034513506713
 68. Girish Rao S, Bhat P, Nagesh KS, Rao GH, Mirlle B, Kharbhari L, Gangaprasad B (2013) Bone regeneration in extraction sockets with autologous platelet rich fibrin gel. *J Maxillofac Oral Surgery* 12:11–16. doi:10.1007/s12663-012-0370-x
 69. Suttapreyasri S, Leepong N (2013) Influence of platelet-rich fibrin on alveolar ridge preservation. *J Craniofac Surg* 24:1088–1094. doi:10.1097/SCS.0b013e31828b6dc3
 70. Hoaglin DR, Lines GK (2013) Prevention of localized osteitis in mandibular third-molar sites using platelet-rich fibrin. *Int J Dent* 2013:875380. doi:10.1155/2013/875380
 71. Hauser F, Gaydarov N, Badoud I, Vazquez L, Bernard JP, Ammann P (2013) Clinical and histological evaluation of postextraction platelet-rich fibrin socket filling: a prospective randomized controlled study. *Implant Dent* 22:295–303. doi:10.1097/ID.0b013e3182906eb3

72. Tajima N, Ohba S, Sawase T, Asahina I (2013) Evaluation of sinus floor augmentation with simultaneous implant placement using platelet-rich fibrin as sole grafting material. *Int J Oral Maxillofac Implants* 28:77–83. doi:10.11607/jomi.2613
73. Mazor Z, Horowitz RA, Del Corso M, Prasad HS, Rohrer MD, Dohan Ehrenfest DM (2009) Sinus floor augmentation with simultaneous implant placement using Choukroun's platelet-rich fibrin as the sole grafting material: a radiologic and histologic study at 6 months. *J Periodontol* 80:2056–2064. doi:10.1902/jop.2009.090252
74. Simonpieri A, Choukroun J, Del Corso M, Sammartino G, Dohan Ehrenfest DM (2011) Simultaneous sinus-lift and implantation using microthreaded implants and leukocyte- and platelet-rich fibrin as sole grafting material: a six-year experience. *Implant Dent* 20:2–12. doi:10.1097/ID.0b013e3181faa8af
75. Inchingolo F, Tatullo M, Marrelli M, Inchingolo AM, Scacco S, Inchingolo AD, Dipalma G, Vermesan D, Abbinante A, Cagiano R (2010) Trial with platelet-rich fibrin and Bio-Oss used as grafting materials in the treatment of the severe maxillary bone atrophy: clinical and radiological evaluations. *Eur Rev Med Pharmacol Sci* 14:1075–1084
76. Tatullo M, Marrelli M, Cassetta M, Pacifici A, Stefanelli LV, Scacco S, Dipalma G, Pacifici L, Inchingolo F (2012) Platelet rich fibrin (P.R.F.) in reconstructive surgery of atrophied maxillary bones: clinical and histological evaluations. *Int J Med Sci* 9: 872–880. doi:10.7150/ijms.5119
77. Zhang Y, Tangl S, Huber CD, Lin Y, Qiu L, Rausch-Fan X (2012) Effects of Choukroun's platelet-rich fibrin on bone regeneration in combination with deproteinized bovine bone mineral in maxillary sinus augmentation: a histological and histomorphometric study. *J Cranio-Maxillo-Facial Surg* 40:321–328. doi:10.1016/j.jcms.2011.04.020
78. Choukroun J, Diss A, Simonpieri A, Girard MO, Schoeffler C, Dohan SL, Dohan AJ, Mouhyi J, Dohan DM (2006) Platelet-rich fibrin (PRF): a second-generation platelet concentrate. Part V: histologic evaluations of PRF effects on bone allograft maturation in sinus lift. *Oral Surg Oral Med Oral Pathol Oral Rad Endod* 101: 299–303. doi:10.1016/j.tripleo.2005.07.012
79. Ali S, Bakry SA, Abd-Elhakam H (2015) Platelet-rich fibrin in maxillary sinus augmentation: a systematic review. *J Oral Implantol* 41:746–753. doi:10.1563/aaid-joi-D-14-00167
80. Roy S, Driggs J, Elgharably H, Biswas S, Findley M, Khanna S, Gnyawali U, Bergdall VK, Sen CK (2011) Platelet-rich fibrin matrix improves wound angiogenesis via inducing endothelial cell proliferation. *Wound Repair Regen* 19:753–766. doi:10.1111/j.1524-475X.2011.00740.x
81. Chen FM, Wu LA, Zhang M, Zhang R, Sun HH (2011) Homing of endogenous stem/progenitor cells for in situ tissue regeneration: promises, strategies, and translational perspectives. *Biomaterials* 32:3189–3209. doi:10.1016/j.biomaterials.2010.12.032
82. Steed DL, Donohoe D, Webster MW, Lindsley L (1996) Effect of extensive debridement and treatment on the healing of diabetic foot ulcers. *Diabetic Ulcer Study Group. J Am Coll Surg* 183: 61–64
83. Wieman TJ, Smiell JM, Su Y (1998) Efficacy and safety of a topical gel formulation of recombinant human platelet-derived growth factor-BB (becaplermin) in patients with chronic neuropathic diabetic ulcers. A phase III randomized placebo-controlled double-blind study. *Diabetes Care* 21:822–827
84. White AP, Vaccaro AR, Hall JA, Whang PG, Friel BC, McKee MD (2007) Clinical applications of BMP-7/OP-1 in fractures, nonunions and spinal fusion. *Int Orthop* 31:735–741. doi:10.1007/s00264-007-0422-x
85. Miron RJ, Zhang YF (2012) Osteoinduction: a review of old concepts with new standards. *J Dent Res* 91:736–744. doi:10.1177/0022034511435260
86. Young CS, Ladd PA, Browning CF, Thompson A, Bonomo J, Shockley K, Hart CE (2009) Release, biological potency, and biochemical integrity of recombinant human platelet-derived growth factor-BB (rhPDGF-BB) combined with Augment(TM) Bone Graft or GEM 21S beta-tricalcium phosphate (beta-TCP). *J Control Release* 140:250–255. doi:10.1016/j.jconrel.2009.06.030
87. Park YJ, Lee YM, Lee JY, Seol YJ, Chung CP, Lee SJ (2000) Controlled release of platelet-derived growth factor-BB from chondroitin sulfate-chitosan sponge for guided bone regeneration. *J Control Release* 67:385–394
88. Wissink MJ, Beemink R, Poot AA, Engbers GH, Beugeling T, van Aken WG, Feijen J (2000) Improved endothelialization of vascular grafts by local release of growth factor from heparinized collagen matrices. *J Control Release* 64:103–114
89. Delgado JJ, Evora C, Sanchez E, Baro M, Delgado A (2006) Validation of a method for non-invasive in vivo measurement of growth factor release from a local delivery system in bone. *J Control Release* 114:223–229
90. Oe S, Fukunaka Y, Hirose T, Yamaoka Y, Tabata Y (2003) A trial on regeneration therapy of rat liver cirrhosis by controlled release of hepatocyte growth factor. *J Control Release* 88:193–200
91. Sculean A, Gruber R, Bosshardt DD (2014) Soft tissue wound healing around teeth and dental implants. *J Clin Periodontol* 41(Suppl 15):S6–22. doi:10.1111/jcpe.12206
92. Adamson R (2009) Role of macrophages in normal wound healing: an overview. *J Wound Care* 18:349–351. doi:10.12968/jowc.2009.18.8.43636
93. Miron RJ, Bosshardt DD (2016) OsteoMacs: key players around bone biomaterials. *Biomaterials* 82:1–19. doi:10.1016/j.biomaterials.2015.12.017
94. Sinder BP, Pettit AR, McCauley LK (2015) Macrophages: their emerging roles in bone. *J Bone Miner Res* 30:2140–2149. doi:10.1002/jbmr.2735
95. Subash D, Shoba K, Aman S, Bharkavi SK (2016) Revitalization of an immature permanent mandibular molar with a necrotic pulp using platelet-rich fibrin: a case report. *J Clin Diagn Res* 10:Zd21–zd23. doi:10.7860/jcdr/2016/21793.8902
96. Rebentish PD, Umashetty G, Kaur H, Doizode T, Kaslekar M, Chowdhury S (2016) Platelet-rich fibrin: a boon in regenerative endodontics. *Minerva Stomatol* 65:385–392
97. Kim JH, Woo SM, Choi NK, Kim WJ, Kim SM, Jung JY (2017) Effect of platelet-rich fibrin on odontoblastic differentiation in human dental pulp cells exposed to lipopolysaccharide. *J Endod* 43:433–438. doi:10.1016/j.joen.2016.11.002
98. Bakhtiar H, Esmaeili S, Fakhr Tabatabayi S, Ellini MR, Nekoofar MH, Dummer PM (2017) Second-generation platelet concentrate (platelet-rich fibrin) as a scaffold in regenerative endodontics: a case series. *J Endod* 43:401–408. doi:10.1016/j.joen.2016.10.016
99. Pradeep K, Kudva A, Narayanamoorthy V, Cariappa KM, Saraswathi MV (2016) Platelet-rich fibrin combined with synthetic nanocrystalline hydroxy apatite granules in the management of radicular cyst. *Niger J Clin Pract* 19:688–691. doi:10.4103/1119-3077.188711
100. Mirkovic S, Djurdjevic-Mirkovic T, Pugkar T (2015) Application of concentrated growth factors in reconstruction of bone defects after removal of large jaw cysts—the two cases report. *Vojnosanit Pregl* 72:368–371
101. Meshram VS, Lambade PN, Meshram PV, Kadu A, Tiwari MS (2015) The autologous platelet rich fibrin: a novel approach in osseous regeneration after cystic enucleation: a pilot study. *Indian J Dent Res* 26:560–564. doi:10.4103/0970-9290.176915
102. Dar M, Hakim T, Shah A, Najjar L, Yaqoob G, Lanker F (2016) Use of autologous platelet-rich fibrin in osseous regeneration after cystic enucleation: a clinical study. *J Oral Biol Craniofac Res* 6: S29–s32. doi:10.1016/j.jobcr.2016.04.004

103. Arunachalam LT, Merugu S, Sudhakar U (2012) A novel surgical procedure for papilla reconstruction using platelet rich fibrin. *Contemp Clin Dent* 3:467–470. doi:[10.4103/0976-237x.107443](https://doi.org/10.4103/0976-237x.107443)
104. Ghanaati S, Booms P, Orłowska A, Kubesch A, Lorenz J, Rutkowski J, Landes C, Sader R, Kirkpatrick C, Choukroun J (2014) Advanced platelet-rich fibrin: a new concept for cell-based tissue engineering by means of inflammatory cells. *J Oral Implantol* 40:679–689. doi:[10.1563/aaid-joi-D-14-00138](https://doi.org/10.1563/aaid-joi-D-14-00138)
105. Choukroun J, Ghanaati S (2017) Reduction of relative centrifugation force within injectable platelet-rich-fibrin (PRF) concentrates advances patients' own inflammatory cells, platelets and growth factors: the first introduction to the low speed centrifugation concept. *Eur J Trauma Emerg Surg*. doi:[10.1007/s00068-017-0767-9](https://doi.org/10.1007/s00068-017-0767-9)
106. El Bagdadi K, Kubesch A, Yu X, Al-Maawi S, Orłowska A, Dias A, Booms P, Dohle E, Sader R, Kirkpatrick CJ, Choukroun J, Ghanaati S (2017) Reduction of relative centrifugal forces increases growth factor release within solid platelet-rich-fibrin (PRF)-based matrices: a proof of concept of LSCC (low speed centrifugation concept). *Eur J Trauma Emerg Surg*. doi:[10.1007/s00068-017-0785-7](https://doi.org/10.1007/s00068-017-0785-7)
107. Kobayashi E, Fluckiger L, Fujioka-Kobayashi M, Sawada K, Sculean A, Schaller B, Miron RJ (2016) Comparative release of growth factors from PRP, PRF, and advanced-PRF. *Clin Oral Investig* 20:2353–2360. doi:[10.1007/s00784-016-1719-1](https://doi.org/10.1007/s00784-016-1719-1)
108. Fujioka-Kobayashi M, Miron RJ, Hernandez M, Kandalam U, Zhang Y, Choukroun J (2017) Optimized platelet-rich fibrin with the low-speed concept: growth factor release, biocompatibility, and cellular response. *J Periodontol* 88:112–121. doi:[10.1902/jop.2016.160443](https://doi.org/10.1902/jop.2016.160443)