

Comment to: Desarda's technique versus Lichtenstein technique for the treatment of primary inguinal hernia: a systematic review and meta-analysis of randomized controlled trials. Emile SH, Elfeki H

J. Li¹ · W. Zhang¹

Received: 2 October 2017 / Accepted: 3 November 2017 / Published online: 10 November 2017
© Springer-Verlag France SAS, part of Springer Nature 2017

Dear Editors:

We read with great interest the recent article by Emile SH. [1], published in *Hernia*. The authors present data from a systematic review and meta-analysis comparing tissue-based repair (Desarda's technique) and Lichtenstein technique for inguinal hernia repair. Based on the analysis of six RCTs comprising 2159 patients [1], they concluded that both Desarda and Lichtenstein technique had similar low recurrence rate; furthermore, Desarda's technique had lower complication rates compared with Lichtenstein technique.

We appreciate for the authors' thorough analysis, and we also very pleased to see the favorable results of the tissue-based inguinal hernia repair technique. Furthermore, their results [1] offered a number of important points to comment, and provided the clues to some important questions in inguinal hernia repair, such as "the ideal layer" and the "ideal tissue" for inguinal hernia repair.

First, it is no doubt that the debate on the benefits of tissue-based repair or prosthetic repair of symptomatic inguinal hernias will continue for some time. Furthermore, the mesh position, either in preperitoneal space or in superficial as Lichtenstein, is still a dispute. Although Lichtenstein technique was strongly recommended by the Europe hernia society (EHS) [2], it is still not the ideal solution due to the consequence after mesh implantation, including stiffness, chronic pain, and foreign body sensation. Thus, the

efforts of searching for non-prosthetic repair never stopped. Next, apart from mesh repair, what is the "ideal" tissue for repair? Most of the conventional tissue-based repairs aim to construct the posterior abdominal wall using patient's deep abdominal wall muscular tissue and fascia, and especially, the transversalis fascia gained much attention [3], such as the Shouldice procedure. However, the Desarda's technique uses the external oblique aponeurosis (EOA), we applauded for the fine and reasonable idea, since the EOA is not only strong, but more superficial and easy to manipulate, which makes the procedure simple. Furthermore, this new established layer, quite resemble the layer of the mesh placement in Lichtenstein procedure, which in part proved the reasonable layer for inguinal hernia repair. Third, most of the meshes for hernia repair are over-weight and much stronger than needed [4]. The use of patients' own tissue of the EOA, although not as strong as a prosthetic mesh, but obviously stronger than the transversalis fascia, thus, the EOA may be a sound alternative tissue material in the majority of inguinal hernia cases, and provides a clue to the path to tissue repair.

Compliance with ethical standards

Funding None.

Conflict of interest No conflict of interest to be declared by the authors.

Ethical approval This article did not require ethical approval of any kind.

Human and animal rights This article does not contain any studies with human participants or animals performed by any of the authors.

Informed consent This article does not include patients, and therefore, informed consent was not applicable.

This comment refers to the article available at doi:[10.1007/s10029-017-1666-z](https://doi.org/10.1007/s10029-017-1666-z).

✉ J. Li
Lijunshenghd@126.com

¹ Department of General Surgery, Affiliated Zhongda Hospital, Southeast University, Nanjing 210009, China

References

1. Emile SH, Elfeki H (2017) Desarda's technique versus Lichtenstein technique for the treatment of primary inguinal hernia: a systematic review and meta-analysis of randomized controlled trials. *Hernia*. <https://doi.org/10.1007/s10029-017-1666-z> (Epub ahead of print)
2. Simons MP, Aufenacker T, Bay-Nielsen M et al (2009) European Hernia Society guidelines on the treatment of inguinal hernia in adult patients. *Hernia* 13:343–403
3. Memon MA, Quinn TH, Cahill DR (1999) Transversalis fascia: historical aspects and its place in contemporary inguinal herniorrhaphy. *J Laparoendosc Adv Surg Tech A* 9(3):267–272
4. Klinge U, Conze J, Limberg W, Brucker C, Ottinger AP, Schumpelick V (1996) Pathophysiology of the abdominal wall. *Chirurg* 67:229–233