



## Correction to: In vitro activity of imipenem-relebactam against non-MBL carbapenemase-producing *Klebsiella pneumoniae* isolated in Greek hospitals in 2015–2016

Irene Galani<sup>1,2</sup>  · Maria Souli<sup>1</sup> · Konstantina Nafplioti<sup>1</sup> · Panagiota Adamou<sup>1</sup> · Ilias Karaiskos<sup>3</sup> · Helen Giamarellou<sup>3</sup> · Anastasia Antoniadou<sup>1</sup> · On Behalf Of The Study Collaborators

Published online: 29 April 2019

© Springer-Verlag GmbH Germany, part of Springer Nature 2019

**Publisher Correction: European Journal of Clinical Microbiology & Infectious Diseases**  
<https://doi.org/10.1007/s10096-019-03517-y>

The publisher regrets that the article has been published online on 01 March 2019 with errors in Table 1.

In the originally published Table 1, the percentage of Imipenem-relebactam susceptibility was incorrectly written as 8 0, while correct data is 98.0. Also, in Meropenem row, column MIC<sub>50</sub> (mg/L), the incorrect data 4 should be 64.

Subsequently, the revised Table 1 with the corrected data is shown below.

The original article has been corrected.

---

The online version of the original article can be found at <https://doi.org/10.1007/s10096-019-03517-y>

---

✉ Irene Galani  
egalani@med.uoa.gr

<sup>1</sup> 4th Department of Internal Medicine, Infectious Diseases Laboratory, Molecular Biology Section, School of Medicine, National and Kapodistrian University of Athens, Athens, Greece

<sup>2</sup> University General Hospital “ATTIKON”, Rimini 1, 124 62 Chaidari, Greece

<sup>3</sup> 1st Internal Medicine & Infectious Diseases Clinic, Hygeia General Hospital, Marousi, Greece

**Table 1** MIC and cumulative percent inhibited distributions for imipenem-relebactam and comparators, in relation to the carbapenemase type produced by the 314 *K. pneumoniae* isolates

Organism /Genotype	Agent	No of isolates / (cumulative % of isolates) inhibited at MIC (mg/L)										MIC <sub>50</sub> (mg/L)	MIC <sub>90</sub> (mg/L)	S (%)
		≤0.25	0.5	1	2	4	8	16	32	64	>64			
<b>KPC-producers</b> (n=295)	Imipenem					2 (0.7)	28 (10.2)	74 (35.3)	47 <b>(51.2)</b>	109 (88.1)	35 (100)	32	>64	0
	Imipenem-relebactam	<b>154</b> (52.2)	<b>91</b> (83.1)	<b>29</b> (92.9)	<b>15</b> (98.0)	6 (100)						0.25	1	98.0
	Meropenem				1 (0.3)		22 (7.8)	41 (21.7)	60 (42.0)	44 (56.9)	127 (100)	64	>64	0.3
	Doripenem				3 (1.0)	28 (10.5)	58 (30.2)	32 (41.0)	39 (54.2)	72 (78.6)	63 (100)	32	>64	0
	Colistin		<b>28</b> (9.5)	<b>133</b> (54.6)	<b>18</b> (60.7)	9 (63.7)	11 (67.5)	18 (73.6)	41 (87.5)	21 (94.6)	16 (100)	1	64	63.4
	Fosfomycin					1 (0.3)	<b>13</b> (4.7)	<b>74</b> (29.8)	<b>87</b> (59.3)	58 (79.0)	62 (100)	32	1024	59.3
	Tigecycline	<b>3</b> (1.0)	<b>37</b> (13.6)	<b>103</b> (48.5)	114 (87.1)	31 (97.6)	5 (99.3)	2 (100)				2	4	48.5
	Gentamicin		<b>6</b> (2.0)	<b>51</b> (19.3)	<b>132</b> (64.1)	41 (78.0)	5 (79.7)	13 (84.1)	4 (85.4)	5 (87.1)	38 (100)	2	>64	64.1
	Ceftazidime-avibactam	<b>20</b> (6.8)	<b>83</b> (34.9)	<b>130</b> (79.0)	<b>52</b> (96.6)	<b>9</b> (99.7)		1 (100)				1	2	99.7
<b>OXA-producers<sup>a</sup></b> (n=19)	Imipenem					6 (31.6)	10 (84.2)	2 (94.7)			1 (100)	8	16	0.0
	Imipenem-relebactam				2 (10.5)	13 (78.9)	2 (89.5)	1 (94.7)	1 (100)			4	16	10.5
	Meropenem						1 (5.3)	9 (52.6)	7 (89.5)	1 (94.7)	1 (100)	16	64	0.0
	Doripenem						9 (47.4)	9 (94.7)			1 (100)	16	16	0.0
	Colistin		<b>1</b> (5.3)	<b>7</b> (42.1)					8 (84.2)	2 (94.7)	1 (100)	32	64	42.1
	Fosfomycin					1 (5.3)	<b>9</b> (52.6)	<b>3</b> (68.4)	1 (73.7)	5 (100)		16	>64	68.4
	Tigecycline	<b>3</b> (15.7)	<b>9</b> (63.2)	4 (84.2)	1 (89.5)	1 (94.7)					1 (100)	1	8	63.2
	Gentamicin			2 (10.5)	2 (21.1)				1 (26.3)	14 (100)		>64	>64	21.1
	Ceftazidime-avibactam	<b>7</b> (36.8)	<b>10</b> (89.5)	<b>2</b> (100)								1	2	100.0

<sup>a</sup>One isolate harboring also *bla*<sub>KPC</sub> is included

Bold indicates susceptible by EUCAST 2018 breakpoint

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.