



Antiproliferative effect of low-level laser/ photobiomodulation on gingival fibroblasts derived from calcium channel blocker-induced gingival overgrowth

Melis özgül slezovic¹ · Işıl Saygun¹ · Vahdi Umut Bengi¹ · Muhittin Serdar² · Alpdogan Kantarci³

Received: 1 March 2024 / Accepted: 24 June 2024
© The Author(s) 2024

Abstract

The aim of this study was to evaluate the antiproliferative properties of low-level laser therapy (LLLT) on gingival fibroblasts obtained from calcium channel blocker-induced gingival overgrowth (GO). Gingival fibroblasts of patients with GO were compared to healthy gingival fibroblasts (H). Both cells were exposed to LLLT (685 nm wavelength, 25mW power, diode laser) and compared to those not treated with LLLT. Cell proliferation and viability were measured with MTT assay at baseline and after 24 and 72 h. TGF- β 1, CTGF, and collagen Type 1 levels were evaluated with Enzyme-Linked Immunosorbent Assay (ELISA). LLLT significantly decreased the proliferation of GO fibroblasts ($p < 0.05$) while leading to a significantly higher proliferation in H fibroblasts compared to the untreated cells ($p < 0.05$). GO cells showed significantly higher CTGF, TGF- β , and collagen Type 1 expression than the H cells ($p < 0.05$). LLLT significantly reduced CTGF levels in GO cells compared to the control group ($p < 0.05$). In H cells, CTGF and TGF- β levels were also significantly decreased in response to LLLT compared to the control group ($p < 0.05$). While LLLT significantly reduced collagen expression in the H group ($p < 0.05$), it did not significantly impact the GO cells. LLLT significantly reduced the synthesis of the growth factors and collagen in both groups with an antiproliferative effect on the gingival fibroblasts from calcium channel blocker-induced GO, suggesting that it can offer a therapeutic approach in the clinical management of drug-induced GO, reversing the fibrotic changes.

Keywords Gingival enlargement · Calcium channel blockers · Low level laser therapy · Photobiomodulation · CTGF · TGF- β · Collagen

Introduction

Gingival overgrowth (GO) develops due to inflammation, drug side effects, and genetic and neoplastic processes [1]. Anticonvulsant, immunosuppressive, and calcium channel blockers (CCB) lead to drug-induced GO [2]. Calcium channel blocker-induced GO is characterized by cell

proliferation, extracellular matrix (ECM) deposition in the gingival connective tissue with varying degrees of inflammation and fibrosis and disruption of collagen metabolism [3] regulated by Transforming Growth Factor- β (TGF- β) [4]. TGF- β controls collagen levels through cellular communication network (CCN) proteins, a family of ECM-related proteins involved in intercellular signaling. Connective tissue growth factor (CTGF), also known as CCN2, is a member of the CCN family; TGF- β 1 rapidly and strongly induces CTGF and contributes to regulating the ECM in GO [5]. CTGF is an important factor in collagen production by regulating the proliferation and differentiation of fibroblasts and stimulating tumor growth and metastasis [6].

Drug-induced GO has a high prevalence, and the fact that many factors such as inflammation, drug dose, agent of the drug used, host response, and genetic predisposition play a role in the pathogenesis makes treatment difficult. Since it

✉ Melis özgül slezovic
melis.ozgulslezovic@sbu.edu.tr

¹ Department of Periodontology, University of Health Sciences, Gulhane Faculty of Dental Medicine, Ankara, Turkey

² Department of Medical Biochemistry, Acıbadem Mehmet Ali Aydınlar University, Ankara, Turkey

³ Forsyth Institute, Cambridge, Massachusetts, USA

is not always possible to discontinue or change the drugs that cause GO, developing treatments to prevent recurrence is critical. The necessity for the development of novel, non-invasive therapeutic approaches becomes apparent.

Low-level laser therapy (LLLT) has been used in many fields of medicine as a non-invasive and non-thermal method for its anti-fibrotic and photobiomodulation effects [7]. While initially expressions such as ‘photobioactivation’ and ‘biostimulation’ were frequently used to describe the stimulation effect of low-level lasers, the term ‘photobiomodulation’ began to be used after the inhibitory effect was observed [8]. The effects of low-level laser therapy (LLLT), also known as photobiomodulation therapy (PBM), in alleviating the negative consequences of inflammation and fibrosis by inducing repair in tissues, have been demonstrated in numerous studies [9, 10]. LLLT or PBM is emerging as a promising new treatment option for fibrosis in different organs. However, the anti-fibrotic potential of this treatment in dentistry needs to be elucidated and the cellular and molecular interactions of the laser need to be clarified. Various studies in dentistry have shown that low-level laser therapy (LLLT) and photobiomodulation (PBM) can accelerate cell proliferation, improve wound healing, and be suitable for treating different oral diseases [11, 12]. However, it is unknown whether LLLT affects GO gingival fibroblasts differently than cells from healthy (non-GO) individuals. Therefore, we tested the hypothesis that LLLT would reduce cell proliferation of GO-derived fibroblasts and affect the levels of TGF- β , CTGF, and collagen, which are involved in fibrotic processes.

Materials and methods

This study was ethically approved by the Health Sciences University Scientific Research Ethics Committee with decision number 2021–371 and supported by the Health Sciences University Scientific Research Projects Unit with project number 2022/003.

Primary cell culture from GO individuals

After obtaining informed consent from a patient with calcium channel blocker-induced GO in need of surgical excision, the excised gingival tissues were transferred to the laboratory by placing the excised gingival tissues in a 5 mm Petri dish containing 14 mm Dulbecco Modified Eagle Medium (DMEM) with 1100 U/ml penicillin and 100 μ g/ml streptomycin. The gingival tissues were cut into small pieces and transferred into a 25-flask with a sterile syringe tip. 10% medium (10% fetal calf serum- 1% penicillin – 1% L-glutamine) was added and placed in an oven containing

5% CO₂ at 37 °C for 72 h. The medium was refreshed every 72 h until all 80% confluency. After trypsinization, the cells were transferred to a large 75 flask. Cells between passages 4 and 7 were used at 3×10^4 cells/well in 24-well plates. Human gingival fibroblasts (HGF-1- ATCC CRL-2014; American Type Culture Collection, Manassas, VA) were used as a healthy control (H). Healthy gingival fibroblasts were incubated in DMEM (Sigma D6429 containing 4500 mg/l glucose, sodium pyruvate, and sodium bicarbonate) medium supplemented with 10% heat-inactivated fetal calf serum FBS (Biochrom S0115) and 100 U/ml penicillin, 100 μ g/ml streptomycin (Biochrom A2213) and incubated at 37 °C in a 95% humidified oven with 5% CO₂. These cells were seeded 3×10^4 cells/well in 24-well plates.

Low-level laser treatment

Each group was divided into LLLT and non-treated control groups. Irradiation was performed in continuous mode with a diode laser (BTL-2000 Benešov Czech Republic) at a wavelength of 685 nm, a power output of 25 mW and with a power density of 2.0 J/cm^2 for 100 s. The laser beam was transmitted by an optical fiber and irradiated to a circular area of 1 cm^2 . Laser irradiation was applied in two doses. The second dose was repeated 3 days after the first dose. Laser application was not applied to the control groups (Fig. 1).


Cell viability and proliferation

MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) (Sigma, St Louis, MO, USA) was used to evaluate the cytotoxicity and cell proliferation of the experimental groups at baseline and after 24 h. Briefly, fibroblasts were seeded in 96-well plates. The media were replaced with fresh media, and MTT (5 mg/mL) solution was added into the wells in one-tenth of the original culture volume, and the incubation was continued at 37 °C in 5% CO₂ for 4 h. Then, the incubation medium was removed, and 100 μ L 0.04 N isopropanol was added to each well. The plate was incubated at room temperature for approximately 30 min. The absorbance was measured at 570 nm in a microplate reader (Biotek Epoch, Germany).

Enzyme-linked immunosorbent assays (ELISA)

Supernatants were frozen at -80 °C until the analysis. TGF- β 1, CTGF, and collagen type I (Col-I) levels were measured using ELISA kits (Fine Test Human TGF- β 1 ELISA Kit, sensitivity level 18.75 pg/ml, Wuhan, China; Fine Test Human CTGF ELISA Kit, sensitivity level 37.5 pg/ml,

Fig. 1 **a.** Initial GO cells (x 4 magnification) **b.** Untreated control GO cells (x10 magnification) **c.** GO cells after LLLT (x10 magnification) **d.** Initial H cells (x10 magnification) **e.** Untreated control H cells (x 10 magnification) **f.** H cells after LLLT (x10 magnification)



Wuhan, China; Fine Test Human Collagen Type I (Col I) ELISA Kit, sensitivity level 0.821 µg/ml, Wuhan, China).

Statistical Analysis Statistical analyses were performed using SPSS (SPSS Inc. Chicago, IL, USA). Shapiro-Wilks test was used to evaluate whether the data were normally distributed. The significance of the difference between the means of the groups in MTT was assessed by the Kruskal-Wallis test. Bonferroni test was used for pairwise post-hoc comparisons. Analysis of Variance (ANOVA) was used to determine whether the growth factor release levels of the study groups differed from each other. Tukey-Kramer test was used to determine which groups differed. Differences below $p < 0.05$ were considered significant.

Results

Effect of LLLT on the proliferation of gingival fibroblasts

The proliferation of GO laser-treated cells was statistically lower than that of control GO cells after the first dose. In contrast, no significant difference was detected between the GO cells after the second dose (Fig. 2a). While a significant difference was seen in the 24th and 72nd hours GO cells in response to LLLT compared to baseline, there was no significant difference between these time points (Figs. 1 and 2d). While there was no significant difference between the baseline and 24 h in H cells in response to LLLT, there was an increase in H cell proliferation after 72 h (Fig. 2b, d). While there was no statistical difference between the baseline viability and proliferation of GO and H cells, LLLT-treated GO cells showed significantly lower proliferation after 24 and 72 h compared to the H group (Fig. 2c).

Effects of LLLT on TGF-β, CTGF and collagen expressions

Untreated control GO cells showed significantly higher CTGF levels than the control H group. LLLT significantly reduced CTGF levels in both GO and H cells (Fig. 3a). Untreated control GO cells showed significantly higher TGF-β1 levels than control H cells. The reduction of TGF-β1 levels in GO cells in response to LLLT was not statistically significant. LLLT resulted in a significant decrease in TGF-β1 levels in H cells (Fig. 3b). Untreated control GO cells showed significantly higher collagen levels than control H cells. LLLT was found to reduce collagen levels in both GO and H cells. However, this effect was only statistically significant in H cells (Fig. 3c). A high positive correlation was found between TGF-β1 and CTGF ($r = 0.798$; $p < 0.0001$). A low positive correlation was found between CTGF and collagen ($r = 0.335$ $p < 0.05$).

Discussion

LLLT suppressed proliferation in cells obtained from CCB-induced GO without showing a cytotoxic effect, while it stimulated proliferation in healthy cells. A similar proliferative effect of LLLT in healthy cells has also been shown in previous studies [13, 14]. In a previous in vitro study, we divided human gingival fibroblast cells into 3 groups. In the first group, the cells were irradiated with a single dose of diode laser, the second group was irradiated with the same dose of laser for two consecutive days, and the third group, which was not irradiated with diode laser, served as the control group. Both irradiated groups showed higher proliferation and viability than the control group [15]. In another in vitro study on healthy gingival fibroblasts obtained from a cell bank, gingival fibroblasts were divided into




Fig. 2 a. Comparison of MTT levels between GO cells b. Comparison of MTT levels between H cells c. Comparison of MTT levels between laser-treated GO and H cells. d. Intra-group MTT comparison by time

points. MTT 0: Baseline MTT, MTT 24: 24th-hour MTT, MTT 72: 72nd-hour MTT. MTT 0: Baseline MTT, MTT 24: 24th hour MTT, MTT 72: 72nd hour MTT *: $p < 0,05$ **: $p < 0,01$ ***: $p < 0,001$

experimental and control groups. The experimental group received LLLT with a diode laser for 3 consecutive days, while the control group received no treatment. A significantly higher cell proliferation was detected in the experimental group at 48 and 72 h [16]. In contrast, some studies did not report cell proliferation [17], possibly due to differences in the characteristics of the laser device, such as dose wavelength, irradiation mode, and the characteristics of the cell type. It is crucial to employ adequate dosimetry in LLLT. A fundamental principle, termed the “biphasic dose response,” has emerged, indicating that larger doses of light tend to be less efficacious than smaller doses [18]. However, no consensus has been reached on the optimal range. In studies, a diode laser with a wavelength ranging from 660 to 940 nm was frequently used in continuous mode, with a large range in output power ranging from 10 to 500 mW, while the energy density range ranged from 1 to 15.8 J/cm² [19]. In our study, LLLT were performed in 2 doses, similar to the parameters of our previous studies in which the photobiomodulation effect of LLLT was demonstrated [15, 20].

LLLT has been reported to be beneficial in reducing fibrosis in various organs [7, 21]. Still, there is no study on the anti-fibrotic effect of LLLT in drug-induced gingival

growths. In our study, while proliferation was suppressed with the first dose of LLLT in cells obtained from CCB-induced gingival growth, no significant difference in cell proliferation value was observed after the second dose of LLLT. However, a decrease in the proliferation level of the control group without LLLT was observed at 72 h. In primary cell cultures, cells have a limited lifespan and proliferation capacity decreases over time [22]. The increase in the number of these cells, which are thought to have high proliferation potential, in the wells may have caused the environmental conditions to be insufficient for cell proliferation and maintenance of cell viability. In contrast, cells of the same character treated with LLLT simultaneously showed a stable level of viability, a remarkable result of the biomodulation property of the low-level laser.

In the pathogenesis of gingival overgrowth, extracellular matrix (ECM) accumulation and fibrotic changes occur by releasing excessive amounts of mediators that contribute to regulating fibrogenic and regenerative signals with increased fibroblasts [23]. In our study, TGF- β levels were significantly higher in cells obtained from CCB-induced growths than in healthy cells, consistent with previous studies [24, 25]. A decrease in TGF- β levels was observed in




Fig. 3 a. Comparison of CTGF levels between groups b. Comparison of TGF- β levels between groups. c. Comparison of collagen levels between groups. *: $p < 0,05$ **: $p < 0,01$ ***: $p < 0,001$

these hyperplastic cells and healthy fibroblasts treated with LLLT compared to non-laser-treated cells. Studies in which the anti-fibrotic effect of LLLT was demonstrated were generally performed in the medical field [26, 27]. No studies in the literature evaluated the anti-fibrotic properties of LLLT on gingival growths and its effect on the expression of growth factors thought to play a role in fibrosis. TGF- β levels of healthy gingival fibroblasts treated with LLLT were shown to be significantly lower than those of untreated cells

[11]. The researchers stated that TGF- β is a versatile growth factor and may play a role in undesirable wound healing models such as fibrosis and scar tissue formation. In the study by Kuo et al. on keloid tissues obtained from patients, the effect of LLLT on gene expression was evaluated, and it was reported that LLLT mediated the blockade of TGF- β expression [28]. It is thought that TGF- β can slow down the degradation of connective tissue matrix by decreasing

the synthesis of matrix metalloproteinases and plasminogen activators [29].

Recent studies have shown that TGF- β controls CTGF expression may have effects on subsequent steps of tissue fibrosis [30]. CTGF is involved in several important biological functions, including cell proliferation, differentiation, adhesion, and angiogenesis, and in controlling multiple pathological processes such as tumor development and tissue fibrosis [31]. The literature shows increased CTGF levels are generally demonstrated in drug-induced growths and patients with hereditary gingival fibromatosis [32, 33]. Our study detected significantly high CTGF levels in cells obtained from CCB-induced growths. These cells showed very high CTGF release even when their proliferation decreased, CTGF is a reliable biomarker in drug-induced gingival overgrowth. In our study, the increased CTGF release detected in the cells obtained from CCB-induced gingival overgrowth was suppressed by LLLT, and the CTGF level in LLLT-treated cells was found to be significantly lower in healthy gingival fibroblasts compared to the control group.

Suppression of excessive synthesis of CTGF in healthy cells, which is a biomarker for drug-induced gingival overgrowth, may provide a prophylactic benefit in terms of the occurrence of gingival hyperplasia. Similar to the results of our study, Yeh et al. showed in vitro that LLLT inhibited the transcriptional activity of CTGF in cells obtained from patients with oral submucous fibrosis [34]. In another study supporting these results, Zhu et al. reported that CTGF mRNA expression decreased with laser irradiation in gingival fibroblasts obtained from keloid tissues [35].

Our study detected increased TGF- β , CTGF and collagen levels in cells obtained from CCB-induced growths. Similarly, Fuji et al. showed increased collagen levels in their study with gingival fibroblasts obtained from patients using nifedipine [36]. Lu et al. reported that fibroblast cultures formed from gingival tissues obtained from patients with nifedipine-induced gingival overgrowth exhibited increased expression of type I collagen compared to fibroblasts obtained from healthy gingiva [37]. Similarly, there are other studies in the literature showing that collagen, TGF- β 1 and CTGF levels increase in drug-induced gingival enlargement and hereditary gingival fibromatosis [38, 39]. Pisoschi et al. reported in their study that the TGF- β -CTGF relationship was activated in gingival growth caused by calcium channel blockers [40]. Another study reported that TGF- β acts as a strong stimulatory signal for connective tissue formation in fibrotic conditions and acts together with CTGF in its growth-stimulating effect [41]. Some authors have reported that the relationship between CTGF and TGF- β may be a mediator that strengthens the fibrogenic function of fibroblasts [42, 43]. In our study, a positive

correlation was observed between TGF- β and CTGF and between collagen and CTGF, similar to previous studies.

Conclusion

This is the first study to demonstrate that LLLT has a modulating effect on cell proliferation in gingival fibroblasts from tissues with gingival hyperplasia due to the use of calcium channel blockers without causing cytotoxicity and can suppress the synthesis of CTGF, TGF- β , and collagen—all of which have been linked to the pathogenesis of gingival hyperplasia. Low-dose laser applications suppress the proliferation of fibroblasts isolated from hyperplastic tissue while stimulating the proliferation of healthy cells. This indicates that LLLT can be used in growth therapies without causing any pathological effects in the mouth.

In conclusion, for LLLT to be included in clinical practice as a new treatment model in treating hyperplasia in dentistry, in vitro studies evaluating biological markers for pathogenesis and randomized controlled clinical trials evaluating long-term recurrence are needed.

Acknowledgements We would like to thank the Health Sciences University Scientific Research Projects Unit for the support of the study. We would also like to thank Pınar Eçi for her contributions to the laboratory stages.

Author contributions Conceptualization: Işıl Saygun. Methodology: Işıl Saygun. Formal analysis: Muhittin Serdar, Vahdi umut Bengi. Investigation: Melis Özgül Slezovic, Işıl Saygun. Writing - original draft preparation: Melis Özgül Slezovic. Writing - review and editing: Işıl Saygun, Alpdoğan Kantarci. Visualization: Melis Özgül Slezovic, Vahdi umut Bengi. Supervision: Alpdoğan Kantarci.

Funding Open access funding provided by the Scientific and Technological Research Council of Türkiye (TÜBİTAK). This study was funded by Health Sciences University Scientific Research Projects Unit.

Open access funding provided by the Scientific and Technological Research Council of Türkiye (TÜBİTAK).

Declarations

Ethical approval Approval was obtained from the ethics committee of Health Sciences University. The procedures used in this study adhere to the tenets of the Declaration of Helsinki.

Consent to participate Informed consent was obtained from all individual participants included in the study.

Financial interests The authors declare they have no financial interests.

Competing interests The authors have no competing interests to declare that are relevant to the content of this article.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Brunet L, Miranda J, Farré M, Berini L, Mendieta C (1996) Gingival Enlargement Induced by drugs. *Drug-Safety* 15(3):219–231. <https://doi.org/10.2165/00002018-199615030-00007>
2. Mavrogiannis M, Ellis JS, Thomason JM, Seymour RA (2006) The management of drug-induced gingival overgrowth. *J Clin Periodontol* 33(6):434–439. <https://doi.org/10.1111/j.1600-051X.2006.00930.x>
3. Seymour RA, Ellis JS, Thomason JM (2000) Risk factors for drug-induced gingival overgrowth. *J Clin Periodontol* 27(4):217–223. <https://doi.org/10.1034/j.1600-051x.2000.027004217.x>
4. Prime SS, Pring M, Davies M, Paterson IC (2004) TGF- β Signal Transduction in oro-facial health and non-malignant disease (part I). *Crit Reviews Oral Biology Med* 15(6):324–336. <https://doi.org/10.1177/154411130401500602>
5. Blom IE, Goldschmeding R, Leask A (2002) Gene regulation of connective tissue growth factor: new targets for antifibrotic therapy? *Matrix Biol* 21(6):473–482. [https://doi.org/10.1016/S0945-053X\(02\)00055-0](https://doi.org/10.1016/S0945-053X(02)00055-0)
6. Zhu X, Zhong J, Zhao Z, Sheng J, Wang J, Liu J et al (2015) Epithelial derived CTGF promotes breast tumor progression via inducing EMT and collagen I fibers deposition. *Oncotarget* 6(28):25320–25338. <https://doi.org/10.18632/oncotarget.4659>
7. Mamalis A, Siegel D, Jagdeo J (2016) Visible Red Light Emitting Diode Photobiomodulation for skin fibrosis: Key Molecular pathways. *Curr Derm Rep* 5(2):121–128. <https://doi.org/10.1007/s13671-016-0141-x>
8. Schindl A, Schindl M, Pernerstorfer-Schön H, Schindl L (2000) Low-intensity laser therapy: a review. *J Investig Med* 48(5):312–326
9. Feliciano RDS, Manchini MT, Atum ALB, da Silva GA, Antônio EL, Serra AJ, Tucci PJF, Andrade de Mello R, Chavantes MC, Baltatu OC, Silva Júnior JA (2022) Photobiomodulation therapy's effects on cardiac fibrosis activation after experimental myocardial infarction. *Lasers Surg Med* 54(6):883–894. <https://doi.org/10.1002/lsm.23544>
10. Chiang MH, Lee KT, Chen CH, Chen KK, Wang YH (2020) Photobiomodulation therapy inhibits oral submucous fibrosis in mice. *Oral Dis* 26(7):1474–1482. <https://doi.org/10.1111/odi.13409>
11. Karoussis IK, Kyriakidou K, Psarros C, Afouxenides P, Vrotsos IA (2021) Dosage effects of an 810 nm Diode Laser on the proliferation and growth factor expression of human gingival fibroblasts. *J Lasers Med Sci* 12:e25. <https://doi.org/10.34172/jlms.2021.25>
12. Vale FA, Moreira MS, Almeida FCS, Ramalho KM (2015) Low-level laser therapy in the treatment of recurrent aphthous ulcers: a systematic review. *Sci World J* 7. <https://doi.org/10.1155/2015/150412>
13. Illescas-Montes R, Melguizo-Rodríguez L, Manzano-Moreno FJ, García-Martínez O, Ruiz C, Ramos-Torrecillas J (2017) Cultured human fibroblast Biostimulation using a 940 nm Diode Laser. *Materials* 10(7):793. <https://doi.org/10.3390/ma10070793>
14. Vinck EM, Cagnie BJ, Cornelissen MJ, Declercq HA, Cambier DC (2003) Increased fibroblast proliferation induced by light emitting diode and low power laser irradiation. *Lasers Med Sci* 18(2):95–99. <https://doi.org/10.1007/s10103-003-0262-x>
15. Saygun I, Karacay S, Serdar M, Ural AU, Sencimen M, Kurtis B (2008) Effects of laser irradiation on the release of basic fibroblast growth factor (bFGF), insulin like growth factor-1 (IGF-1), and receptor of IGF-1 (IGFBP3) from gingival fibroblasts. *Lasers Med Sci* 23(2):211–215. <https://doi.org/10.1007/s10103-007-0477-3>
16. Frozanfar A, Ramezani M, Rahpeyma A, Khajehahmadi S, Arbab HR (2013) The effects of low-level laser therapy on the expression of collagen type I gene and proliferation of human gingival fibroblasts (Hgf3-Pi 53): in vitro study. *Iran J Basic Med Sci* 16(10):1071–1074
17. Gholami L, Parsamanesh G, Shahabi S, Jazaeri M, Baghaei K, Fekrazad R (2021) The Effect of Laser Photobiomodulation on Periodontal Ligament Stem cells. *Photochem Photobiol* 97(4):851–859. <https://doi.org/10.1111/php.13367>
18. Huang YY, Sharma SK, Carroll J, Hamblin MR (2011) Biphasic dose response in low level light therapy - an update. *Dose Response* 9(4):602–618. <https://doi.org/10.2203/dose-response.11-009.Hamblin>
19. Ren C, McGrath C, Jin L et al (2016) Effect of diode low-level lasers on fibroblasts derived from human periodontal tissue: a systematic review of in vitro studies. *Lasers Med Sci* 31:1493–1510. <https://doi.org/10.1007/s10103-016-2026-4>
20. Saygun I, Nizam N, Ural AU, Serdar MA, Avcu F, Tözüm TF (2012) Low-level laser irradiation affects the release of basic fibroblast growth factor (bFGF), insulin-like growth factor-I (IGF-I), and receptor of IGF-I (IGFBP3) from osteoblasts. *Photomed Laser Surg* 30(3):149–154. <https://doi.org/10.1089/pho.2011.3079>
21. Alves AN, Fernandes KPS, Melo CaV, Yamaguchi RY, França CM, Teixeira DF et al (2014) Modulating effect of low level-laser therapy on fibrosis in the repair process of the tibialis anterior muscle in rats. *Lasers Med Sci* 29(2):813–821. <https://doi.org/10.1007/s10103-013-1428-9>
22. Butler M, *Animal Cell Culture and Technology* (2003) 2nd edition, London. <https://doi.org/10.4324/9780203427835>
23. Hallmon WW, Rossmann JA (1999) The role of drugs in the pathogenesis of gingival overgrowth. A collective review of current concepts. *Periodontol* 2000 21(1):176–196
24. Nițulescu EA, Crăitoiu MM, Baniță MI, Drăghici E, Crăitoiu Ș (2012) The involvement of TGF- β 1 and CTGF in regional gingival overgrowth. *Rom J Morphol Embryol* 53(1):143–150
25. Vahabi S, Salman BN, Rezazadeh F, Namdari M (2014) Effects of cyclosporine and phenytoin on biomarker expressions in gingival fibroblasts of children and adults: an in vitro study. *J Basic Clin Physiol Pharmacol* 25(2):167–173. <https://doi.org/10.1515/jbcpp-2013-0003>
26. Ahrabi B, Bahrami M, Moghadasali R, Zamanian-Azodi M, Khoramgah MS, Tabatabaei Mirakabad FS et al (2020) The effect of low-power laser therapy on the TGF/ β signaling pathway in chronic kidney disease: a review. *J Lasers Med Sci* 11(2):220–225. <https://doi.org/10.34172/jlms.2020.36>
27. Brochetti RA, Leal MP, Rodrigues R, da Palma RK, de Oliveira LVF, Horliana ACRT et al (2017) Photobiomodulation therapy improves both inflammatory and fibrotic parameters in experimental model of lung fibrosis in mice. *Lasers Med Sci* 32(8):1825–1834. <https://doi.org/10.1007/s10103-017-2281-z>
28. Kuo YR, Wu WS, Wang FS (2007) Flashlamp pulsed-dye laser suppressed TGF-beta1 expression and proliferation in cultured

- keloid fibroblasts is mediated by MAPK pathway. *Lasers Surg Med* 39(4):358–364. <https://doi.org/10.1002/lsm.20489>
29. Javelaud D, Mauviel A (2004) Mammalian transforming growth factor- β s: smad signaling and physio-pathological roles. *Int J Biochem Cell Biol* 36(7):1161–1165. [https://doi.org/10.1016/S1357-2725\(03\)00255-3](https://doi.org/10.1016/S1357-2725(03)00255-3)
 30. Holmes A, Abraham DJ, Sa S, Shiwen X, Black CM, Leask A (2001) CTGF and SMADs, maintenance of scleroderma phenotype is independent of SMAD signaling. *J Biol Chem* 276(14):10594–10601. <https://doi.org/10.1074/jbc.M010149200>
 31. Barreto SC, Ray A, Ag Edgar P (2016) Biological characteristics of CCN proteins in tumor development. *J BUON* 21(6):1359–1367
 32. Anand AJ, Gopalakrishnan S, Karthikeyan R, Mishra D, Mohapatra S (2018) Immunohistochemical analysis of the role connective tissue growth factor in drug-induced Gingival Overgrowth in response to Phenytoin, Cyclosporine, and Nifedipine. *J Int Soc Prev Community Dent* 8(1):12–20. https://doi.org/10.4103/jispcd.JISPCD_403_17
 33. Kantarci A, Black S, Xydias C, Murawel P, Uchida Y, Yucekal-Tuncer B et al (2006) Epithelial and connective tissue cell CTGF/CCN2 expression in gingival fibrosis. *J Pathol* 210(1):59–66. <https://doi.org/10.1002/path.2000>
 34. Yeh MC, Chen KK, Chiang MH, Chen CH, Chen PH, Lee HE et al (2017) Low-power laser irradiation inhibits arecoline-induced fibrosis: an in vitro study. *Int J Oral Sci* 9(1):38–42. <https://doi.org/10.1038/ijos.2016.49>
 35. Zhu R, Yue B, Yang Q, Ma Y, Huang G, Guan M et al (2015) The effect of 595 nm pulsed dye laser on connective tissue growth factor (CTGF) expression in cultured keloid fibroblasts. *Lasers Surg Med* 47(2):203–209. <https://doi.org/10.1002/lsm.22334>
 36. Fujii A, Matsumoto H, Nakao S, Teshigawara H, Akimoto Y (1994) Effect of calcium-channel blockers on cell proliferation, DNA synthesis and collagen synthesis of cultured gingival fibroblasts derived from human nifedipine responders and non-responders. *Arch Oral Biol* 39(2):99–104. [https://doi.org/10.1016/0003-9969\(94\)90103-1](https://doi.org/10.1016/0003-9969(94)90103-1)
 37. Lu HK, Tseng CC, Lee YH, Li CL, Wang LF (2010) Flutamide inhibits nifedipine- and interleukin-1 β -induced collagen overproduction in gingival fibroblasts. *J Periodontol Res* 45(4):451–457. <https://doi.org/10.1111/j.1600-0765.2009.01255.x>
 38. Chung Y, Fu E, Chin YT, Tu HP, Chiu HC, Shen EC et al (2015) Role of shh and TGF in cyclosporine-enhanced expression of collagen and α -SMA by gingival fibroblast. *J Clin Periodontol* 42(1):29–36. <https://doi.org/10.1111/jcpe.12333>
 39. Yamazaki K, Igarashi-Takeuchi H, Numabe Y (2022) Hepatocyte growth factor exhibits anti-fibrotic effects in an in vitro model of nifedipine-induced gingival overgrowth. *J Oral Sci* 64(1):99–104. <https://doi.org/10.2334/josnusd.21-0419>
 40. Pisoschi CG, Stănculescu CE, Andrei AM, Berbecaru-Iovan A, Munteanu C, Popescu F et al (2014) Role of transforming growth factor β -connective tissue growth factor pathway in dihydropyridine calcium channel blockers-induced gingival overgrowth. *Rom J Morphol Embryol* 55(2):285–290
 41. Kothapalli D, Frazier KS, Welply A, Segarini PR, Grotendorst GR (1997) Transforming growth factor beta induces anchorage-independent growth of NRK fibroblasts via a connective tissue growth factor-dependent signaling pathway. *Cell Growth Differ* 8(1):61–68
 42. Trackman PC, Kantarci A (2004) Connective tissue metabolism and Gingival Overgrowth. *Crit Reviews Oral Biology Med* 15(3):165–175. <https://doi.org/10.1177/154411130401500305>
 43. Heng ECK, Huang Y, Black SA Jr., Trackman PC (2006) CCN2, connective tissue growth factor, stimulates collagen deposition by gingival fibroblasts via module 3 and α 6- and β 1 integrins. *J Cell Biochem* 98(2):409–420. <https://doi.org/10.1002/jcb.20810>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.