

An alternating linearization bundle method for convex optimization and nonlinear multicommodity flow problems

K. C. Kiwiel

Received: 8 August 2008 / Accepted: 24 September 2009 / Published online: 10 November 2009
© Springer and Mathematical Programming Society 2009

Abstract We give a bundle method for minimizing the sum of two convex functions, one of them being known only via an oracle of arbitrary accuracy. Each iteration involves solving two subproblems in which the functions are alternately represented by their linearizations. Our approach is motivated by applications to nonlinear multicommodity flow problems. Encouraging numerical experience on large scale problems is reported.

Keywords Nondifferentiable optimization · Convex programming · Proximal bundle methods · Approximate subgradients · Network flow problem

Mathematics Subject Classification (2000) 65K05 · 90C25 · 90C27

1 Introduction

We give a bundle method for the structured convex minimization problem

$$\theta_* := \inf\{\theta(\cdot) := \sigma(\cdot) + \pi(\cdot)\}, \quad (1.1)$$

where $\sigma : \mathbb{R}^m \rightarrow (-\infty, \infty]$ and $\pi : C \rightarrow \mathbb{R}$ are closed proper convex functions, and $C := \text{dom } \sigma := \{u : \sigma(u) < \infty\}$ is the effective domain of σ . Such problems may appear via duality when the primal has a certain structure. For instance, consider the

Research supported by the Polish Ministry of Science and Higher Education Grant No. N N514 408736.

K. C. Kiwiel (✉)
Systems Research Institute, Polish Academy of Sciences,
Newelska 6, 01-447 Warsaw, Poland
e-mail: kiwiel@ibspan.waw.pl

two equivalent minimization problems

$$f_* := \inf\{f(Ax) : x \in X\} = \inf\{f(y) : y = Ax, x \in X\}, \quad (1.2)$$

where $X \subset \mathbb{R}^n$ and A is an $m \times n$ matrix. For the Lagrangian $L(x, y; u) := f(y) + \langle u, Ax - y \rangle$, minimization over $(x, y) \in X \times \mathbb{R}^m$ yields (1.1) as a dual problem with

$$\sigma(u) := f^*(u) := \sup_y \{\langle u, y \rangle - f(y)\} \quad \text{and} \quad \pi(u) := \sup\{\langle -A^T u, x \rangle : x \in X\}. \quad (1.3)$$

We assume that σ is “simple” in the sense that minimizing σ plus a separable convex quadratic function is “easy”. On the other hand, π is known only via an oracle, which at any query point $u \in C$ delivers an affine minorant of π (e.g., $\langle -Ax, \cdot \rangle$) for a possibly inexact maximizer x in (1.3).

Our method is an approximate version of the proximal point algorithm [18,21] which generates a sequence

$$\hat{u}^{k+1} = \arg \min \sigma(\cdot) + \pi(\cdot) + \frac{1}{2t_k} |\cdot - \hat{u}^k|^2 \quad \text{for } k = 1, 2, \dots, \quad (1.4)$$

starting from a point $\hat{u}^1 \in C$, where $|\cdot|$ is the Euclidean norm and $t_k > 0$ are step-sizes. It combines two basic ideas: *bundling* from the proximal bundle methods [7,9, Sect. XV.3], and their extensions [12,13] to inexact oracles, and *alternating linearization* (AL for short) from [11,13,16]. Here bundling means replacing π in (1.4) by its polyhedral model $\check{\pi}_k \leq \pi$ derived from the past oracle answers. Since the resulting subproblem may still be too difficult, we follow the AL approach in which a subproblem involving the sum of two functions (here σ and $\check{\pi}_k$) is replaced by two subproblems in which the functions are alternately represented by linear models. Thus, (1.4) is replaced by the two easier subproblems

$$\check{u}^{k+1} := \arg \min \bar{\sigma}_{k-1}(\cdot) + \check{\pi}_k(\cdot) + \frac{1}{2t_k} |\cdot - \hat{u}^k|^2, \quad (1.5)$$

$$u^{k+1} := \arg \min \sigma(\cdot) + \bar{\pi}_k(\cdot) + \frac{1}{2t_k} |\cdot - \hat{u}^k|^2. \quad (1.6)$$

The first subproblem (1.5) employs a linearization $\bar{\sigma}_{k-1} \leq \sigma$ obtained at the previous iteration. Its solution yields by the usual optimality condition a linearization $\bar{\pi}_k \leq \check{\pi}_k$ which may a posteriori replace $\check{\pi}_k$ in (1.5) without changing its optimal value and solution. Similarly, the solution of (1.6) provides a linearization $\bar{\sigma}_k \leq \sigma$ which may a posteriori replace σ in (1.6).

Our method coincides with that of [13] in the special case of σ being the indicator function i_C of C ($i_C(u) = 0$ if $u \in C$, ∞ otherwise). Then u^{k+1} in (1.6) is the projection onto C of $\hat{u}^k - t_k \nabla \bar{\pi}_k$; this projection is straightforward if the set C is “simple”. For more difficult cases, it is crucial to allow for approximate solutions in (1.6). We show (cf. Sect. 4.2) that such solutions can be obtained by solving the Fenchel dual of (1.6) approximately; this is conceptually related to the use of Fenchel’s duality in [7, Proposition XV.2.4.3 and p. 306].

For dual applications, we restrict our attention to the setup of (1.2)–(1.3) with f closed proper convex and X compact and convex (since other examples of [16] could be treated in similar ways). As in [13], even when the dual has no solutions, our method can still asymptotically find ε_π -optimal primal solutions, where ε_π is an upper bound on the oracle’s errors; in fact only the asymptotic oracle errors matter, as discussed in [13, Sect. 4.2].

Actually, our theoretical contributions outlined above were motivated by applications to nonlinear multicommodity flow problems (NMFP for short); more concretely, by the good experimental results of [1], where the analytic center cutting plane method (ACCPM for short) exploited “nice” second-order properties of σ . This gave tremendous improvements over an earlier version of ACCPM [6] which used a first-order oracle for σ . We show that our method can exploit such properties as well, obtaining significant speedups with respect to standard bundle on most instances used in [1]. The alternative approach of [17] for adapting standard bundle to NMFP is promising, but has not been tested on large instances (see Sect. 8.3 for rough comparisons with our AL). Finally, we note that the ballstep subgradient method of [14] is quite efficient only for fairly low accuracy requirements.

As for the state-of-the-art in NMFP, we refer the reader to [1] for the developments subsequent to the review of [19].

The paper is organized as follows. In Sect. 2 we present our method. Its convergence is analyzed in Sect. 3. Useful modifications, including approximate solutions of (1.6), are given in Sect. 4. Application to the Lagrangian relaxation of (1.2) is studied in Sect. 5. Specializations to NMFP are given in Sect. 6. Implementation issues are discussed in Sect. 7. Numerical benchmarks on the instances of [1] and comparisons with standard bundle and the method of [17] are given in Sect. 8.

2 The alternating linearization bundle method

We first explain our use of approximate objective values in (1.5), (1.6). Our method generates a sequence of *trial points* $\{u^k\}_{k=1}^\infty \subset C$ at which the oracle is called. We assume that for a fixed *accuracy tolerance* $\varepsilon_\pi \geq 0$, at each $u^k \in C$ the oracle delivers an *approximate value* π_u^k and an *approximate subgradient* g_π^k of π that produce the *approximate linearization* of π :

$$\pi_k(\cdot) := \pi_u^k + \langle g_\pi^k, \cdot - u^k \rangle \leq \pi(\cdot) \quad \text{with} \quad \pi_k(u^k) = \pi_u^k \geq \pi(u^k) - \varepsilon_\pi. \quad (2.1)$$

Thus $\pi_u^k \in [\pi(u^k) - \varepsilon_\pi, \pi(u^k)]$, whereas g_π^k lies in the ε_π -subdifferential of π at u^k

$$\partial_{\varepsilon_\pi} \pi(u^k) := \left\{ g_\pi : \pi(\cdot) \geq \pi(u^k) - \varepsilon_\pi + \langle g_\pi, \cdot - u^k \rangle \right\}.$$

Then $\theta_u^k := \sigma_u^k + \pi_u^k$ is the approximate value of θ at u^k , where $\sigma_u^k := \sigma(u^k)$.

At iteration $k \geq 1$, the current *prox* (or *stability*) center $\hat{u}^k := u^{k(l)} \in C$ for some $k(l) \leq k$ has the value $\theta_{\hat{u}}^k := \theta_u^{k(l)}$ (usually $\theta_{\hat{u}}^k = \min_{j=1}^k \theta_u^j$); note that, by (2.1),

$$\theta_{\hat{u}}^k \in [\theta(\hat{u}^k) - \varepsilon_{\pi}, \theta(\hat{u}^k)]. \quad (2.2)$$

If $\pi_{\hat{u}}^k < \bar{\pi}_k(\hat{u}^k)$ in (1.6) due to evaluation errors, the *predicted descent*

$$v_k := \theta_{\hat{u}}^k - \left[\sigma(u^{k+1}) + \bar{\pi}_k(u^{k+1}) \right] \quad (2.3)$$

may be nonpositive; hence, if necessary, t_k is increased and (1.5)–(1.6) are solved again until $v_k \geq |u^{k+1} - \hat{u}^k|^2/2t_k$ as in [12, 13, 15]. A *descent* step to $\hat{u}^{k+1} := u^{k+1}$ is taken if

$$\theta_{\hat{u}}^{k+1} \leq \theta_{\hat{u}}^k - \kappa v_k \quad (2.4)$$

for a fixed $\kappa \in (0, 1)$. Otherwise, a *null* step $\hat{u}^{k+1} := \hat{u}^k$ occurs; then $\bar{\pi}_k$ and the new linearization π_{k+1} are used to produce a better model $\check{\pi}_{k+1} \geq \max\{\bar{\pi}_k, \pi_{k+1}\}$.

Specific rules of our method will be discussed after its formal statement below.

Algorithm 2.1

Step 0 (Initiation). Select $u^1 \in C$, a *descent parameter* $\kappa \in (0, 1)$, a *stepsize bound* $t_{\min} > 0$ and a *stepsize* $t_1 \geq t_{\min}$. Call the oracle at u^1 to obtain π_u^1 and g_{π}^1 of (2.1). Set $\bar{\pi}_0 := \pi_1$ by (2.1), and $\bar{\sigma}_0(\cdot) := \sigma(u^1) + \langle p_{\sigma}^0, \cdot - u^1 \rangle$ with $p_{\sigma}^0 \in \partial\sigma(u^1)$. Set $\hat{u}^1 := u^1$, $\theta_{\hat{u}}^1 := \theta_u^1 := \sigma_u^1 + \pi_u^1$ with $\sigma_u^1 := \sigma(u^1)$, $i_t^1 := 0$, $k := k(0) := 1$, $l := 0$ ($k(l) - 1$ will denote the iteration of the l th descent step).

Step 1 (Model selection). Choose $\check{\pi}_k : \mathbb{R}^m \rightarrow \mathbb{R}$ convex and such that

$$\max\{\bar{\pi}_{k-1}, \pi_k\} \leq \check{\pi}_k \leq \pi. \quad (2.5)$$

Step 2 (Solving the π -subproblem). Find \check{u}^{k+1} of (1.5) and the *aggregate linearization* of $\check{\pi}_k$

$$\bar{\pi}_k(\cdot) := \check{\pi}_k(\check{u}^{k+1}) + \langle p_{\pi}^k, \cdot - \check{u}^{k+1} \rangle \quad \text{with} \quad p_{\pi}^k := (\hat{u}^k - \check{u}^{k+1})/t_k - p_{\sigma}^{k-1}. \quad (2.6)$$

Step 3 (Solving the σ -subproblem). Find u^{k+1} of (1.6) and the *aggregate linearization* of σ

$$\bar{\sigma}_k(\cdot) := \sigma(u^{k+1}) + \langle p_{\sigma}^k, \cdot - u^{k+1} \rangle \quad \text{with} \quad p_{\sigma}^k := (\hat{u}^k - u^{k+1})/t_k - p_{\pi}^k. \quad (2.7)$$

Compute v_k of (2.3), and the aggregate subgradient and linearization error of θ

$$p^k := (\hat{u}^k - u^{k+1})/t_k \quad \text{and} \quad \varepsilon_k := v_k - t_k |p^k|^2. \tag{2.8}$$

Step 4 (Stopping criterion). If $\max\{|p^k|, \varepsilon_k\} = 0$, stop ($\theta_{\hat{u}}^k \leq \theta_*$).

Step 5 (Noise attenuation). If $v_k < -\varepsilon_k$, set $t_k := 10t_k$, $i_t^k := k$ and go back to Step 2.

Step 6 (Oracle call). Call the oracle at u^{k+1} to obtain π_u^{k+1} and g_{π}^{k+1} of (2.1).

Step 7 (Descent test). If the descent test (2.4) holds with $\theta_u^{k+1} := \sigma(u^{k+1}) + \pi_u^{k+1}$, set $\hat{u}^{k+1} := u^{k+1}$, $\theta_{\hat{u}}^{k+1} := \theta_u^{k+1}$, $i_t^{k+1} := 0$, $k(l+1) := k+1$ and increase l by 1 (descent step); otherwise, set $\hat{u}^{k+1} := \hat{u}^k$, $\theta_{\hat{u}}^{k+1} := \theta_{\hat{u}}^k$, and $i_t^{k+1} := i_t^k$ (null step).

Step 8 (Stepsize updating). If $k(l) = k+1$ (i.e., after a descent step), select $t_{k+1} \geq t_{\min}$; otherwise, either set $t_{k+1} := t_k$, or choose $t_{k+1} \in [t_{\min}, t_k]$ if $i_t^{k+1} = 0$.

Step 9 (Loop). Increase k by 1 and go to Step 1.

Several comments on the method are in order. Step 1 may choose the simplest model $\check{\pi}_k = \max\{\bar{\pi}_{k-1}, \pi_k\}$. More efficient choices are discussed in [13, Sect. 4.4] and [15, Sect. 2.3]. For polyhedral models, Step 2 may use the QP methods of [3, 8, 10], which can handle efficiently sequences of subproblems (1.5).

We now use the relations of Steps 2 and 3 to derive an optimality estimate, which involves the aggregate linearization $\bar{\theta}_k := \bar{\sigma}_k + \bar{\pi}_k$ and the optimality measure

$$V_k := \max \left\{ |p^k|, \varepsilon_k + \langle p^k, \hat{u}^k \rangle \right\}. \tag{2.9}$$

Lemma 2.2 (1) *The vectors p_{π}^k and p_{σ}^k of (2.6) and (2.7) are in fact subgradients:*

$$p_{\pi}^k \in \partial \check{\pi}_k(\check{u}^{k+1}) \quad \text{and} \quad p_{\sigma}^k \in \partial \sigma(u^{k+1}), \tag{2.10}$$

and the linearizations $\bar{\pi}_k$ and $\bar{\sigma}_k$ of (2.6) and (2.7) provide the minorizations

$$\bar{\pi}_k \leq \check{\pi}_k, \quad \bar{\sigma}_k \leq \sigma \quad \text{and} \quad \bar{\theta}_k := \bar{\pi}_k + \bar{\sigma}_k \leq \theta. \tag{2.11}$$

(2) *The aggregate subgradient p^k of (2.8) and the linearization $\bar{\theta}_k$ above satisfy*

$$p^k = p_{\pi}^k + p_{\sigma}^k = (\hat{u}^k - u^{k+1})/t_k, \tag{2.12}$$

$$\bar{\theta}_k(\cdot) = \bar{\theta}_k(u^{k+1}) + \langle p^k, \cdot - u^{k+1} \rangle. \tag{2.13}$$

(3) *The predicted descent v_k of (2.3) and the aggregate error ε_k of (2.8) satisfy*

$$v_k = \theta_{\hat{u}}^k - \bar{\theta}_k(u^{k+1}) = t_k |p^k|^2 + \varepsilon_k \quad \text{and} \quad \varepsilon_k = \theta_{\hat{u}}^k - \bar{\theta}_k(\hat{u}^k). \tag{2.14}$$

(4) The aggregate linearization $\bar{\theta}_k$ is expressed in terms of p^k and ε_k as follows:

$$\theta_{\hat{u}}^k - \varepsilon_k + \langle p^k, \cdot - \hat{u}^k \rangle = \bar{\theta}_k(\cdot) \leq \theta(\cdot). \tag{2.15}$$

(5) The optimality measure V_k of (2.9) satisfies $V_k \leq \max\{|p^k|, \varepsilon_k\}(1 + |\hat{u}^k|)$ and

$$\theta_{\hat{u}}^k \leq \theta(u) + V_k(1 + |u|) \text{ for all } u. \tag{2.16}$$

(6) We have $v_k \geq -\varepsilon_k \Leftrightarrow t_k|p^k|^2/2 \geq -\varepsilon_k \Leftrightarrow v_k \geq t_k|p^k|^2/2$. Moreover, $v_k \geq \varepsilon_k, -\varepsilon_k \leq \varepsilon_\pi$ and

$$v_k \geq \max\{t_k|p^k|^2/2, |\varepsilon_k|\} \quad \text{if } v_k \geq -\varepsilon_k, \tag{2.17}$$

$$V_k \leq \max\{(2v_k/t_k)^{1/2}, v_k\} (1 + |\hat{u}^k|) \quad \text{if } v_k \geq -\varepsilon_k, \tag{2.18}$$

$$V_k < (2\varepsilon_\pi/t_k)^{1/2} (1 + |\hat{u}^k|) \quad \text{if } v_k < -\varepsilon_k. \tag{2.19}$$

Proof (1) Let $\phi_\pi^k, \phi_\sigma^k$ denote the objectives of (1.5), (1.6). By (2.6), the optimality condition $0 \in \partial\phi_\pi^k(\check{u}^{k+1})$ for (1.5) with $\nabla\bar{\sigma}_{k-1} = p_\sigma^{k-1}$ by Step 0 and (2.7), i.e.,

$$0 \in \partial\phi_\pi^k(\check{u}^{k+1}) = \partial\check{\pi}_k(\check{u}^{k+1}) + p_\sigma^{k-1} + (\check{u}^{k+1} - \hat{u}^k)/t_k = \partial\check{\pi}_k(\check{u}^{k+1}) - p_\pi^k,$$

and $\bar{\pi}_k(\check{u}^{k+1}) = \check{\pi}_k(\check{u}^{k+1})$ yield $p_\pi^k \in \partial\check{\pi}_k(\check{u}^{k+1})$ and $\bar{\pi}_k \leq \check{\pi}_k$. Similarly, by (2.7),

$$0 \in \partial\phi_\sigma^k(u^{k+1}) = p_\pi^k + \partial\sigma(u^{k+1}) + (u^{k+1} - \hat{u}^k)/t_k = \partial\sigma(u^{k+1}) - p_\sigma^k$$

(using $\nabla\bar{\pi}_k = p_\pi^k$) and $\bar{\sigma}_k(u^{k+1}) = \sigma(u^{k+1})$ give $p_\sigma^k \in \partial\sigma(u^{k+1})$ and $\bar{\sigma}_k \leq \sigma$. Combining both minorizations, we obtain that $\bar{\pi}_k + \bar{\sigma}_k \leq \check{\pi}_k + \sigma \leq \theta$ by (2.5) and (1.1).

- (2) Use the linearity of $\bar{\theta}_k := \bar{\pi}_k + \bar{\sigma}_k$, (2.6), (2.7) and (2.8).
- (3) Rewrite (2.3), using the fact that $\bar{\theta}_k(\hat{u}^k) = \bar{\theta}_k(u^{k+1}) + t_k|p^k|^2$ by (2).
- (4) We have $\theta_{\hat{u}}^k - \varepsilon_k = \bar{\theta}_k(\hat{u}^k)$ by (3), $\bar{\theta}_k$ is affine by (2) and minorizes θ by (1).
- (5) Using the Cauchy–Schwarz inequality in the definition (2.9) gives

$$\begin{aligned} V_k &\leq \max\{|p^k|, \varepsilon_k + |p^k||\hat{u}^k|\} \leq \max\{|p^k|, \varepsilon_k\} + |p^k||\hat{u}^k| \\ &\leq \max\{|p^k|, \varepsilon_k\} (1 + |\hat{u}^k|). \end{aligned}$$

Since $|a||b| + c \leq \max\{|a|, c\}(1 + |b|)$ for any scalars a, b, c , in (2.15) we have

$$\begin{aligned} -\langle p^k, u \rangle + \varepsilon_k + \langle p^k, \hat{u}^k \rangle &\leq |p^k||u| + \varepsilon_k + \langle p^k, \hat{u}^k \rangle \\ &\leq \max\{|p^k|, \varepsilon_k + \langle p^k, \hat{u}^k \rangle\}(1 + |u|). \end{aligned}$$

(6) The equivalences follow from the expression of $v_k = t_k|p^k|^2 + \varepsilon_k$ in (3); in particular, $v_k \geq \varepsilon_k$. Next, by (2.14), (2.11) and (2.2), we have

$$-\varepsilon_k = \bar{\theta}_k(\hat{u}^k) - \theta_{\hat{u}}^k \leq \theta(\hat{u}^k) - \theta_{\hat{u}}^k \leq \varepsilon_{\pi}.$$

Finally, to obtain the bounds (2.17)–(2.19), use the equivalences together with the facts that $v_k \geq \varepsilon_k$, $-\varepsilon_k \leq \varepsilon_{\pi}$ and the bound on V_k from assertion (5). \square

The optimality estimate (2.16) justifies the stopping criterion of Step 4: $V_k = 0$ yields $\theta_{\hat{u}}^k \leq \inf \theta = \theta_*$; thus, the point \hat{u}^k is ε_{π} -optimal, i.e., $\theta(\hat{u}^k) \leq \theta_* + \varepsilon_{\pi}$ by (2.2). If the oracle is exact ($\varepsilon_{\pi} = 0$), we have $v_k \geq \varepsilon_k \geq 0$ by Lemma 2.2(6), and Step 5 is redundant. When inexactness is discovered at Step 5 via $v_k < -\varepsilon_k$ and the stepsize t_k is increased, the *stepsize indicator* $i_t^k \neq 0$ prevents Step 7 from decreasing t_k after null steps until the next descent step occurs (cf. Step 6). At Step 6, we have $u^{k+1} \in C$ and $v_k > 0$ (by (2.17), since $\max\{|p^k|, \varepsilon_k\} > 0$ at Step 4), so that $\hat{u}^{k+1} \in C$ and $\theta_{\hat{u}}^{k+1} \leq \theta_{\hat{u}}^k$.

3 Convergence

With Lemma 2.2 replacing [13, Lemma 3.2], it is easy to check that the convergence results of [13, Sect.3] will hold once we prove [13, Lemma 3.2] for our method. To this end, as usual, we assume that the oracle’s subgradients are *locally bounded*:

$$\{g_{\pi}^k\} \text{ is bounded if } \{u^k\} \text{ is bounded.} \tag{3.1}$$

Further, as in [13], we assume that the model subgradients p_{π}^k in (2.10) satisfy

$$\{p_{\pi}^k\} \text{ is bounded if } \{u^k\} \text{ is bounded.} \tag{3.2}$$

Remark 3.1 Note that (3.1) holds if $C = \mathbb{R}^m$ or if π can be extended to become finite-valued on a neighborhood of C , since $g_{\pi}^k \in \partial_{\varepsilon_{\pi}} \pi(u^k)$ by (2.1), whereas the mapping $\partial_{\varepsilon_{\pi}} \pi$ is locally bounded on C in both cases [7, Sect. XI.4.1]. As discussed in [13, Remark 4.4], typical models $\check{\pi}_k$ satisfy condition (3.2) automatically when (3.1) holds.

A suitable modification of the proof of [13, Lemma 3.2] follows.

Lemma 3.2 *Suppose there exists \bar{k} such that for all $k \geq \bar{k}$, only null steps occur and Step 5 doesn’t increase t_k . Then $V_k \rightarrow 0$.*

Proof Let ϕ_{π}^k and ϕ_{σ}^k denote the objectives of subproblems (1.5) and (1.6). First, using partial linearizations of these subproblems, we show that their optimal values $\phi_{\pi}^k(\check{u}^{k+1}) \leq \phi_{\sigma}^k(u^{k+1})$ are nondecreasing and bounded above for $k \geq \bar{k}$.

Fix $k \geq \bar{k}$. By the definitions in (1.5) and (2.6), we have $\bar{\pi}_k(\check{u}^{k+1}) = \check{\pi}_k(\check{u}^{k+1})$ and

$$\check{u}^{k+1} = \arg \min \left\{ \bar{\phi}_{\pi}^k(\cdot) := \bar{\pi}_k(\cdot) + \bar{\sigma}_{k-1}(\cdot) + \frac{1}{2t_k} |\cdot - \hat{u}^k|^2 \right\} \tag{3.3}$$

from $\nabla \bar{\phi}_\pi^k(\check{u}^{k+1}) = 0$. Since $\bar{\phi}_\pi^k$ is quadratic and $\bar{\phi}_\pi^k(\check{u}^{k+1}) = \phi_\pi^k(\check{u}^{k+1})$, by Taylor’s expansion

$$\bar{\phi}_\pi^k(\cdot) = \phi_\pi^k(\check{u}^{k+1}) + \frac{1}{2t_k} |\cdot - \check{u}^{k+1}|^2. \tag{3.4}$$

Similarly, by the definitions in (1.6) and (2.7), we have $\bar{\sigma}_k(u^{k+1}) = \sigma(u^{k+1})$,

$$u^{k+1} = \arg \min \left\{ \bar{\phi}_\sigma^k(\cdot) := \bar{\pi}_k(\cdot) + \bar{\sigma}_k(\cdot) + \frac{1}{2t_k} |\cdot - \hat{u}^k|^2 \right\}, \tag{3.5}$$

$$\bar{\phi}_\sigma^k(\cdot) = \phi_\sigma^k(u^{k+1}) + \frac{1}{2t_k} |\cdot - u^{k+1}|^2. \tag{3.6}$$

Next, to bound the objective values of the linearized subproblems (3.3) and (3.5) from above, we use the minorizations $\bar{\pi}_k \leq \pi$ and $\bar{\sigma}_{k-1}, \bar{\sigma}_k \leq \sigma$ of (2.11) for $\theta := \pi + \sigma$:

$$\phi_\pi^k(\check{u}^{k+1}) + \frac{1}{2t_k} |\check{u}^{k+1} - \hat{u}^k|^2 = \bar{\phi}_\pi^k(\hat{u}^k) \leq \theta(\hat{u}^k), \tag{3.7a}$$

$$\phi_\sigma^k(u^{k+1}) + \frac{1}{2t_k} |u^{k+1} - \hat{u}^k|^2 = \bar{\phi}_\sigma^k(\hat{u}^k) \leq \theta(\hat{u}^k), \tag{3.7b}$$

where the equalities stem from (3.4) and (3.6). Due to the minorization $\bar{\sigma}_{k-1} \leq \sigma$, the objectives of (3.3) and (1.6) satisfy $\bar{\phi}_\pi^k \leq \phi_\sigma^k$. On the other hand, since $\hat{u}^{k+1} = \hat{u}^k$, $t_{k+1} \leq t_k$ (cf. Step 7), and $\bar{\pi}_k \leq \bar{\pi}_{k+1}$ by (2.5), the objectives of (3.5) and the next subproblem (1.5) satisfy $\bar{\phi}_\sigma^k \leq \phi_\pi^{k+1}$. Altogether, by (3.4) and (3.6), we see that

$$\phi_\pi^k(\check{u}^{k+1}) + \frac{1}{2t_k} |u^{k+1} - \check{u}^{k+1}|^2 = \bar{\phi}_\pi^k(u^{k+1}) \leq \phi_\sigma^k(u^{k+1}), \tag{3.8a}$$

$$\phi_\sigma^k(u^{k+1}) + \frac{1}{2t_k} |\check{u}^{k+2} - u^{k+1}|^2 = \bar{\phi}_\sigma^k(\check{u}^{k+2}) \leq \phi_\pi^{k+1}(\check{u}^{k+2}). \tag{3.8b}$$

In particular, the inequalities $\phi_\pi^k(\check{u}^{k+1}) \leq \phi_\sigma^k(u^{k+1}) \leq \phi_\pi^{k+1}(\check{u}^{k+2})$ imply that the non-decreasing sequences $\{\phi_\pi^k(\check{u}^{k+1})\}_{k \geq \bar{k}}$ and $\{\phi_\sigma^k(u^{k+1})\}_{k \geq \bar{k}}$, which are bounded above by (3.7) with $\hat{u}^k = \hat{u}^{\bar{k}}$ for all $k \geq \bar{k}$, must have a common limit, say $\phi_\infty \leq \theta(\hat{u}^{\bar{k}})$. Moreover, since $t_k \leq t_{\bar{k}}$ for all $k \geq \bar{k}$, we deduce from the bounds (3.7)–(3.8) that

$$\phi_\pi^k(\check{u}^{k+1}), \phi_\sigma^k(u^{k+1}) \uparrow \phi_\infty, \quad \check{u}^{k+2} - u^{k+1} \rightarrow 0, \tag{3.9}$$

and the sequences $\{\check{u}^{k+1}\}$ and $\{u^{k+1}\}$ are bounded. Then the sequences $\{g_\pi^k\}$ and $\{p_\pi^k\}$ are bounded by (3.1) and (3.2).

We now show that the approximation error $\bar{e}_k := \pi_u^{k+1} - \bar{\pi}_k(u^{k+1})$ vanishes. Using the form (2.1) of π_{k+1} , the minorization $\pi_{k+1} \leq \bar{\pi}_{k+1}$ of (2.5), the Cauchy–Schwarz inequality, and the optimal values of subproblems (1.5) and (1.6) with $\hat{u}^k = \hat{u}^{\bar{k}}$ for

$k \geq \bar{k}$, we estimate

$$\begin{aligned} \bar{\varepsilon}_k &:= \pi_{u^{k+1}} - \bar{\pi}_k(u^{k+1}) = \pi_{k+1}(\check{u}^{k+2}) - \bar{\pi}_k(u^{k+1}) + \langle g_\pi^{k+1}, u^{k+1} - \check{u}^{k+2} \rangle \\ &\leq \check{\pi}_{k+1}(\check{u}^{k+2}) - \bar{\pi}_k(u^{k+1}) + |g_\pi^{k+1}| |u^{k+1} - \check{u}^{k+2}| \\ &= \phi_\pi^{k+1}(\check{u}^{k+2}) - \phi_\sigma^k(u^{k+1}) + \Delta_u^k + \Delta_\sigma^k + |g_\pi^{k+1}| |u^{k+1} - \check{u}^{k+2}|, \end{aligned} \tag{3.10}$$

where $\Delta_u^k := |u^{k+1} - \hat{u}^k|^2/2t_k - |\check{u}^{k+2} - \hat{u}^k|^2/2t_{k+1}$ and $\Delta_\sigma^k := \sigma_u^{k+1} - \bar{\sigma}_k(\check{u}^{k+2})$; in fact, $\Delta_\sigma^k = -\langle p_\sigma^k, \check{u}^{k+2} - u^{k+1} \rangle$ by (2.7). To see that $\Delta_u^k \rightarrow 0$, note that

$$|\check{u}^{k+2} - \hat{u}^k|^2 = |u^{k+1} - \hat{u}^k|^2 + 2\langle \check{u}^{k+2} - u^{k+1}, u^{k+1} - \hat{u}^k \rangle + |\check{u}^{k+2} - u^{k+1}|^2,$$

$|u^{k+1} - \hat{u}^k|^2$ is bounded, $\check{u}^{k+2} - u^{k+1} \rightarrow 0$ by (3.9), and $t_{\min} \leq t_{k+1} \leq t_k$ for $k \geq \bar{k}$ by Step 7. These properties also give $\Delta_\sigma^k \rightarrow 0$, since by (2.7) and the Cauchy–Schwarz inequality,

$$|\Delta_\sigma^k| \leq |p_\sigma^k| |\check{u}^{k+2} - u^{k+1}| \quad \text{with} \quad |p_\sigma^k| \leq |u^{k+1} - \hat{u}^k|/t_k + |p_\pi^k|,$$

where $\{p_\pi^k\}$ is bounded. Hence, using (3.9) and the boundedness of $\{g_\pi^{k+1}\}$ in (3.10) yields $\lim_k \bar{\varepsilon}_k \leq 0$. On the other hand, $\bar{\varepsilon}_k = \theta_u^{k+1} - \bar{\theta}_k(u^{k+1})$ from $\bar{\sigma}_k(u^{k+1}) = \sigma_u^{k+1}$ in (2.7), while for $k \geq \bar{k}$ the null step condition $\theta_u^{k+1} > \theta_{\hat{u}}^k - \kappa v_k$ gives

$$\bar{\varepsilon}_k = \left[\theta_u^{k+1} - \theta_{\hat{u}}^k \right] + \left[\theta_{\hat{u}}^k - \bar{\theta}_k(u^{k+1}) \right] > -\kappa v_k + v_k = (1 - \kappa)v_k \geq 0$$

by (2.14), where $\kappa < 1$ by Step 0; we conclude that $\bar{\varepsilon}_k \rightarrow 0$ and $v_k \rightarrow 0$. Finally, since $v_k \rightarrow 0$, $t_k \geq t_{\min}$ (cf. Step 7) and $\hat{u}^k = \hat{u}^{\bar{k}}$ for $k \geq \bar{k}$, we have $V_k \rightarrow 0$ by (2.18). □

Our principal result on the asymptotic objective value $\theta_{\hat{u}}^\infty := \lim_k \theta_{\hat{u}}^k$ follows.

Theorem 3.3 (1) We have $\theta_{\hat{u}}^k \downarrow \theta_{\hat{u}}^\infty \leq \theta_*$, and additionally $\lim_k V_k = 0$ if $\theta_* > -\infty$.
 (2) $\theta_* \leq \underline{\lim}_k \theta(\hat{u}^k) \leq \overline{\lim}_k \theta(\hat{u}^k) \leq \theta_{\hat{u}}^\infty + \varepsilon_\pi$.

Proof Use the proof of [13, Theorem 3.5], with obvious modifications. □

4 Modifications

4.1 Looping between subproblems

To obtain a more accurate solution to subproblem (1.4) with π replaced by $\check{\pi}_k$, we may cycle between subproblems (1.5) and (1.6), updating their data as if null steps occurred without changing the model $\check{\pi}_k$. Specifically, for a given subproblem accuracy threshold $\check{\kappa} \in (0, 1)$, suppose that the following step is inserted after Step 5.

Step 5' (*Subproblem accuracy test*). If

$$\sigma(u^{k+1}) + \check{\pi}_k(u^{k+1}) > \theta_{\check{u}}^k - \check{\kappa}v_k, \tag{4.1}$$

set $\bar{\sigma}_{k-1}(\cdot) := \bar{\sigma}_k(\cdot)$, $p_{\sigma}^{k-1} := p_{\sigma}^k$ and go back to Step 2.

The main aim of this modification is to avoid “unnecessary” null steps. Namely, if the test (4.1) holds with $\check{\kappa} \leq \kappa$ and the oracle is exact enough to deliver $\pi_u^{k+1} \geq \check{\pi}_k(u^{k+1})$, then the descent test (2.4) can't hold and a null step must occur, which is bypassed by Step 5'.

When the oracle is expensive, the optional use of Step 5' with $\check{\kappa} \in [\kappa, 1)$ gives room for deciding whether to continue working with the current model $\check{\pi}_k$ before calling the oracle.

Convergence for this modification can be analyzed as in [13, Remark 4.1]. Omitting details for brevity, here we just observe that for the test (4.1) written as (cf. (2.14))

$$\check{\epsilon}_k := \check{\pi}_k(u^{k+1}) - \bar{\pi}_k(u^{k+1}) > (1 - \check{\kappa})v_k,$$

the $\check{\epsilon}_k$ above may play the role of $\bar{\epsilon}_k$ in (3.10).

4.2 Solving the σ -subproblem approximately

For a given tolerance $\kappa_N \in (0, 1 - \kappa)$, suppose Step 3 is replaced by the following. Step 3' (*Solving the σ -subproblem approximately*). Find a linearization $\bar{\sigma}_k \leq \sigma$ s.t.

$$\phi_{\pi}^k(\check{u}^{k+1}) \leq \bar{\phi}_{\sigma}^k(u^{k+1}), \tag{4.2}$$

$$\sigma(u^{k+1}) - \bar{\sigma}_k(u^{k+1}) \leq \kappa_N v_k, \tag{4.3}$$

for u^{k+1} given by (3.5) and v_k by (2.14). Set p^k and ϵ_k by (2.8), and $p_{\sigma}^k := \nabla \bar{\sigma}_k$.

Before discussing implementations, we show that Step 3' does not spoil convergence. In Sect. 2, $\bar{\sigma}_k(u^{k+1})$ replaces $\sigma(u^{k+1})$ in (2.3), (2.7) and (2.10). In Sect. 3, it suffices to validate Lemma 3.2.

Lemma 4.1 *Lemma 3.2 still holds for Step 3 replaced by Step 3' above.*

Proof We only sketch how to modify the proof of Lemma 3.2. First, referring to (3.5) instead of (1.6), replace ϕ_{σ}^k by $\bar{\phi}_{\sigma}^k$ throughout, and (3.8a) by (4.2). Second, let $\Delta_{\sigma}^k := \bar{\sigma}_k(u^{k+1}) - \bar{\sigma}_k(\check{u}^{k+2})$ in (3.10). Third, by (4.3), the null step condition yields

$$\bar{\sigma}_k(u^{k+1}) + \pi_u^{k+1} > \theta_{\check{u}}^k - \kappa v_k + \bar{\sigma}_k(u^{k+1}) - \sigma(u^{k+1}) \geq \theta_{\check{u}}^k - \tilde{\kappa}v_k$$

for $\tilde{\kappa} := \kappa + \kappa_N < 1$, and hence

$$\bar{\epsilon}_k = \bar{\sigma}_k(u^{k+1}) + \pi_u^{k+1} - \bar{\theta}_k(u^{k+1}) > (1 - \tilde{\kappa})v_k \geq 0,$$

so that the proof may finish as before. □

Step 3' can be implemented by solving the Fenchel dual of (1.6) approximately. Indeed, using the representation $\sigma(\cdot) = \sup_z \{ \langle z, \cdot \rangle - \sigma^*(z) \}$ in (1.6), consider the Lagrangian

$$L(u, z) := \langle z, u \rangle - \sigma^*(z) + \bar{\pi}_k(u) + \frac{1}{2t_k} |u - \hat{u}^k|^2, \tag{4.4}$$

and associate with each dual point $z \in \text{dom } \sigma^*$ the following quantities:

$$\bar{u}(z) := \arg \min_u L(u, z) = \hat{u}^k - t_k \left(p_\pi^k + z \right), \tag{4.5}$$

$$\bar{\sigma}(\cdot; z) := \langle z, \cdot \rangle - \sigma^*(z), \tag{4.6}$$

$$\varepsilon(z) := \sigma(\bar{u}(z)) - \bar{\sigma}(\bar{u}(z); z) = \sigma(\bar{u}(z)) + \sigma^*(z) - \langle z, \bar{u}(z) \rangle, \tag{4.7}$$

$$v(z) := \theta_{\hat{u}}^k - [\bar{\pi}_k(\bar{u}(z)) + \bar{\sigma}(\bar{u}(z); z)], \tag{4.8}$$

where $\bar{u}(z)$ is the Lagrangian solution (with $p_\pi^k = \nabla \bar{\pi}_k$), $\bar{\sigma}(\cdot; z)$ is the linearization of σ , $\varepsilon(z)$ is its linearization error at $\bar{u}(z)$, and $v(z)$ is the predicted descent. Maximizing $L(\bar{u}(z), z)$ or minimizing $w(z) = -L(\bar{u}(z), z)$ leads to the following dual problem:

$$w_* := \min_z \left\{ w(z) := \sigma^*(z) + \frac{t_k}{2} |p_\pi^k + z|^2 - \langle z, \hat{u}^k \rangle - \bar{\pi}_k(\hat{u}^k) \right\}, \tag{4.9}$$

with a unique solution z^* giving $u^* := \bar{u}(z^*)$ such that $u^* \in \partial \sigma^*(z^*)$, $z^* \in \partial \sigma(u^*)$ and

$$\sigma(u^*) + \sigma^*(z^*) - \langle z^*, u^* \rangle = 0; \tag{4.10}$$

not suprisingly, u^* is the exact solution of (1.6) and z^* is the corresponding p_σ^k in (2.7). Note that (4.9) can be restricted to the set $D := \text{dom } \partial \sigma^* := \{z : \partial \sigma^*(z) \neq \emptyset\}$, which contains z^* .

Now, suppose we have a method for solving (4.9) with the following properties:

- (1) It starts from the point $z^1 := p_\sigma^{k-1} \in D$ such that $\sigma_{k-1}(\cdot) = \langle z^1, \cdot \rangle - \sigma^*(z^1)$; thus, by (3.3), (3.4) and (4.4)–(4.6), $w(z^1) = -\phi_\pi^k(\check{u}^{k+1})$ from $w(z^1) = -L(\bar{u}(z^1), z^1)$.
- (2) It generates points $z^i \in D$ with $w(z^i) \leq w(z^1)$ such that $z^i \rightarrow z^*$, $\sigma^*(z^i) \rightarrow \sigma^*(z^*)$ and $\sigma(\bar{u}(z^i)) \rightarrow \sigma(u^*)$, where $\bar{u}(z^i) \rightarrow u^*$ by (4.5).

Then $\varepsilon(z^i) \rightarrow 0$ by (4.7) and (4.10), whereas $v(z^i) \rightarrow v(z^*)$ by (4.8). Thus, if $v(z^*) > 0$, we will eventually have $\varepsilon(z^i) \leq \kappa_N v(z^i)$. Then the method may stop with $u^{k+1} := \bar{u}(z^i)$, $v_k := v(z^i)$, $\bar{\sigma}_k(\cdot) := \bar{\sigma}(\cdot; z^i)$ and $p_\sigma^k := z^i$ to meet the requirements of Step 3', with (4.2) following from $-\phi_\sigma^k(u^{k+1}) = w(z^i) \leq w(z^1) = -\phi_\pi^k(\check{u}^{k+1})$; see (1) above and (3.5).

As for the assumptions in (2) above, note that $\sigma^*(z^i) \rightarrow \sigma^*(z^*)$ if σ^* is continuous on $D := \text{dom } \partial \sigma^*$ (e.g., in Sect. 6.3). Similarly, $\sigma(\bar{u}(z^i)) \rightarrow \sigma(u^*)$ holds if σ is continuous on $\text{dom } \partial \sigma$ and $\bar{u}(z^i) \in \text{dom } \partial \sigma$ for large i .

5 Lagrangian relaxation

We now consider the application of our method to (1.2) treated as the *primal problem*

$$\varphi_* := \sup \{ \varphi(y) := -f(y) \} \quad \text{s.t.} \quad \psi(x, y) := y - Ax = 0, \quad x \in X, \quad (5.1)$$

assuming that f is closed proper convex and the set $X \neq \emptyset$ is compact and convex. In view of (1.3) and (2.1), suppose that, at each $u^k \in C$, the oracle delivers

$$g_\pi^k := -Ax^k \quad \text{and} \quad \pi_k(\cdot) := \langle -Ax^k, \cdot \rangle \quad \text{for some } x^k \in X. \quad (5.2)$$

For simplicity, let Step 1 retain only selected past linearizations for its k th model

$$\check{\pi}_k(\cdot) := \max_{j \in J_k} \pi_j(\cdot) \quad \text{with} \quad k \in J_k \subset \{1, \dots, k\}. \quad (5.3)$$

Then (see (2.10) and [13, Sect. 4.4]) there are convex weights $v_j^k \geq 0$ such that

$$(\bar{\pi}_k, p_\pi^k, 1) = \sum_{j \in \hat{J}_k} v_j^k (\pi_j, g_\pi^j, 1) \quad \text{with} \quad \hat{J}_k := \{j \in J_k : v_j^k > 0\}, \quad (5.4)$$

and for convergence it suffices to choose $J_{k+1} \supset \hat{J}_k \cup \{k+1\}$. Using these weights and (2.7), we may estimate a solution to (5.1) via the *aggregate primal solution* (\hat{x}^k, \hat{y}^k) :

$$\hat{x}^k := \sum_{j \in J_k} v_j^k x^j \quad \text{and} \quad \hat{y}^k := p_\sigma^k. \quad (5.5)$$

We first derive useful expressions of $\varphi(\hat{y}^k)$ and $\psi(\hat{x}^k, \hat{y}^k)$.

Lemma 5.1 *We have $\hat{x}^k \in X$, $\varphi(\hat{y}^k) = \theta_u^k - \varepsilon_k - \langle p^k, \hat{u}^k \rangle$ and $\psi(\hat{x}^k, \hat{y}^k) = p^k$.*

Proof First, $\hat{x}^k \in \text{co}\{x^j\}_{j \in \hat{J}_k} \subset X$, $\bar{\pi}_k(\cdot) = \langle -A\hat{x}^k, \cdot \rangle$ and $p_\pi^k = -A\hat{x}^k$ by convexity of X , (5.2), (5.4) and (5.5). Then $p^k = \hat{y}^k - A\hat{x}^k = \psi(\hat{x}^k, \hat{y}^k)$ by (2.12), (5.1) and (5.5). Next, by [20, Theorem 23.5], the inclusion $\hat{y}^k := p_\sigma^k \in \partial\sigma(u^{k+1})$ of (2.10) with $\sigma := f^*$ in (1.3) yields $\sigma(u^{k+1}) = \langle u^{k+1}, \hat{y}^k \rangle - f(\hat{y}^k)$; thus $\varphi(\hat{y}^k) := -f(\hat{y}^k) = \bar{\sigma}_k(0)$ by (5.1) and (2.7). Since $\bar{\pi}_k(0) = 0$ in (2.11), (2.15) gives $\bar{\sigma}_k(0) = \bar{\theta}_k(0) = \theta_u^k - \varepsilon_k + \langle p^k, \hat{u}^k \rangle$, as required. \square

In terms of the optimality measure V_k of (2.9), Lemma 5.1 says that

$$\hat{x}^k \in X \quad \text{with} \quad \varphi(\hat{y}^k) \geq \theta_u^k - V_k, \quad |\psi(\hat{x}^k, \hat{y}^k)| \leq V_k. \quad (5.6)$$

We now show that $\{(\hat{x}^k, \hat{y}^k)\}$ has cluster points in the set of ε_π -optimal solutions of (5.1)

$$Z_{\varepsilon_\pi} := \{ (x, y) \in X \times \mathbb{R}^m : \varphi(y) \geq \varphi_* - \varepsilon_\pi, \psi(x, y) = 0 \}, \quad (5.7)$$

unless $\varphi_* = -\infty$, i.e., the primal problem is infeasible. Note that (5.2) with X compact and (5.4) yield (3.1)–(3.2), as required for Theorem 3.3.

Theorem 5.2 *Either $\theta_* = -\infty$ and $\theta_{\hat{u}}^k \downarrow -\infty$, in which case the primal problem (5.1) is infeasible, or $\theta_* > -\infty$, $\theta_{\hat{u}}^k \downarrow \theta_{\hat{u}}^\infty \in [\theta_* - \varepsilon_\pi, \theta_*]$, $\overline{\lim}_k \theta(\hat{u}^k) \leq \theta_{\hat{u}}^\infty + \varepsilon_\pi$ and $\underline{\lim}_k V_k = 0$. In the latter case, let $K \subset \mathbb{N}$ be a subsequence such that $V_k \xrightarrow{K} 0$. Then:*

- (1) $\{(\hat{x}^k, \hat{y}^k)\}_{k \in K}$ is bounded and all its cluster points lie in the set $X \times \mathbb{R}^m$.
- (2) Let $(\hat{x}^\infty, \hat{y}^\infty)$ be a cluster point of the sequence $\{(\hat{x}^k, \hat{y}^k)\}_{k \in K}$. Then $(\hat{x}^\infty, \hat{y}^\infty) \in Z_{\varepsilon_\pi}$.
- (3) $d_{Z_{\varepsilon_\pi}}((\hat{x}^k, \hat{y}^k)) := \inf_{(x,y) \in Z_{\varepsilon_\pi}} |(\hat{x}^k, \hat{y}^k) - (x, y)| \xrightarrow{K} 0$.
- (4) If $\varepsilon_\pi = 0$, then $\theta_{\hat{u}}^k \downarrow \theta_*$, $\varphi(\hat{y}^k) \xrightarrow{K} \varphi_* = \theta_*$, and $\psi(\hat{x}^k, \hat{y}^k) \xrightarrow{K} 0$.

Proof The first assertion follows from Theorem 3.3 (since $\theta_* = -\infty$ implies primal infeasibility by weak duality). In the second case, using $\theta_{\hat{u}}^k \downarrow \theta_{\hat{u}}^\infty \geq \theta_* - \varepsilon_\pi$ and $V_k \xrightarrow{K} 0$ in the bounds of (5.6) yields $\underline{\lim}_{k \in K} \varphi(\hat{y}^k) \geq \theta_* - \varepsilon_\pi$ and $\lim_{k \in K} \psi(\hat{x}^k, \hat{y}^k) = 0$.

- (1) By (5.6), $\{\hat{x}^k\}$ lies in the compact X , and $\{\hat{y}^k\}_{k \in K}$ is bounded by (5.1), (5.6).
- (2) We have $\hat{x}^\infty \in X$, $\varphi(\hat{y}^\infty) \geq \theta_* - \varepsilon_\pi$ and $\psi(\hat{x}^\infty, \hat{y}^\infty) = 0$ by closedness of φ and continuity of ψ . Since $\theta_* \geq \varphi_*$ by weak duality (cf. (1.1), (1.3), (5.1)), we get $\varphi(\hat{y}^\infty) \geq \varphi_* - \varepsilon_\pi$. Thus $(\hat{x}^\infty, \hat{y}^\infty) \in Z_{\varepsilon_\pi}$ by the definition (5.7).
- (3) This follows from (1), (2) and the continuity of the distance function $d_{Z_{\varepsilon_\pi}}$.
- (4) In the proof of (2), $\theta_* \geq \varphi_* \geq \varphi(\hat{y}^\infty) \geq \theta_*$ yields $\varphi_* = \varphi(\hat{y}^\infty) = \theta_*$, and for $K' \subset K$ such that $\hat{y}^k \xrightarrow{K'} \hat{y}^\infty$ we have $\varphi(\hat{y}^\infty) \geq \underline{\lim}_{k \in K'} \varphi(\hat{y}^k) \geq \underline{\lim}_{k \in K'} \varphi(\hat{y}^k) \geq \theta_*$, i.e., $\varphi(\hat{y}^k) \xrightarrow{K'} \varphi_*$. So considering convergent subsequences in (1) gives $\varphi(\hat{y}^k) \xrightarrow{K} \varphi_*$. □

6 Application to multicommodity network flows

6.1 The nonlinear multicommodity flow problem

Let $(\mathcal{N}, \mathcal{A})$ be a directed graph with $N := |\mathcal{N}|$ nodes and $m := |\mathcal{A}|$ arcs. Let $E \in \mathbb{R}^{N \times m}$ be its node-arc incidence matrix. There are n commodities to be routed through the network. For each commodity i there is a *required flow* $r_i > 0$ from its *source* node o_i to its *sink* node d_i . Let s_i be the *supply* N -vector of commodity i , having components $s_{io_i} = r_i, s_{id_i} = -r_i, s_{il} = 0$ if $l \neq o_i, d_i$. Our *nonlinear multicommodity flow problem* (NMFP for short) is:

$$\min f(y) := \sum_{j=1}^m f_j(y_j) \tag{6.1a}$$

$$\text{s.t. } y = \sum_{i=1}^n x_i, \tag{6.1b}$$

$$x_i \in X_i := \{x_i : Ex_i = s_i, 0 \leq x_i \leq \bar{x}_i\}, \quad i = 1 : n, \tag{6.1c}$$

where each *arc cost* function f_j is closed proper convex, y is the *total flow* vector, x_i is the *flow vector* of commodity i , and \bar{x}_i is a fixed positive vector of *flow bounds*.

Our assumptions seem to be weaker than those used in the literature. We add that if $\text{dom } f^* \subset \mathbb{R}_+^m$, then the bounds \bar{x}_i are not needed in (6.1c): Even if they are absent, our algorithm will proceed as if we had $\bar{x}_{ij} = r_i$ for all i and j ; cf. [14, Sect. 7.2].

6.2 Primal recovery

We may treat problem (6.1) as (5.1) with $Ax = \sum_{i=1}^n x_i$, $X = \prod_{i=1}^n X_i$, and the oracle solving shortest path problems to evaluate $\pi(u^k) = -\sum_{i=1}^n \min\{\langle u^k, x_i \rangle : x_i \in X_i\}$ at each u^k . Thus the results of Sect. 5 hold. Yet, as in [14, Sect. 7.3], for stopping criteria it is useful to employ another aggregate solution (\hat{x}^k, \check{y}^k) with \hat{x}^k given by (5.5) and

$$\check{y}^k := A\hat{x}^k = \sum_{i=1}^n \hat{x}_i^k, \quad (6.2)$$

which satisfies the constraints of (6.1). Thus $f(\check{y}^k) \geq f_*$, where the optimal value f_* of (6.1) satisfies $-f_* = \varphi_* \leq \theta_*$ by weak duality. Hence, if the oracle is exact, $\theta_u^k \geq \theta_*$ implies that the method may stop when $f(\check{y}^k) + \theta_u^k \leq \varepsilon$ for a given tolerance $\varepsilon > 0$, in which case (\hat{x}^k, \check{y}^k) is an ε -solution of (6.1). This stopping criterion will be met for some k under conditions similar to those in [14, Proposition 7.1].

Proposition 6.1 *Suppose problem (6.1) is feasible and has a unique optimal total flow y^* (e.g., f is strictly convex on $\mathbb{R}_+^m \cap \text{dom } f$) that satisfies $y^* \in [0, c) \subset \text{dom } f$ for some $c \in \mathbb{R}_+^m$. Further, let $\varepsilon_\pi = 0$ (i.e., the oracle is exact), and let $K \subset \mathbb{N}$ be a subsequence such that $V_k \xrightarrow{K} 0$. Then $\check{y}^k \xrightarrow{K} y^*$, $f(\check{y}^k) \xrightarrow{K} f_* = -\theta_*$ and $f(\check{y}^k) + \theta_u^k \xrightarrow{K} 0$.*

Proof By Theorem 5.2(3) and the uniqueness of y^* , $\hat{y}^k \xrightarrow{K} y^*$. Hence $\check{y}^k \xrightarrow{K} y^*$ from $\hat{y}^k - \check{y}^k = \psi(\hat{x}^k, \hat{y}^k) \xrightarrow{K} 0$ (cf. Theorem 5.2(4)), where $\check{y}^k \geq 0$ by (6.2) with $\hat{x}^k \in X$ (Lemma 5.1). Consequently, $y^* \in [0, c)$ gives $\check{y}^k \in [0, c)$ for all large $k \in K$. Since each function f_j in (6.1a) is continuous on $\text{dom } f_j \supset [0, c_j)$, we have $f(\check{y}^k) \xrightarrow{K} f(y^*) = f_*$. The conclusion follows from Theorem 5.2(4) with $\theta_* = \varphi_* = -f_*$. \square

An extension to the case where some arc costs are linear follows.

Proposition 6.2 *Let problem (6.1) be feasible. Suppose that the first \check{m} components of any optimal total flow y^* are unique (e.g., f_j are strictly convex on $\mathbb{R}_+^m \cap \text{dom } f_j$ for $j \leq \check{m}$) and satisfy $y_j^* \in [0, c_j) \subset \text{dom } f_j$ for some $c_j > 0$, whereas the costs f_j are linear for $j > \check{m}$. Further, let $\varepsilon_\pi = 0$ (i.e., the oracle is exact), and let $K \subset \mathbb{N}$ be a subsequence such that $V_k \xrightarrow{K} 0$. Then $\check{y}_j^k \xrightarrow{K} y_j^*$ for $j \leq \check{m}$, $f(\check{y}^k) \xrightarrow{K} f_* = -\theta_*$ and $f(\check{y}^k) + \theta_u^k \xrightarrow{K} 0$.*

Proof The proof of Proposition 6.1 gives $\hat{y}_j^k, \check{y}_j^k \xrightarrow{K} y_j^*$ and $f_j(\hat{y}_j^k), f_j(\check{y}_j^k) \xrightarrow{K} f_j(y_j^*)$ for $j \leq \check{m}$, since $\hat{y}_j^k \in \text{dom } f$ by (5.6). For $j > \check{m}$, $f_j(y_j) = \alpha_j y_j$ for some $\alpha_j \in \mathbb{R}$;

thus $\sigma_j(u_j) := f_j^*(u_j) = i_{\{\alpha_j\}}(u_j)$. Then $u_j^{k+1} = \hat{u}_j^k = \alpha_j$ in (1.6) yields $p_j^k = 0$ in (2.8), so $\psi_j(\hat{x}^k, \hat{y}^k) = 0$ by Lemma 5.1; since $\hat{y}^k - \check{y}^k = \psi(\hat{x}^k, \hat{y}^k)$, we have $\hat{y}_j^k = \check{y}_j^k$ for $j > \check{m}$. Therefore, by (6.1a), $f(\check{y}^k) = f(\hat{y}^k) + \sum_{j \leq \check{m}} [f_j(\check{y}_j^k) - f_j(\hat{y}_j^k)]$, where the sum vanishes as $k \xrightarrow{K} \infty$; Theorem 5.2(4) with $\varphi := -f$ gives the conclusion. \square

6.3 Specific arc costs

For specific arc costs, as in [1, 14], we shall consider Kleinrock’s average delays

$$f_j(y_j) := \begin{cases} \infty & \text{if } y_j \geq c_j, \\ y_j/(c_j - y_j) & \text{if } y_j \in [0, c_j), \\ y_j/c_j & \text{if } y_j < 0, \end{cases} \tag{6.3a}$$

$$f_j^*(u_j) := \begin{cases} (\sqrt{c_j u_j} - 1)^2 & \text{if } u_j \geq 1/c_j, \\ \infty & \text{if } u_j < 1/c_j, \end{cases} \tag{6.3b}$$

with arc capacities $c_j > 0$, the BPR (Bureau of Public Roads) nonlinear delays

$$f_j(y_j) := \begin{cases} \alpha_j y_j + \beta_j y_j^{\gamma_j} & \text{if } y_j \geq 0, \\ \alpha_j y_j & \text{if } y_j < 0, \end{cases} \tag{6.4a}$$

$$f_j^*(u_j) := \begin{cases} \frac{\gamma_j - 1}{\gamma_j} (u_j - \alpha_j)^{\gamma_j/(\gamma_j - 1)} / (\beta_j \gamma_j)^{1/(\gamma_j - 1)} & \text{if } u_j \geq \alpha_j, \\ \infty & \text{if } u_j < \alpha_j, \end{cases} \tag{6.4b}$$

with parameters $\alpha_j \geq 0, \beta_j > 0, \gamma_j \geq 2$, as well as BPR linear delays with $\alpha_j \geq 0$:

$$f_j(y_j) := \alpha_j y_j \quad \text{for all } y_j, \tag{6.5a}$$

$$f_j^*(u_j) := \begin{cases} 0 & \text{if } u_j = \alpha_j, \\ \infty & \text{if } u_j \neq \alpha_j. \end{cases} \tag{6.5b}$$

Our costs are linearly extrapolated versions of the “standard” costs used in [14], where $f_j(y_j)$ is set to ∞ for $y_j < 0$, so that $f_j^*(u_j)$ becomes 0 instead of ∞ for $u_j < f_j'(0)$. Note that the value of f_j at $y_j < 0$ does not matter for (6.1), where the constraints yield $y_j \geq 0$. Further, if (6.1) is feasible, the assumptions of Propositions 6.1 and 6.2 hold for our Kleinrock and nonlinear BPR costs, and for a mixture of our nonlinear and linear BPR costs, respectively. Finally, since $\text{dom } \sigma = \text{dom } f^* \subset \mathbb{R}_+^m$ for our costs, the oracle has to solve shortest path problems with nonnegative arc lengths u^k only; hence, we may assume that $\varepsilon_\pi = 0$.

6.4 Solving the σ -subproblem for specific arc costs

We now specialize the results of Sect. 4.2 with $\sigma^* := f$ for the costs of Sect. 6.3. Since σ^* is separable, we may handle (4.9) by solving m one-dimensional subproblems to

determine components of an approximate solution, say \tilde{z} . Thus we need a stopping criterion for each subproblem. To this end, we replace the criterion $\varepsilon(z^i) \leq \kappa_N v(z^i)$ by $\varepsilon(\tilde{z}) \leq \kappa_N \bar{v}(\tilde{z})$ for

$$\bar{v}(z) := \sigma_{\hat{u}}^k - \bar{\sigma}(\hat{u}^k; z) + t_k |p_{\pi}^k + z|^2 = v(z) - \left[\pi_{\hat{u}}^k - \bar{\pi}_k(\hat{u}^k) \right], \tag{6.6}$$

where the second equality follows from (4.5), (4.6) and (4.8) with $\theta_{\hat{u}}^k = \sigma_{\hat{u}}^k + \pi_{\hat{u}}^k$. Moreover, $\sigma_{\hat{u}}^k - \bar{\sigma}(\hat{u}^k; z) \geq 0$ yields $\bar{v}(z) \geq 0$, whereas by the results of Sect. 4.2, $\bar{v}(z) = 0$ only if $z = z^* = -p_{\pi}^k$; since checking if $\bar{v}(-p_{\pi}^k) = 0$ is easy, we may assume that $\bar{v}(z^*) > 0$. Finally, $\bar{v}(z) \leq v(z)$ from $\varepsilon_{\pi} = 0$. The resulting ‘‘natural’’ subproblem criteria are discussed below.

To simplify notation, we assume $m = 1$, drop the subscript j in (6.3)–(6.5) and let $t := t_k$ in (4.5). We first consider the Kleinrock and nonlinear BPR costs in (6.3)–(6.4). For finding an approximate solution \tilde{z} , we exploit the following properties:

- $f(z) = f'(0)z$ for $z \leq 0$ with $f'(0) \geq 0$;
- $f''(z) > 0$ for $z > 0$ in $F := \text{dom } f = (-\infty, c)$, with $c := \infty$ in the BPR case;
- $\sigma^* = f$ is continuous on F with $\text{dom } \partial\sigma^* = F$;
- $\sigma := f^*$ is continuous on $\text{dom } \sigma = [f'(0), \infty)$ with $\text{dom } \partial\sigma = \text{dom } \sigma$;
- $w'(z) = f'(z) - \bar{u}(z)$ and $w''(z) = f''(z) + t$ for $z \in F$ in (4.9) by (4.5).

If $w'(0) \geq 0$, then $\tilde{z} := -w'(0)/t$ is optimal ($w'(\tilde{z}) = 0$), $\varepsilon(\tilde{z}) = 0$ and $\bar{u}(\tilde{z}) = f'(0)$.

If $w'(0) < 0$, then $z^* \in (0, -w'(0)/t)$, since for $z \geq -w'(0)/t$, $f'(z) > f'(0)$ yields

$$w'(z) = f'(z) - \bar{u}(z) > f'(0) - \bar{u}(z) = w'(0) + tz \geq 0.$$

Further, $z^* \in (0, z^{\text{up}})$ for $z^{\text{up}} := \min\{-w'(0)/t, c\}$ from $z^* \in F$, and $\bar{u}(z) \in \text{dom } \sigma$ for $z \in (0, z^{\text{up}})$, since $\bar{u}(z) > f'(0)$ if $z < -w'(0)/t$. These properties and the results of Sect. 4.2 yield the following. Suppose we minimize w over $(0, z^{\text{up}})$ via a descent method, starting from $z^1 := p_{\sigma}^{k-1}$ if $p_{\sigma}^{k-1} \in (0, z^{\text{up}})$ or any $z^1 \in (0, z^{\text{up}})$ otherwise, which generates points $z^i \in (0, z^{\text{up}})$ such that $z^i \rightarrow z^*$. Then $\varepsilon(z^i) \rightarrow 0$ and $\bar{v}(z^i) \rightarrow \bar{v}(z^*) > 0$ in (6.6) imply that we will eventually have $\varepsilon(z^i) \leq \kappa_N \bar{v}(z^i)$, in which case the method may stop with $\tilde{z} := z^i$.

Next, for the linear BPR costs in (6.5) with $w'(z) = f'(0) - \bar{u}(z)$, $\tilde{z} := -w'(0)/t$ is optimal ($w'(\tilde{z}) = 0$), $\varepsilon(\tilde{z}) = 0$ and $\bar{u}(\tilde{z}) = f'(0)$ (as in the case of $w'(0) \geq 0$ above).

For $m > 1$, expressing $\varepsilon(z)$ in (4.7), $w(z)$ in (4.9) and $\bar{v}(z)$ in (6.6) as sums of $\varepsilon_j(z_j)$, $w_j(z_j)$ and $\bar{v}_j(z_j)$ respectively over $j = 1, \dots, m$, for each j we may find \tilde{z}_j as above so that $\varepsilon_j(\tilde{z}_j) \leq \kappa_N \bar{v}_j(\tilde{z}_j)$, and $w(\tilde{z}) \leq w(p_{\sigma}^{k-1})$; since $\bar{v}(z) \leq v(z)$ in (6.6), we also have $\varepsilon(\tilde{z}) \leq \kappa_N v(\tilde{z})$. Thus, as in Sect. 4.2, we may set $u^{k+1} := \bar{u}(\tilde{z})$, $v_k := v(\tilde{z})$, $\bar{\sigma}_k(\cdot) := \bar{\sigma}(\cdot; \tilde{z})$ and $p_{\sigma}^k := \tilde{z}$.

7 Implementation issues

We now describe the main issues in our implementation of each step of Algorithm 2.1 for the network applications of Sect. 6. We also highlight aspects where our implementation could be improved; this is left for future work.

7.1 Initial settings

In the Kleinrock case of (6.3), the initial $u_j^1 := (1 - \rho_*)^{-2}/c_j$ for all j , with $\rho_* := \frac{1}{4}$ estimating the maximum traffic intensity $\max_j y_j^*/c_j$ as in [5, 14]; then $p_\sigma^0 := \nabla\sigma(u^1)$. In the BPR case of (6.4)–(6.5), $u_j^1 := \alpha_j$ for all j , and we let $p_\sigma^0 := 0$.

As usual in bundle methods, we use the descent parameter $\kappa = 0.1$ in (2.4). We set the initial stepsize to $t_1 := 1$, corresponding to the inverse of the initial proximal coefficient of [1], and let $t_{\min} := 10^{-20}t_1$.

7.2 Subproblem solution

For the models $\check{\pi}_k$ of (5.3), subproblem (1.5) is solved by the QP routine of [10]. This routine has at least two drawbacks. First, being designed for bound-constrained problems, it employs data structures that are not efficient in the unconstrained case. Second, its linear algebra is behind the current state of the art; it could benefit from tuned versions of LAPACK like the MATLAB implementation of [1].

The one-dimensional subproblems of Sect. 6.4 are solved for the tolerance $\kappa_N = 10^{-3}$ by Newton's method with Armijo's backtracks for a descent tolerance of 10^{-6} , where at each iteration the initial unit stepsize is reduced if necessary to 0.9 times the maximum feasible stepsize, and the stepsize is divided by 2 for each Armijo's failure. This works quite well, but implementations based on self-concordant ideas (as in [1]) could be more efficient.

The looping Step 5' of Sect. 4.1 employs the tolerance $\check{\kappa} = 0.2$, but the number of loops at any iteration is limited to 30.

7.3 Shortest-path oracle

Let $S \leq n$ be the number of common sources (different source nodes) in (6.1). To evaluate $\pi(u^{k+1})$, we call S times subroutine L2QUE of [4], which finds shortest paths from a given source to all other nodes. We chose L2QUE simply because it performed well in our earlier work [14]; most probably, faster routines exist.

7.4 Termination criterion

In view of Sect. 6.2, we stop when the relative optimality gap is small enough:

$$\gamma_{\text{rel}}^k := \left(f_{\text{up}}^k - f_{\text{low}}^k \right) / \max\{ f_{\text{low}}^k, 1 \} \leq \varepsilon_{\text{opt}}, \quad (7.1)$$

where $\varepsilon_{\text{opt}} = 10^{-5}$ as in [1], whereas f_{up}^k and f_{low}^k are the best upper and lower bounds on f_* obtained so far. Specifically, $f_{\text{low}}^k := -\min_{j \leq k+1} \theta_u^j$, whereas f_{up}^k is the minimum of $f(\check{y}^j)$ over iterations $j \leq k$, $j = 10, 20, \dots$, at which $f(\check{y}^j)$ is computed. A more frequent computation of $f(\check{y}^j)$ could save work on small instances.

7.5 Step size updating

Our implementation of Step 8 uses the following procedure, in which γ_{rel}^k is the gap of (7.1), $\gamma_k := f_{\text{up}}^k - f_{\text{low}}^k$ is the absolute gap, l_k is the number of loops made on iteration k , and n_k counts descent or null steps since the latest change of t_k , with $n_1 := 1$.

Procedure 7.6 (Step size updating)

- (1) Set $t_k + 1 := t_k$.
- (2) If $\hat{u}^{k+1} = \hat{u}^k$ or $l_k > 0$ go to (5).
- (3) If $n_k \geq 10$, or $v_k < \gamma_k/2$ and $\gamma_{\text{rel}}^k \leq 0.01$, set $t_{k+1} := 2t_k$.
- (4) Set $n_{k+1} := \max\{n_k + 1, 1\}$. If $t_{k+1} \neq t_k$, set $n_{k+1} := 1$. Exit.
- (5) If $i_t^{k+1} = 0$, $n_k \leq -10$, and either $v_k > \gamma_k/2$ or $\gamma_{\text{rel}}^k > 0.01$, set $t_{k+1} := \max\{t_k/5, t_{\min}\}$. Set $n_{k+1} := \min\{n_k - 1, -1\}$. If $t_{k+1} \neq t_k$, set $n_{k+1} := -1$. Exit.

The counter n_k introduces some inertia, which smooths out the step size updating. In general, t_k should be increased (respectively decreased) if “too many” descent (respectively null) steps are occurring, but v_k should be of order γ_k , since descent steps with $v_k \ll \gamma_k$ bring little. Of course, our procedure is just an example and there is still room for improvement.

8 Numerical illustrations

To get a feeling for the practical merits and drawbacks of our approach, we first benchmark our AL implementation on the test problems of [1].

8.1 Test problems of Babonneau and Vial

We used the four sets of test problems of [1]. Their features are given in Table 1, where N is the number of nodes, m is the number of arcs, n is the number of commodities, S is the number of common sources, and $f_*^{\text{Kleinrock}}$ and f_*^{BPR} are the optimal values of (6.1) for the Kleinrock and BPR costs respectively, with relative optimality gaps of at most 10^{-5} . Table 1 corrects some values of [1, Table 2]; see [2] and below.

For the first two sets of planar and grid problems¹, the cost functions are generated as in [1, Sect. 8.1]; we add that problem planar150 is missing in [1].

The third set of telecommunication problems includes a corrected version of problem ndo22 [2]; the BPR costs are generated as in [1].

¹ Available at <http://www.di.unipi.it/di/groups/optimize/Data/MMCF.html>.

Table 1 Test problems of Babonneau and Vial

Problem	N	m	n	S	$f_{*}^{\text{Kleinrock}}$	f_{*}^{BPR}
<i>Planar problems</i>						
planar30	30	150	92	29	40.5668	4.44549×10^7
planar50	50	250	267	50	109.478	1.21236×10^8
planar80	80	440	543	80	232.321	1.81906×10^8
planar100	100	532	1,085	100	226.299	2.29114×10^8
planar150	150	850	2,239	150	715.309	5.27985×10^8
planar300	300	1,680	3,584	300	329.120	6.90748×10^8
planar500	500	2,842	3,525	500	196.394	4.83309×10^9
planar800	800	4,388	12,756	800	354.008	1.16952×10^9
planar1,000	1,000	5,200	20,026	1,000	1,250.92	3.41859×10^9
planar2,500	2,500	12,990	81,430	2,500	3,289.05	1.23827×10^{10}
<i>Grid problems</i>						
grid1	25	80	50	23	66.4002	8.33599×10^5
grid2	25	80	100	25	194.512	1.72689×10^6
grid3	100	360	50	40	84.5618	1.53241×10^6
grid4	100	360	100	63	171.331	3.05543×10^6
grid5	225	840	100	83	236.699	5.07921×10^6
grid6	225	840	200	135	652.877	1.05075×10^7
grid7	400	1,520	400	247	776.566	2.60669×10^7
grid8	625	2,400	500	343	1,542.15	4.21240×10^7
grid9	625	2,400	1,000	495	2,199.83	8.36394×10^7
grid10	625	2,400	2,000	593	2,212.89	1.66084×10^8
grid11	625	2,400	4,000	625	1,502.75	3.32475×10^8
grid12	900	3,480	6,000	899	1,478.93	5.81488×10^8
grid13	900	3,480	12,000	900	1,760.53	1.16933×10^9
grid14	1,225	4,760	16,000	1,225	1,414.39	1.81297×10^9
grid15	1,225	4,760	32,000	1,225	1,544.15	3.61568×10^9
<i>Telecommunication problems</i>						
ndo22	14	22	23	5	103.412	1.86767×10^3
ndo148	61	148	122	61	151.926	1.40233×10^5
904	106	904	11,130	106	33.4931	1.29197×10^7
<i>Transportation problems</i>						
Sioux-Falls	24	76	528	24	600.679	4.23133×10^6
Winnipeg	1,067	2,836	4,344	135	1,527.41	8.25673×10^5
Barcelona	1,020	2,522	7,922	97	845.872	1.22856×10^6
Chicago-sketch	933	2,950	93,135	386	614.726	1.67484×10^7
Chicago-region	12,982	39,018	2,296,227	1,771	3,290.49	2.58457×10^7
Philadelphia	13,389	40,003	1,149,795	1,489	2,557.42	2.24926×10^8

The fourth set of transportation problems² uses original BPR costs, and Kleinrock costs generated as in [1]. To clarify the description of [1], we add that in the Kleinrock case the demands are divided by 2 for Sioux-Falls, 2000 for Winnipeg, 5100 for Barcelona, 2.5 for Chicago-sketch, 6 for Chicago-region, and 7 for Philadelphia. We also observe that although [1, Table 2] gives wrong Kleinrock values for Chicago-sketch, Chicago-region and Philadelphia, their entries in [1, Table 5] are apparently correct. In contrast, for the BPR versions of Barcelona and Philadelphia, [1, Table 6] must be corrected as in [2].

8.2 Numerical results for the test problems of Babonneau and Vial

Tables 2 and 3 give our results for the problems of Sect. 8.1. In these tables,

- k and l are the numbers of iterations and descent steps respectively;
- Σ is the average number of subproblems solved at Step 3 per iteration;
- N is the average number of Newton's iterations for the one-dimensional subproblems solved approximately at Step 3 (cf. Sect. 7.2);
- CPU is the total CPU time in seconds;
- $\%Si$ is the percentage of CPU time spent on the subproblems of Step 3;
- $\%Or$ is the percentage of CPU time spent in the shortest-path oracle.

We used a Dell M60 notebook (Pentium M 755 2 GHz, 1.5 GB RAM) under MS Windows XP and Fortran 77, with SPECint2000 of 1541 and SPECfp2000 of 1088. Our machine was comparable with that of [1] (Pentium 4 2.8 GHz, 2 GB RAM, with SPECint2000 of 1254 and SPECfp2000 of 1327). Yet we refrain from comparing the CPU times, as they could depend on many other factors. Here our main message is that AL can solve all the instances of [1] in reasonable time.

Table 4 gives our results for standard bundle (cf. Sect. 8.3). In this table,

- $T_{SB/AL}$ is the ratio of the CPU times of standard bundle (SB for short) and AL, with the times increased to 0.1 if necessary.

Moreover, to avoid too long run times, we imposed an iteration limit of 9,999 (thus $T_{SB/AL}$ does not mean much for runs with $k = 9,999$), and skipped some largest instances. AL is significantly faster than SB on all but the smallest instances.

8.3 Numerical comparisons with disaggregate bundle

For comparing AL with SB and the method of [17] we also used the small and medium sized test problems of [17] with Kleinrock costs. Their features are given in Table 5; problems p1 and p4 are called ndo22 and ndo148 in Table 1.

Standard bundle replaces (1.5)–(1.6) by the single subproblem

$$u^{k+1} := \arg \min \left\{ \check{\sigma}_k(u) + \check{\tau}_k(u) + \frac{1}{2l_k} |u - \hat{u}^k|^2 : u \in C \right\}, \quad (8.1)$$

² Available at <http://www.bgu.ac.il/bargera/tntp/>.

Table 2 Performance of AL for Kleinrock costs

Problem	k	l	Sigma	Newton	CPU	%Si	%Or
planar30	125	62	4.7	1.9	0.1	60	0
planar50	214	73	3.2	2.2	0.2	31	10
planar80	308	80	3.0	2.2	0.6	28	28
planar100	312	75	3.9	2.4	0.8	24	28
planar150	979	95	1.7	2.1	12.2	3	17
planar300	303	84	6.4	2.7	4.7	27	46
planar500	253	77	8.3	2.6	9.7	23	55
planar800	341	82	7.7	2.7	28.1	16	69
planar1000	648	104	4.1	3.0	74.8	8	73
planar2500	1,530	103	2.5	2.6	1,092.1	2	86
grid1	92	65	8.2	2.3	0.1	40	0
grid2	185	62	2.9	2.4	0.0	50	50
grid3	222	74	6.7	2.2	0.4	37	13
grid4	247	79	5.3	2.7	0.4	45	30
grid5	290	82	5.5	2.3	1.3	35	19
grid6	453	89	2.9	2.5	2.4	15	28
grid7	646	98	3.0	2.4	8.4	12	33
grid8	940	102	2.1	2.3	21.0	7	41
grid9	900	99	2.2	2.4	24.4	7	48
grid10	730	100	2.8	2.7	22.1	9	54
grid11	424	85	5.6	3.3	14.1	18	52
grid12	458	96	5.8	3.4	27.0	17	59
grid13	423	94	6.4	3.7	26.1	18	58
grid14	470	106	7.1	3.9	49.5	17	63
grid15	451	102	7.7	4.1	49.7	19	62
ndo22	374	290	9.5	1.9	0.1	38	0
ndo148	91	56	2.7	2.0	0.0	33	0
904	216	57	8.3	3.0	1.5	45	16
Sioux-Falls	300	61	2.8	2.5	0.1	11	11
Winnipeg	1,149	303	4.6	1.8	104.4	4	11
Barcelona	3044	314	5.4	1.7	397.6	3	6
Chicago-sketch	280	80	8.8	2.4	13.3	19	62
Chicago-region	303	73	7.7	2.1	901.0	4	88
Philadelphia	433	89	8.4	3.2	1431.3	5	85

where $\check{\sigma}_k \leq \sigma$ is a polyhedral approximation built from linearizations obtained from a first-order oracle for σ similarly to $\check{\pi}_k \leq \pi$. Since $\sigma(u) = \sum_{j=1}^m f_j^*(u_j)$ for f_j^* given by (6.3), constructing an exact first-order oracle for σ is simple. Further, given an integer $n_\sigma \leq m$, we may treat σ as the sum of n_σ functions, say $\sigma = \sum_{i=1}^{n_\sigma} \sigma_i$, where $\sigma_1(u) = \sum_{j=1}^{\lfloor m/n_\sigma \rfloor} f_j^*(u_j)$, etc., using a richer model $\check{\sigma}_k := \sum_{i=1}^{n_\sigma} \check{\sigma}_{ik}$ in (8.1), where each $\check{\sigma}_{ik} \leq \sigma_i$ stems from past linearizations delivered by an oracle for σ_i . Of course, richer models may speed up convergence, but the QP work in solving (8.1) may grow.

Table 3 Performance of AL for BPR costs

Problem	k	l	Sigma	Newton	CPU	%Si	%Or
planar30	75	69	1.3	1.1	0.0	66	33
planar50	105	64	1.4	1.3	0.0	66	33
planar80	150	59	1.1	1.3	0.2	8	73
planar100	108	44	1.4	1.3	0.2	20	54
planar150	194	52	1.1	1.5	0.9	12	67
planar300	97	31	1.3	1.2	1.4	8	86
planar500	50	23	1.7	1.0	3.3	4	92
planar800	108	33	1.9	1.2	25.4	2	94
planar1000	209	41	1.4	1.3	32.6	2	88
planar2500	264	52	1.3	1.6	411.8	0	97
grid1	48	29	3.6	2.2	0.0	100	0
grid2	61	27	1.7	2.2	0.0	100	0
grid3	43	23	2.5	1.3	0.0	25	50
grid4	59	26	1.8	2.2	0.1	77	11
grid5	86	28	2.1	1.7	0.3	38	38
grid6	150	33	2.0	2.0	0.6	44	33
grid7	108	31	2.1	2.3	1.0	34	50
grid8	143	36	1.6	2.3	2.3	25	56
grid9	183	37	1.7	2.4	4.0	16	62
grid10	200	34	2.3	2.5	5.5	19	59
grid11	120	32	4.2	3.2	4.1	36	49
grid12	122	31	5.8	3.4	8.8	38	48
grid13	140	30	5.5	3.6	10.1	39	50
grid14	111	28	8.0	4.0	15.9	43	46
grid15	115	26	8.0	4.3	16.9	44	47
ndo22	11	8	2.2	2.2	0.0	0	0
ndo148	14	11	2.4	2.1	0.0	0	100
904	116	32	1.2	2.8	0.5	32	57
Sioux-Falls	105	37	6.3	2.6	0.1	85	0
Winnipeg	127	31	8.4	1.8	4.5	51	39
Barcelona	92	24	14.3	3.0	5.6	74	18
Chicago-sketch	129	32	7.0	2.2	7.2	34	57
Chicago-region	300	51	3.6	2.6	891.0	5	89
Philadelphia	671	62	2.7	1.9	3,239.7	2	94

Since our AL is implemented on top of SB, they share the same QP routine, primal recovery, etc. In particular, SB uses the descent test (2.4) with $\kappa = 0.1$ and $v_k := \theta_{\tilde{u}}^k - [\check{\sigma}_k(u^{k+1}) + \check{\pi}_k(u^{k+1})]$, and the stopping criterion (7.1) with $\varepsilon_{\text{opt}} = 10^{-5}$.

The *Newton-cutting-plane* (NCP for short) method of [17] replaces (8.1) by

$$\tilde{u}^{k+1} := \arg \min \left\{ \bar{\sigma}_k(u) + \check{\pi}_k(u) + \frac{1}{2} |u - \hat{u}^k|_{H_k}^2 : u \in C \right\}, \quad (8.2)$$

Table 4 Performance of standard bundle

Problem	Kleinrock costs				BPR costs			
	k	CPU	%Or	$T_{SB/AL}$	k	CPU	%Or	$T_{SB/AL}$
planar30	333	0.6	0	5.6	291	0.2	4	2.2
planar50	553	2.0	7	9.9	103	0.3	11	2.7
planar80	1,498	5.9	17	9.8	218	1.8	12	9.2
planar100	2,210	10.6	19	13.3	113	0.9	14	4.5
planar150	3,435	54.3	15	4.5	890	38.1	8	42.3
planar300	3,870	153.1	18	32.6	234	37.6	7	26.9
planar500	4,613	1,183.2	8	122.0	151	80.3	11	24.3
planar800	5,630	2,132.3	14	75.9	307	569.7	11	22.4
planar1000	9,999 ^a	2,709.8	31	36.2	578	1,365.4	5	41.9
planar2500	9,999 ^a	13,399.5	46	12.3	–	–	–	–
grid1	184	0.1	0	1.0	50	0.0	0	1.0
grid2	134	0.1	0	1.0	56	0.0	0	1.0
grid3	1,150	5.7	4	14.3	59	0.2	6	1.6
grid4	1,189	3.8	10	9.6	60	0.2	4	2.4
grid5	1,729	32.1	6	24.7	72	1.0	9	3.2
grid6	2,127	30.8	11	12.8	105	1.7	6	2.9
grid7	4,712	226.7	9	27.0	85	4.9	7	4.9
grid8	9,691	1,502.6	6	71.6	119	25.0	4	10.9
grid9	6,705	965.6	9	39.6	139	34.2	5	8.6
grid10	9,999 ^a	1,383.1	12	62.6	135	39.3	5	7.1
grid11	5,008	700.4	12	49.7	94	32.3	5	7.9
grid12	9,999 ^a	2,782.7	13	103.1	81	74.5	3	8.5
grid13	6,983	2,061.8	12	79.0	91	85.9	3	8.5
grid14	8,582	5,157.5	11	104.2	75	169.7	2	10.7
grid15	9,999 ^a	6,049.1	11	121.7	73	162.0	3	9.6
ndo22	727	0.2	0	1.6	22	0.0	100	1.0
ndo148	218	0.2	17	2.3	36	0.0	0	1.0
904	2,030	87.5	3	58.3	1,504	285.9	1	571.8
Sioux-Falls	860	0.3	11	3.5	117	0.1	9	1.0
Winnipeg	9,999 ^a	1,247.9	8	12.0	443	140.9	4	31.3
Barcelona	9,999 ^a	1,031.7	8	2.6	2,743	2,541.3	1	453.8
Chicago-sketch	9,999 ^a	4,971.6	6	373.8	490	217.9	7	30.3

^a Failure to obtain required accuracy

where $\bar{\sigma}_k(\cdot) := \sigma(\hat{u}^k) + \langle \sigma'(\hat{u}^k), \cdot - \hat{u}^k \rangle$ is the linearization of σ at \hat{u}^k and $|\cdot|_{H_k} := \langle H_k \cdot, \cdot \rangle^{1/2}$ is the norm generated by a symmetric positive definite matrix H_k which approximates the Hessian $\sigma''(\hat{u}^k)$. Exploiting the structure of $\pi = \sum_{i=1}^n \pi_i$ with $\pi_i(\cdot) = -\min_{x_i \in X_i} \langle \cdot, x_i \rangle$, NCP employs the disaggregated model $\tilde{\pi}_k := \sum_{i=1}^n \tilde{\pi}_{ik}$ with $\tilde{\pi}_{ik}(\cdot) = \max_{j=1}^k \langle g_{\pi_i}^j, \cdot \rangle$ and $g_{\pi_i}^j \in \partial \pi_i(u^j)$. For the search direction $d^k := \bar{u}^{k+1} - \hat{u}^k$, a backtracking search finds a stepsize $t_k \in (0, 1]$ and a point $u^{k+1} :=$

Table 5 Test problems of Lemaréchal et al.

Problem	N	m	n	S	$f_*^{\text{Kleinrock}}$
p1	14	22	23	5	103.4120
p2	19	68	30	15	8.994992
p3	60	280	100	48	53.08077
p4	61	148	122	61	151.9269
p5	20	64	133	20	39.63546
p6	122	332	162	45	276.3214
p7	100	600	200	88	84.96748
p8	30	72	335	20	36.45172
p9	21	68	420	21	68.83896
p10	100	800	500	99	139.0965
p11	67	170	761	20	109.8956
p12	34	160	946	34	19.56668
p13	300	2,000	1,000	293	304.3895
p14	48	198	1,583	47	135.4632
p15	81	188	2,310	66	41.79184
p16	122	342	2,881	102	242.7148

$\hat{u}^k + t_k d^k$ such that either $\hat{u}^{k+1} := u^{k+1}$ if θ is reduced significantly or $\hat{u}^{k+1} := \hat{u}^k$; see [17] for details.

Table 6 gives the AL and NCP results for the problems of Table 5. In this table,

- $\#Or$ is the number of oracle calls made by NCP from [17, Table 1];
- $T_{\text{NCP/AL}}$ is the ratio of the CPU times of NCP from [17, Table 1] and our AL, with our times increased to 0.01 if necessary.

As for CPU comparisons, [17] used a desktop PC (Xeon 2.4 GHz 2 cores, 1.5 GB RAM) under Linux, CPLEX 10.0 for solving QPs and C for the shortest path computation via Dijkstra's method, with SPECint2000 of 2564 and SPECfp2000 of 2522. Thus our machine was about twice slower, but the QP and shortest-path solvers were different. In CPU times, AL is substantially faster than NCP on most instances. Here two points should be noted. First, NCP's CPU times would probably change substantially with the use of a specialized QP solver such as [3, 8, 10]. Second, without implementing primal recovery, NCP had to rely on an "artificial" stopping criterion instead of (7.1), possibly spending more work than necessary to meet (7.1) with $\varepsilon_{\text{opt}} = 10^{-5}$.

Table 7 reports the SB results for several values of the disaggregation parameter n_σ (the oracle percentages $\%Or$ were marginal: at most 16 for $n_\sigma = 1$, and 3 for $n_\sigma = 20$). Clearly, AL is much faster than SB in CPU times. This is mostly due to SB spending more time on its QP subproblems, since the iteration counts do not increase so much except for problems p15 and p16. Note that increasing n_σ may help for some problems (e.g., p15 and p16), but not for others (e.g., p13).

The interested readers might compare our results with those given in [17] for two other standard bundle variants using $n_\sigma = m$ or 1, as well as full disaggregation for π just like NCP; neither variant was competitive with NCP.

Table 6 Performance of AL and NCP for small and medium Kleinrock problems

Pb	AL							NCP			
	k	l	Sigma	Newton	CPU	%Si	%Or	k	#Or	CPU	$T_{NC/AL}$
p1	374	290	9.5	1.9	0.06	33	0	12	100	0.03	0.5
p2	40	34	2.5	1.7	0.00	0	0	5	11	0.02	2.0
p3	92	57	2.3	1.2	0.11	0	54	7	102	0.20	1.8
p4	91	56	2.7	2.0	0.10	49	10	7	15	1.07	10.7
p5	88	54	5.1	2.1	0.07	28	14	7	15	0.35	5.0
p6	139	99	1.7	1.4	0.11	36	18	9	99	0.61	5.5
p7	92	56	4.0	1.1	0.16	6	43	7	103	0.78	4.9
p8	104	41	20.6	1.8	0.10	29	0	7	111	0.13	1.3
p9	112	59	13.9	2.7	0.06	66	0	7	15	5.62	93.7
p10	174	65	4.1	0.9	0.44	20	38	10	314	9.60	21.8
p11	86	57	3.3	2.0	0.05	79	0	8	141	4.84	96.8
p12	83	47	10.1	1.6	0.06	66	33	5	11	1.03	17.2
p13	208	65	4.6	1.1	2.61	13	59	11	330	73.37	28.1
p14	167	67	3.1	1.9	0.14	64	14	9	89	13.09	93.5
p15	119	37	22.8	1.3	0.24	33	16	3	7	2.44	10.2
p16	310	211	4.8	1.3	0.55	20	41	9	82	311.85	567.0

Table 7 Performance of disaggregate bundle for small and medium Kleinrock problems

Pb	$n_\sigma = 1$		$n_\sigma = 3$		$n_\sigma = 5$		$n_\sigma = 10$		$n_\sigma = 20$	
	k	CPU	k	CPU	k	CPU	k	CPU	k	CPU
p1	727	0.1	71	0.0	47	0.0	33	0.0	21	0.0
p2	118	0.1	96	0.1	112	0.2	87	0.2	70	0.2
p3	180	0.5	235	1.4	224	2.3	233	2.9	181	3.2
p4	218	0.2	169	0.4	150	0.5	166	1.0	136	1.4
p5	149	0.1	197	0.3	166	0.3	132	0.4	105	0.3
p6	334	0.9	302	1.5	320	2.0	330	2.8	261	3.6
p7	294	2.4	305	5.0	298	11.4	315	13.1	275	12.7
p8	546	0.4	244	0.3	149	0.3	80	0.3	84	0.3
p9	276	0.1	265	0.3	238	0.4	185	0.6	161	0.6
p10	390	6.4	378	15.0	471	22.5	386	27.4	399	38.3
p11	147	0.2	180	0.7	172	1.0	163	1.5	129	2.0
p12	386	0.6	436	1.7	320	1.9	216	1.9	159	1.7
p13	479	36.3	567	73.2	588	101.7	507	264.3	673	380.8
p14	262	0.6	272	1.1	269	1.3	307	1.9	277	2.8
p15	5,610	31.4	4,962	57.8	3,153	41.2	1,320	22.0	501	10.4
p16	3,282	21.0	2,559	31.5	1,010	21.3	1,424	31.3	520	18.4

Acknowledgments I would like to thank F. Babonneau, H. Bar-Gera and J.-P. Vial for numerous discussions and help with the test problems. An associate editor and two referees helped in improving the presentation of numerical results.

References

1. Babonneau, F., Vial, J.P.: ACCPM with a nonlinear constraint and an active set strategy to solve nonlinear multicommodity flow problems. *Math. Program.* **120**, 179–210 (2009)
2. Babonneau, F., Vial, J.P.: ACCPM with a nonlinear constraint and an active set strategy to solve nonlinear multicommodity flow problems: A corrigendum. *Math. Program.* **120**, 211–212 (2009)
3. Frangioni, A.: Solving semidefinite quadratic problems within nonsmooth optimization algorithms. *Comput. Oper. Res.* **23**, 1099–1118 (1996)
4. Gallo, G., Pallottino, S.: Shortest path algorithms. *Ann. Oper. Res.* **13**, 3–79 (1988)
5. Goffin, J.L.: The ellipsoid method and its predecessors. In: Contesse, L., Correa, R., Weintraub, A. *Recent Advances in System Modelling and Optimization*, Lecture Notes in Control and Information Sciences 87, pp. 127–141. Springer, Berlin (1987)
6. Goffin, J.L., Gondzio, J., Sarkissian, R., Vial, J.P.: Solving nonlinear multicommodity flow problems by the analytic center cutting plane method. *Math. Program.* **76**, 131–154 (1996)
7. Hiriart-Urruty, J.B., Lemaréchal, C.: *Convex Analysis and Minimization Algorithms*. Springer, Berlin (1993)
8. Kiwiel, K.C.: A method for solving certain quadratic programming problems arising in nonsmooth optimization. *IMA J. Numer. Anal.* **6**, 137–152 (1986)
9. Kiwiel, K.C.: Proximity control in bundle methods for convex nondifferentiable minimization. *Math. Program.* **46**, 105–122 (1990)
10. Kiwiel, K.C.: A Cholesky dual method for proximal piecewise linear programming. *Numer. Math.* **68**, 325–340 (1994)
11. Kiwiel, K.C.: A projection-proximal bundle method for convex nondifferentiable minimization. In: Théra, M., Tichatschke, R. *Ill-posed Variational Problems and Regularization Techniques*, Lecture Notes in Economics and Mathematical Systems 477, pp. 137–150. Springer, Berlin (1999)
12. Kiwiel, K.C.: A proximal bundle method with approximate subgradient linearizations. *SIAM J. Optim.* **16**, 1007–1023 (2006)
13. Kiwiel, K.C.: A proximal-projection bundle method for Lagrangian relaxation, including semidefinite programming. *SIAM J. Optim.* **17**, 1015–1034 (2006)
14. Kiwiel, K.C., Larsson, T., Lindberg, P.O.: Lagrangian relaxation via ballstep subgradient methods. *Math. Oper. Res.* **32**, 669–686 (2007)
15. Kiwiel, K.C., Lemaréchal, C.: An inexact bundle variant suited to column generation. *Math. Program.* **118**, 177–206 (2009)
16. Kiwiel, K.C., Rosa, C.H., Ruszczyński, A.: Proximal decomposition via alternating linearization. *SIAM J. Optim.* **9**, 668–689 (1999)
17. Lemaréchal, C., Ouerou, A., Petrou, G.: A bundle-type algorithm for routing in telecommunication data networks. *Comput. Optim. Appl.* (2007). doi:[10.1007/s10589-007-9160-7](https://doi.org/10.1007/s10589-007-9160-7)
18. Martinet, B.: Régularisation d'inéquations variationnelles par approximations successives. *RAIRO Rech. Opér.* **4**(R3), 154–158 (1970)
19. Ouerou, A., Mahey, P., Vial, J.P.: A survey of algorithms for convex multicommodity flow problems. *Manag. Sci.* **46**, 126–147 (2000)
20. Rockafellar, R.T.: *Convex Analysis*. Princeton University Press, Princeton (1970)
21. Rockafellar, R.T.: Monotone operators and the proximal point algorithm. *SIAM J. Control Optim.* **14**, 877–898 (1976)