



# Development of a Method for Fucoxanthin Production Using the Haptophyte Marine Microalga *Pavlova* sp. OPMS 30543

Akihiko Kanamoto<sup>1,2</sup> · Yuichi Kato<sup>3</sup> · Erina Yoshida<sup>1</sup> · Tomohisa Hasunuma<sup>1,3</sup> · Akihiko Kondo<sup>1,3,4</sup>

Received: 17 August 2020 / Accepted: 1 March 2021 / Published online: 13 March 2021  
© The Author(s), under exclusive licence to Springer Science+Business Media, LLC, part of Springer Nature 2021

## Abstract

The natural pigment fucoxanthin has attracted global attention because of its superior antioxidant properties. The haptophyte marine microalgae *Pavlova* spp. are assumed to be promising industrial fucoxanthin producers as their lack of a cell wall could facilitate the commercialization of cultured cells as a whole food. This study screened promising *Pavlova* strains with high fucoxanthin content to develop an outdoor cultivation method for fucoxanthin production. Initial laboratory investigations of *P. pinguis* NBRC 102807, *P. lutheri* NBRC 102808, and *Pavlova* sp. OPMS 30543 identified OPMS 30543 as having the highest fucoxanthin content. The culture conditions were optimized for OPMS 30543. Compared with f/2 and Walne's media, the use of Daigo's IMK medium led to the highest biomass production and highest fucoxanthin accumulation. The presence of seawater elements in Daigo's IMK medium was necessary for the growth of OPMS 30543. OPMS 30543 was then cultured outdoors using acrylic pipe photobioreactors, a plastic bag, an open tank, and a raceway pond. Acrylic pipe photobioreactors with small diameters enabled the highest biomass production. Using an acrylic pipe photobioreactor with 60-mm diameter, a fucoxanthin productivity of 4.88 mg/L/day was achieved in outdoor cultivation. Thus, this study demonstrated the usefulness of *Pavlova* sp. OPMS 30543 for fucoxanthin production in outdoor cultivation.

**Keywords** Fucoxanthin · Marine microalgae · Outdoor cultivation · *Pavlova*

## Introduction

Fucoxanthin is synthesized by brown algae and diatoms as a major photosynthetic pigment; thus, it is the most abundant marine carotenoid and is widely distributed in nature (Dembitsky

and Maoka 2007). Fucoxanthin has attracted considerable attention for use in the pharmaceutical, nutraceutical, and cosmetic industries because of its superior antioxidant properties (Peng et al. 2011). Fucoxanthin has also been studied for its anti-cancer activity in human cells (Hosokawa et al. 1999; Kotake-Nara et al. 2001), anti-type 2 diabetes and anti-obesity effects in mice and human cells (Gammone and d'Orazio 2015; Maeda et al. 2007), in vitro anti-cholesterol activity (Kawee-ai et al. 2013), anti-inflammatory effects in rats (Shiratori et al. 2005), anti-angiogenic effects in human cells (Sugawara et al. 2006), anti-malarial effects against *Plasmodium falciparum* (Afolayan et al. 2008), and anti-hypertensive effects in rats (Ikeda et al. 2003; Sivagnanam et al. 2015), as well as for the treatment of Alzheimer's disease (Kawee-ai et al. 2013). Currently, fucoxanthin is produced commercially from brown algae such as *Laminaria* spp. and *Undaria pinnatifida* and diatoms such as *Phaedactylum tricorutum* (Gayen et al. 2019). Algatechnologies Inc. supplies Fucoxital™, which is manufactured from *P. tricorutum*, and this was the first fucoxanthin food ingredient product approved by the US Food and Drug Administration (NDI 1048, 2017). Fucoxanthin obtained from diatoms such as *Chaetoceros gracilis* and *Odontella aurita* also have potential industrial applications

A. Kanamoto was a CEO of OP Bio Factory at the time this study was conducted. A. Kanamoto participated in the experiments as a representative of OP Bio Factory. The corresponding author has full access to all the data in the study and is completely responsible for the data and its accuracy.

✉ Tomohisa Hasunuma  
hasunuma@port.kobe-u.ac.jp

- 1 Graduate School of Innovation, Science and Technology, Kobe University, 1-1 Rokkodai, Nada, Kobe 657-8501, Japan
- 2 OP Bio Factory Co., Ltd., 5-8 Aza-Suzaki, Uruma, Okinawa 904-2234, Japan
- 3 Engineering Biology Research Center, Kobe University, 1-1 Rokkodai, Nada, Kobe 657-8501, Japan
- 4 RIKEN Center for Sustainable Resource Science, 1-7-22 Suehiro, Tsurumi, Yokohama, Kanagawa 230-0045, Japan

(Tokushima et al. 2016; Xia et al. 2018). Culture conditions such as light and nutrients have been reported to affect microalgal fucoxanthin production (Xia et al. 2013; Gómez-Loredo et al. 2016; Lu et al. 2018; Yang and Wei 2020). In *O. aurita*, cultivation in a high nitrate medium led to high fucoxanthin content and volumetric fucoxanthin production (Xia et al. 2013). In *P. tricornutum*, tryptone and urea were examined as supplemental nitrogen sources, and tryptone was found to improve cell growth and fucoxanthin production (Yang and Wei 2020).

In addition to brown algae and diatoms, haptophyte microalgae of *Pavlova* spp., such as *P. lutheri* and *P. pinguis*, can produce fucoxanthin (Hiller et al. 1988; Lananan et al. 2013). The marine microalga *P. lutheri*, which can produce considerable amounts of polyunsaturated fatty acids (PUFAs), is commonly employed as a larval feed in aquaculture (Brown et al. 1997; Guihéneuf and Stengel 2013), and its PUFA yield is increased via random mutagenesis (Meireles et al. 2003). *P. pinguis* contains abundant docosapentaenoic acid (Milke et al. 2008). As *Pavlova* spp. do not have a cell wall (Green 1980); they can be commoditized as whole foods without the need to extract intracellular fucoxanthin. Thus, *Pavlova* spp. are considered valuable fucoxanthin producers. However, there are no quantitative reports regarding fucoxanthin production by *Pavlova* spp.

In the present study, screening of several *Pavlova* spp. to identify a strain with high fucoxanthin content revealed that *Pavlova* sp. OPMS 30543 is a promising producer. Culture conditions for OPMS 30543 were examined and optimized, and factors affecting biomass and fucoxanthin production were investigated in laboratory experiments. Large-scale and outdoor cultivation of OPMS 30543 was also conducted using various culture facilities.

## Materials and Methods

### Strains and Laboratory-Scale Cultivation

*Pavlova pinguis* NBRC 102807 and *P. lutheri* NBRC 102808 were obtained from the National Biological Resource Center (NBRC) of the National Institute of Technology and Evaluation. *Pavlova* sp. OPMS 30543 was isolated from brackish water from Okinawa Main Island, Japan. Microalgae were photoautotrophically cultivated in artificial seawater (Marine Art SF-1, Tomita Pharmaceutical, Tokushima, Japan) enriched with either Daigo's IMK (FUJIFILM Wako Pure Chemical Corp., Osaka, Japan), f/2 (Guillard and Ryther 1962), or Walne's (Walne 1970) elements (Table 1). Culture conditions were as follows, unless otherwise noted in the figure legends: 800 mL of medium in 1-L sterilized bottles, illumination with white fluorescent lamps at an intensity of 150  $\mu\text{mol photons/m}^2/\text{s}$  with a 12-h:12-h light/dark cycle, and continuous aeration of 0.25 mL/mL/min. Cells were harvested using 0.7- $\mu\text{m}$  pore size glass fiber filter paper GF/F (Cytiva, Tokyo, Japan), washed with distilled water, and dried at 120 °C for 2 h before measurement of dry cell weight (DCW). To examine alternative nitrogen sources for Daigo's IMK, media were prepared as shown in Table 2.

### Pigment Analysis

Approximately 10 mg of dried cells was suspended in 1 mL of acetonitrile, mixed by vortexing for 1 min, and disrupted by sonication for 10 min. After centrifugation at 10,000 $\times g$  for 2 min, the supernatant was analyzed by high-performance liquid chromatography (Shimadzu, Kyoto, Japan) under the following conditions: reverse-phase column, COSMOSIL 5C<sub>18</sub>-AR-II, 4.6 mm I.D.  $\times$  150 mm (Nacalai Tesque, Kyoto, Japan); column oven temperature, 40 °C; mobile phase, 80%

**Table 1** Nutrients in seawater media (mg/L)

1 $\times$ Daigo's IMK		f/2		Walne's	
NaNO <sub>3</sub>	200	NaNO <sub>3</sub>	75	NaNO <sub>3</sub>	100
Na <sub>2</sub> HPO <sub>4</sub>	1.4	NaH <sub>2</sub> PO <sub>4</sub> · 2H <sub>2</sub> O	6	NaH <sub>2</sub> PO <sub>4</sub> · 2H <sub>2</sub> O	20
K <sub>2</sub> HPO <sub>4</sub>	5	-	-	-	-
NH <sub>4</sub> Cl	2.68	-	-	-	-
Fe-EDTA	5.2	FeCl <sub>3</sub> · 6H <sub>2</sub> O	3.16	FeCl <sub>3</sub> · 6H <sub>2</sub> O	1.3
Mn-EDTA	0.332	MnCl <sub>2</sub> · 4H <sub>2</sub> O	0.18	MnCl <sub>2</sub> · 4H <sub>2</sub> O	0.36
Na <sub>2</sub> -EDTA	37.2	Na <sub>2</sub> -EDTA	4.4	Na <sub>2</sub> -EDTA	45
ZnSO <sub>4</sub> · 7H <sub>2</sub> O	0.023	ZnSO <sub>4</sub> · 7H <sub>2</sub> O	0.021	ZnCl <sub>2</sub>	0.021
CoSO <sub>4</sub> · 7H <sub>2</sub> O	0.014	CoSO <sub>4</sub> · 7H <sub>2</sub> O	0.012	CoCl <sub>2</sub> · 6H <sub>2</sub> O	0.02
Na <sub>2</sub> MoO <sub>4</sub> · 2H <sub>2</sub> O	0.0073	Na <sub>2</sub> MoO <sub>4</sub> · 2H <sub>2</sub> O	0.007	(NH <sub>4</sub> ) <sub>6</sub> Mo <sub>7</sub> O <sub>24</sub> · 4H <sub>2</sub> O	0.009
CuSO <sub>4</sub> · 5H <sub>2</sub> O	0.0025	CuSO <sub>4</sub> · 5H <sub>2</sub> O	0.007	CuSO <sub>4</sub> · 5H <sub>2</sub> O	0.02
H <sub>2</sub> SeO <sub>3</sub>	0.0017	-	-	-	-
-	-	Na <sub>2</sub> SiO <sub>3</sub> · 9H <sub>2</sub> O	10	-	-
-	-	-	-	H <sub>3</sub> BO <sub>3</sub>	33.6
Thiamine-HCl	0.2	Thiamine-HCl	0.1	Thiamine-HCl	0.01
Biotin	0.0015	Biotin	0.0005	Biotin	0.0002
Vitamin B12	0.0015	Vitamin B12	0.0005	Vitamin B12	0.01

**Table 2** Nutrients in modified IMK (mIMK) media (mg/L)

	1× Daigo's IMK	mIMK (NaNO <sub>3</sub> )	mIMK (KNO <sub>3</sub> )	mIMK (CO[NH <sub>2</sub> ] <sub>2</sub> )	mIMK (NH <sub>4</sub> Cl)
NaNO <sub>3</sub>	200	200	-	-	-
KNO <sub>3</sub>	-	-	200	-	-
CO(NH <sub>2</sub> ) <sub>2</sub>	-	-	-	200	-
NH <sub>4</sub> Cl	2.68	-	-	-	200
Na <sub>2</sub> HPO <sub>4</sub>	1.4	-	-	-	-
K <sub>2</sub> HPO <sub>4</sub>	5	5	5	5	5
Fe-EDTA	5.2	-	-	-	-
Mn-EDTA	0.332	-	-	-	-
Na <sub>2</sub> -EDTA	37.2	37.2	37.2	37.2	37.2
ZnSO <sub>4</sub> ·7H <sub>2</sub> O	0.023	0.023	0.023	0.023	0.023
CoSO <sub>4</sub> ·7H <sub>2</sub> O	0.014	-	-	-	-
Na <sub>2</sub> MoO <sub>4</sub> ·2H <sub>2</sub> O	0.0073	-	-	-	-
CuSO <sub>4</sub> ·5H <sub>2</sub> O	0.0025	0.0025	0.0025	0.0025	0.0025
H <sub>2</sub> SeO <sub>3</sub>	0.0017	-	-	-	-
Thiamine-HCl	0.2	-	-	-	-
Biotin	0.0015	-	-	-	-
Vitamin B12	0.0015	-	-	-	-

acetonitrile aqueous containing 0.1% formic acid; flow rate, 1 mL/min; and detection, 450 nm using a photodiode array detector. Fucoxanthin signals were identified and quantified using a standard curve generated using the fucoxanthin standard (FUJIFILM Wako Pure Chemical Corp.).

## Large-Scale Cultivation

OPMS 30543 was cultivated outdoors under natural sunlight using the following common cultivation systems: (1) 60-mm outer diameter and 5-mm thickness acrylic pipe photobioreactor (PBR), (2) 114-mm outer diameter and 5-mm thickness acrylic pipe PBR, (3) 216-mm outer diameter and 5-mm thickness acrylic pipe PBR, (4) 267-mm outer diameter and 5-mm thick acrylic pipe PBR, (5) 450-mm outer diameter and 0.1-mm thickness plastic bag, (6) 200-L polycarbonate open tank, and (7) 500-L raceway pond, in 50% artificial seawater containing 2× Daigo's IMK elements described above (Table 1). Agitation was performed by aeration at 0.25 mL/min for (1) and (2), and 0.1 mL/min for (3), (4), (5), and (6) except for the raceway pond, in which the flow rate was adjusted to 0.5 m/s by stirring with a paddle. During cultivation, the pH was adjusted to 8 by supplying 100% CO<sub>2</sub>.

## Results


### Screening of *Pavlova* Strains for Fucoxanthin Production

To develop a fucoxanthin production method using *Pavlova* spp., three strains (i.e., *P. pinguis* NBRC 102807, *P. lutheri* NBRC 102808, and *P. sp.* OPMS 30543) were examined in

this study (Fig. 1a). The strains were cultured in 50% seawater containing 2× Daigo's IMK at 25 °C to identify a promising strain with high fucoxanthin production. Strain NBRC 102808 exhibited the lowest biomass production, whereas NBRC 102807 exhibited the highest biomass production, 1.54 g DCW/L at day 12 (Fig. 1b). In contrast, among these *Pavlova* strains, strain NBRC 102807 exhibited the lowest fucoxanthin content (2.06 mg/g DCW, day 3) (Fig. 1c). OPMS 30543 exhibited measurable biomass production of 0.85 g DCW/L over 12 days and achieved the highest fucoxanthin content, 12.88 mg/g DCW at day 9. Fucoxanthin production (calculated by multiplying the biomass and fucoxanthin content) of 9.01 mg/L at day 9 was achieved by OPMS 30543, which was higher than that of strains NBRC 102807 (2.32 mg/L, day 12) and NBRC 102808 (0.61 mg/L, day 9) (Fig. 1d). Thus, OPMS 30543 was identified as a promising *Pavlova* strain for fucoxanthin production.

### Examination of Culture Medium for OPMS 30543

To determine the optimal medium for fucoxanthin production, biomass and fucoxanthin content were investigated using OPMS 30543 grown in 50% seawater enriched with either 2× Daigo's IMK, f/2 (Guillard and Ryther 1962) or Walne's (Walne 1970) elements (Table 1). Among these conditions, cultivation in 2× Daigo's IMK medium resulted in higher biomass (0.92 g DCW/L) relative to f/2 (0.55 g DCW/L) and Walne's (0.56 g DCW/L) media after 14 days of cultivation (Fig. 2a). In addition, the fucoxanthin content of OPMS 30543 grown in 2× Daigo's IMK medium was significantly higher (2.62 mg/g DCW, day 14) than that of cells grown in f/2 (1.48 mg/g DCW, day 7) or Walne's (1.39 mg/g DCW, day 7) media (Fig. 2b).




**Fig. 1** Comparison of three *Pavlova* strains. **a** Microscopic images of *Pavlova* cells. Scale bars: 50 µm. **b** Biomass. **c** Fucoxanthin content. **d** Fucoxanthin production

Fucoxanthin production of 1.51 mg/L on day 14 was achieved by culturing cells in 2× Daigo's IMK medium, which was double the production of cells grown in medium containing f/2 (0.73 mg/L, day 7) or Walne's (0.79 mg/L) elements (Fig. 2c). Thus, these data suggest that the use of 2× Daigo's IMK was the most suitable for maximizing OPMS 30543 biomass and fucoxanthin production.

### Examination of Culture Conditions for OPMS 30543

To improve the biomass production of OPMS 30543, various culture conditions (i.e., seawater concentration, pH, and temperature) were examined. When cultivated in 2×

Daigo's IMK with different concentrations of seawater, biomass production was observed only in the presence of seawater; OPMS 30543 did not grow in 0% seawater medium (Fig. 3a). The highest biomass of 6.16 g DCW/L on day 14 was achieved in the medium with 50% seawater. The effect of varying the culture pH by supplying CO<sub>2</sub> gas to the medium was also examined (Fig. 3b). OPMS 30543 biomass production was reduced when the pH was adjusted to 6, whereas the highest biomass of 3.78 g DCW/L on day 6 was observed when pH was adjusted to 8. Culture temperature was investigated over the range of 15–35 °C (Fig. 3c). Within this temperature range, OPMS 30543 produced higher biomass at higher temperatures,



**Fig. 2** Comparison of different media for OPMS 30543 cultivation. **a** Biomass. **b** Fucoxanthin content. **c** Fucoxanthin production. Cells were statically cultivated in 200 mL Erlenmeyer flasks with a 100-mL

working volume of 50% seawater containing either 2x Daigo's IMK, f/2, or Walne's elements

and cultivation at 35 °C resulted in the highest biomass production of 3.32 g DCW/L on day 6. Thus, cultivation in 50% seawater medium at 35 °C and pH 8 was determined to be the optimal condition for OPMS 30543 biomass production.


### Modification of IMK Medium by Replacing Nitrogen Sources and Adding Carbon Sources

To further improve OPMS 30543 biomass production and fucoxanthin content, the effect of varying the nitrogen source in the medium was examined. The modified IMK medium was prepared by replacing  $\text{NaNO}_3$  in 1x Daigo's IMK with either  $\text{NaNO}_3$ ,  $\text{KNO}_3$ ,  $\text{CO}(\text{NH}_2)_2$ , or  $\text{NH}_4\text{Cl}$  (Table 2). After 9 days of cultivation, cells cultured in the modified IMK medium containing  $\text{KNO}_3$  exhibited the highest biomass of 1.8 g DCW/L (Fig. 4a). Both urea  $\text{CO}(\text{NH}_2)_2$  and  $\text{NH}_4\text{Cl}$  were found to be available as nitrogen sources for OPMS 30543 cultivation, and biomass production of 1.58 and 0.82 g DCW/L at 10 days was observed, respectively. Use of  $\text{NaNO}_3$ -containing medium resulted in higher fucoxanthin content (12.74 mg/g DCW) than in media with  $\text{KNO}_3$  (5.57 mg/g DCW),  $\text{CO}(\text{NH}_2)_2$

(8.38 mg/g DCW), or  $\text{NH}_4\text{Cl}$  (7.80 mg/g DCW) (Fig. 4b). Fucoxanthin production was the highest when  $\text{NaNO}_3$  was used as the nitrogen source (Fig. 4c). Fucoxanthin production of OPMS 30543 grown in modified IMK medium containing  $\text{NaNO}_3$ ,  $\text{KNO}_3$ ,  $\text{CO}(\text{NH}_2)_2$ , or  $\text{NH}_4\text{Cl}$  was 17.84, 10.03, 13.24, and 6.40 mg/L, respectively. Thus, these data suggest that  $\text{NaNO}_3$  is the best nitrogen source for maximizing OPMS 30543 fucoxanthin production.

The effect of adding various carbon sources to the medium was also examined to enhance biomass and fucoxanthin production. Modified IMK medium was prepared by adding either glucose, methanol, sodium acetate, or sodium bicarbonate to 50% seawater enriched with 1x Daigo's IMK. Each of the additional carbon sources increased biomass production compared with that with the normal 1x Daigo's IMK (Fig. 4d). After 4 days of cultivation, OPMS 30543 grown in medium with sodium acetate exhibited the highest biomass of 1.79 g DCW/L, whereas OPMS 30543 biomass in medium containing glucose, methanol, and sodium bicarbonate was 1.19, 0.71, and 1.28 g DCW/L, respectively. Use of medium containing methanol resulted in the highest fucoxanthin content (7.26 mg/g DCW) relative to medium containing glucose





**Fig. 3** Comparison of culture conditions for OPMS 30543. **a** Seawater concentration in medium. **b** pH, adjusted by supplying CO<sub>2</sub> gas to the culture. Cultures were illuminated with red, blue, and white LEDs

at a total intensity of 300  $\mu\text{mol photons/m}^2/\text{s}$  with a 12-h:12-h light/dark cycle. **c** Culture temperature

(4.25 mg/g DCW), sodium acetate (4.11 mg/g DCW), or sodium bicarbonate (2.99 mg/g DCW) (Fig. 4e). Fucoxanthin production was the highest when sodium acetate was added to the medium (Fig. 4f). Fucoxanthin production by OPMS 30543 grown with glucose, methanol, sodium acetate, and sodium bicarbonate was 5.06, 5.15, 7.36, and 3.83 mg/L, respectively. Thus, sodium acetate was suggested as the optimal carbon source for enhancing fucoxanthin production.


### Large-Scale Outdoor Cultivation of OPMS 30543

A large-scale outdoor OPMS 30543 cultivation test was performed to evaluate the potential of fucoxanthin production outdoors. Acrylic pipe PBRs (5-mm thickness with different outer diameters of 114, 216, and 267 mm), a plastic bag (0.1-mm thickness with 450-mm outer diameter), a 200-L polycarbonate open tank, and a 500-L raceway pond were used for cultivation (Fig. 5). Six days of cultivation outdoors in acrylic pipe PBRs with 114-, 216-, and 267-mm outer diameter produced biomass of 0.73, 0.39, and 0.31 g DCW/L, respectively (Fig. 6a). Cultivation using a plastic bag, a 200-L polycarbonate open tank, and a 500-L raceway pond produced 0.24, 0.26, and 0.10 g DCW/L, respectively,

on day 6. Thus, the acrylic pipe PBRs with smaller outer diameters achieved higher biomass production than the plastic bag, open tank, or raceway pond. To further examine these results, OPMS 30543 was cultivated using an acrylic pipe PBR with a 60-mm outer diameter. Biomass of 1.82 g DCW/L and 2.20 g DCW/L were observed on days 6 and 8, respectively (Fig. 6b), both of which were higher than the biomass production achieved using the acrylic pipe PBR with a 114-mm outer diameter. The fucoxanthin content on day 8 was 20.86 mg/g DCW, which was higher than that achieved with any of the laboratory-scale cultivations in this study. Using a PBR with a 60-mm outer diameter, biomass productivity of 0.23 g DCW/L/day and fucoxanthin productivity of 4.88 mg/L/day were demonstrated in large-scale outdoor cultivation.

### Discussion

In previous studies, *P. lutheri* and *P. pinguis* were examined as aquatic feed producers that accumulate high levels of  $\omega$ -3 fatty acids, including docosahexaenoic acid and eicosapentaenoic acid (Guihéneuf and Stengel 2013; Guihéneuf et al. 2015; Fernandes et al. 2020). However, these organisms




**Fig. 4** Examination of alternative nitrogen sources and additional carbon sources. **a** Biomass, **b** fucoxanthin content, and **c** fucoxanthin production of cells grown in 50% seawater enriched with modified IMK and different nitrogen sources. **d** Biomass, **e** fucoxanthin con-

tent, and **f** fucoxanthin production of cells grown in 50% seawater enriched with 2× Daigo's IMK with additional carbon sources, illuminated with red, blue, and white LEDs at a total intensity of 300  $\mu\text{mol photons/m}^2/\text{s}$  with a 12-h:12-h light/dark cycle

have not been studied extensively for their use as fucoxanthin producers, despite several reports describing fucoxanthin production by *P. lutheri* (Hiller et al. 1988; Lananan et al. 2013) and the advantages of the lack of a cell wall in *Pavlova* spp. (Green 1980). To develop a useful fucoxanthin production method, this study first compared fucoxanthin production in three *Pavlova* strains and identified *Pavlova* sp. OPMS 30543 as a promising strain owing to its significantly higher fucoxanthin production than that of *P. pinguis* NBRC 102807 and *P. lutheri* NBRC 102808 (Fig. 1d).

To determine the optimal conditions for OPMS 30543 cultivation, three types of media were examined. The use of 2× Daigo's IMK medium resulted in higher fucoxanthin production than with either f/2 or Walne's medium (Fig. 2c). A likely reason is that 2× Daigo's IMK contains a much higher level of nitrate (400 mg/L NaNO<sub>3</sub>) than f/2 (75 mg/L NaNO<sub>3</sub>) or Walne's (100 mg/L NaNO<sub>3</sub>) (Table 1). Nitrate supplementation has been reported to increase fucoxanthin production in the diatoms *Phaeodactylum tricornutum* and *O. aurita* (Xia et al. 2013; McClure et al. 2018). Nitrogen supplementation with tryptone improved fucoxanthin

production in *P. tricornutum* (Yang and Wei 2020). This study also investigated different nitrogen sources with which to modify 2× Daigo's IMK and found that the use of NaNO<sub>3</sub> resulted in the highest fucoxanthin accumulation (Fig. 4c). Microalgae growth and fucoxanthin generally show a positive relationship, except under some conditions such as nitrogen depletion, under which fucoxanthin content decreases (Xia et al. 2018). In this study, the modified IMK medium containing KNO<sub>3</sub> led to the highest biomass (Fig. 4a), although the fucoxanthin content was the lowest (Fig. 4b). This might be because the nitrogen source was depleted in the KNO<sub>3</sub> medium owing to the highest cell growth. The effect of the nitrogen source on fucoxanthin production has not been examined in detail in previous studies. Absorption and assimilation of different nitrogen sources were investigated in Pelagophyceae *Aureococcus anophagefferens*, which also accumulates fucoxanthin (Ou et al. 2018). Different from the results of this study, cultivation using urea resulted in the highest fucoxanthin content in this microalga compared with cultivation with NaNO<sub>3</sub>, NH<sub>4</sub>Cl, or glutamic acid. Although the effects differ among




**Fig. 5** Facilities used for outdoor cultivation. **a** Acrylic pipe photobioreactors (5-mm thickness with outer diameters of 114, 216, and 267 mm) and a plastic bag (0.1-mm thickness with a 450-mm outer

diameter). **b** 200-L polycarbonate open tank. **c** 500-L raceway pond. **d** Acrylic pipe photobioreactor (60-mm outer diameter)


algae species, these results suggest that supplementation and type of nitrogen source are important factors affecting fucoxanthin accumulation.

Among the *Pavlova* strains tested in this study, *P. pinguis* NBRC 102807 exhibited the highest biomass production

(Fig. 1b). In contrast, *Pavlova* sp. OPMS 30543 could grow under a wide range of seawater concentrations, ranging from 25 to 100%, with similar biomass productivity (Fig. 3a). This robustness toward salinity is a valuable characteristic for seawater cultivation. OPMS 30543 did not produce



**Fig. 6** Large-scale outdoor cultivation of OPMS 30543. **a** Biomass of OPMS 30543 cultivated using natural light in acrylic pipe PBRs (5-mm thickness with different outer diameters of 114, 216, and 267 mm), a plastic bag (0.1-mm thickness with a 450-mm outer diameter), 200-L polycarbonate open tank, and 500-L raceway pond. **b** Biomass of OPMS 30543 cultivated outdoors under natural light in



an acrylic pipe PBR with a 60-mm outer diameter. In these experiments, 50% seawater enriched with 2× Daigo's IMK was used as the medium. Aeration was provided except for the raceway pond. In the raceway pond, cells were stirred using a paddle. During cultivation, the pH was adjusted to 8 by blowing CO<sub>2</sub>.



**Table 3** Summary of fucoxanthin content in microalgae and macroalgae

	Species	Cell wall	Fucoxanthin content (mg/g DCW)	References
Haptophytes	<i>Pavlova</i> sp.	Negative	20.86	This study
	<i>Isochrysis</i> aff. <i>galbana</i>	Negative	18.23	Kim et al. (2012)
	<i>Isochrysis galbana</i>	Negative	15.8	Sun et al. (2019)
	<i>Tisochrysis lutea</i>	Negative	16.39	Gao et al. (2020)
Diatoms	<i>Chaetoceros gracilis</i>	Positive	2.24	Kim et al. (2012)
	<i>Cylindrotheca closterium</i>	Positive	25.5	Wang et al. (2018)
	<i>Nitzschia laevis</i>	Positive	12.0	Lu et al. (2018)
	<i>Nitzschia</i> sp.	Positive	4.92	Kim et al. (2012)
	<i>Odontella aurita</i>	Positive	18.47	Xia et al. (2013)
	<i>Phaeodactylum tricornutum</i>	Positive	59.2	McClure et al. (2018)
	<i>Thalassiosira weissflogii</i>	Positive	9.5	Marella and Tiwari (2020)
	Chrysophytes	<i>Mallomonas</i> sp.	Positive	26.6
Brown algae	<i>Cystoseira hakodatensis</i>	Positive	2.01	Susanto et al. (2016)
	<i>Cystoseira indica</i>	Positive	3.56	Fariman et al. (2016)
	<i>Nizamuddinina zanardinii</i>	Positive	1.65	Fariman et al. (2016)
	<i>Padina</i> sp.	Positive	1.97	Dang et al. (2017)
	<i>Sargassum horneri</i>	Positive	2.12	Susanto et al. (2016)
	<i>Sargassum linearifolium</i>	Positive	1.76	Dang et al. (2017)
	<i>Sargassum siliquastrum</i>	Positive	1.99	Susanto et al. (2016)
	<i>Sphaerotrichia divaricata</i>	Positive	1.15	Maeda et al. (2018)
	<i>Undaria pinnatifida</i>	Positive	0.73	Xiao et al. (2012)

biomass when cultured in medium with 0% seawater, possibly because Daigo's IMK medium depends upon supplementation of  $Mg^{2+}$  and  $Ca^{2+}$  in seawater (Table 1). Of the three media examined, 2× Daigo's IMK provided the highest OPMS 30543 biomass production (Fig. 2a), probably because it contained more nitrate than either f/2 or Walne's media (Table 1). The effects of an additional carbon source were also examined. This analysis revealed that the addition of glucose, sodium acetate, or sodium bicarbonate to 2× Daigo's IMK medium enhanced OPMS 30543 biomass production (Fig. 4d). In haptophyte *Isochrysis galbana*, glycerol was found to be the best additional carbon source to enhance biomass production, whereas acetate had no effect and glucose only slightly enhanced the growth rate (Alkhamis and Qin 2013). Overall, these data suggest that the addition of a suitable carbon is a promising approach for enhancing the biomass production of microalgae, including OPMS 30543.

In the large-scale outdoor cultivation experiment, the acrylic pipe PBRs demonstrated higher biomass production than the open tank or raceway pond (Fig. 6a). A possible reason for this result is that the open tank and raceway pond were highly contaminated with bacteria, fungi, and protozoa (data not shown). Among the acrylic pipe PBRs examined, those with a smaller diameter produced higher biomass, most likely because the higher surface area-to-volume ratio contributes to more efficient illumination. Using the 60-mm diameter acrylic pipe PBR, a fucoxanthin content of 20.86

mg/g DCW and fucoxanthin productivity of 4.88 mg/L/day was obtained after 8 days of cultivation (Fig. 6b). Fucoxanthin content in various microalgae and macroalgae has been reported in previous studies (Table 3). Microalgae such as haptophytes, diatoms, and chrysophytes generally show higher fucoxanthin content than macroalgae. In diatoms, *P. tricornutum* and *Cylindrotheca closterium* were reported to achieve 59.2 mg/g DCW and 25.5 mg/g DCW fucoxanthin content, respectively (McClure et al. 2018; Wang et al. 2018). Chrysophytes *Mallomonas* sp. also showed a high fucoxanthin content of 26.6 g/g DCW (Petrushkina et al. 2017). For commercialization of cultured cells as a whole food, however, these microalgae would not be favorable because they have a cell wall. In this study, as a cell wall-lacking microalga, *Pavlova* sp. OPMS 30543 achieved a fucoxanthin content of 20.86 mg/g DCW, which is higher than that achieved with *Isochrysis* aff. *galbana* (Kim et al. 2012). Thus, *Pavlova* sp. OPMS 30543 is a promising feedstock for fucoxanthin, characterized by both a high fucoxanthin content and the absence of cell wall. With the development of a large-scale outdoor cultivation method for OPMS 30543 fucoxanthin production as demonstrated in this study, the utilization of *Pavlova* cells as whole foods has taken a step toward successful commercialization.

**Acknowledgements** The authors thank Dr. Takeshi Fujiwara, Dr. Takafumi Watanabe, and Ms. Yuko Koizumi for their technical assistance.

We would like to thank NBRC for supplying *Pavlova pinguis* NBRC 102807 and *Pavlova lutheri* NBRC 102808.

**Author Contribution** A. Kanamoto designed the study, conducted the experiments, and drafted the manuscript. Y. K., E. Y., T. H., and A. Kondo commented on the study, helped interpret results, and revised the manuscript. All authors approved the final version of the manuscript.

**Data Availability** The data supporting the findings of this study are available within this article or from the corresponding author upon reasonable request. *Pavlova pinguis* NBRC 102807 and *Pavlova lutheri* NBRC 102808 can be obtained from the National Biological Resource Center (NBRC).

## Declarations

**Competing Interests** The authors declare that they have no competing interests.

## References

- Afolayan AF, Bolton JJ, Lategan CA, Smith PJ, Beukes DR (2008) Fucoxanthin, tetraprenylated toluquinone and toluhydroquinone metabolites from *Sargassum heterophyllum* inhibit the in vitro growth of the malaria parasite *Plasmodium falciparum*. *Z Naturforsch C J Biosci* 63:848–852
- Alkhamis Y, Qin JG (2013) Cultivation of *Isochrysis galbana* in phototrophic, heterotrophic, and mixotrophic conditions. *Biomed Res Int* 2013:983465
- Brown MR, Jeffrey SW, Volkman JK, Dunstan GA (1997) Nutritional properties of microalgae for mariculture. *Aquaculture* 151:315–331
- Dang TT, Bowyer MC, Van Altena IA, Scarlett CJ (2017) Comparison of chemical profile and antioxidant properties of the brown algae. *Int J Food Sci Technol* 53:174–181
- Dembitsky VM, Maoka T (2007) Allenic and cumulenyl lipids. *Prog Lipid Res* 46:328–375
- Fariman GA, Shastan SJ, Zahedi MM (2016) Seasonal variation of total lipid, fatty acids, fucoxanthin content, and antioxidant properties of two tropical brown algae (*Nizamuddinina zardinii* and *Cystoseira indica*) from Iran. *J Appl Phycol* 28:1323–1331
- Fernandes T, Martel A, Cordeiro N (2020) Exploring *Pavlova pinguis* chemical diversity: a potentially novel source of high value compounds. *Sci Rep* 10:339
- Gammone MA, d'Orazio N (2015) Anti-obesity activity of the marine carotenoid fucoxanthin. *Mar Drugs* 13:2196–2214
- Gao F, Teles Cabanelas I, Wijffels RH, Barbosa MJ (2020) Process optimization of fucoxanthin production with *Tisochrysis lutea*. *Bioresour Technol* 315:123894
- Gayen K, Bhowmick TK, Maity SK (2019) Sustainable Downstream Processing of Microalgae for Industrial Application. CRC Press, Boca Raton
- Gómez-Loredo A, Benavides J, Rito-Palmares M (2016) Growth kinetics and fucoxanthin production of *Phaeodactylum tricoratum* and *Isochrysis galbana* cultures at different light and agitation conditions. *J Appl Phycol* 28:849–860
- Green JC (1980) The fine structure of *Pavlova pinguis* Green and a preliminary survey of the order Pavloales (Prymnesiophyceae). *Br Phycol J* 15:151–191
- Guihéneuf F, Stengel DB (2013) LC-PUFA-enriched oil production by microalgae: accumulation of lipid and triacylglycerols containing n-3 LC-PUFA is triggered by nitrogen limitation and inorganic carbon availability in the marine haptophyte *Pavlova lutheri*. *Mar Drugs* 11:4246–4266
- Guihéneuf F, Mimouni V, Tremblin G, Ulmann L (2015) Light intensity regulates LC-PUFA incorporation into lipids of *Pavlova lutheri* and the final desaturase and elongase activities involved in their biosynthesis. *J Agric Food Chem* 63:1261–1267
- Guillard RRL, Ryther JH (1962) Studies of marine planktonic diatoms: I. *Cyclotella nana* Hustedt, and *Detonula confervacea* (Cleve) Gran. *Can J Microbiol* 8:229–239
- Hiller RG, Larkum AWD, Wrench PM (1988) Chlorophyll proteins of the prymnesiophyte *Pavlova lutheri* (Droop) comb. nov.: identification of the major light-harvesting complex. *Biochimica et Biophysica Acta - Bioenergetics* 932:223–231
- Hosokawa M, Wanezaki S, Miyauchi K, Kurihara H, Kohno H, Kawabata J, Odashima S, Takahashi K (1999) Apoptosis-inducing effect of fucoxanthin on human leukemia cell line HL-60. *Food Sci Technol Res* 5:243–246
- Ikeda K, Kitamura A, Machida H, Watanabe M, Negishi H, Hiraoka J, Nakano T (2003) Effect of *Undaria pinnatifida* (Wakame) on the development of cerebrovascular diseases in stroke-prone spontaneously hypertensive rats. *Clin Exp Pharmacol Physiol* 30:44–48
- Kawee-ai A, Kuntiya A, Kim SM (2013) Anticholinesterase and antioxidant activities of fucoxanthin purified from the microalga *Phaeodactylum tricoratum*. *Nat Prod Commun* 8:1381–1386
- Kim SM, Kang SW, Kwon ON, Chung D, Pan CH (2012) Fucoxanthin as a major carotenoid in *Isochrysis* aff. *galbana*: Characterization of extraction for commercial application. *J Korean Soc Appl Biol Chem* 55:477–483
- Kotake-Nara E, Kushirom M, Zhang H, Sugawara T, Miyashita K, Nagao A (2001) Carotenoids affect proliferation of human prostate cancer cells. *J Nutr* 131:3303–3306
- Lananan F, Jusoh A, Ali N, Lam SS, Endut A (2013) Effect of Conway medium and f/2 medium on the growth of six genera of South China Sea marine microalgae. *Bioresour Technol* 141:75–82
- Lu X, Sun H, Zhao W, Cheng KW, Chen F, Liu B (2018) A hetero-photoautotrophic two-stage cultivation process for production of fucoxanthin by the marine diatom *Nitzschia laevis*. *Mar Drugs* 16:219
- Maeda H, Fukuda S, Izumi H, Saga N (2018) Anti-oxidant and fucoxanthin contents of brown alga *Ishimozuku* (*Sphaerotrichia divaricata*) from the West Coast of Aomori Japan. *Mar Drugs* 16:255
- Maeda H, Hosokawa M, Sashima T, Miyashita K (2007) Dietary combination of fucoxanthin and fish oil attenuates the weight gain of white adipose tissue and decreases blood glucose in obese/diabetic KK-A<sup>y</sup> mice. *J Agric Food Chem* 55:7701–7706
- Marella TK, Tiwari A (2020) Marine diatom *Thalassiosira weissflogii* based biorefinery for co-production of eicosapentaenoic acid and fucoxanthin. *Bioresour Technol* 307:123245
- McClure DD, Luiz A, Gerber B, Barton GW, Kavanagh JM (2018) An investigation into the effect of culture conditions on fucoxanthin production using the marine microalgae *Phaeodactylum tricoratum*. *Algal Res* 29:41–48
- Meireles LA, Guedes C, Malcata FX (2003) Increase of the yields of eicosapentaenoic and docosahexaenoic acids by the microalga *Pavlova lutheri* following random mutagenesis. *Biotechnol Bioeng* 81:50–55
- Milke LM, Bricelj VM, Parrish CC (2008) Biochemical characterization and nutritional value of three *Pavlova* spp. in unialgal and mixed diets with *Chaetoceros muelleri* for postlarval sea scallops *Placopecten magellanicus*. *Aquaculture* 276:130–142
- Ou L, Cai Y, Jin W, Wang Z, Lu S (2018) Understanding the nitrogen uptake and assimilation of the Chinese strain of *Aureococcus anophagefferens* (Pelagophyceae). *Algal Res* 34:182–190
- Peng J, Yuan JP, Wu CF, Wang JH (2011) Fucoxanthin, a marine carotenoid present in brown seaweeds and diatoms: metabolism and bioactivities relevant to human health. *Mar Drugs* 9:1806–1828

- Petrushkina M, Gusev E, Sorokin B, Zotko N, Mamaeva A, Filimonova A, Kulikovskiy M, Maltsev Y, Yampolsky I, Guglya E, Vinokurov V, Namsaraev Z, Kuzmin D (2017) Fucoxanthin production by heterokont microalgae. *Algal Res* 24:387–393
- Shiratori K, Ohgami K, Ilieva I, Jin XH, Koyama Y, Miyashita K, Yoshida K, Kase S, Ohno S (2005) Effects of fucoxanthin on lipopolysaccharide-induced inflammation in vitro and in vivo. *Exp Eye Res* 81:422–428
- Sivagnanam SP, Yin S, Choi JH, Park YB, Woo HC, Chun BS (2015) Biological properties of fucoxanthin in oil recovered from two brown seaweeds using supercritical CO<sub>2</sub> extraction. *Mar Drugs* 13:3422–3442
- Sugawara T, Matsubara K, Akagi R, Mori M, Hirata T (2006) Antiangiogenic activity of brown algae fucoxanthin and its deacetylated product, fucoxanthinol. *J Agric Food Chem* 54:9805–9810
- Sun Z, Wang X, Liu J (2019) Screening of *Isochrysis* strains for simultaneous production of docosahexaenoic acid and fucoxanthin. *Algal Res* 41:101545
- Susanto E, Fahmi AS, Abe M, Hosokawa M, Miyashita K (2016) Lipids, fatty acids, and fucoxanthin content from temperate and tropical brown seaweeds. *Aquat Procedia* 7:66–75
- Tokushima H, Inoue-Kashino N, Nakazato Y, Masuda A, Ifuku K, Kashino Y (2016) Advantageous characteristics of the diatom *Chaetoceros gracilis* as a sustainable biofuel producer. *Biotechnol Biofuels* 9:235
- Walne PR (1970) Studies on the food value of nineteen genera of algae to juvenile bivalves of the genera *Ostrea*, *Crassostrea*, *Mercenaria*, and *Mytilus*. *Fish Invest* 26:1–62
- Wang S, Verma SK, Said IH, Thomsen L, Ullrich MS, Kuhnert N (2018) Changes in the fucoxanthin production and protein profiles in *Cylindrotheca closterium* in response to blue light-emitting diode light. *Microb Cell Fact* 17:110
- Xia S, Gao B, Fu J, Xiong J, Zhang C (2018) Production of fucoxanthin, chrysolaminarin, and eicosapentaenoic acid by *Odontella aurita* under different nitrogen supply regimes. *J Biosci Bioeng* 126:723–729
- Xia S, Wang K, Wan L, Li A, Hu Q, Zhang C (2013) Production, characterization, and antioxidant activity of fucoxanthin from the marine diatom *Odontella aurita*. *Mar Drugs* 11:2667–2681
- Xiao X, Si X, Yuan Z, Xu X, Li G (2012) Isolation of fucoxanthin from edible brown algae by microwave-assisted extraction coupled with high-speed countercurrent chromatography. *J Sep Sci* 35:2313–2317
- Yang R, Wei D (2020) Improving fucoxanthin production in mixotrophic culture of marine diatom *Phaeodactylum tricornutum* by LED light shift and nitrogen supplementation. *Front Bioeng Biotechnol* 8:820

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.