



Correction to: Efficacy of superficial temporal artery-middle cerebral artery double bypass in patients with hemorrhagic moyamoya disease: surgical effects for operated hemispheric sides

Taichi Ishiguro¹ · Yoshikazu Okada² · Tatsuya Ishikawa¹ · Koji Yamaguchi¹ · Akitsugu Kawashima³ · Takakazu Kawamata¹

Published online: 4 January 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Correction to: Neurosurgical Review
<https://doi.org/10.1007/s10143-018-01059-z>

Table 2 of the original version of this article contained an added data. Correct Table 2 is presented here.

The original article has been corrected.

Table 2 Perioperative outcomes (within 30 days) in all 52 operated sides

Perioperative outcomes	<i>n</i> = 52 (operated sides)	%
Hyperperfusion (estimated by our method)	11	21.2
Transient neurological deficit	6	11.5
Cerebral infarction	0	0
Partial seizure	4	7.7
Surgical site infection	1	1.9
Focal vasogenic edema	7	13.5
Subcortical hemorrhage at the bypassed site	4	7.7
Morbidity and mortality	<i>n</i> = 36 (patients)	%
Deterioration of mRS score at discharge	2	5.6
Death	0	0

The online version of the original article can be found at <https://doi.org/10.1007/s10143-018-01059-z>

✉ Takakazu Kawamata
tkawamata@twmu.ac.jp

¹ Department of Neurosurgery, Tokyo Women's Medical University, 8-1 Kawada-cho, Shinjuku-ku, Tokyo 162-8666, Japan

² Department of Neurosurgery, St. Luke's International Hospital, Tokyo, Japan

³ Department of Neurosurgery, Tokyo Women's Medical University Yachiyo Medical Center, Chiba, Japan