



# Effect of humic acids, nitrate and oxygen on the photodegradation of the flubendiamide insecticide: identification of products

M. Bavcon Kralj<sup>1</sup> · H. Divanović<sup>2</sup> · S. Košenina<sup>1</sup> · M. Kete<sup>3</sup> · A. T. Lebedev<sup>4</sup> · V. B. Artaev<sup>5</sup> · P. Trebše<sup>1</sup>

Received: 17 July 2017 / Accepted: 12 December 2017 / Published online: 21 December 2017  
© Springer International Publishing AG, part of Springer Nature 2017

## Abstract

Flubendiamide is a ryanodine insecticide that shows a strong insecticidal activity and is relatively safe for non-target organisms. Actually only flubendiamide and its product desiodo-flubendiamide have been studied during catalytic degradation using TiO<sub>2</sub> and ZnO. Therefore, here we tested the photocatalytic removal of flubendiamide in the presence of nitrates or humic acids. Degradation kinetics were monitored using high-performance liquid chromatography ultraviolet–visible detector. Product identification was done using a high-resolution time-of-flight mass spectrometer coupled to a gas chromatograph (GC-HRMS). Results show that the addition of humic acids at 10 mg l<sup>-1</sup> increased the removal of flubendiamide more than five times. The addition of nitrate ions at 10 mg l<sup>-1</sup> had no influence. The removal of flubendiamide was more than ten times faster in experiments with oxygen purging. Fourteen degradation products were identified, which can be classified into three groups: phthalimide and related phthalic acid derivatives, fluorinated species related to the second amide moiety, and advanced transformation products.

**Keywords** TiO<sub>2</sub> photocatalysis · Flubendiamide · Degradation products

## Introduction

Advanced oxidation processes (AOPs), although making use of different reacting systems (H<sub>2</sub>O<sub>2</sub> and/or O<sub>3</sub>; Fenton system: addition of H<sub>2</sub>O<sub>2</sub> to Fe<sup>2+</sup> salts, semiconductors as TiO<sub>2</sub>, ozone chemistry O<sub>3</sub>/H<sub>2</sub>O<sub>2</sub> and O<sub>3</sub>/UV and other combinations of oxidants and UV light), are all defined by the same chemical feature: production of ·OH radicals (Andreozzi et al. 1999).

Photocatalysis is an important AOP, making use of a semiconductor–metal oxides as a catalyst, because of their

ability to create photogenerated conduction band electrons ( $e_{cb}^-$ ) and valence band holes ( $h_{vb}^+$ ), which enable redox reactions with adsorbed aqueous species. Although several catalysts have been exploited so far, only TiO<sub>2</sub> in the anatase form expressed sufficiently high stability, good performance and low cost. That is why it is one of the most important materials used in oxidation of organic molecules (Castellote and Bengtsson 2011). It is well known that photocatalytic activity of TiO<sub>2</sub> depends on its electronic properties and structure, doping TiO<sub>2</sub> with cationic and anionic metals, and coupling TiO<sub>2</sub> with other semiconductors rises its capability of degrading pollutants (Daghrir et al. 2013).

Conventional photocatalytic processes have been exploited in large reactors of a wide variety of designs (e.g. parabolic trough, inclined plate, fixed/fluidised bed, packed bed, corrugated plate and falling film photoreactors). However, the commercial use of photocatalytic technologies implies still high efforts to promote the contact between solar irradiation and the surface of the photocatalyst in order to achieve an efficient photodegradation of organic pollutants (Braham and Harris 2009). However, it has been frequently observed in water purification systems (Barreiro and Pratt 1992) that some pollutants are not amenable to biological treatments and express a high chemical stability and/

✉ M. Bavcon Kralj  
mojca.kralj@zf.uni-lj.si

<sup>1</sup> Faculty of Health Sciences, University of Ljubljana, 1000 Ljubljana, Slovenia

<sup>2</sup> Faculty of Pharmacy, University of Sarajevo, Sarajevo, Bosnia and Herzegovina

<sup>3</sup> Laboratory for Environmental Research, University of Nova Gorica, 5000 Nova Gorica, Slovenia

<sup>4</sup> Organic Chemistry Department, Lomonosov Moscow State University, Moscow, Russia

<sup>5</sup> LECO Corporation, 3000 Lakeview Avenue, St. Joseph, MI, USA

or a strong resistance to be completely mineralised. In these cases, it is necessary to define which degradation products are formed during the catalytic process.

Photocatalytic reactors used for water treatment can be classified not only regarding their designs, but also according to the catalysts' application. There are designs with the photocatalyst suspended in the aqueous medium or with the photocatalyst immobilised on the surface. The most practical and frequently used immobilisation methods to produce thick highly active layers for air/water treatment are sol–gel, powder modified titania sol and sol–pray (Kete et al. 2014). In this study, we proved that the hybrid sol suspension method can be successfully used to prepare stable, thick and highly active photocatalytic layers used for flubendiamide degradation.

Flubendiamide is one of RI (ryanodine insecticides), developed by Nihon Nohyaku Co., Ltd. (Tokyo, Japan). Its mode of action involves activation of ryanodine-sensitive calcium release channels (ryanodine receptors; RyRs) in insects. The insect RyRs remain in an open state in a species-specific manner and affect the  $\text{Ca}^{2+}$  pump activity (Masaki et al. 2005; Masaki 2008). The chemical name of flubendiamide is *N*<sup>2</sup>-[1,1-dimethyl-2-(methylsulphonyl)ethyl]-3-iodo-*N*<sup>1</sup>-[2-methyl-4-[1,2,2,2-tetrafluoro-1-(trifluoromethyl)ethyl]phenyl]-1,2-benzenedicarboxamide. The uniqueness of the structure results from three parts with novel substituents: heptafluoroisopropyl group in the amide moiety, a sulphonylalkyl group in the aliphatic amide moiety and an iodine atom at the 3-position of the phthalic acid moiety (Tohnishi et al. 2005). The compound shows an extremely strong insecticidal activity and relative safety for non-target organisms.

However, there was a need of additional peer review for flubendiamide carried out by the competent authority of the rapporteur Member State Greece 3 years ago. The European Food Safety Authority (EFSA) concluded that flubendiamide is appropriate for use in regulatory risk. In a laboratory sterile aqueous photolysis experiment, flubendiamide formed two major transformation products named desiodo-flubendiamide (NNI-0001-desiodo (A1)) and hydroxy-perfluoroalkyl-flubendiamide [NNI-0001-3-OH-hydroxy-perfluoroalkyl (A10)]. EFSA concluded that the risk assessment for aquatic invertebrates from metabolite A10 could not be finalised (EFSA 2013).

Flubendiamide and its major transformation product desiodo-flubendiamide have been studied in different plants (Buddidathia et al. 2016; Mohapatra et al. 2010, 2011) and soils. The calculated half-life values for flubendiamide ( $t_{1/2}$ ) ranged from 37.62 to 50.17, 43–50.17 to 50.17–60.21 days for new alluvial, red/laterite and coastal soil (Paramasivam and Banerjee 2012) to 155.1 and 130.8 days in field capacity and submerged soils. In the later case, 2.5% of organic manure was added (Das and Mukherjee 2012).

In aqueous solutions (tap, leaching and watercourse), the photocatalytic degradation of flubendiamide using ZnO and  $\text{TiO}_2$  was enhanced by the addition of an electron acceptor, such as  $\text{Na}_2\text{S}_2\text{O}_8$ . In addition, the main phototransformation product desiodo-flubendiamide was also successfully photodegraded under the same conditions (Fenoll et al. 2015). To our knowledge, there is a lack of information regarding photocatalytic removal of ryanodine insecticides, in the presence of nitrates or dissolved organic matter in general as naturally the most widespread oxidisers. Many examples summarised by Remucal (2014) show that dissolved organic matter, nitrites and nitrates can be an important source of  $\cdot\text{OH}$  radicals and therefore may play a significant role in the elimination of pollutants. On the other side, it is known that nitrate ions undergo photocatalytic degradation during photocatalytic processes using different photocatalytic materials (Remucal 2014).

Within this research, we have studied the photocatalytic degradation of flubendiamide using  $\text{TiO}_2$  films in order to identify its transformation products and propose its degradation pathway. Additionally, we have investigated how the addition of nitrates or humic acids influences the photocatalytic degradation of flubendiamide.

## Experimental

### Materials

The analytical standard of flubendiamide (98.8% purity) and organic solvents: acetone, acetonitrile, salt  $\text{NaNO}_3$  and humic substances, were all from Sigma-Aldrich (St. Louis, Missouri, USA).

### Photocatalytic experiment

The reactor cell consisted of a Durand glass tube (280 mm, inner diameter 80 mm) with the effective volume of 1.5 L. The reactor cell was then placed in the photoreactor chamber equipped with ultraviolet A (UVA) lamps (15 W, 265 mm × 16 mm, Philips Cleo; broad maximum at 355 nm) mounted on the reflective surface of polished aluminium which was placed behind the lamps. Twelve glass sheets with immobilised catalyst  $\text{TiO}_2$  (Kete et al. 2014) were fastened around the axis of a special Teflon holder immersed in the centre of the reactor cell.

For the photocatalytic experiments, 5 mg of flubendiamide in 50 mL of acetone ( $100 \text{ mg l}^{-1}$ ) was diluted with distilled water up to 1000 mL. The solution was then filtered and transferred into the reactor cell. The Teflon holder with 12 glass sheets with immobilised  $\text{TiO}_2$  was immersed in the centre of the cell. A gentle purging of oxygen ( $1\text{--}2 \text{ mL min}^{-1}$ ) allowed the mixing of the solution and

the contact of the pesticide with the catalysts. The reaction time was set to 3 h. At different time intervals (5, 15, 30, 45, 60, 90, 120 and 180 min), the samples of 1 mL were taken for HPLC analyses. After 3 h of photocatalytic irradiation, the final solution was extracted with solid-phase extraction (SPE) C<sub>18</sub> cartridges (Supelco, Discovery DSC-18), washed with 5 mL of acetonitrile, dried to dryness, redissolved in dichloromethane and analysed by gas chromatography–mass spectrometry (GC–MS).

### Photocatalytical degradation of flubendiamide in the presence of humic acids and nitrate

To study the influence of nitrates and humic acids on flubendiamide photodegradation, 10 mg of NO<sub>3</sub><sup>-</sup> and 10 mg of humic acids were added to the solution of flubendiamide (10 mg l<sup>-1</sup> in 5% acetone solution). The solutions were then placed in the reactor cell and photocatalytically degraded as described in the paragraph above (photocatalytic experiments).

### Analytical methods

#### Kinetic studies with high-performance liquid chromatography with ultraviolet–visible detector (HPLC–UV–Vis)

Solutions of flubendiamide were analysed by Agilent 1100 HPLC–UV–Vis chromatograph (Agilent Technologies, Palo Alto, CA, USA). The separation was done using Supelco Ascentis<sup>®</sup> Express 5 μm C18 column (5 μm, 150 mm × 4.6 mm). The temperature was set at 25 °C, the injection volume was 20 μL, the mobile phase consisted of 60% acetonitrile and 40% distilled water with the flow rate of 1.0 mL min<sup>-1</sup> and the wavelength was set at 230 nm. For the quantification purposes, the calibration curve ranged from 0.04 to 50.0 mg l<sup>-1</sup> with seven different concentrations in triplicates. The correlation coefficient (*r*<sup>2</sup>) was above 0.99.

#### Product identification with gas chromatography–mass spectrometry (GC–MS)

All data were obtained using high resolution Folded Flight Path (FFP<sup>™</sup>) multiple reflecting geometry time-of-flight (TOF) mass spectrometer Pegasus<sup>®</sup> GC-HRT (LECO Corporation, Saint Joseph, MI, USA) coupled to an Agilent 7890A gas chromatograph (Agilent Technologies, Palo Alto, CA, USA). The system was controlled by the ChromaTOF-HRT<sup>®</sup> software (Version 1.91, LECO Corporation), which was also used for spectra collection and data processing. The data were acquired using ten full (10–800 *m/z* range) spectra per second in high-resolution mode (25,000 or above at *m/z* 218.9851), with high mass accuracy (< 1 ppm), reliably

determining elemental composition of all ions of interest in mass spectra. The multi-point mass calibration of mass spectra with FC-43 (perfluorotributylamine) was performed before running the samples as a part of the automated tuning routine. The mass spectrometer's hardware and acquisition software allows minimising mass drift during data collection. The electron ionisation source temperature was kept at 270 °C, while the electron energy was 70 eV. Chromatographic separation of the samples was performed using an Rxi-5SilMS 30 m × 0.25 mm (id) × 0.25 μm (*df*) (Restek Corporation, Bellefonte, PA) column with a constant helium flow of 1 mL min<sup>-1</sup>. All injection volumes were 1 μL, splitless for 60 s, and thereafter purged with 20 mL min<sup>-1</sup> flow. The septum purge flow was 3 mL min<sup>-1</sup>. The injector and the transfer line temperature were set at 270 and 320 °C, respectively. The oven program was as follows: 0.5 min isothermal at 50 °C, then 10 °C min<sup>-1</sup> ramping to 320 °C and 8 min isothermal hold at 320 °C.

### Results and discussion

The photocatalytic experiments with flubendiamide were carried out by applying gas purging with oxygen and without gas purging. Besides, the influence of nitrates and humic acids on the process was studied.


Generally, the application of gas enables a better mixing of solution and a better contact between the catalyst and the compound that should be degraded. The role of nitrate in the photodegradation of organic pesticides involves the absorption of solar light between 290 and 400 nm and the generation of highly reactive hydroxyl radicals (Méallier 1999). Many examples (benzoylcyclohexanedione herbicides and phenyl urea herbicides), summarised by Remucal (2014), showed that nitrates, besides the dissolved organic matter, can play an important role in the pesticides' removal in nitrate rich natural waters. Our results indicate that the degradation of flubendiamide is more than ten times faster in experiments which included the gas purging, compared to those when the gas purging was turned off (Table 1).

Table 1 shows the first-order reaction rate constants ( $[k] = \text{min}^{-1}$ ) describing the photocatalysis of flubendiamide without and with oxygen purging in solutions with addition of nitrates and humic acids (both 10 mg l<sup>-1</sup>). The correlation coefficients (second column) indicate good correlation between experimental results and fitted curves of the first-order reaction.

Surprisingly, the photocatalysis of flubendiamide under oxygen flow in the presence of nitrates was the same as without added nitrates. During the experiment, the concentration of added nitrates remained the same ( $\pm 5\%$ ). No difference in degradation kinetics was observed (Fig. 1). The same observation was found in the research of Rabindranathan

**Table 1** Kinetic parameters describing the photocatalysis of flubendiamide under oxygen purging, in solution of nitrates and humic acids (HA) (both 10 mg l<sup>-1</sup>) and without gas purging

	Corr. coef.	k (first-order decay)
Purging with O <sub>2</sub>	0.978	0.047 ± 0.016
No gas purging	0.951	0.004 ± 0.001
Addition of NO <sub>3</sub> <sup>-</sup> + O <sub>2</sub> purging	0.998	0.050 ± 0.009
Addition of HA + O <sub>2</sub> purging	0.989	0.267 ± 0.141

**Fig. 1** The degradation curve of flubendiamide under oxygen purging with addition of humic acids enhancing the degradation and addition of nitrate ions, demonstrating no effect

et al. (2003), who studied the photocatalytic degradation of phosphamidon, where in the presence of  $1.5 \times 10^{-4}$  M NO<sub>3</sub><sup>-</sup>, no significant effect under the examined conditions

**Table 2** Identified flubendiamide products of the photocatalytic experiment

Peak #	Molecular formula	R.T. (s)	Area (a.u.)
1	C <sub>11</sub> H <sub>6</sub> F <sub>7</sub> NO	604.008	464,672
2	C <sub>10</sub> H <sub>8</sub> F <sub>7</sub> N	680.448	119,861,554
3	C <sub>10</sub> H <sub>9</sub> F <sub>6</sub> N	711.192	8,493,067
4	C <sub>10</sub> H <sub>8</sub> F <sub>5</sub> N	714.888	4,169,189
5	C <sub>10</sub> H <sub>6</sub> F <sub>7</sub> NO	756.048	296,952
6	C <sub>11</sub> H <sub>8</sub> F <sub>7</sub> NO	883.496	129,024
7	C <sub>8</sub> H <sub>3</sub> IO <sub>3</sub>	1099.86	61,523
8	C <sub>8</sub> H <sub>4</sub> INO <sub>2</sub>	1214.76	1,226,026
9	C <sub>18</sub> H <sub>10</sub> F <sub>7</sub> NO <sub>2</sub>	1260.72	1,184,771
10	C <sub>11</sub> H <sub>9</sub> INO <sub>2</sub>	1468.54	155,226
11	C <sub>18</sub> H <sub>9</sub> IF <sub>7</sub> NO <sub>2</sub>	1475.26	4,817,429
11	C <sub>18</sub> H <sub>9</sub> IF <sub>7</sub> NO <sub>2</sub> —Isomer 11	1520.78	928,348
12	C <sub>13</sub> H <sub>14</sub> INSO <sub>4</sub>	1620.41	24,444,942
13	C <sub>18</sub> H <sub>11</sub> F <sub>7</sub> NO <sub>2</sub>	1756.49	22,981,705
14	C <sub>18</sub> H <sub>10</sub> IF <sub>7</sub> NO <sub>2</sub>	1943.81	10,224,000

**Fig. 2** Proposed degradation pathway of flubendiamide under photocatalytic conditions. Identified products are divided into three groups: Group 1: phthalimide and related phthalic acid derivatives, Group 2: fluorinated species and Group 3: advanced transformation products marked in brackets by numbers of each compound found in Table 2


was observed. Authors reported the decrease in degradation rate which was attributed to the competition of ions for active sites on the TiO<sub>2</sub> surface (Rabindranathan et al. 2003).

On the other hand, the addition of humic acids in photodegradation processes induced more than five times faster transformation of flubendiamide. The irradiation of dissolved organic matter results in arising of different reactive species including hydroxyl, hydroperoxyl and superoxide radical anions, singlet oxygen, excited triplet states of dissolved organic matter, hydrated electrons, carbonate radicals and other radicals (Richard and Canonica 2005). In fact, dissolved organic matter enhanced the photodegradation of the fungicides carboxin and oxycarboxin (Hustert et al. 1999). On the other hand, there are also examples where dissolved organic matter had no effect (anthracene; Bertilsson and Widenfalk 2002) or even an inhibition effect in case of carbofuran (Bachman and Patterson 1999).

Photochemical study, performed by Chen et al. (2010), examined the effect of humic acids (0–25 mg l<sup>-1</sup>) as possible photosensitiser on the photocatalytic mineralisation of dimethoate. Upon 60 min irradiation, the mineralisation efficiency is shown to increase from 63.73 to 78.88% as the humic acids concentration increases from 0 to 5 mg/L. A further increase in humic acids led to a decrease in photocatalytic efficiency. This was attributed to the reduced light transmittance due to the presence of excessive humic acids in reactive solution and the competitive adsorption of humic acids for the active sites on the catalyst surface (Chen et al. 2010).

The degradation pathway in flubendiamide photocatalysis suggested the amide moiety being the most susceptible part in the molecule for the hydroxyl attack. The most abundant degradation products are listed in Table 2. Although the majority of the identified compounds were not present in the available mass spectrometry library (MS library, National Institute of Standards and Technology, NIST14), the identification was rather straightforward, simple and reliable using the accurate mass measurements feature of the Pegasus-HRT instrument (Lebedev et al. 2013) and the known pathways of the transformation of organic compounds under electron ionisation (Lebedev 2015; McLafferty and Turecek 1993).

Table 2 shows the identified flubendiamide photocatalytic products. Consecutive numbers of identified compounds in the chromatogram are presented in the first column, the molecular formulas of identified compounds are represented in the second column, the retention times of identified compounds are mentioned in the third column, while the peak



areas expressed in the arbitrary units (a.u.) are shown in the last column.

The transformation products may be roughly divided into three groups (Fig. 2). The first one consists of the phthalimide and related phthalic acid derivatives (9, 11, 12, 13, 14) (Kushwaha and Kaushik 2016). Fluorinated species related to the second amide moiety represent the second group (1–6). The third one involves advanced transformation products (7, 8, 10). It is worth mentioning that dialkylsulphonyl chain appears to be unstable in reaction conditions, and the corresponding products are not detected in the reaction mixture, possibly degrading rapidly to polar or volatile small molecules.

The process then involves the cleavage of the phthalimide cycle with the formation of aldehydes 13 and 14 or anchimeric assistance of the second nitrogen atom and the formation of phthalimide derivatives. It is worth mentioning a secondary process dealing with deiodination, i.e. substitution of iodine atom in the aromatic ring for a hydrogen atom (compounds 9 and 13). The loss of the second amide moiety (or its part) results in the formation of the advanced transformation products 7, 8, 10. A group of fluorinated alkylbenzene derivatives (1–6) includes iodocyanate, amines, nitroso-derivative, aldehyde and compounds with a shortened heptafluoroisopropyl group.

## Conclusion

It is the first report on the photocatalytic degradation of flubendiamide, novel registered ryanodine insecticide. Flubendiamide transforms into several degradation products. GC–MS proved to be an optimal tool for the identification of the products formed under photocatalytic conditions in oxygen flow, with the addition of nitrates or humic acids. Fourteen novel compounds were identified, and the degradation pathways were proposed. One group of identified products belong to the phthalimide and related phthalic acid derivatives, which are known to express pharmacological activities, and should not be released uncontrolled to the environment.

**Acknowledgements** The authors acknowledge the financial support from the Slovenian Research Agency [research core funding No. P3-0388, Mechanisms of health protection (Mehanizmi varovanja zdravja)]. We acknowledge financial support from Scholarship of the Republic of Slovenia scheme Mobility Grant (CMEPIUS).

## References

Andreozzi R, Caprio V, Insola A, Marotta R (1999) Advanced oxidation processes (AOP) for water purification and recovery. *Catal Today* 53(1):51–59. [https://doi.org/10.1016/S0920-5861\(99\)00102-9](https://doi.org/10.1016/S0920-5861(99)00102-9)

- Bachman J, Patterson HH (1999) Photodecomposition of the carbamate pesticide carbofuran: kinetics and the influence of dissolved organic matter. *Environ Sci Technol* 33(6):874–881. <https://doi.org/10.1021/es9802652>
- Barreiro R, Pratt JR (1992) Toxic effects of chemicals on microorganisms. *Water Environ Res* 64(4):632–641 (ISSN 1061-4303)
- Bertilsson S, Widenfalk A (2002) Photochemical degradation of PAHs in freshwaters and their impact on bacterial growth: influence of water chemistry. *Hydrobiologia* 469:23–32. <https://doi.org/10.1023/A:1015579628189>
- Braham RJ, Harris AT (2009) Review of major design and scale-up considerations for solar photocatalytic reactors. *Ind Eng Chem Res* 48(19):8890–8905. <https://doi.org/10.1021/ie900859z>
- Buddidathia R, Mohapatra S, Siddamallaiah L, Manikrao G, Shankara Hebbar S (2016) Dissipation pattern of flubendiamide residues on capsicum fruit (*Capsicum annum* L.) under field and controlled environmental conditions. *J Environ Sci Health B* 51(1):44–51. <https://doi.org/10.1080/03601234.2015.1080496>
- Castellote M, Bengtsson N (2011) Principles of TiO<sub>2</sub> photocatalysis. In: Ohama Y, Van Gemert D (eds) *Application of titanium dioxide photocatalysis to construction materials*, 1st edn. Springer, Berlin, pp 5–9
- Chen J, Hub Z, Wang D, Gao C, Ji R (2010) Photocatalytic mineralization of dimethoate in aqueous solutions using TiO<sub>2</sub>: parameters and by-products analysis. *Desalination* 258(1–3):28–33. <https://doi.org/10.1016/j.desal.2010.03.053>
- Daghrir R, Drogué P, Didier R (2013) Modified TiO<sub>2</sub> for environmental photocatalytic applications: a review. *Ind Eng Chem Res* 52(10):3581–3599. <https://doi.org/10.1021/ie303468t>
- Das SK, Mukherjee I (2012) Effect of moisture and organic manure on persistence of flubendiamide in soil. *Bull Environ Contam Toxicol* 88:515–520. <https://doi.org/10.1007/s00128-012-0551-9>
- European Food Safety Authority (2013) Conclusion on the peer review of the pesticide risk assessment of the active substance flubendiamide. *EFSA J* 11(9):3298–3360. <https://doi.org/10.2903/j.efs.2013.3298>
- Fenoll J, Vela N, Garrido I, Navarro G, Perez-Lucas G, Navarro S (2015) Reclamation of water polluted with flubendiamide residues by photocatalytic treatment with semiconductor oxides. *Photochem Photobiol* 91:1088–1094. <https://doi.org/10.1111/php.12479>
- Hustert K, Moza PN, Kettrup A (1999) Photochemical degradation of carboxin and oxycarboxin in the presence of humic substances and soil. *Chemosphere* 38(14):3423–3429. [https://doi.org/10.1016/S0045-6535\(98\)00555-4](https://doi.org/10.1016/S0045-6535(98)00555-4)
- Kete M, Pavlica E, Fresno F, Bratina G, Lavrenčič Štangar U (2014) Highly active photocatalytic coatings prepared by a low-temperature method. *Environ Sci Pollut Res* 21:11238–11249. <https://doi.org/10.1007/s11356-014-3077-3>
- Kushwaha N, Kaushik D (2016) Recent advances and future prospects of phthalimide derivatives. *J Appl Pharm Sci* 6(03):159–171. <https://doi.org/10.7324/JAPS.2016.60330>
- Lebedev AT (2015) Mass spectrometry in organic chemistry. *Tehnosfera, Moscow*
- Lebedev AT, Polyakova OV, Mazur DM, Artaev VB (2013) The benefits of high resolution mass spectrometry in environmental analysis. *Analyst* 138:6946–6953. <https://doi.org/10.1039/C3AN01237A>
- Linsebigler AL, Lu G, Yates JT (1995) Photocatalysis on TiO<sub>2</sub> surfaces: principles, mechanisms, and selected results. *Chem Rev* 95(3):735–758. <https://doi.org/10.1021/cr00035a013>
- Masaki T (2008) Study on the mechanism of insecticidal activity through disruption of intracellular calcium homeostasis. *J Pestic Sci* 33(3):271–272. <https://doi.org/10.1584/jpestics.a330301>
- Masaki T, Tohnishi M, Nishimatsu T, Tsubata K, Inoue K, Motoba K, Hirooka T (2005) Flubendiamide, a novel Ca<sup>2+</sup> channel

- modulator, reveals evidence for functional cooperation between  $\text{Ca}^{2+}$  pumps and  $\text{Ca}^{2+}$  release. *Mol Pharmacol* 69(5):1733–1739. <https://doi.org/10.1124/mol.105.020339>
- McLafferty FW, Turecek F (1993) Interpretation of mass spectra. University Science Books, Mill Valley
- Méallier P (1999) Phototransformation of pesticides in aqueous solution. In: Boule P (ed) The handbook of environmental chemistry, Part L: environmental photochemistry, vol 2, 1st edn. Springer, Berlin, pp 241–261
- Mohapatra S, Ahuja AK, Deepa M, Sharma D, Jagadish GK, Rashmi N (2010) Persistence and dissipation of flubendiamide and des-iodo flubendiamide in cabbage (*Brassica oleracea* Linne) and soil. *Bull Environ Contam Toxicol* 85(3):352–356. <https://doi.org/10.1007/s00128-010-0063-4>
- Mohapatra S, Ahuja AK, Deepa M, Jagadish GK, Rashmi N, Sharma D (2011) Development of an analytical method for analysis of flubendiamide, des-iodo flubendiamide and study of their residue persistence in tomato and soil. *Environ Sci Health B* 46(1):264–271. <https://doi.org/10.1080/03601234.2011.540536>
- Paramasivam M, Banerjee H (2012) Degradation dynamics of flubendiamide in different types of soils. *Bull Environ Contam Toxicol* 88:511–514. <https://doi.org/10.1007/s00128-012-0552-8>
- Rabindranathan S, Devipriya S, Yesodharan S (2003) Photocatalytic degradation of phosphamidon on semiconductor oxides. *J Hazard Mater* 102(2):217–229. [https://doi.org/10.1016/S0304-3894\(03\)00167-5](https://doi.org/10.1016/S0304-3894(03)00167-5)
- Remucal CK (2014) The role of indirect photochemical degradation in the environmental fate of pesticides: a review. *Environ Sci Process Impacts* 16:628–653. <https://doi.org/10.1039/c3em00549f>
- Richard C, Canonica S (2005) Aquatic phototransformation of organic contaminants induced by coloured dissolved organic matter. In: Boule P, Bahnemann DW, Robertson PKJ (eds) Environmental photochemistry part II, the handbook of environmental chemistry, 2nd edn. Springer, Berlin, pp 299–323
- Tohnishi M, Nakao H, Furuya T, Seo A, Odama H, Tsubata K, Fujioka S, Kodama S, Hirooka T, Nishimatsu T (2005) Flubendiamide, a novel insecticide highly active against Lepidopterous insect. *Pests J Pestic Sci* 30(4):354–360. <https://doi.org/10.1584/jpestics.30.354>