PUBLISHER CORRECTION



Publisher Correction: Usefulness of ultrasound in diagnosing long head of the biceps tendon malposition in patients with rotator cuff tears

Yoshiko Fujiwara¹ · Syuichi Yamamoto¹ · Yumi Kato¹ · Shimpei Kurata² · Shuhei Fujii² · Kazuya Inoue² · Takashi Inoue³ · Takamitsu Mondori² · Yoshiyuki Nakagawa² · Yasuhito Tanaka⁴

Published online: 15 April 2022

© The Author(s), under exclusive licence to The Japan Society of Ultrasonics in Medicine 2022

Correction to: Journal of Medical Ultrasonics

https://doi.org/10.1007/s10396-022-01200-y

The Publisher regrets the following error in Fig. 1 where the label of the figure was published incorrectly. The correct Fig. 1 is given in this correction.

The original article has been corrected.




Fig. 1 Ultrasonic view of the long head of the biceps tendon (LHBT) with relation to the bicipital groove. a Normal anatomy of the LHBT in the bicipital groove, b Subluxation of the LHBT (\rightarrow) over the

lesser tubercle, **c** Dislocation of the LHBT with complete deviation from the bicipital groove distal LHBT (\rightarrow) , **d** Tear (\rightarrow) of the LHBT

The original article can be found online at https://doi.org/10.1007/ $\,$ s10396-022-01200-y.

- Department of Clinical Laboratory, Uda City Hospital, 815 Haibarahagihara, Uda, Nara 633-0298, Japan
- Department of Orthopaedic Surgery, Uda City Hospital, 815 Haibarahagihara, Uda, Nara 633-0298, Japan
- Department of Evidence-Based Medicine, Institute for Clinical and Translational Science, Nara Medical University Hospital, 840 Shijo-cho, Kashihara, Nara 634-8522, Japan
- Department of Orthopaedic Surgery, Nara Medical University, 840 Shijo-cho, Kashihara, Nara 634-8522, Japan

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

