



Generalized Analytic Integrability of a Class of Polynomial Differential Systems in \mathbb{C}^2

Jaume Llibre¹ · Yuzhou Tian²

Received: 25 April 2020 / Accepted: 21 April 2021 / Published online: 30 April 2021
© The Author(s), under exclusive licence to Springer Nature B.V. 2021

Abstract

This paper study the type of integrability of differential systems with separable variables $\dot{x} = h(x)f(y)$, $\dot{y} = g(y)$, where h , f and g are polynomials. We provide a criterion for the existence of generalized analytic first integrals of such differential systems. Moreover we characterize the polynomial integrability of all such systems.

In the particular case $h(x) = (ax + b)^m$ we provide necessary and sufficient conditions in order that this subclass of systems has a generalized analytic first integral. These results extend known results from Giné et al. (Discrete Contin. Dyn. Syst. 33:4531–4547, 2013) and Llibre and Valls (Discrete Contin. Dyn. Syst., Ser. B 20:2657–2661, 2015). Such differential systems of separable variables are important due to the fact that after a blow-up change of variables any planar quasi-homogeneous polynomial differential system can be transformed into a special differential system of separable variables $\dot{x} = xf(y)$, $\dot{y} = g(y)$, with f and g polynomials.

Mathematics Subject Classification (2010) Primary 37C10 · Secondary 34A34 · 37J35

Keywords Polynomial systems · Generalized analytic integrability · Polynomial first integrals · Residue

1 Introduction and the main results

Planar polynomial differential systems play an important role in the qualitative theory of dynamical systems due to their many applications in physics, chemist, biology, economics, Nowadays the qualitative theory has gained wide development for polynomial systems. For a planar differential system, the existence of a first integral determines completely its global dynamical behavior. So a natural problem arises: Given a polynomial differential system in

✉ Y. Tian
tianyhz3@mail2.sysu.edu.cn

J. Llibre
jllibre@mat.uab.cat

¹ Departament de Matemàtiques, Universitat Autònoma de Barcelona, 08193 Bellaterra, Barcelona, Catalonia, Spain

² School of Mathematics (Zhuhai), Sun Yat-sen University, Zhuhai 519082, P.R. China

\mathbb{R}^2 or \mathbb{C}^2 , how to decide if this system has a first integral? For general polynomial differential systems this problem is very difficult to solve. During the past three decades many mathematicians investigated the integrability of different classes of polynomial differential systems, such as Liénard systems [5, 13, 15, 16], Lotka-Volterra systems [7, 8, 10–12], and quasi-homogeneous polynomial systems [1–4, 6, 17], etc.

Let \mathbb{C} be the set of complex numbers and $\mathbb{C}[x]$ be the ring of all complex polynomials in the variable x . We consider the following complex polynomial differential systems

$$\dot{x} = h(x) f(y), \quad \dot{y} = g(y), \tag{1}$$

where $h \in \mathbb{C}[x]$ and $f, g \in \mathbb{C}[y]$ are coprime. The associated vector field of this system is

$$\mathcal{X} = h(x) f(y) \frac{\partial}{\partial x} + g(y) \frac{\partial}{\partial y}. \tag{2}$$

The integer $d = \max \{ \deg hf, \deg g \}$ is the degree of the vector field \mathcal{X} .

Let U be an open set of \mathbb{C}^2 . A non-locally constant function $H : U \rightarrow \mathbb{C}$ is called a *first integral* of system (1) if it is constant along any solution curve of system (1) contained in U . If $H(x, y)$ is differentiable, then H is a first integral of system (1) if and only if

$$\mathcal{X}H = h(x) f(y) \frac{\partial H}{\partial x} + g(y) \frac{\partial H}{\partial y} = 0 \tag{3}$$

in U .

System (1) has an *analytic first integral* if there exists a first integral $H(x, y)$ which is an analytic function in the variables x and y . A function of the form $\varphi(y) = a \prod_{i=1}^k (y - \alpha_i)^{\gamma_i}$ is called a *product function* with $\alpha_j, \gamma_j \in \mathbb{C}$ and $a \in \mathbb{C} \setminus \{0\}$. The polynomial function $\varphi(y)$ is *square-free* if it can be written as $\varphi(y) = a \prod_{i=1}^k (y - \alpha_i)$ with $a \in \mathbb{C} \setminus \{0\}$, $\alpha_i \neq \alpha_j$ for $i, j = 1, \dots, k$ and $i \neq j$. We say that system (1) has a *generalized analytic first integral* if there exists a first integral $H(x, y)$ which is an analytic function in the variable x whose coefficients are product functions in the variable y .

Let

$$F(z) = \sum_{n=-\infty}^{\infty} a_n (z - z_0)^n$$

be a Laurent expansion at a point z_0 . The coefficient $a_{-1} = \text{Res}[F(z), z_0]$ is the *residue* of $F(z)$ at z_0 .

Differential system (1) of separable variables has a lot of applications. For example Giné et al. in Lemma 2.2 of [6] proved that any planar quasi-homogeneous polynomial differential system can be transformed into a polynomial differential system (1) of the form

$$\dot{x} = xf(y), \quad \dot{y} = g(y), \tag{4}$$

with $f(y), g(y) \in \mathbb{C}[y]$. Hence the study of the type of integrability of the quasi-homogeneous polynomial systems can be reduced to study the type of integrability of their corresponding polynomial systems (4). Note that the polynomial differential systems (4) is a subclass of polynomial differential systems (1). In this paper we generalize some known facts for the systems (4) to systems (1), and provide other new results.

First we present a necessary condition for the existence of generalized analytic first integrals of system (1).

Theorem 1 Assume that $\alpha_1, \dots, \alpha_k$ are the different roots of the polynomial $g(y)$. If system (1) has a generalized analytic first integral, then it must satisfy one of the following two conditions.

- (a) The polynomials $h(x)$ and $g(y)$ are square-free, and $\deg f < \deg g$.
- (b) The roots of the polynomial $h(x)$ are not simple and the $\text{Res}[f(y)/g(y), \alpha_i] = 0$ for all $i = 1, \dots, k$.

The following result is due to Llibre and Valls, see Theorem 1.1 of [14].

Theorem 2 Let $h(x) = x$. Then the polynomial differential system (1) has a generalized analytic first integral if and only if $g(y)$ is square-free, and $\deg f < \deg g$.

In the next theorem we generalize Theorem 2 when $h(x) = (ax + b)^m$, where $a \in \mathbb{C} \setminus \{0\}$ and $m \in \mathbb{N}$. As usual \mathbb{N} denotes the set of positive integers.

Theorem 3 Let $h(x) = (ax + b)^m$ with $a \in \mathbb{C} \setminus \{0\}$ and $m \in \mathbb{N}$. Assume that $\alpha_1, \dots, \alpha_k$ are the different roots of the polynomial $g(y)$. The following statements hold.

- (a) If $m = 1$, then system (1) has a generalized analytic first integral if and only if $g(y)$ is square-free, and $\deg f < \deg g$.
- (b) If $m \geq 2$, then system (1) has a generalized analytic first integral if and only if $\text{Res}[f(y)/g(y), \alpha_i] = 0$ for all $i = 1, \dots, k$.

System (4) has a polynomial first integral if and only if $g(y)$ is square-free, $\deg f < \deg g$ and $\text{Res}[f(y)/g(y), \alpha_j] \in \mathbb{Q}^-$ for $j = 1, \dots, k$, see statement (viii) of Lemma 2.4 of [6]. For the more general polynomial differential system (1) we provide necessary and sufficient conditions for its polynomial integrability in the following theorem.

Theorem 4 Let $\alpha_1, \dots, \alpha_k$ be different roots of the polynomial $g(y)$. System (1) has a polynomial first integral if and only if the two following conditions hold.

- (a) The polynomials $h(x) = ax + b$ and $g(y)$ is square-free, and $\deg f < \deg g$.
- (b) $a \text{Res}[f(y)/g(y), \alpha_j] \in \mathbb{Q}^-$ for $j = 1, \dots, k$.

This paper is organized as follows. We present some preliminary results in Sect. 2. The proofs of Theorems 1, 3 and 4 are given in Sect. 3. In Sect. 4 we illustrate our results with some examples.

2 Preliminaries

In this section we introduce some necessary lemmas for the proof of Theorems 1, 3 and 4. The following lemma can be found in many textbooks, as for instance in [9].

Lemma 5 Assume that $F, G \in \mathbb{C}[y]$ are coprime with $\deg F < \deg G = w$. Let p be the coefficient of the monomial y^w of the polynomial $G(y)$ and q the one of the monomial y^{w-1} of the polynomial $F(y)$.

(a) If y_1, y_2, \dots, y_s are the distinct roots of $G(y)$ with multiplicity n_1, n_2, \dots, n_s , respectively, then

$$\frac{F(y)}{G(y)} = \sum_{i=1}^s \sum_{j=1}^{n_i} \frac{t_{i,j}}{(y - y_i)^j}, \tag{5}$$

where $t_{i,1} = \text{Res}[F(y)/G(y), y_i]$ and $t_{i,n_i} \neq 0$ for $i = 1, \dots, s$.

(b) If $G(y)$ is square-free that is $G(y) = \prod_{i=1}^w (y - y_i)$, then

$$\frac{F(y)}{G(y)} = \sum_{i=1}^w \frac{t_i}{y - y_i}, \tag{6}$$

where

$$t_i = \text{Res}[F(y)/G(y), y_i] \text{ for } i = 1, \dots, w \quad \text{and} \quad \sum_{i=1}^w t_i = q/p. \tag{7}$$

The rational function $F(y)/G(y)$ is a square-free rational function if it satisfies statement (b) of Lemma 5.

Lemma 6 The function $\varphi(y)$ is a product function if and only if $\varphi'(y)/\varphi(y)$ is a square-free rational function.

Proof Necessity. Assume that $\varphi(y)$ is the product function $\varphi(y) = a \prod_{i=1}^k (y - \alpha_i)^{\gamma_i}$. Then

$$\ln \varphi(y) = \ln a + \sum_{i=1}^k \gamma_i \ln (y - \alpha_i). \tag{8}$$

Derivating equation (8) with respect to y , we get that

$$\frac{\varphi'(y)}{\varphi(y)} = \sum_{i=1}^k \frac{\gamma_i}{y - \alpha_i} \tag{9}$$

is a square-free rational function. Hence necessity is proved.

Sufficiency. Since $\varphi'(y)/\varphi(y)$ is a square-free rational function we have equation (9). Integrating equation (9) we get

$$\varphi(y) = a \prod_{i=1}^k (y - \alpha_i)^{\gamma_i},$$

where a is an integration constant. The proof of Lemma 6 is completed. □

Consider $h(x) = \sum_{i=0}^{m-n} h_{n+i} x^{n+i}$ with $m \geq n \geq 1$, $h_{n+i} \in \mathbb{C}$ and $h_m h_n \neq 0$. If system (1) has a generalized analytic first integral $H(x, y)$, then $H(x, y)$ can be written as a power series in x of the form

$$H(x, y) = \sum_{j \geq 0} a_j(y) x^j, \tag{10}$$

where the coefficients $a_j(y)$ are product functions in the variable y . From equation (3) we obtain

$$\begin{aligned} \mathcal{X}H &= \sum_{j \geq 0} j f(y) a_j(y) \left(\sum_{i=0}^{m-n} h_{n+i} x^{n+i} \right) x^{j-1} + \sum_{j \geq 0} g(y) a'_j(y) x^j \\ &= f(y) \sum_{i=0}^{m-n} \sum_{j \geq 0} j h_{n+i} a_j(y) x^{n+i+j-1} + g(y) \sum_{j \geq 0} a'_j(y) x^j = 0. \end{aligned} \tag{11}$$

The equation

$$\sum_{i=0}^{m-n} \sum_{j \geq 0} j h_{n+i} a_j(y) x^{n+i+j-1}$$

can be decomposed into sum of the following equations:

for $i = m - n$, $\sum_{j \geq 0} j h_m a_j(y) x^{m+j-1}$;

for $i = m - n - 1$, $\sum_{j \geq 0} j h_{m-1} a_j(y) x^{m+j-2} = \sum_{j \geq 0} (j + 1) h_{m-1} a_{j+1}(y) x^{m+j-1}$;

for $i = m - n - 2$,

$$\begin{aligned} \sum_{j \geq 0} j h_{m-2} a_j(y) x^{m+j-3} &= \sum_{j \geq 0} (j + 2) h_{m-2} a_{j+2}(y) x^{m+j-1} \\ &+ \sum_{j=0}^1 j h_{m-2} a_j(y) x^{m+j-3}; \end{aligned}$$

for $i = m - n - 3$,

$$\begin{aligned} \sum_{j \geq 0} j h_{m-3} a_j(y) x^{m+j-4} &= \sum_{j \geq 0} (j + 3) h_{m-3} a_{j+3}(y) x^{m+j-1} \\ &+ \sum_{j=0}^2 j h_{m-3} a_j(y) x^{m+j-4}; \end{aligned}$$

for $i = m - n - 4$,

$$\begin{aligned} \sum_{j \geq 0} j h_{m-4} a_j(y) x^{m+j-5} &= \sum_{j \geq 0} (j + 4) h_{m-4} a_{j+4}(y) x^{m+j-1} \\ &+ \sum_{j=0}^3 j h_{m-4} a_j(y) x^{m+j-5}; \end{aligned}$$

⋮

for $i = 2$,

$$\sum_{j \geq 0} j h_{n+2} a_j(y) x^{n+j+1} = \sum_{j \geq 0} (m - n + j - 2) h_{n+2} a_{m-n+j-2}(y) x^{m+j-1} + \sum_{j=0}^{m-n-3} j h_{n+2} a_j(y) x^{n+j+1};$$

for $i = 1$,

$$\sum_{j \geq 0} j h_{n+1} a_j(y) x^{n+j} = \sum_{j \geq 0} (m - n + j - 1) h_{n+1} a_{m-n+j-1}(y) x^{m+j-1} + \sum_{j=0}^{m-n-2} j h_{n+1} a_j(y) x^{n+j};$$

for $i = 0$,

$$\sum_{j \geq 0} j h_n a_j(y) x^{n+j-1} = \sum_{j \geq 0} (m - n + j) h_n a_{m-n+j}(y) x^{m+j-1} + \sum_{j=0}^{m-n-1} j h_n a_j(y) x^{n+j-1};$$

and
$$\sum_{j \geq 0} g(y) a'_j(y) x^j = \sum_{j \geq 0} g(y) a'_{m+j-1}(y) x^{m+j-1} + \sum_{j=0}^{m-2} g(y) a'_j(y) x^j.$$

Then equating the coefficients of x^j in (11) we get the equations that $a_j(y)$ must satisfy:

$$\begin{aligned} a'_j(y) &= 0 \quad \text{for } j = 0, \dots, n - 1; \\ a'_j(y) &= -\frac{f(y)}{g(y)} \sum_{i=1}^{j-n+1} i h_{j-i+1} a_i(y) \quad \text{for } j = n, \dots, m - 2; \\ a'_j(y) &= -\frac{f(y)}{g(y)} \sum_{i=0}^{m-n} (j + i - m + 1) h_{m-i} f(y) a_{j+i-m+1}(y) \quad \text{for } j \geq m - 1. \end{aligned} \tag{12}$$

Remark 7 Note that $a_0(y)$ is a constant. In the following we can assume $a_0(y) = 0$, because a first integral does not depend on the sum of an additional constant.

The solutions of equations (12) are characterized by the following two lemmas.

Lemma 8 Let $h(x) = \sum_{i=0}^{m-n} h_{n+i} x^{n+i}$ with $h_{n+i} \in \mathbb{C}$ and $h_m h_n \neq 0$. Assume that $n = 1$ and that the differential polynomial system (1) has a generalized analytic first integral (10). Then the following statements hold.

(a) There exist polynomials $F_j(u)$ with $\deg F_j = j$ and $F_j(0) = 0$ such that

$$a_j(y) = F_j \left(\exp \left(-h_1 \int \frac{f(y)}{g(y)} dy \right) \right) \quad \text{for all } j \in \mathbb{N}.$$

(b) The polynomial $g(y)$ is square-free and $\deg f < \deg g$.

Proof (a) When $n = 1$ equations (12) can be rewritten as

$$a'_j(y) = -jh_1 \frac{f(y)}{g(y)} a_j(y) - \frac{f(y)}{g(y)} \sum_{i=1}^{j-1} ih_{j-i+1} a_i(y) \quad \text{for } j = 1, \dots, m-2, \quad (13)$$

and

$$a'_j(y) = -jh_1 \frac{f(y)}{g(y)} a_j(y) - \frac{f(y)}{g(y)} \sum_{i=0}^{m-2} (j+i-m+1) h_{m-i} f(y) a_{j+i-m+1}(y), \quad (14)$$

for $j \geq m-1$.

For $m = 2$ we only need to study equation (14) that is

$$a'_j(y) = -jh_1 \frac{f(y)}{g(y)} a_j(y) - (j-1)h_2 \frac{f(y)}{g(y)} a_{j-1}(y), \quad (15)$$

with $j \geq 1$.

When $j = 1$ the solution of equation (15) is

$$a_1(y) = C_1 \exp\left(-h_1 \int \frac{f(y)}{g(y)} dy\right), \quad (16)$$

where C_1 is an integration constant. Obviously $F_1(u) = C_1u$. Hence for $j = 1$ statement (a) holds.

Assume that there exists polynomial $F_j(u)$ with $\deg F_j = j$ and $F_j(0) = 0$ such that

$$a_j(y) = F_j\left(\exp\left(-h_1 \int \frac{f(y)}{g(y)} dy\right)\right).$$

By the induction hypothesis and equation (15) we have

$$a'_{j+1}(y) = -(j+1)h_1 \frac{f(y)}{g(y)} a_{j+1}(y) - jh_2 \frac{f(y)}{g(y)} F_j\left(\exp\left(-h_1 \int \frac{f(y)}{g(y)} dy\right)\right). \quad (17)$$

The solution of the linear differential equation (17) is

$$\begin{aligned} a_{j+1}(y) &= \exp\left(- (j+1)h_1 \int \frac{f(y)}{g(y)} dy\right) \\ &\times \left(C_{j+1} - jh_2 \int \frac{f(y)}{g(y)} \exp\left((j+1)h_1 \int \frac{f(y)}{g(y)} dy\right) dy\right) \\ &\times F_j\left(\exp\left(-h_1 \int \frac{f(y)}{g(y)} dy\right)\right). \end{aligned} \quad (18)$$

Let $u = \exp\left(-h_1 \int \frac{f(y)}{g(y)} dy\right)$. Then

$$\frac{du}{u} = -h_1 \frac{f(y)}{g(y)} dy. \quad (19)$$

So equation (18) can be written as

$$a_{j+1}(y) = F_{j+1}(u) = u^{j+1} \left(C_{j+1} + \frac{j h_2}{h_1} \int \frac{1}{u^{j+2}} F_j(u) du \right). \tag{20}$$

Using the induction hypothesis $\deg F_j = j$ and $F_j(0) = 0$, we get $\deg F_{j+1} = j + 1$ and $F_{j+1}(0) = 0$. The induction is proved and statement (a) follows for $m = 2$.

For $m \geq 3$ we need to consider equations (13) and (14), that is

$$a'_j(y) = -j h_1 \frac{f(y)}{g(y)} a_j(y) - \frac{f(y)}{g(y)} \sum_{i=1}^{j-1} i h_{j+1-i} a_i(y) \quad \text{for } j = 1, \dots, m-2, \tag{21}$$

and

$$a'_j(y) = -j h_1 \frac{f(y)}{g(y)} a_j(y) - \frac{f(y)}{g(y)} \sum_{i=0}^{m-2} (j+i-m+1) h_{m-i} a_{j+i-m+1}(y), \tag{22}$$

for $j \geq m-1$.

For $j = 1$ equation (21) becomes

$$\frac{a'_1(y)}{a_1(y)} = -h_1 \frac{f(y)}{g(y)}. \tag{23}$$

It is easy to get that

$$a_1(y) = C_1 \exp \left(-h_1 \int \frac{f(y)}{g(y)} dy \right),$$

where C_1 is an integration constant. Let $F_1(u) = C_1 u$. So statement (a) holds for $j = 1$.

Assume that for $j = 1, \dots, l$ there exist polynomials $F_j(u)$ with $\deg F_j = j$ such that

$$a_j(y) = F_j \left(\exp \left(-h_1 \int \frac{f(y)}{g(y)} dy \right) \right).$$

Next we consider $j = l + 1$. If $l + 1 \leq m - 2$, then

$$a'_{l+1}(y) = -(l+1) h_1 \frac{f(y)}{g(y)} a_{l+1}(y) - \frac{f(y)}{g(y)} \sum_{i=1}^l i h_{l+2-i} F_i(u), \tag{24}$$

with $u = \exp \left(-h_1 \int \frac{f(y)}{g(y)} dy \right)$. The solution of the linear differential equation (24) is

$$a_{l+1}(y) = u^{l+1} \left(C_{l+1} - \sum_{i=1}^l i h_{l+2-i} \int \frac{f(y)}{g(y)} \frac{F_i(u)}{u^{l+1}} dy \right), \tag{25}$$

with $u = \exp \left(-h_1 \int \frac{f(y)}{g(y)} dy \right)$. From equation (19) it follows that

$$a_{l+1}(y) = F_{l+1}(u) = u^{l+1} \left(C_{l+1} + \sum_{i=1}^l \frac{i h_{l+2-i}}{h_1} \int \frac{F_i(u)}{u^{l+2}} du \right).$$

By the induction hypothesis $\deg F_j = j$ and $F_j(0) = 0$ for $j = 1, \dots, l$, we obtain $\deg F_{l+1} = l + 1$ and $F_{l+1}(0) = 0$.

If $l + 1 \geq m - 1$, then

$$a'_{l+1}(y) = -(l + 1)h_1 \frac{f(y)}{g(y)} a_{l+1}(y) - \frac{f(y)}{g(y)} \sum_{i=0}^{m-2} (l + i - m + 2) h_{m-i} F_{l+i-m+2}(u), \tag{26}$$

with $u = \exp\left(-h_1 \int \frac{f(y)}{g(y)} dy\right)$. By the same arguments as above one can get that

$$a_{l+1}(y) = F_{l+1}(u) = u^{l+1} \left(C_{l+1} + \sum_{i=0}^{m-2} \frac{(l + i - m + 2) h_{m-i}}{h_1} \int \frac{F_{l+i-m+2}(u)}{u^{l+1}} du \right). \tag{27}$$

Applying the induction hypothesis $\deg F_j = j$ and $F_j(0) = 0$ for $j = 1, \dots, l$, we have $\deg F_{l+1} = l + 1$ and $F_{l+1}(0) = 0$. The proof of statement (a) is done.

(b) Let $a_j(y) = \text{constant} = C_j$ for $j \in \mathbb{N}$. From statement (a) we know that there exists a polynomial $F_j(u)$ such that

$$a_j(y) = F_j(u) = C_j \text{ with } u = \exp\left(-h_1 \int \frac{f(y)}{g(y)} dy\right) \text{ and } F_j(0) = 0.$$

Thus $C_j = 0$. Since the first integral $H(x, y)$ is a non-locally constant function, there exists a positive integer j_0 such that $a_{j_0}(y)$ is not a constant and $a_i(y) = 0$ for $i = 1, \dots, j_0 - 1$. Using equations (13) and (14) we have

$$a'_{j_0}(y) = -j_0 h_1 \frac{f(y)}{g(y)} a_{j_0}(y).$$

Consequently

$$a_{j_0}(y) = C_{j_0} \exp\left(-j_0 h_1 \int \frac{f(y)}{g(y)} dy\right), \tag{28}$$

with constant $C_{j_0} \neq 0$. From Lemma 6 we get that $a_{j_0}(y)$ if and only if $f(y)/g(y)$ is a square-free rational function. So $g(y)$ is square-free and $\deg f < \deg g$. This completes the proof of this lemma. □

Lemma 9 Let $h(x) = \sum_{i=0}^{m-n} h_{n+i} x^{n+i}$ with $h_{n+i} \in \mathbb{C}$ and $h_m h_n \neq 0$. Assume that $n \geq 2$ and that the polynomial differential system (1) has a generalized analytic first integral (10), and $\alpha_1, \dots, \alpha_k$ are the different roots of the polynomial $g(y)$. The following statements hold.

(a) There exist polynomials $F_j(u)$ such that

$$a_j(y) = F_j\left(\int \frac{f(y)}{g(y)} dy\right),$$

for $j \geq n$, and $a_j(y)$ are constants for $j = 1, \dots, n - 1$.

(b) Then $\text{Res}[f(y)/g(y), \alpha_i] = 0$ for all $i = 1, \dots, k$.

Proof (a) From equations (12) we have that

$$a'_j(y) = 0 \quad \text{for } j = 1, \dots, n - 1; \tag{29}$$

$$a'_j(y) = -\frac{f(y)}{g(y)} \sum_{i=1}^{j-n+1} i h_{j-i+1} a_i(y) \quad \text{for } j = n, n+1, \dots, m-2; \tag{30}$$

and

$$a'_j(y) = -\frac{f(y)}{g(y)} \sum_{i=0}^{m-n} (j+i-m+1) h_{m-i} a_{j+i-m+1}(y) \quad \text{for } j \geq m-1. \tag{31}$$

Then $a_j(y) = \text{constant} = C_j$ for $j = 1, \dots, n-1$.

For $j = n$ equation (30) can be written as

$$a'_n(y) = -h_n \frac{f(y)}{g(y)} a_1(y) = -h_n C_1 \frac{f(y)}{g(y)}.$$

We get

$$a_n(y) = -h_n C_1 \int \frac{f(y)}{g(y)} dy + C_n,$$

where C_n is an integration constant. Let $F_n(u) = -h_n C_1 u + C_n$. Thus statement (a) holds for $j = n$.

The constants $a_j(y) = C_j$ for $j = 1, \dots, n-1$ can be regarded as polynomials of degree 0. Assume that for $j = 1, \dots, l$ there exist polynomials $F_j(u)$ such that

$$a_j(y) = F_j \left(\int \frac{f(y)}{g(y)} dy \right).$$

If $n \leq l+1 \leq m-2$, then function $a_{l+1}(y)$ satisfy

$$a'_{l+1}(y) = -\frac{f(y)}{g(y)} \sum_{i=1}^{l-n+2} i h_{l-i+2} F_i \left(\int \frac{f(y)}{g(y)} dy \right). \tag{32}$$

Let $u = \int (f(y)/g(y)) dy$. Note that $du = (f(y)/g(y)) dy$. Therefore

$$\begin{aligned} a_{l+1}(y) &= F_{l+1}(u) \\ &= -\sum_{i=1}^{l-n+2} i h_{l-i+2} \int \frac{f(y)}{g(y)} F_i \left(\int \frac{f(y)}{g(y)} dy \right) dy \\ &= -\sum_{i=1}^{l-n+2} i h_{l-i+2} \int F_i(u) du. \end{aligned}$$

If $l+1 \geq m-1$, then

$$a'_{l+1}(y) = -\frac{f(y)}{g(y)} \sum_{i=0}^{m-n} (l+i-m+2) h_{m-i} F_{l+i-m+2}(u).$$

Using similar arguments we obtain

$$a_{l+1}(y) = F_{l+1}(u) = -\sum_{i=0}^{m-n} (l+i-m+2) h_{m-i} \int F_{l+i-m+2}(u) du.$$

Therefore statement (a) is proved.

(b) Suppose that $a_j(y) = \text{constant} = C_j$ for all $j \geq n$. Then the first integral $H(x, y)$ is independent of the variable y . This implies that $\dot{x} = 0$, which is a contradiction. Therefore there exists a positive integer $j_0 \geq n$ such that $a_{j_0}(y)$ is not a constant and $a_i(y) = \text{constant} = C_i$ for $i = 1, \dots, j_0 - 1$. From equations (30) and (31) we obtain that

$$a'_{j_0}(y) = C \frac{f(y)}{g(y)}, \tag{33}$$

where $C = -\sum_{i=1}^{j-n+1} i h_{j-i+1} C_i$ or $\sum_{i=0}^{m-n} (j+i-m+1) h_{m-i} C_{j+i-m+1}$. Hence

$$a_{j_0}(y) = C \int \frac{f(y)}{g(y)} dy + C_{j_0}, \tag{34}$$

with constant $C \neq 0$. Since $a_{j_0}(y)$ is a product function, by Lemma 6, we get that

$$\frac{C}{C \int \frac{f(y)}{g(y)} dy + C_{j_0}} \frac{f(y)}{g(y)}$$

is a square-free rational function. This implies that

$$\int \frac{f(y)}{g(y)} dy \tag{35}$$

is a rational function.

We know that there exist two polynomials $p(y), r(y) \in \mathbb{C}[y]$ such that

$$f(y) = p(y)g(y) + r(y) \quad \text{and} \quad \deg r < \deg g.$$

The polynomial $r(y)$ cannot be zero due to the fact that $f(y)$ and $g(y)$ are coprime. Consequently

$$\int \frac{f(y)}{g(y)} dy = Q(y) + \int \frac{r(y)}{g(y)} dy, \tag{36}$$

with $Q'(y) = p(y)$. Assume that $\alpha_1, \dots, \alpha_k$ are the distinct roots of $g(y)$ with multiplicity n_1, n_2, \dots, n_k , respectively. Using Lemma 5 $r(y)/g(y)$ can be expressed as

$$\frac{r(y)}{g(y)} = \sum_{i=1}^k \sum_{j=1}^{n_i} \frac{c_{i,j}}{(y - \alpha_i)^j}, \tag{37}$$

where $c_{i,n_i} \neq 0$ for $i = 1, \dots, k$. Thus

$$\int \frac{r(y)}{g(y)} dy = \ln \left(\prod_{i=1}^k (y - \alpha_i)^{c_{i,1}} \right) + \sum_{i=1}^k \sum_{j=2}^{n_i} \int \frac{c_{i,j}}{(y - \alpha_i)^j} dy. \tag{38}$$

Since equation (35) is a rational function and $Q(y)$ (see equation (36)) is polynomial, equation (38) is also a rational function. Note that $j \geq 2$ in equation (38). This implies that

$$\int \frac{c_{i,j}}{(y - \alpha_i)^j} dy$$

is a rational function. Then $c_{i,1}$ must be 0, that is $\text{Res}[f(y)/g(y), \alpha_i] = 0$ for all $i = 1, \dots, k$. The proof is done. \square

3 Proofs of Theorems 1, 3 and 4

The main purpose of this section is to prove Theorems 1, 3 and 4.

Proof of Theorem 1 We claim that if $h(x)$ simultaneously has simple roots and multiple roots, then system (1) has no generalized analytic first integral.

Let β_1 and β_2 be a simple root and a root of multiplicity n of $h(x)$ with $n \geq 2$, respectively. Assume that system (1) has a generalized analytic first integral. Doing the change of variables $(x, y, t) \mapsto (x + \beta_1, y, t)$, system (1) becomes

$$\dot{x} = \tilde{h}(x) f(y), \quad \dot{y} = g(y), \tag{39}$$

where $\tilde{h}(x) = \sum_{i=1}^m \tilde{h}_i x^i$ with $\tilde{h}_i \in \mathbb{C}$ and $\tilde{h}_m \tilde{h}_1 \neq 0$. Since system (1) has a generalized analytic first integral, system (39) also has a generalized analytic first integral. By Lemma 8 we get that $g(y)$ is square-free and $\deg f < \deg g$. This means that $\text{Res}[f(y)/g(y), \alpha_i] \neq 0$ for all $i = 1, \dots, k$.

Under the transformation $(x, y, t) \mapsto (x + \beta_2, y, t)$ system (1) changes to

$$\dot{x} = \bar{h}(x) f(y), \quad \dot{y} = g(y), \tag{40}$$

where $\bar{h}(x) = \sum_{i=n}^m \bar{h}_i x^i$ with $\bar{h}_i \in \mathbb{C}$ and $\bar{h}_m \bar{h}_n \neq 0$. From Lemma 9 it follows that

$$\text{Res}[f(y)/g(y), \alpha_i] = 0$$

for all $i = 1, \dots, k$. This is in contradiction with $\text{Res}[f(y)/g(y), \alpha_i] \neq 0$ for all $i = 1, \dots, k$. So the claim is proved.

In summary, the polynomial $h(x)$ is square-free or it has no simple roots. If $h(x)$ is square-free, using Lemma 8, we obtain statement (a). If $h(x)$ has no simple roots, by Lemma 9, statement (b) holds. This completes the proof of the theorem. \square

Proof of Theorem 3 Doing the change of variables $(x, y, t) \mapsto ((x - b)/a, y, t/a)$, system (1) becomes

$$\dot{x} = x^m f(y), \quad \dot{y} = \frac{1}{a} g(y). \tag{41}$$

(a) From Theorem 2 it follows that statement (a) holds.

(b) *Necessity.* Using statement (a) of Theorem 1 the necessity is obvious.

Sufficiency. It is sufficient to show that system (41) has a generalized analytic first integral. Assume that $\alpha_1, \dots, \alpha_k$ are the distinct roots of $g(y)$ with multiplicity n_1, n_2, \dots, n_k , respectively. There exist two polynomials $p(y), r(y) \in \mathbb{C}[y]$ such that

$$f(y) = p(y) g(y) + r(y) \quad \text{and} \quad \deg r < \deg g.$$

The polynomial $r(y)$ cannot be zero due to the fact that $f(y)$ and $g(y)$ are coprime.

By Lemma 5 we have

$$\frac{f(y)}{g(y)} = p(y) + \frac{r(y)}{g(y)} = p(y) + \sum_{i=1}^k \sum_{j=1}^{n_i} \frac{c_{i,j}}{(y - \alpha_i)^j}, \tag{42}$$

where $c_{i,n_i} \neq 0$ for $i = 1, \dots, k$. Since $\text{Res}[f(y)/g(y), \alpha_i] = 0$ that is $c_{i,1} = 0$ for all $i = 1, \dots, k$, we obtain

$$P(y) := \int \frac{f(y)}{g(y)} dy = \int p(y) dy + \sum_{i=1}^k \sum_{j=2}^{n_i} \int \frac{c_{i,j}}{(y - \alpha_i)^j} dy. \tag{43}$$

Note that $j \geq 2$ in equation (43). Thus $P(y)$ is a rational function, that is, a product function. Now we show that

$$H(x, y) = \frac{(m - 1)x^{m-1}}{1 + a(m - 1)x^{m-1}P(y)} \tag{44}$$

is a generalized analytic first integral of system (41). Doing simple computations we have

$$\frac{\partial H}{\partial x} = \frac{(m - 1)^2 x^{m-2}}{(1 + a(m - 1)x^{m-1}P(y))^2} \quad \text{and} \quad \frac{\partial H}{\partial y} = -\frac{a(m - 1)^2 x^{2m-2}}{(1 + a(m - 1)x^{m-1}P(y))^2} \frac{f(y)}{g(y)}.$$

Therefore

$$\mathcal{X}H = x^m f(y) \frac{\partial H}{\partial x} + \frac{1}{a} g(y) \frac{\partial H}{\partial y} = 0.$$

Moreover $H(x, y)$ can be written as a power series in x

$$H(x, y) = \frac{(m - 1)x^{m-1}}{1 - a(1 - m)x^{m-1}P(y)} = (m - 1)x^{m-1} \sum_{j \geq 0} a^j (1 - m)^j P^j(y) x^{(m-1)j}.$$

This completes the proof of the theorem. □

Proof of Theorem 4 Necessity. We claim that $h(x)$ is square-free.

Let β_1, \dots, β_l be different roots of the polynomial $h(x)$. Suppose that β is an arbitrary root of the polynomial $h(x)$ with multiplicity n . By changing the variables $(x, y, t) \mapsto (x + \beta, y, t)$, system (1) is equivalent to

$$\dot{x} = \tilde{h}(x) f(y), \quad \dot{y} = g(y), \tag{45}$$

where $\tilde{h}(x) = \sum_{i=n}^m \tilde{h}_i x^i$ with $\tilde{h}_i \in \mathbb{C}$ and $\tilde{h}_m \tilde{h}_n \neq 0$. Note that $\tilde{h}_n = h^{(n)}(\beta)/n!$. Since system (1) has a polynomial first integral, system (45) also has a polynomial first integral, that is

$$H(x, y) = \sum_{j \geq 0} a_j(y) x^j, \tag{46}$$

where $a_j(y)$ are polynomials. Obviously $H(x, y)$ is a generalized analytic first integral. From the proof of Lemmas 8 and 9 we know that there exists a positive integer j_0 such that $a_{j_0}(y)$ is not a constant, and $a_i(y) = \text{constant} = C_i$ for $i = 1, \dots, j_0 - 1$.

Assume that $n \geq 2$. From the proof of statement (b) of Lemma 9, we get

$$a_{j_0}(y) = C \int \frac{f(y)}{g(y)} dy + C_{j_0}, \tag{47}$$

with constant $C \neq 0$ (see equation (34)). By equation (38) $a_{j_0}(y)$ is not a polynomial. Thus β is a simple root of $h(x)$, that is $n = 1$. Using Theorem 1 we obtain that the polynomials $h(x)$ and $g(y)$ are square-free, and $\deg f < \deg g$. Hence the claim is proved.

From the proof of statement (b) of Lemma 8 we have

$$a_{j_0}(y) = C_{j_0} \exp\left(-j_0 \tilde{h}_1 \int \frac{f(y)}{g(y)}\right), \tag{48}$$

with $\tilde{h}_1 = h'(\beta)$ and the constant $C_{j_0} \neq 0$ (see equation (28)).

Applying Lemma 5 $f(y)/g(y)$ can be expressed as

$$\frac{f(y)}{g(y)} = \sum_{j=1}^k \frac{\mu_j}{y - \alpha_j}, \tag{49}$$

where $\mu_j = \text{Res}[f(y)/g(y), \alpha_j]$ for $j = 1, \dots, k$. Therefore

$$a_{j_0}(y) = C_{j_0} \prod_{j=1}^k (y - \alpha_j)^{-j_0 \tilde{h}_1 \mu_j}. \tag{50}$$

Since $a_{j_0}(y)$ is a polynomial we have $\tilde{h}_1 \mu_j \in \mathbb{Q}^-$ for all $j = 1, \dots, k$. Note that β is an arbitrary root of the polynomial $h(x)$. Thus

$$h'(\beta_i) \cdot \text{Res}[f(y)/g(y), \alpha_j] \in \mathbb{Q}^- \text{ for } i = 1, \dots, l \text{ and } j = 1, \dots, k. \tag{51}$$

This means that $h'(\beta_1)/h'(\beta_i) \in \mathbb{Q}^+$ for $i = 1, \dots, l$.

Assume that $l \geq 2$. Using statement (b) of Lemma 5 $1/h(x)$ can be written as

$$\frac{1}{h(x)} = \sum_{i=1}^l \frac{t_i}{x - \beta_i},$$

with $t_i = \text{Res}[1/h(x), \beta_i] = 1/h'(\beta_i) \neq 0$. From equation (7) we obtain

$$\sum_{i=1}^l t_i = \sum_{i=1}^l \frac{1}{h'(\beta_i)} = 0. \tag{52}$$

One can get

$$\sum_{i=1}^l \frac{h'(\beta_1)}{h'(\beta_i)} = 0,$$

which is in contradiction with $h'(\beta_1)/h'(\beta_i) \in \mathbb{Q}^+$ for $i = 1, \dots, l$. So $l = 1$, that is $h(x) = ax + b$ with $a \in \mathbb{C} \setminus \{0\}$. Then equation (51) becomes

$$a \text{Res}[f(y)/g(y), \alpha_j] \in \mathbb{Q}^- \text{ for } j = 1, \dots, k.$$

This proves the necessity.

Sufficiency. Let $\mu_j = \text{Res} [f(y)/g(y), \alpha_j]$ and consider

$$\tilde{H}(x, y) = \left(x + \frac{b}{a}\right)^{\frac{1}{a}} \left(\prod_{j=1}^k (y - \alpha_j)^{-\mu_j}\right). \tag{53}$$

Since $a\mu_j \in \mathbb{Q}^-$ there exists a positive integer N such that

$$H(x, y) = (\tilde{H}(x, y))^{aN} = \left(x + \frac{b}{a}\right)^N \left(\prod_{j=1}^k (y - \alpha_j)^{-\mu_j aN}\right) \tag{54}$$

is a polynomial. Next we show that polynomial (54) is a first integral of system (1). In fact it is sufficient to prove that $\tilde{H}(x, y)$ is a first integral of system (1).

Straightforward computations show that

$$\frac{\partial \tilde{H}}{\partial x} = \frac{\tilde{H}(x, y)}{ax + b}, \tag{55}$$

and

$$\frac{\partial \tilde{H}}{\partial y} = -\tilde{H}(x, y) \left(\sum_{j=1}^k \frac{\mu_j}{y - \alpha_j}\right). \tag{56}$$

The polynomial $g(y)$ is square-free with $\deg f < \deg g$. Using Lemma 5 we have

$$\frac{f(y)}{g(y)} = \sum_{j=1}^k \frac{\mu_j}{y - \alpha_j}. \tag{57}$$

Equation (56) can be written as

$$\frac{\partial \tilde{H}}{\partial y} = -\tilde{H}(x, y) \frac{f(y)}{g(y)}.$$

Thus

$$x\tilde{H} = (ax + b) f(y) \frac{\partial \tilde{H}}{\partial x} + g(y) \frac{\partial \tilde{H}}{\partial y} = 0.$$

That is $\tilde{H}(x, y)$ is a first integral of system (1). This completes the proof of the theorem. \square

4 Examples

In this section we present some applications of our results.

Example 10 Consider the differential system

$$\dot{x} = (x - 1)(x - 2)^2 y, \quad \dot{y} = y + 1. \tag{58}$$

It has a first integral

$$H(x, y) = \frac{x-2}{(x-1)(y+1)} \exp\left(y + \frac{1}{x-2}\right).$$

By Theorem 1 system (58) has no generalized analytic first integral, because $h(x) = (x-1)(x-2)^2$ simultaneously has simple roots and multiple roots.

Example 11 Consider the differential system

$$\dot{x} = (2x-1)^m (y^3 - 8y^2 + 29y - 26), \quad \dot{y} = (y-3)^3 (y+1)^2, \quad (59)$$

with $m \in \mathbb{N}$ and $m \geq 2$. For this system we have $f(y) = y^3 - 8y^2 + 29y - 26$, $g(y) = (y-3)^3 (y+1)^2$, $\alpha_1 = -1$ and $\alpha_2 = 3$. So $\text{Res}[f(y)/g(y), \alpha_1] = \text{Res}[f(y)/g(y), \alpha_2] = 0$. Applying Theorem 3 system (59) has the generalized analytic first integral (see equation (44))

$$H(x, y) = \frac{2(m-1)(y-1)(y-3)^2(2x-1)^{m-1}}{(y+1)(y-3)^2 - (m-1)(2y^2 - 11y + 19)(2x-1)^{m-1}}.$$

Example 12 Consider the differential system

$$\dot{x} = (5x-1) \left(3\sqrt[3]{3} + 2\sqrt{2} - 5y\right), \quad \dot{y} = 6 \left(y - \sqrt{2}\right) \left(y - \sqrt[3]{3}\right). \quad (60)$$

Using the notations of Theorem 4 we get that $g(y) = 6(y - \sqrt{2})(y - \sqrt[3]{3})$ is square-free, $\alpha_1 = \sqrt{2}$, $\alpha_2 = \sqrt[3]{3}$, $h(x) = 5x - 1$ and $f(y) = 3\sqrt[3]{3} + 2\sqrt{2} - 5y$. For this system we have

$$5\text{Res}[f(y)/g(y), \alpha_1] = -\frac{5}{2} \quad \text{and} \quad 5\text{Res}[f(y)/g(y), \alpha_2] = -\frac{5}{3}.$$

By Theorem 4 system (60) has the polynomial first integral

$$H(x, y) = \left(x - \frac{1}{5}\right)^6 \left(y - \sqrt{2}\right)^{15} \left(y - \sqrt[3]{3}\right)^{10}.$$

Acknowledgements The first author is partially supported by the Ministerio de Ciencia, Innovación y Universidades, Agencia Estatal de Investigación grants M TM2016-77278-P (FEDER), the Agència de Gestió d'Ajuts Universitaris i de Recerca grant 2017SGR1617, and the H2020 European Research Council grant MSCA-RISE-2017-777911. The second author is partially supported by the National Natural Science Foundation of China (No. 11971495 and No. 11801582), China Scholarship Council (No. 201906380022) and Guangdong Basic and Applied Basic Research Foundation (No. 2019A1515011239).

References

1. Algaba, A., Freire, E., Gamero, E., García, C.: Monodromy, center-focus and integrability problems for quasi-homogeneous polynomial systems. *Nonlinear Anal.* **72**, 1726–1736 (2010)
2. Algaba, A., García, C., Giné, J.: Analytic integrability of some examples of degenerate planar vector fields. *Acta Appl. Math.* **141**, 1–15 (2016)
3. Algaba, A., García, C., Reyes, M.: Rational integrability of two-dimensional quasi-homogeneous polynomial differential systems. *Nonlinear Anal.* **73**, 1318–1327 (2010)

4. García, B., Llibre, J., Pérez del Río, J.S.: Planar quasi-homogeneous polynomial differential systems and their integrability. *J. Differ. Equ.* **255**, 3185–3204 (2013)
5. Gasull, A., Giné, J.: Integrability of Liénard systems with a weak saddle. *Z. Angew. Math. Phys.* **68**(13), 7 (2017)
6. Giné, J., Grau, M., Llibre, J.: Polynomial and rational first integrals for planar quasi-homogeneous polynomial differential systems. *Discrete Contin. Dyn. Syst.* **33**, 4531–4547 (2013)
7. Giné, J., Romanovski, V.G.: Integrability conditions for Lotka-Volterra planar complex quintic systems. *Nonlinear Anal., Real World Appl.* **11**, 2100–2105 (2010)
8. Gravel, S., Thibault, P.: Integrability and linearizability of the Lotka-Volterra system with a saddle point with rational hyperbolicity ratio. *J. Differ. Equ.* **184**, 20–47 (2002)
9. Lang, S.: *Complex Analysis*, fourth edition. Graduate Texts in Mathematics, vol. 103. Springer, New York (1999).
10. Liu, C., Chen, G., Li, C.: Integrability and linearizability of the Lotka-Volterra systems. *J. Differ. Equ.* **198**, 301–320 (2004)
11. Liu, C., Li, Y.: The integrability of a class of cubic Lotka-Volterra systems. *Nonlinear Anal., Real World Appl.* **19**, 67–74 (2014)
12. Llibre, J., Ramírez, R., Ramírez, V.: Integrability of a class of N -dimensional Lotka-Volterra and Kolmogorov systems. *J. Differ. Equ.* **269**(13), 2503–2531 (2020). <https://doi.org/10.1016/j.jde.2020.02.001>
13. Llibre, J., Valls, C.: On the local analytic integrability at the singular point of a class of Liénard analytic differential systems. *Proc. Am. Math. Soc.* **138**, 253–261 (2010)
14. Llibre, J., Valls, C.: Analytic integrability of a class of planar polynomial differential systems. *Discrete Contin. Dyn. Syst., Ser. B* **20**, 2657–2661 (2015)
15. Llibre, J., Valls, C.: Liouvillian first integrals for a class of generalized Liénard polynomial differential systems. *Proc. R. Soc. Edinb., Sect. A* **146**, 1195–1210 (2016)
16. Llibre, J., Valls, C.: On the analytic integrability of the Liénard analytic differential systems. *Discrete Contin. Dyn. Syst., Ser. B* **21**, 557–573 (2016)
17. Llibre, J., Zhang, X.: Polynomial first integrals for quasi-homogeneous polynomial differential systems. *Nonlinearity* **15**, 1269–1280 (2002)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.