



EIGENPAIRS OF A FAMILY OF TRIDIAGONAL MATRICES: THREE DECADES LATER

C. M. DA FONSECA^{1,2,*} and V. KOWALENKO³

¹Kuwait College of Science and Technology, Doha District, Block 4, P.O. Box 27235, Safat 13133, Kuwait

e-mail: c.dafonseca@kcst.edu.kw

²University of Primorska, FAMNIT, Glagoljska 8, 6000 Koper, Slovenia

e-mail: carlos.dafonseca@famnit.upr.si

³Department of Mathematics and Statistics, University of Melbourne, Victoria 3010, Australia

e-mail: vkowa@unimelb.edu.au

(Received April 3, 2019; revised May 13, 2019; accepted May 13, 2019)

Abstract. This survey paper summarizes the more important recent applications of the eigenpairs formulas for a family of tridiagonal matrices based on Losonczi's seminal work of almost thirty years ago, which not only seems to have been largely ignored, but has also been re-cast or re-discovered in alternative guises by various authors since. In the course of presenting these applications, we also make contact with earlier more specific applications where Losonczi's work could have been applied to yield the results more quickly. Many of the recent applications in physics and engineering cite less general work, which followed Losonczi more than a decade later.

1. Introduction

For almost a century there has been an active interest/fascination in tridiagonal matrices. For the first 75 years since the pioneering paper of Egerváry and Szász [13], the interest was primarily mathematical where the aim was to develop the spectral theory, viz. the determinants, eigenvalues and eigenvectors, for increasingly more general and complicated tridiagonal matrices. This culminated in the most general class of these matrices studied by Losonczi in [33]. As we shall observe here, over the last two decades there has been an explosion in applications of tridiagonal matrices to important topics in applied mathematics, physics and technology/engineering. In fact, we cannot do due justice to all these applications in this brief survey.

* Corresponding author.

Key words and phrases: tridiagonal matrix, characteristic polynomial, eigenvalue, Chebyshev polynomial of second kind.

Mathematics Subject Classification: 15A18, 15B05.

Thus we have been restricted to presenting only a sample of them. Unfortunately, most of these applications refer to the more recent work of Yueh [52] without recognizing the more general and powerful work of Losonczi that preceded a decade earlier. Even Yueh did not cite Losonczi's work. The aim of the present paper is to highlight the issue in order to give Losonczi the recognition he justly deserves.

2. A class of tridiagonal matrices

Based on an earlier work [34], Losonczi [33] in 1990 introduced and studied the following general family of $n \times n$ complex tridiagonal matrices:

$$(1) \quad M_{n,k} = \begin{pmatrix} a & & & & & & & & c \\ & \ddots & & & & & & & \ddots \\ & & a & & & & & & \ddots \\ \hline d & & & & & & & & 0 \\ & \ddots & & & & & & & \ddots \\ & & \ddots & & & & & & 0 \\ & & & & & & & & c \\ & & & & & & & & \ddots \\ & & & & & & & & \ddots \\ & & & & & & & & d \\ & & & & & & & & \ddots \\ & & & & & & & & \ddots \\ & & & & & & & & b \\ & & & & & & & & \ddots \\ & & & & & & & & \ddots \\ & & & & & & & & b \end{pmatrix}_{n \times n} .$$

In this form where $n - 2k \geq 0$, the upper left and bottom right corners represent $k \times k$ principal submatrices. For the case of $n - 2k < 0$, the matrices took the following form:

$$(2) \quad M_{n,k} = \begin{pmatrix} a & & & & & & & & c \\ & \ddots & & & & & & & \ddots \\ & & a & & & & & & c \\ \hline & & & a + b & & & & & \\ & & & & \ddots & & & & \\ & & & & & a + b & & & \\ \hline d & & & & & & & & b \\ & \ddots & & & & & & & \ddots \\ & & d & & & & & & \ddots \\ & & & & & & & & \ddots \\ & & & & & & & & \ddots \\ & & & & & & & & b \end{pmatrix}_{n \times n} ,$$

where the principal submatrices in the upper left and lower right corners are now $(n - k) \times (n - k)$. The main reason for studying these matrices is

that they arise: (i) in determining the discrete quadratic inequalities of the Wirtinger type [15] and (ii) when estimating specific coefficients in trigonometric polynomials [13,21,43]. From this study Losonczi was able to derive general formulas for the eigenpairs of $M_{n,k}$, resulting in explicit formulas for several special cases. In what follows we shall without loss of generality only consider matrices of the first form, viz. (1). The corresponding results for matrices of the form in (2) follow naturally and are left as an exercise for the reader.

If we put $n = qk + r$, with $r \in \{0, 1, \dots, k - 1\}$, and let $P_{n,k}(x)$ represent the characteristic polynomial of $M_{n,k}$, then according to Theorem 1 in [33], we find that

$$(3) \quad P_{n,k}(x) = P_{q+1,1}(x)^r P_{q,1}(x)^{k-r}.$$

This means that the eigenvalues of $M_{n,k}$ can be found from the spectral analysis of the following matrix:

$$(4) \quad M_{\ell,1} = \begin{pmatrix} a & c & & & & \\ d & 0 & c & & & \\ & d & \ddots & \ddots & & \\ & & \ddots & \ddots & c & \\ & & & d & 0 & c \\ & & & & d & b \end{pmatrix}_{\ell \times \ell}.$$

Moreover, from (7) in Theorem 2 of the same reference, we have

$$(5) \quad P_{n,1}(x) = (\sqrt{cd})^n \left(U_n \left(\frac{x}{2\sqrt{cd}} \right) - \frac{a+b}{\sqrt{cd}} U_{n-1} \left(\frac{x}{2\sqrt{cd}} \right) + \frac{ab}{cd} U_{n-2} \left(\frac{x}{2\sqrt{cd}} \right) \right).$$

where $\{U_n\}_{n \geq 0}$ are Chebyshev polynomials of the second kind. For $n \geq 1$, these orthogonal polynomials, which have been studied extensively since their discovery in the 1850's (for example, see [10,42]), satisfy the three-term recurrence relation given by

$$(6) \quad 2xU_n(x) = U_{n+1}(x) + U_{n-1}(x), \quad \text{for } n \geq 1,$$

subject to the initial values $U_0(x) = 1$ and $U_1(x) = 2x$. One of the most famous formulas for $U_n(x)$, valid for all values of n , is

$$U_n(x) = \frac{\sin(n + 1)\theta}{\sin \theta},$$

where $x = \cos \theta$ and $0 \leq \theta < \pi$.

In regard to eigenvectors, if $\lambda = 2\sqrt{cd}\cos\theta$ represents an eigenvalue of $M_{n,1}$, then according to Theorem 3 of [33] the corresponding eigenvectors can be expressed as $u = (u_0, u_1, \dots, u_{n-1})$, where

$$u_\ell = \left(\frac{\sqrt{c}}{\sqrt{d}}\right)^\ell \left(\sin(\ell+1)\theta + \frac{a}{b}\sin\ell\theta\right), \quad \text{for } \ell = 0, 1, \dots, n-1.$$

We present here a survey of some of the most important recent papers that deal with the applications and theory of these tridiagonal matrices, which continue to this day to fascinate scientists in engineering and physics. As part of this survey, we shall discuss important references preceding Losonczi's seminal work as far back as 1928, which deal with more specific or less general matrices than those studied by Losonczi. At the same time we shall cite applications up to the present day in order to demonstrate that these matrices represent a hive of activity in different fields of science and technology. It should be emphasised that the list of publications provided in this survey is by no means exhaustive, but should give the reader a taste of just how important Losonczi's work is.

In the following sections we shall survey several earlier papers containing similar results to Losonczi, then present several other versions of his work, where the authors were unaware of [33] and finally discuss the most recent applications, where the results of [33] emerge in different fields of activity, but are not duly acknowledged. Before doing so, however, we need to make some preliminary remarks to help the reader to understand the later material.

3. Preliminary remarks

First, it is obvious that spectral theory when applied to (1) and especially (4) reduces them to the symmetric case where b and c are replaced by \sqrt{bc} . Moreover, if we regard a tridiagonal matrix as the adjacency matrix of a path, then the off-diagonal entries become the weights of each edge. Similarly, one can regard (1) as the (weight) adjacency matrix of two types of path. For more details, the reader is referred to [19].

Second, when we replace d by 1 and c by cd in (1), the resulting matrix is known in the theory of orthogonal polynomials theory as a monic Jacobi matrix [4]. Obviously, the characteristic polynomial is invariant, while the signs of c and d are completely redundant. Nevertheless, they are retained in many examples here for convenience.

4. Before Losonczi

In 1945 Rutherford [44] derived (5) for the case of $c = d = 1$ with a and b , arbitrary. As a consequence, he was able to obtain the eigenvalues for

$$(7) \quad \begin{pmatrix} 1 & 1 & & & \\ & 1 & 0 & \ddots & \\ & & \ddots & \ddots & 1 \\ & & & & 1 & 0 \end{pmatrix}_{n \times n}.$$

These are in turn given by

$$\lambda_\ell = -2 \cos\left(\frac{2\ell\pi}{2n+1}\right),$$

where $\ell = 1, 2, \dots, n$. For the matrix

$$(8) \quad \begin{pmatrix} 1 & 1 & & & \\ & 1 & 0 & 1 & \\ & & 1 & \ddots & \ddots \\ & & & \ddots & \ddots & 1 \\ & & & & 1 & 0 & 1 \\ & & & & & 1 & 1 \end{pmatrix}_{n \times n}$$

he found that the eigenvalues are given by

$$\lambda_\ell = 2 \cos\left(\frac{\ell\pi}{n}\right), \quad \text{for } \ell = 1, 2, \dots, n.$$

Several years later, in his Master's thesis [14] Elliott not only re-derived Rutherford's results by providing a more detailed exposition, but also included the matrix

$$(9) \quad \begin{pmatrix} 1 & 1 & & & \\ & 1 & 0 & 1 & \\ & & 1 & \ddots & \ddots \\ & & & \ddots & \ddots & 1 \\ & & & & 1 & 0 & 1 \\ & & & & & 1 & -1 \end{pmatrix}_{n \times n},$$

whose eigenvalues were found to be

$$\lambda_\ell = 2 \cos\left(\frac{(2\ell-1)\pi}{2n}\right), \quad \text{for } \ell = 1, 2, \dots, n.$$

Although outside the scope of this work, it should be pointed out that the matrices studied by Rutherford in [44] were motivated by various problems in physics and chemistry and have since had a tremendous impact in modern research areas such as nanotechnology.

Another motivation for Losonczi's study of his more general tridiagonal matrices was the pioneering paper of E. Egerváry and O. Szász [13] on multivariable operator theory, free pluriharmonic functions and trigonometric polynomials. In this work the authors related their study to the case of $a = b = 0$ in $M_{n,k}$. As a consequence, they derived (3) by rearranging the rows and columns.

Losonczi was able to generalize all the above-mentioned results via one elegant and powerful tridiagonal matrix. Moreover, in addition to calculating the eigenvalues for the matrix, he was able to derive formulas for the corresponding eigenvectors. In the following sections we shall turn our attention to more recent literature, where his results have emerged, but have not been cited. It should be mentioned that the list of applications presented here is by no means exhaustive.

5. Applications

Matrices of the form of (7) are structurally simple but are, nevertheless, rich in applications. Our survey of important examples begins with the description of a major result in [5]. There the authors determine the singular eigenvalues of a Jordan block denoted by $J_n(\mu)$. To determine these values, they present the product of Jordan blocks, one of which is the complex conjugate. Specifically, they consider

$$J_n^*(\mu)J_n(\mu) = \begin{pmatrix} r^2 & r & & & \\ r & 1+r^2 & \ddots & & \\ & \ddots & \ddots & r & \\ & & r & 1+r^2 & \\ & & & & \ddots \end{pmatrix}_{n \times n},$$

where $\mu = re^{i\theta}$. The singular eigenvalues of $J_n(1)$, which represent the main result of their paper, are later found to be

$$\lambda_n = \sqrt{2 + 2 \cos\left(\frac{2k\pi}{2n+1}\right)} = 2 \cos\left(\frac{k\pi}{2n+1}\right), \quad \text{for } k = 1, \dots, n.$$

This result appears in Theorem 3.3 of [5].

Capparelli and Maroscia also study the generating functions for the determinants of these matrices, but this is outside the scope of the present paper. However, it should be noted that the polynomials associated with

matrices of the form given by (4) with $b = 0$, are referred to as co-recursive in the literature, and their properties, particularly orthogonality, have been studied for a long time since Chihara [11]. A recent example is the application of Caputo–Fabrizio fractional operators in various physical problems [9]. The above matrix for $\mu = 1$, viz. $J_n^*(1)J_n(1)$, also appears in many different areas such as civil engineering construction [46,48,49], mechanical engineering [28], or in the analysis of multiconductor transmission lines [25].

Similar matrices also appear in spectral graph theory [1] where we have

$$\begin{pmatrix} 3 & 1 & & & \\ 1 & 2 & \ddots & & \\ & \ddots & \ddots & \ddots & 1 \\ & & & 1 & 2 \end{pmatrix}.$$

Such matrices emerge in deterministic port-based teleportation (dPBT) protocol schemes, where a quantum state is guaranteed to be transferred to another system without unitary correction as in [36]. Moreover, very similar matrices arise in mechanical engineering processes such as freeze-form extrusion fabrication processes [32].

6. Other applications

In the process of linearizing stochastic resonance problems, which arises in a very wide class of dynamical systems, one encounters the following matrix:

$$\begin{pmatrix} \lambda - D & D & & & & & \\ D & \lambda - 2D & D & & & & \\ & & D & \ddots & \ddots & & \\ & & & \ddots & \ddots & D & \\ & & & & D & \lambda - 2D & D \\ & & & & D & & \lambda - D \end{pmatrix}_{n \times n}.$$

The above result is referred to as a tridiagonal *Toeplitz* matrix in [38]. Via spectral theory, the above matrix becomes the type of matrix given by (8). In addition, almost identical matrices can be found in: (i) the theory of experiments in fluids [45] with $D = 1$, (ii) the fast two-dimensional smoothing with the discrete cosine transform in information science [35], where $D = 1$ and $\lambda = 0$, and (iii) the stability of the matrix formulations for the Crank–Nicolson finite difference method for time-dependent diffusion on a staggered grid [39]. In the last example it is found that the above matrix arises when implicit boundary conditions are applied resulting in (4.4) of their paper,

which in turn corresponds to $D = 1/2$ and $\lambda = 2$ in the above matrix. To solve the eigenvalue problem these authors use the paper by Yueh [52], which is discussed in the following section, but is itself a particular case of Losonczi’s more universal work.

Another application where the above tridiagonal matrices arise is in the study of powers of them such as (8) or (9). This was first carried out by Rimas in [40,41], but continues to fascinate many researchers to this day. Because the knowledge of the eigenpairs of these matrices is vital in these efforts, it means that Losonczi’s results in [33] are continually being rediscovered and used. However, he is often not credited for the solutions because the authors are unaware of his seminal work. Instead they only cite more recent references, which also do not cite him properly. In fact, there is a host of papers with applications where the above matrices appear. Some recent examples are [2,7,8,27,30]. Yet again, none of these papers cites Losonczi’s work.

7. More general cases

One of the most famous realizations of the general case (4), bearing in mind that it is a simplification of the Losonczi class given by (1), is the Kac–Murdock–Szegő, KMS for short, or Markovian matrix, which first appeared in [29]. This matrix is a square Toeplitz matrix and is defined as

$$K_n(\rho) = (\rho^{|i-j|})_{i,j=1,\dots,n},$$

where ρ is a real number. The above class of matrices represents a very fertile area of research and consequently, they continue to be much studied up to the present day. For example, see [12,50] or Problem 7.2 on pp. 12–13 of [26]. The determinants for these matrices are known to be given by $\det K_n(\rho) = (1 - \rho^2)^{n-1}$. Furthermore, for $\rho \neq \pm 1$, they reduce to

$$K_n^{-1}(\rho) = \frac{1}{1 - \rho^2} \begin{pmatrix} 1 & -\rho & & & & & \\ -\rho & 1 + \rho^2 & -\rho & & & & \\ & -\rho & \ddots & \ddots & & & \\ & & \ddots & \ddots & -\rho & & \\ & & & -\rho & 1 + \rho^2 & -\rho & \\ & & & & -\rho & 1 & \end{pmatrix}_{n \times n}.$$

By discarding the factor outside the above matrix and setting $\rho = 1$, one obtains the matrix $D^T D$, which arises in the theory of signal processing as exemplified by [16].

The general matrix given by (4) is fundamental in the analysis of the KMS matrices. For example, the following tridiagonal matrix appears in [50]:

$$\begin{pmatrix} \frac{2-\gamma}{1-\gamma} & -1 & & & & \\ -1 & 2 & -1 & & & \\ & -1 & \ddots & \ddots & & \\ & & \ddots & \ddots & -1 & \\ & & & -1 & 2 & -1 \\ & & & & -1 & 1 \end{pmatrix}_{n \times n},$$

as does the inverse of the matrix

$$L_n = (\min(i - j) - \gamma)_{i,j=1,\dots,n}.$$

A similar matrix also appears in [17].

At the beginning of this section we presented the type of matrix that emerges in the linearization of stochastic resonance problems given by (6). In their study of the stochastic resonance problem modelled as a one-dimensional array of nonlinear spatially coupled subsystems in the mean field limit, Nicolis and Nicolis [37] obtain the following matrix:

$$\begin{pmatrix} \lambda - D & D & & & & \\ D & -2\lambda - D & D & & & \\ & D & \ddots & \ddots & & \\ & & \ddots & \ddots & D & \\ & & & D & -2\lambda - D & D \\ & & & D & D & \lambda - D \end{pmatrix}_{n \times n}.$$

In determining the eigenvalues for this matrix they cite once again the work of Yueh [52], which as have stated previously did not reference the earlier and more general work by Losonczi.

Occasionally, due to awkward notation, it is not obvious that simple tridigonal matrices of the above form emerge. For example, in the diagonalization of the Ising model with a transverse magnetic field in the presence of a local field defect at one edge, Francica et al. obtain by introducing the nonlocal Jordan-Wigner tranformation the following matrix in Appendix A

of [20]:

$$\begin{pmatrix} 1 + h^2\mu^2 & -h & & & \\ -h & 1 + h^2 & \ddots & & \\ & \ddots & \ddots & \ddots & \\ & & \ddots & 1 + h^2 & -h \\ & & & -h & h^2 \end{pmatrix}.$$

Next the authors obtain a version of (5) for their specific application. Again, rather than cite Losonczi, they use the Yueh's approach to derive (A12) in their paper. Interestingly, if one sets $\mu = 0$ in the above matrix, then one obtains (B3) in [22], which deals with the study of freeze-in dark matter models where the "Clockwork" mechanism is used to suppress dark matter couplings. Again, Yueh's paper is cited in their derivation of the eigenvalues for the fermionic Clockwork matrix.

Another physical example of this type of matrix is (6) in [3], which takes the following form:

$$\begin{pmatrix} -2\Delta_L & 2J_L & & & \\ 2J_L & -4\Delta_L & \ddots & & \\ & \ddots & \ddots & \ddots & \\ & & \ddots & -4\Delta_L & 2J_L \\ & & & 2J_L & -2\Delta_L \end{pmatrix}.$$

Balachandran et al. refer to this as a Toeplitz bordered matrix. It arises from studying two spin chains of the Heisenberg XXZ model of condensed matter physics, each with a different anisotropy parameter. To explain the spin current rectification arising from the different anisotropy in both half-chains, they consider the case of coupling two chains, one of which is completely polarized and the other is at infinite temperature. By studying the case where the half-chain with non-zero anisotropy is polarized, they derive the above matrix determining its eigenvalues, which represents the magnon excitation spectrum for the half-chain.

In a similar fashion Guo and Poletti [23,24] also obtain an $n \times n$ tridiagonal Toeplitz bordered matrix in their solution for a large class of Lindblad master equations for noninteracting particles (both fermions and bosons are considered) on n sites. For the fermionic case the tridiagonal matrix is

given by

$$\begin{pmatrix} -ih_z - \Gamma_1/2 & -iJ/2 & & & \\ & -iJ/2 & -ih_z & \ddots & \\ & & \ddots & \ddots & \ddots \\ & & & \ddots & -ih_z & -iJ/2 \\ & & & & -iJ/2 & -ih_z - \Gamma_n/2 \end{pmatrix}.$$

Once again, the eigenvalues are determined by referring to the more recent work of Yueh rather than Losonczi’s work.

8. Extensions

As described in [19], the so-called k -tridiagonal matrices denoted as $T_n^{(k)}(a_*, b_*, c_*)$ with $a_* \doteq \{a_1, \dots, a_n\}$, $b_* \doteq \{b_1, \dots, b_{n-k}\}$ and $c_* \doteq \{c_{k+1}, \dots, c_n\}$ can be interpreted as a matrix, whose graph is a set of disjoint paths or a forest with r paths of length $\ell - 1$ and $k - r$ paths of length ℓ , where $n = k\ell - r$. The authors then show that these matrices can be expressed as a direct sum of tridiagonal matrices denoted by T_n . Consequently, the application of spectral theory, i.e., solving for the determinant, eigenvalues and eigenvectors, of these more general matrices than those discussed in Sections 4 to 6 can be achieved by using Egerváry and Szász [13] or the more general results of Losonczi [33].

There are two interesting extensions of these matrices. The first is when the main diagonal is comprised of zero entries and both subdiagonals are not in symmetric positions. Such matrices are studied in [31]. The other has no such restriction on the main diagonal, but possesses a quasi-symmetry in the sense that the upper subdiagonal begins horizontally at position $k + 1$, and the lower subdiagonal begins vertically at $k + 2$ as described in [47]. There they are referred to $(k, k + 1)$ tridiagonal matrices of order n and are denoted by $T_n^{(k,k+1)}$ with $n \leq 2k$.

The second type of generalization includes matrices of the form

$$\begin{pmatrix} a & c & & & \\ c & 0 & e & & \\ & e & \ddots & \ddots & \\ & & \ddots & \ddots & e \\ & & & e & 0 & c \\ & & & & c & b \end{pmatrix}.$$

The above matrices are studied in detail in [6,18] with many specific cases of this extension together with their spectra. The eigenvalues are derived by

relying on the frequently cited [51,52] which we have indicated are simply replicating Losonczi's results.

9. Conclusion

In this brief survey, we have presented a great number of important applications/problems in mathematics, physics and technology involving tridiagonal matrices that fall into the class given by (1). The spectral theory for these matrices and those given by (2) was developed by Losonczi in the early 1990's [33]. Yet this work has been largely ignored. In fact, many of the more recent problems cited here have been solved with reference to Yueh's work [52], which appeared more than a decade later and is less general than Losonczi's paper. It is hoped that this article has redressed this issue so that Losonczi's work receives the due recognition it deserves.

References

- [1] S. Akbari, W.H. Haemers, H.R. Maimani and L.P. Majd, Signed graphs cospectral with the path, *Linear Algebra Appl.*, **553** (2018), 104–116.
- [2] H. Al Ba'ba'a, M. Nouh and T. Singh, Formation of local resonance band gaps in finite acoustic metamaterials: A closed-form transfer function model, *J. Sound Vib.*, **410** (2017), 429–446.
- [3] V. Balachandran, G. Benenti, E. Pereira, G. Casati and D. Poletti, Perfect diode in quantum spin chains, *Phys. Rev. Lett.*, **120** (2018), 200603.
- [4] M.I. Bueno and F. Marcellán, Darboux transformation and perturbation of linear functionals, *Linear Algebra Appl.*, **384** (2004), 215–242.
- [5] S. Capparelli and P. Maroscia, On two sequences of orthogonal polynomials related to Jordan blocks, *Mediterr. J. Math.*, **10** (2010), 1609–1630.
- [6] H.C. Chang, S.E. Liu and R. Burrige, Exact eigensystems for some matrices arising from discretizations, *Linear Algebra Appl.*, **430** (2009), 999–1006.
- [7] A. Chatterjee, Sensitivity and error analysis of a coupled micro-resonator array for ultra-sensitive mass detection using matrix perturbation theory, *IEEE Sens. J.*, **16** (2016), 1304–1309.
- [8] A. Chatterjee, Lumped parameter modelling of turbine blade packets for analysis of modal characteristics and identification of damage induced mistuning, *Appl. Math. Model.*, **40** (2016), 2119–2133.
- [9] M. Chen, J. Shi and W. Deng, High order algorithms for Fokker–Planck equation with Caputo–Fabrizio fractional derivative, arXiv:1809.03263.
- [10] T.S. Chihara, *An Introduction to Orthogonal Polynomials*, Gordon and Breach (New York, 1978).
- [11] T.S. Chihara, On co-recursive orthogonal polynomials, *Proc. Amer. Math. Soc.*, **8** (1957), 899–905.
- [12] Z. Dong, J. Zhang and H. Chen, Average SEP-optimal precoding for correlated massive MIMO with ZF detection: An asymptotic analysis, arXiv:1901.00313
- [13] E. Egerváry and O. Szász, Einige Extremalprobleme im Bereiche der trigonometrischen Polynome, *Math. Z.*, **27** (1928), 641–652.
- [14] J.F. Elliott, *The characteristic roots of certain real symmetric matrices*, Master's thesis, University of Tennessee (1953).

- [15] K. Fan, O. Taussky and J. Todd, Discrete analogs of inequalities of Wirtinger, *Monatsh. Math.*, **59** (1955), 73–79.
- [16] A. Fasano and V. Villani, Baseline wander removal for bioelectrical signals by quadratic variation reduction, *Signal Process.*, **99** (2014), 48–57.
- [17] C.M. da Fonseca, On the eigenvalues of some tridiagonal matrices, *J. Comput. Appl. Math.*, **200** (2007), 283–286.
- [18] C.M. da Fonseca, S. Kouachi, D.A. Mazilu and I. Mazilu, A multi-temperature kinetic Ising model and the eigenvalues of some perturbed Jacobi matrices, *Appl. Math. Comput.*, **259** (2015), 205–211.
- [19] C.M. da Fonseca and F. Yilmaz, Some comments on k -tridiagonal matrices: determinant, spectra, and inversion, *Appl. Math. Comput.*, **270** (2015), 644–647.
- [20] G. Francica, T.J.G. Apollaro, N. Lo Gullo and F. Plastina, Local quench, majorana zero modes, and disturbance propagation in the Ising chain, *Phys. Rev. B*, **94** (2016), 245103.
- [21] Ya.L. Geronimus, Refinement of estimates of van der Corput, Visser, Fejes and Boas for the coefficients of trigonometric polynomials, *Dokl. Akad. Nauk SSSR (N.S.)*, **63** (1948), 479–482 (in Russian).
- [22] A. Goudelis, K.A. Mohan and D. Sengupta, Clockworking FIMPs, *J. High Energy Phys.*, **2018** (2018), 14.
- [23] C. Guo and D. Poletti, Analytical solutions for a boundary driven XY chain, *Phys. Rev. A*, **98** (2018), 052126.
- [24] C. Guo and D. Poletti, Solutions for dissipative quadratic open systems: part I – bosons, *Phys. Rev. A*, **95** (2017), 052107.
- [25] B. Honarbakhsh and S. Asadi, Analysis of multiconductor transmission lines using the CN-FDTD method, *IEEE T. Electromagnet. C.*, **59** (2017), 184–192.
- [26] R.A. Horn and C.R. Johnson, *Matrix Analysis*, Second Edition, Cambridge University Press (New York, 2013).
- [27] J. Hong, X. Li, D. Zhou, S. Liu, X. Liu and D.F. Wang, Localization in coupled systems: Part III – Vibration characteristics analysis in a mode-localized four cantilever array with and without mass perturbation, 2018 Symposium on Design, Test, Integration & Packaging of MEMS and MOEMS.
- [28] S. Hovda, Gibbs-like phenomenon inherent in a lumped element model of a rod, *Adv. Mech. Eng.*, **9** (2017), 1–12.
- [29] M. Kac, W.L. Murdock and G. Szegő, On the eigenvalues of certain Hermitian forms, *J. Rational Mech. Anal.*, **2** (1953), 767–800.
- [30] V. Katewa, F. Pasqualetti and V. Gupta, On privacy vs cooperation in multi-agent systems, *Int. J. Control*, **91** (2018), 1693–1707.
- [31] T. McMillen, On the eigenvalues of double band matrices, *Linear Algebra Appl.*, **431** (2009), 1890–1897.
- [32] M. Li, Modeling, analysis, and simulation for aqueous-based ceramic pastes in freeze-form extrusion fabrication process, Doctoral Dissertations, 2481, Missouri University of Science and Technology (2016).
- [33] L. Losonczi, Eigenvalues and eigenvectors of some tridiagonal matrices, *Acta Math. Hungar.*, **60** (1992), 309–332.
- [34] L. Losonczi, On some discrete quadratic inequalities, *Int. Ser. Numer. Math.*, **80** (1987), 73–85.
- [35] P. Lyubin and E. Shchetinin, Fast two-dimensional smoothing with discrete cosine transform, *Commun. Comput. Inf. Sci.*, **678** (2016), 646–656.
- [36] M. Mozrzymas, M. Studziński, S. Strelchuk and M. Horodecki, Optimal port-based teleportation, *New J. Phys.*, **20** (2018), 053006.
- [37] C. Nicolis and G. Nicolis, Coupling-enhanced stochastic resonance, *Phys. Rev. E*, **96** (2017), 042214.

- [38] S. Noschese and L. Reichel, Eigenvector sensitivity under general and structured perturbations of tridiagonal Toeplitz-type matrices, *Numer. Linear Algebra Appl.*, doi.org/10.1002/nla.2232.
- [39] C.M. Oishi, J.Y. Yuan, J.A. Cuminato and D.E. Stewart, Stability analysis of Crank-Nicolson and Euler schemes for time-dependent diffusion equations, *BIT Numer. Math.*, **55** (2015), 487–513.
- [40] J. Rimas, On computing of arbitrary positive integer powers for tridiagonal matrices with elements $1, 0, 0, \dots, 0, 1$ in principal and $1, 1, 1, \dots, 1$ in neighbouring diagonals – I, *Appl. Math. Comput.*, **186** (2007), 1254–1257.
- [41] J. Rimas, On computing of arbitrary positive integer powers for tridiagonal matrices with elements $-1, 0, 0, \dots, 0, 1$ in principal and $1, 1, 1, \dots, 1$ in neighbouring diagonals – II, *Appl. Math. Comput.*, **188** (2007), 634–637.
- [42] T.J. Rivlin, *Chebyshev Polynomials: From Approximation Theory to Algebra and Number Theory*, 2nd Ed., Wiley (New York, 1990).
- [43] W.W. Rogosinski and G. Szegő, Extremum problems for non-negative sine polynomials, *Acta Sci. Math. (Szeged)*, **12** (1950), 112–124.
- [44] D.E. Rutherford, Some continuant determinants arising in physics and chemistry, *Proc. Roy. Soc. Edinburgh Sect. A*, **62** (1947), 229–239.
- [45] C. Tang, W. Sun, H. He, H. Li and E. Li, Spurious PIV vector detection and correction using a penalized leastsquares method with adaptive order differentials, *Exp. Fluids*, **57** (2017), 81.
- [46] G. Tong and Y. Tang, Buckling of parallel purlins inter-braced by sag-rods, *J. Constr. Steel Res.*, **139** (2017), 123–134.
- [47] S. Takahira, T. Sogabe and T.S. Usuda, Bidiagonalization of $(k, k + 1)$ -tridiagonal matrices, *Spec. Matrices*, **7** (2019), 20–26.
- [48] Y. Tang, G. Tong and L. Zhang, Buckling of parallel purlins in standing seam or screw-fastened roofs, *Thin Walled Struct.*, **132** (2018), 136–150.
- [49] Y. Tang, G. Tong and L. Zhang, Effectiveness of sag rods as lateral supports for parallel purlins and girts, *J. Struct. Eng.*, **144** (2018), 04018151
- [50] W.F. Trench, Properties of some generalizations of Kac–Murdock–Szegő matrices, *Contemp. Math.*, **281** (2001), 233–245.
- [51] A.R. Willms, Analytic results for the eigenvalues of certain tridiagonal matrices, *SIAM J. Matrix Anal. Appl.*, **30** (2008), 639–656.
- [52] W.C. Yueh, Eigenvalues of several tridiagonal matrices, *Appl. Math. E-Notes*, **5** (2005), 66–74.