



Facet separation for disjunctive constraints with network flow representation

Péter Dobrovocski^{1,2} · Tamás Kis^{1,2}

Received: 28 April 2023 / Accepted: 30 August 2024
© The Author(s) 2024

Abstract

We present a novel algorithm for separating facet-inducing inequalities for the convex-hull of the union of polytopes representing a disjunctive constraint of special structure. It is required that the union of polytopes admit a certain network flow representation. The algorithm is based on a new, graph theoretic characterization of the facets of the convex-hull. Moreover, we characterize the family of polytopes that are representable by the networks under consideration. We demonstrate the effectiveness of our approach by computational results on a set of benchmark problems.

Keywords Disjunctive programming · Embedding formulations · Branch-and-cut

1 Introduction

Modeling with disjunctive constraints is a standard technique in the theory and practice of mathematical programming. The concept was introduced by Balas (1979), and has become a flourishing research area with many applications (Balas, 2018).

A central question is the following: given a finite set of polyhedra $P_1, \dots, P_m \subseteq \mathbb{R}_{\geq 0}^n$, how can we express the constraint

$$x \in \bigcup_{i=1}^m P_i \quad (1)$$

in a linear program? Balas had the idea to lift this problem into a higher dimensional space. Suppose $P_i = \{x \in \mathbb{R}^n \mid A^i x \leq b^i\}$ for $i \in [m]$. Let $\Gamma = [m]$, and $\Gamma^* = \{i \in \Gamma \mid P_i \neq \emptyset\}$. Let $\mathcal{S}(Q^*)$ be the set of those vectors $x \in \mathbb{R}^n$ such that there exist vectors $(y^i, \lambda_i)_{i \in \Gamma^*}$ satisfying

✉ Tamás Kis
kis.tamas@sztaki.hu

Péter Dobrovocski
peter.dobrovocski@sztaki.hu

¹ HUN-REN Institute for Computer Science and Control, Kende str. 13-17, Budapest 1111, Hungary

² Department of Operations Research, Eötvös Loránd University, Pázmány Péter sétány 1/A, Budapest 1117, Hungary

$$\begin{aligned}
x - \sum_{i \in \Gamma^*} y^i &= 0, \\
A^i y^i - b^i \lambda_i &\leq 0, \quad \forall i \in \Gamma^*, \\
\sum_{i \in \Gamma^*} \lambda_i &= 1, \\
\lambda_i &\geq 0, \quad \forall i \in \Gamma^*.
\end{aligned} \tag{2}$$

Theorem 1 (Balas (1985)) $\text{cl conv}(\bigcup_{i=1}^m P_i) = \mathcal{S}(\Gamma^*)$.

It may not be straightforward to determine the subset Γ^* of Γ . Balas has given necessary and sufficient conditions for replacing Γ^* by Γ in (2). For our purposes, the following condition suffices.

Theorem 2 (Balas (1985)) *If for every $i \in \Gamma$, some subset of the set of inequalities $A^i x \leq b^i$ defines a bounded nonempty polyhedron, then $\text{cl conv}(\bigcup_{i=1}^m P_i) = \mathcal{S}(\Gamma)$.*

From now on, we assume that each P_i is a polytope.

An important property of the linear system (2) is that for every basic (or extreme) solution, there exists an index $i \in \Gamma$ such that $x = y^i$ and $\lambda_i = 1$, while $y^{i'} = 0$ and $\lambda_{i'} = 0$ for all $i' \in \Gamma \setminus \{i\}$, see Balas (1998).

A drawback of the extended formulation (2) is that the x variables are copied m times, which may considerably increase the size of the LP formulation. To alleviate this burden, Vielma introduced the concept of *embedding formulations* (Vielma, 2018). Let $P_i^{\text{emb}} = P_i \times \epsilon_i$ be the embedding of polytope P_i into \mathbb{R}^{n+m} , where $\epsilon_i \in \mathbb{R}^m$ has a 1 in position i , and 0 otherwise. Then

$$P^{\text{emb}} = \text{conv} \left(\bigcup_{i=1}^m P_i^{\text{emb}} \right), \tag{3}$$

is the *Cayley Embedding* of the union of the P_i (Huber et al., 2000; Vielma, 2018).

In Kis and Horváth (2022) the concept of *network-flow representable polytope* has been defined, and explored to characterize the facets of (3). We recapitulate the fundamental definitions and key properties in Sect. 2.

Recall the notion of ideal, and non-extended formulation from Vielma (2018; 2019), Huchette and Vielma (2019a; 2019b). Let $Ax + By + Cz \leq b$, $z \in \mathbb{Z}^k$ be a formulation of (3) and Q denote the polyhedron determined by its LP relaxation. We call the formulation *ideal* if z is integral in the extreme points of Q , and *non-extended*, if it contains no y variables (otherwise extended). Our formulation of (3) is ideal, since it gives the convex hull of feasible points, and non-extended, since beside the original problem variables, there are only m new binary variables to select the P_i . Note that Balas' formulation (2) is ideal, and extended.

In the present paper we complement the results of Kis and Horváth (2022) by efficient separation algorithms that can be used in branch-and-cut solvers. To demonstrate the effectiveness of the new cuts, and to test the separation procedures, we summarize our computational experiments on a set of benchmark problems.

The structure of the paper is as follows. Section 2 describes the necessary background and some basic results to be used throughout the paper. We define the special cases P_{\leq}^{emb} and $P_{=}^{\text{emb}}$ of P^{emb} here as well. We characterize those disjunctive constraints that admit the proposed network flow representation in Sect. 3. The affine hull of P^{emb} is determined in Sect. 4. In Sect. 5 we recall the characterization of facets of P_{\leq}^{emb} from Kis and Horváth (2022), and

give a new characterization of the non-trivial facets of P_{\equiv}^{emb} , which is more general than the result in Kis and Horváth (2022). Separation procedures are provided along with proofs of their correctness in Sect. 6. In Sect. 7 we describe a general test problem for the separation algorithm and summarize our computational results. We also provide a comparison to other approaches, including Balas’ extended formulation.

2 Background and preliminary results

A network representation for P^{emb} consists of a network $G = (V, E)$ along with a capacity function $c_{x,\lambda}$ on the arcs, where G is of special structure. The set of nodes V comprises a source node $s \in V$ and a unique sink node $t \in V$. Let $V_t = \{v_1, \dots, v_n\}$ denote the neighboring nodes of t , and we will call them the *terminal nodes*. We assume that $G \setminus \{t\}$ can be decomposed into m subgraphs G_1, \dots, G_m . Each G_i is a directed tree rooted at s with all leaves in V_t . In addition, we assume that for each node $v_j \in V_t$ there exists at least one subgraph G_i containing v_j . Moreover, for $i \neq i'$, $V(G_i) \cap V(G_{i'})$ comprises node s , and possibly some nodes from V_t , but they have no other nodes in common. This implies that each $v \in V(G_i) \setminus \{s\}$ has a unique in-arc in G_i , denoted by $e_i(v)$, and a unique parent node $p_i(v)$, where $e_i(v) = (p_i(v), v)$.

The *capacity function* $c_{x,\lambda}$ has the following properties: if $e \in E(G_i)$, then $c_{x,\lambda}(e) = \beta_e \lambda_i$, where $\beta_e > 0$ is a constant. The sum of the capacities of the arcs emanating s in subgraph G_i is denoted by $\alpha_i \lambda_i$. The capacity of the arc (v_j, t) is $c_{x,\lambda}(v_j, t) = x_j$. The sum of the capacities of those arcs entering t is $\sum_{j=1}^n x_j$.

For simplicity, for a fixed graph G we will denote the network by $N(x, \lambda)$, where x and λ are the parameters of the capacity function. We say that $N(x, \lambda)$ *represents* P^{emb} , if for each i , when setting $\lambda_i = 1$, and the other coordinates of λ to 0, and for any $x \geq 0$, $(x, \lambda) \in P^{\text{emb}}$ if and only if the network N parametrized by x and λ as above, admits a feasible $s - t$ flow of value $\sum_{j=1}^n x_j$, and (x, λ) satisfies the valid equations for P^{emb} .

Throughout the paper, we will consider two special cases of the polyhedron P^{emb} . Either the only valid equation for P^{emb} is

$$\sum_{i=1}^m \lambda_i = 1, \tag{4}$$

or all $(x, \lambda) \in P^{\text{emb}}$ also satisfy

$$\sum_{i=1}^m \alpha_i \lambda_i = \sum_{j=1}^n x_j. \tag{5}$$

We distinguish these two cases by P_{\leq}^{emb} and P_{\equiv}^{emb} , respectively. Moreover, P_{\star}^{emb} will denote either of these two polytopes. Let $R_{\leq} := \{(x, \lambda) \in \mathbb{R}_{\geq 0}^{n+m} \mid (4)\}$, $R_{=} := \{(x, \lambda) \in R_{\leq} \mid (5)\}$, and R_{\star} will denote either of them.

We continue with a couple of definitions and notations used throughout the paper. The set U will denote a subset of the terminal nodes V_t , and \bar{U} will always denote $V_t \setminus U$. For an arc $e = (u, v)$ we call u the *tail of the arc* and v the *head of the arc*. In graph G , for any $X \subset V(G)$, we denote the set of those arcs (u, v) such that $v \in X$ and $u \in V \setminus X$ by $\rho(X)$ and the set of those arcs (u, v) such that $u \in X$, and $v \in V \setminus X$ by $\delta(X)$. Note that for $v \in V(G_i) \setminus (\{s\} \cup V_t)$, $\delta(v)$ and $\rho(v)$ consist of arcs in $E(G_i)$ only. For any node

$w \in V(G)$, let $\sigma(w)$ denote the subset of nodes in $V(G) \setminus \{t\}$ consisting of w and all those nodes reachable from w by a directed path.

An $s-t$ cut in this network is a partitioning $[S, \bar{S}]$ of V such that $s \in S$ and $t \in \bar{S} := V \setminus S$. We call S the *source side* of the cut and \bar{S} the *sink side* of the cut. Let $E(S, \bar{S})$ be the set of *cut-arcs*, with tail in S and head in \bar{S} , and let $E_i(S, \bar{S}) = E(G_i) \cap E(S, \bar{S})$. For any subset of nodes $S \subset V(G)$, let $E(S)$ be the set of those arcs from $E(G)$ with both end-points in S , and let $E_i(S) = E(G_i) \cap E(S)$. For $w \in S \cap V(G_i)$, let $c_w^i(S) = \sum_{e \in E_i(S \cap V(\sigma(w)), \bar{S})} \beta_e$. When the $s-t$ cut is clear from the context, we simply put c_w^i . For $w \in V_t$, $c_w^i(S) = 0$.

An $s-t$ cut $[S, \bar{S}]$ in G induces the following inequality:

$$\sum_{e \in E(S, \bar{S})} c_{x, \lambda}(e) \geq \sum_{j=1}^n x_j. \quad (6)$$

Consider a network $N(x, \lambda)$. We define $Q_{\leq}(N)$ as the set of vectors (x, λ) that satisfy (4) and all the inequalities (6):

$$Q_{\leq}(N) = \left\{ (x, \lambda) \in \mathbb{R}_{\geq 0}^{n+m} \left| \begin{array}{l} \sum_{i=1}^m \lambda_i = 1, \\ \sum_{e \in E(S, \bar{S})} c_{x, \lambda}(e) \geq \sum_{j=1}^n x_j, \forall s-t \text{ cut } [S, \bar{S}] \text{ in } G \end{array} \right. \right\}. \quad (7)$$

We define $Q_{=}(N)$ by requiring also (5):

$$Q_{=}(N) = \left\{ (x, \lambda) \in Q_{\leq}(N) \left| \sum_{i=1}^m \alpha_i \lambda_i = \sum_{j=1}^n x_j \right. \right\}. \quad (8)$$

If the network is understood from the context, then we use the notation Q_{\leq} and $Q_{=}$. For the sake of brevity, let Q_{\star} denote either Q_{\leq} or $Q_{=}$. An important property of these polyhedra is that in their vertices $\lambda = \epsilon_i$ for some $i \in \llbracket m \rrbracket$, see Kis and Horváth (2022).

Theorem 3 (Corollary 1 of Kis and Horváth (2022)) *Network N represents the polytope P_{\star}^{emb} if and only if $P_{\star}^{\text{emb}} = Q_{\star}(N)$.*

We continue with reduction rules for the capacity function $c_{x, \lambda}$. We call a capacity function *reduced* if for all $i \in \llbracket m \rrbracket$, and $v \in V(G_i) \setminus (\{s\} \cup V_t)$ the following holds:

$$\begin{aligned} \beta_e &\leq \beta_{e_i(v)} \quad \forall e \in \delta(v), \\ \beta_{e_i(v)} &< \sum_{e \in \delta(v)} \beta_e. \end{aligned} \quad (9)$$

Proposition 1 *A network $N = (G, c_{x, \lambda})$ with non-reduced capacity function can always be transformed to a network $N' = (G', c'_{x, \lambda})$ with a reduced capacity function such that $P_{\star}^{\text{emb}}(N) = P_{\star}^{\text{emb}}(N')$.*

The proofs of Propositions 1–6 can be found in Appendix A. For an $s-t$ cut $[S, \bar{S}]$, let k_i denote the sum of coefficients of the cut arcs that belong to $E(G_i)$, i.e.

$$k_i = \sum_{e \in E_i(S, \bar{S})} \beta_e \quad (10)$$

For a subset $U \subset V_t$ an $s - t$ cut with respect to U is an $s - t$ cut $[S, \bar{S}]$ such that $U \subseteq S$, and $V_t \setminus U = \bar{U} \subseteq \bar{S}$. There may be several $s - t$ cuts w.r.t. U , and we will be interested only in those of smallest capacity.

Definition 1 (Dominating cut) A dominating $s - t$ cut w.r.t. U is an $s - t$ cut such that k_i is minimal for each $i \in [m]$, where the minimum is taken over all $s - t$ cuts w.r.t. U .

It can be shown that there always exists a dominating $s - t$ cut for any $U \subset V_t$, see Kis and Horváth (2022). The dominating $s - t$ cuts w.r.t. U are not unique in general. We will denote the set of dominating $s - t$ cuts w.r.t. U by $\mathcal{C}_{\min}(U)$. All $s - t$ cuts in $\mathcal{C}_{\min}(U)$ have the same capacity by definition. We can also define a partial order on $\mathcal{C}_{\min}(U)$ based on the set inclusion relation.

Definition 2 (Minimal/maximal dominating cut) An $s - t$ cut $[S, \bar{S}]$ is *minimal dominating* w.r.t. U if

$$S \subseteq S' \quad \forall [S', \bar{S}'] \in \mathcal{C}_{\min}(U)$$

and it is *maximal dominating* if

$$S' \subseteq S \quad \forall [S', \bar{S}'] \in \mathcal{C}_{\min}(U).$$

The minimal and the maximal dominating $s - t$ cuts w.r.t. U are unique as proved in Proposition 4 of Kis and Horváth (2022). Throughout this paper, $[S^-, \bar{S}^-]$ and $[S^+, \bar{S}^+]$ will denote the unique minimal and maximal dominating $s - t$ cut w.r.t. U , respectively.

Next, we give a constructive characterization of these cuts. To this end, we define a special capacity function c^U on the arcs of G as follows:

$$c_{uv}^U = \begin{cases} \beta_{uv} & \text{if } v \neq t \\ 0 & \text{if } u \in U \\ M & \text{if } u \in \bar{U} \end{cases}, \tag{11}$$

where M is a very large number, for example $M = \sum_{i=1}^m \alpha_i + 1$. With the new arc capacities c_{uv}^U , in any minimum capacity $s - t$ cut, the arcs (v_j, t) are always cut-arcs for all $v_j \in U$ and the arcs $(v_{j'}, t)$ are never cut-arcs for any $v_{j'} \in \bar{U}$. Thus for any minimum capacity $s - t$ cut $[S, \bar{S}]$ we have $U \subset S$, and $\bar{U} \subset \bar{S}$. Let f denote a maximum $s - t$ flow in the network with arc capacities c^U . We call arc e *tight* if $\beta_e = f_e$. It is well-known that in any minimum capacity $s - t$ cut, the cut-arcs are tight. Observe that tight arcs occur only on paths from s to \bar{U} by the definition of c^U . Our goal is to find sets S^-, S^+ such that $s \in S^- \cap S^+$, S^- is minimal, S^+ is maximal and all arcs leaving S^- or S^+ are tight.

Proposition 2 Let $[S^-, \bar{S}^-]$ be the minimal dominating $s - t$ cut w.r.t. U . Let f be an $s - t$ flow of maximum value in the network (G, c^U) . The set of cut-arcs $E^- := E(S^-, \bar{S}^-)$ has the following properties:

1. All arcs from U to t belong to E^- ,
2. No arc from \bar{U} to t is in E^- ,
3. The arcs in E^- are tight and cover all paths from s to \bar{U} ,
4. E^- is minimal in the sense that no proper subset of it covers all paths from s to t ,
5. For all $uv \in E^-$ with $v \neq t$, there is no tight arc on the unique $s - u$ path in G .

In a similar manner, we can derive the key properties of the maximal dominating $s - t$ cuts w.r.t. U .

Proposition 3 Let $[S^+, \overline{S^+}]$ be a maximal dominating $s - t$ cut w.r.t. U . Let f be a maximum flow in the network (G, c^U) . The set of cut-arcs $E^+ := E(S^+, \overline{S^+})$ has the following properties:

1. All arcs from U to t are in E^+ ,
2. No arc from \overline{U} to t belong to E^+ ,
3. The arcs in E^+ are tight and cover all paths from s to \overline{U} ,
4. E^+ is maximal in the sense that no proper subset of it covers all paths from s to t ,
5. For all $uv \in E^+$ with $v \neq t$, there exists a path from v to \overline{U} with no tight arc.

The minimal and the maximal dominating $s - t$ cuts, respectively, satisfy the following properties:

Proposition 4 For $U \subset U' \subseteq V_t$ and minimal dominating $s - t$ cuts $[S^-, \overline{S^-}], [Z^-, \overline{Z^-}]$ w.r.t. U and U' , respectively, we have $S^- \subset Z^-$.

Proposition 5 For $U \subset U' \subseteq V_t$ and maximal dominating $s - t$ cuts $[S^+, \overline{S^+}], [Z^+, \overline{Z^+}]$ w.r.t. U and U' , respectively, we have $S^+ \subset Z^+$.

Finally, consider S^+ and S^- , where $[S^+, \overline{S^+}]$ and $[S^-, \overline{S^-}]$ are maximal and minimal dominating $s - t$ cuts w.r.t. $U \subseteq V_t$.

Proposition 6 The connected components of the subgraph spanned by $S^+ \setminus S^-$ are rooted trees. If $v \in S^+ \setminus S^-$ is the root node of such a tree, then $e_i(v)$ emanates from S^- , and $\beta_{e_i(v)} = c_v^i(S^+)$.

Corollary 1 Let $[S^+, \overline{S^+}], [S^-, \overline{S^-}]$ be the maximal and minimal dominating $s - t$ cuts w.r.t. U , respectively. For all $i \in [m]$, we have

$$\sum_{e \in E_i(S^+ \setminus S^-, \overline{S^+})} \beta_e = \sum_{e \in E_i(S^-, S^+ \setminus S^-)} \beta_e. \quad (12)$$

Now we consider the inequalities induced by dominating $s - t$ cuts. For any $U \subset V_t$, the dominating $s - t$ cuts in $\mathcal{C}_{\min}(U)$ uniquely induce the inequality

$$\sum_{i=1}^m k_i \lambda_i + \sum_{j: v_j \in U} x_j \geq \sum_{j=1}^n x_j. \quad (13)$$

We can simplify this by subtracting $\sum_{j: v_j \in U} x_j$ from both sides:

$$\sum_{i=1}^m k_i \lambda_i \geq \sum_{j: v_j \in \overline{U}} x_j. \quad (14)$$

For any subset $U \subset V_t$, let F_U be the set of vectors $(x, \lambda) \in P_{\star}^{\text{emb}}$ that satisfy the equation

$$\sum_{i=1}^m k_i \lambda_i = \sum_{j: v_j \in \overline{U}} x_j. \quad (15)$$

Note that (15) is obtained from (13) or (14) by turning the inequality into an equation. We call $U \subset V_t$ facet inducing for P_{\star}^{emb} if F_U is a facet of P_{\star}^{emb} .

In the P_{\leq}^{emb} polytope, the points of F_U also satisfy

$$\sum_{i=1}^m (\alpha_i - k_i) \lambda_i = \sum_{j: v_j \in U} x_j, \tag{16}$$

which can be derived by subtracting (15) from (5).

Let $(x, \lambda) \in P_{\star}^{\text{emb}}$ be a vector that satisfies an inequality induced by a dominating $s - t$ cut $[S, \bar{S}]$ (for some $U \subset V_t$) at equality. Let ϕ be a maximum $s - t$ flow in $N(x, \lambda)$. Then $\phi_e = c_{x,\lambda}(e)$ for all $e \in E(S, \bar{S})$ and $\phi_e = 0$ for all $e \in E(\bar{S}, S)$. For polytope P_{\leq}^{emb} we also have $\phi_{v_j t} = x_j$ for all $j \in \llbracket n \rrbracket$.

3 Network flow representable polytopes

In this section we characterize the family of those polytopes, that admit a network flow representation. Recall, that a family of sets S is called *laminar*, if

$$X \cap Y \neq \emptyset \implies X \subseteq Y \text{ or } Y \subseteq X \tag{17}$$

holds for all $X, Y \in \mathcal{S}$, see e.g., section 13.4 of Schrijver (2004).

Theorem 4 For $i \in \llbracket m \rrbracket$, let \mathcal{L}_i be a laminar family on $\llbracket n \rrbracket$ and for each $L \in \mathcal{L}_i$, b_L a positive number, and define a polytope

$$P_i = \left\{ x \in \mathbb{R}_{\geq 0}^n \mid \sum_{j \in L} x_j \leq b_L, \forall L \in \mathcal{L}_i \right\}, \tag{18}$$

for each of these families. Then $P_{\leq}^{\text{emb}} = \text{conv}(\cup_{i=1}^m P_i^{\text{emb}})$ is network flow representable, where $P_i^{\text{emb}} = P_i \times \epsilon_i$.

Conversely, given a network flow representable polytope $P_{\leq}^{\text{emb}} = \text{conv}(\cup_{i=1}^m P_i^{\text{emb}})$, we can find laminar families \mathcal{L}_i , $i \in \llbracket m \rrbracket$ and $b_L > 0$ for each $L \in \mathcal{L}_i$, such that the P_i satisfy (18).

Proof To prove the statement, we first show that for a polytope P_{\leq}^{emb} or $P_{=}^{\text{emb}}$, we can construct a network N representing it. Then we show that from a network N , we can retrieve the systems of inequalities of (18).

Recall that a laminar family \mathcal{L}_i admits a rooted tree representation, see e.g., Edmonds and Giles (1977). Let the graph G_i be the rooted tree representation of \mathcal{L}_i . Note that every inequality $\sum_{j \in L} x_j \leq b_L$ is represented by a node of G_i , and every leaf v_j of G_i corresponds to a variable x_j . Let G be the graph we get by joining the graphs G_i for all i on their leaves. Then we add source node s and sink node t , and add arcs from every leaf to t and from s to every node with no in-arcs. Define capacity function $c_{x,\lambda}$ as $c_{x,\lambda}(u, v) = b_L \lambda_i$ if node v corresponds to $\sum_{j \in L} x_j \leq b_L$ and $L \in \mathcal{L}_i$. If $v = v_j$ is a leaf, and there is no explicit upper bound on it defined by (18), then the $c_{x,\lambda}(u, v_j) = b_L \lambda_i$ where L is the minimal set in \mathcal{L}_i that contains j . For leaf node v_j , we set $c_{x,\lambda}(v_j t) = x_j$. See Fig. 1 for an illustration. Let $N = (G, c_{x,\lambda})$ be the network flow representation.




Fig. 1 Example for polytope P_i , the corresponding laminar system \mathcal{L}_i and the rooted tree representation G_i along with capacity function $c_{x,\lambda}$

We argue that for any $(x, \lambda) \in P_i^{\text{emb}}$ with $\lambda_i = 1$, we have $\lambda_{i'} = 0$ for all $i' \neq i$, and there exists an $s - t$ -flow f in N with flow value $\sum_{j=1}^n x_j$, that is,

$$f_e = \begin{cases} x_j & \text{if } e = (v_j, t) \\ \sum_{j \in L} x_j & \text{if } e = (u, v), \text{ and } v \text{ represents the inequality } \sum_{j \in L} x_j \leq b_L \end{cases} \quad (19)$$

for all arc e of G_i and $f_e = 0$ if e is not in G_i . Now since (x, λ) satisfies the inequalities (18), it is guaranteed that flow values do not exceed capacity on any arcs, i.e., f is feasible $s - t$ -flow.

For the other direction, we use again the fact that there is a one-to-one correspondence between laminar families and rooted trees. Let $N = (G, c_{x,\lambda})$ be the network flow representation of polytope P_i^{emb} . Let G_i be the subgraph of N , where the capacity of the arcs depend on λ_i . For any node v of G_i , let L_v denote the indices of terminal nodes that can be reached from v on a directed path. Notice that $(L_v)_{v \in V(G_i)}$ is a laminar family. Let b_v denote the coefficient of λ_i on the in-arc of node $v \in V(G_i)$. We have to show that $P_i = \left\{ x \in \mathbb{R}_{\geq 0}^n \mid \sum_{j \in L_v} x_j \leq b_v, \forall v \in V(G_i) \right\}$. Consider $P_i^{\text{emb}} = P_i \times \epsilon_i$, and fix $\lambda_i = 1$ and $\lambda_{i'} = 0$ for all $i' \neq i$ and some $x \in \mathbb{R}_+^n$ such that there is a feasible $s - t$ flow in $N(x, \lambda)$ of value $\sum_{j=1}^n x_j$. Then $(x, \lambda) \in P_i^{\text{emb}}$, which implies $\left\{ x \in \mathbb{R}_{\geq 0}^n \mid \sum_{j \in L_v} x_j \leq b_v, \forall v \in V(G_i) \right\} \subseteq P_i$. The converse inclusion is obvious by the definition of network flow representability. \square

Theorem 4 can be straightforwardly extended to the polytopes

$$P_i^- = \left\{ x \in \mathbb{R}_{\geq 0}^n \mid \sum_{j=1}^n x_j = \alpha_i, \sum_{j \in L} x_j \leq b_L, \forall L \in \mathcal{L}_i \right\}, \quad i \in \llbracket m \rrbracket,$$

where $\alpha_i > 0$ for $i \in \llbracket m \rrbracket$, since the defining equations $\sum_{j=1}^n x_j = \alpha_i$ have no impact on the laminar family induced by the inequalities.




Fig. 2 Illustration of the a_i^ℓ values

The depth of the laminar family \mathcal{L} is the length of the longest chain, i.e., the number of nodes of the longest path from s to t in the network minus 2. In Fig. 1, a longest chain is $\{(x_3), (x_3, x_4), (x_3, x_4, x_5), (x_1, x_2, x_3, x_4, x_5)\}$ with length 4, while the longest path from s to t has 6 nodes.

4 The affine hull of P_{\star}^{emb}

Observe that the only valid equation for P_{\leq}^{emb} is (4) provided that all terminal nodes in V_t are connected to the source node s of the network. On the other hand, the structure of the network may imply some additional valid equations for P_{\leq}^{emb} beside (4) and (5).

Let κ be the number of the connected components (in the undirected sense) of the graph $G \setminus \{s, t\}$ and T_1, \dots, T_κ be the components. By definition, there are no arcs between T_p and T_q for any $p \neq q$. Let U_ℓ be the subset of nodes of T_ℓ that belong to V_t . Note that $\cup_{\ell=1}^\kappa U_\ell = V_t$. Let a_i^ℓ denote the sum of the coefficients of those edges of subgraph G_i that start in s and end in T_ℓ , i.e.

$$a_i^\ell = \sum_{e \in E_i(s, T_\ell)} \beta_e. \tag{20}$$

See Fig. 2 for an illustration.

Proposition 7 *The equations*

$$\sum_{i=1}^m a_i^\ell \lambda_i = \sum_{j: v_j \in U_\ell} x_j \quad \forall \ell \in \llbracket \kappa \rrbracket \tag{21}$$

are all valid for P_{\leq}^{emb} .

Proof We have to prove that all $(x, \lambda) \in P_{\leq}^{\text{emb}}$ satisfy all of the Eq. (21). For any $(x, \lambda) \in P_{\leq}^{\text{emb}}$ we have

$$\sum_{\ell=1}^\kappa \sum_{i=1}^m a_i^\ell \lambda_i = \sum_{i=1}^m \alpha_i \lambda_i = \sum_{j=1}^n x_j = \sum_{\ell=1}^\kappa \sum_{j: v_j \in U_\ell} x_j,$$

since $V_t = \cup_{\ell=1}^\kappa U_\ell$ and $\sum_{\ell=1}^\kappa \alpha_i^\ell = \alpha_i$ for all $i \in \llbracket m \rrbracket$. We will prove that for any $\ell \in \llbracket \kappa \rrbracket$, for all $(x, \lambda) \in P_{\leq}^{\text{emb}}$ the sum of the capacity of edges that enter T_ℓ is exactly the sum of

the capacity of the edges that leave T_ℓ . In any $s - t$ flow in $N(x, \lambda)$, the arcs that leave s and enter T_ℓ are saturated, otherwise Eq. (5) is not satisfied. Also, the arcs leaving T_ℓ and entering t are saturated, for the same reason. No flow that enters T_ℓ enters any other T_k , and the conservation flow implies that the sum of the capacities of the arcs entering T_ℓ is equal to the capacity of those arcs leaving T_ℓ and entering t , which is the statement we wanted to prove. \square

Observation 1 Consider any face F of P_{\leq}^{emb} . Then F consists of those points $(x, \lambda) \in P_{\leq}^{\text{emb}}$ that satisfy (5), and some other Eq. (15) for distinct subsets $U \subset V_t$.

Note that a face F_U may be equal to the intersection of several $F_{U'}$ for distinct subsets U' of V_t .

Proposition 8 The system of equations consisting of (4), and (21) constitute a maximal linearly independent equation system for P_{\leq}^{emb} .

Proof By observation (1), P_{\leq}^{emb} consists of those points of P_{\leq}^{emb} that satisfy (5) and some equations of the form (15) for distinct subsets $U \subset V_t$. Suppose there exists a subset U' of V_t other than $(U_\ell)_{\ell=1}^k$ such that all $(x, \lambda) \in P_{\leq}^{\text{emb}}$ satisfy the equation

$$\sum_{i=1}^m k'_i \lambda_i = \sum_{j: v_j \in U'} x_j, \quad (22)$$

where k'_i is the coefficient of λ_i in the inequality (15) for U' . We can distinguish two cases: i) there exists some index ℓ such that $U' \subset U_\ell$, and ii) the set U' intersects multiple U_ℓ sets. We will prove that case i) is impossible and we will reduce case ii) to case i). Suppose that case i) holds. By subtracting (22) from (21) we obtain

$$\sum_{i=1}^m (a_i^\ell - k'_i) \lambda_i = \sum_{j: v_j \in U'} x_j. \quad (23)$$

Now, since T_ℓ is connected, there exists G_i and $v \in V(G_i) \cap V(T_\ell)$ such that the parent node of v is s and at least one node from both U' and $U_\ell \setminus U'$ is reachable from v on a directed path. Let p_1 and p_2 be two directed paths connecting v with nodes $v_{j'} \in U'$, and $v_{j''} \in U_\ell \setminus U'$, respectively.

We fix $\lambda_i = 1$, and $\lambda_k = 0$ for all $k \neq i$. We construct a feasible flow ϕ in the network as follows. We set $\phi_{su} = \beta_{su}$ for $su \in E(G_i)$, otherwise $\phi_{su} = 0$. Then, for each $w \in V(G_i) \setminus (V_t \cup \{s\})$, we split the flow entering w among the out-arcs of w proportionally to their capacities, i.e.,

$$\phi_e = \frac{\beta_e}{\sum_{e' \in \delta(w)} \beta_{e'}} \phi_{e_i(v)}, \quad \forall e \in \delta(w).$$

We let x_j equal the flow entering v_j , and also let $\phi_{v_j t} = \phi_{e_i(v_j)}$. Finally, we let $\phi_e = 0$ for all other arcs. This defines a feasible flow in $N(x, \lambda)$. Since $\sum_{j=1}^n x_j = \alpha_i$ by construction, we have $(x, \lambda) \in P_{\leq}^{\text{emb}}$. Hence, (x, λ) must satisfy (22).

Observe that $\phi_e = \beta_e$ for all $e \in \delta_{G_i}(s)$ by construction (i.e., all the out-arcs of s are saturated). Moreover, for each $w \in V(G_i) \setminus \{s\}$, $0 < \phi_e < \beta_e$ for all $e \in \delta(w)$. Hence, $0 < \phi < \beta_e$ for all arcs of p_1 and p_2 . Let $\varepsilon = \min\{\phi_e, \beta_e - \phi_e \mid e \in E(p_1) \cup E(p_2)\}$.

We perturb ϕ as follows: we decrease the value of ϕ on the arcs of p_2 by ε and increase ϕ on the arcs of p_1 by ε :

$$\phi'_e = \begin{cases} \phi_e + \varepsilon & e \in p_1 \\ \phi_e - \varepsilon & e \in p_2 \\ \phi_e & \text{otherwise} \end{cases} \tag{24}$$

Clearly, ϕ' is a feasible flow. Now let $(x', \lambda) \in P_{\leq}^{\text{emb}}$ a point such that

$$x'_j = \begin{cases} x_{j'} + \varepsilon & \text{if } j = j' \\ x_{j''} - \varepsilon & \text{if } j = j'' \\ x_j & \text{otherwise.} \end{cases} \tag{25}$$

Note that $\sum_{j=1}^n x_j = \sum_{j=1}^n x'_j$, but $\sum_{j \in U'} x_j \neq \sum_{j \in U'} x'_j$. Since ϕ' is a feasible flow in $N_{\leq}(x', \lambda)$ of value $\sum_{j=1}^n x'_j$, (x', λ) belongs to P_{\leq}^{emb} . However (22) is not satisfied by (x', λ) , since $\sum_{j \in U'} x_j \neq \sum_{j \in U'} x'_j$. Hence, Eq. (22) is not valid for all points of P_{\leq}^{emb} , a contradiction.

As for case (ii), we can assume that U' intersects exactly two of the sets U_1, \dots, U_κ (otherwise it can be reduced to this case by induction), and that the two sets are U_1 and U_2 (by possibly re-labeling U_1, \dots, U_ℓ). Let $[S, \bar{S}]$ be a dominating $s - t$ cut w.r.t. $V_t \setminus U'$. Let $T'_1 = \bar{S} \cap T_1$, $T'_2 = \bar{S} \cap T_2$ and $U'_1 = U_1 \cap U'$, $U'_2 = U_2 \cap U'$. Let k_i^1 denote the sum of the coefficients on those cut-arcs that enter T'_1 and belong to G_i , and define k_i^2 similarly for T'_2 . Since there are no arcs between T'_1 and T'_2 in any direction, we can decompose Eq. (22) the following way:

$$\sum_{i=1}^m k_i^1 \lambda_i + \sum_{i=1}^m k_i^2 \lambda_i = \sum_{j: v_j \in U'_1} x_j + \sum_{j: v_j \in U'_2} x_j, \tag{26}$$

which is exactly the sum of the equations

$$\sum_{i=1}^m k_i^1 \lambda_i = \sum_{j: v_j \in U'_1} x_j \tag{27a}$$

$$\sum_{i=1}^m k_i^2 \lambda_i = \sum_{j: v_j \in U'_2} x_j. \tag{27b}$$

This means that there is an implied valid equation acting on U'_1 , which is a proper subset of U_1 , thus reducing it to case (i). □

The above propositions give a simple method to determine a maximal system of linearly independent valid equations for P_{\leq}^{emb} .

5 Characterization of facet inducing $s - t$ cuts of P_{\leq}^{emb} and $P_{=}^{\text{emb}}$

In this section we give new characterizations of the facet inducing $s - t$ cuts for the P_{\leq}^{emb} and $P_{=}^{\text{emb}}$ polytopes, respectively.

Theorem 5 Let U be an arbitrary subset of the terminal nodes, $[S^-, \overline{S^-}]$ and $[S^+, \overline{S^+}]$ the minimal and the maximal dominating $s - t$ cut w.r.t. U , respectively. U induces a facet of P_{\leq}^{emb} if and only if

1. The subgraph of G induced by $\overline{S^+} \setminus \{t\}$ is connected in the undirected sense, and
2. The subgraph of G induced by S^- is connected in the undirected sense.

Proof These conditions are equivalent to those of Theorem 1 in Kis and Horváth (2022). \square

Now we turn to P_{\leq}^{emb} . In order to characterize its facets, we may assume that a maximal, linearly independent system of valid equations for P_{\leq}^{emb} consists of (4) and (5), see Kis and Horváth (2022), which is equivalent to the condition that $G \setminus \{s, t\}$ is connected (by Proposition 8). In Kis and Horváth (2022), the characterization of non-trivial facets of P_{\leq}^{emb} implicitly assumes that the node-set of the network can be partitioned as $\{s, t\} \cup V_s \cup V_t$, where V_s and V_t are the neighbors of the source s , and sink t , respectively, and the three subsets, namely, V_s , V_t , and $\{s, t\}$, are pairwise disjoint. Below we give a new characterization which is valid for any network.

Theorem 6 Suppose that the only valid equations for P_{\leq}^{emb} are (4) and (5). Fix some $U \subset V_t$, and let $[S^-, \overline{S^-}]$ and $[S^+, \overline{S^+}]$ be the minimal and the maximal dominating $s - t$ cut w.r.t. U , respectively. U induces a facet of P_{\leq}^{emb} if and only if

1. the subgraph of G induced by $\overline{S^+} \setminus \{t\}$ is connected in the undirected sense and
2. the subgraph of G induced by $S^- \setminus \{s\}$ is connected in the undirected sense.

For the proof of Theorems 6, see Appendix B.

These characterizations lead to polynomial-time separation algorithms, which is the topic of the next section.

6 Separation algorithms

The separation algorithms for both of P_{\leq}^{emb} and P_{\leq}^{emb} aim to solve the following problem: given a point $(x, \lambda) \in R_*$, decide if $(x, \lambda) \in P_{\leq}^{\text{emb}}$, and if not, then determine a violated facet inducing inequality. Both algorithms are based on the network flow representation of P_{\leq}^{emb} , and use the $s - t$ cuts of the network.

We can decide if a vector $(x, \lambda) \in R_*$ belongs to P_{\leq}^{emb} by computing the capacity C of a minimum capacity $s - t$ cut in $N(x, \lambda)$ and compare it to $\sum_{i=1}^n x_i$. If

$$C < \sum_{i=1}^n x_i, \quad (28)$$

then (x, λ) is not contained in the polytope, otherwise it is, see Sect. 1.

At first, we describe two methods to find a maximal and a minimal dominating $s - t$ cut, respectively w.r.t. $U \subset V_t$ in Sect. 6.1. We will need them in the separation algorithms for P_{\leq}^{emb} , and P_{\leq}^{emb} , respectively, which are presented in Sects. 6.2 and 6.3, respectively, along with a proof of correctness.

6.1 Finding minimal and maximal dominating $s - t$ cuts w.r.t. a given set U

We will use the properties described in Propositions 2 and 3 for finding minimal and maximal dominating $s - t$ cuts, respectively. The algorithm for finding the minimal dominating $s - t$ cut w.r.t. U is depicted in Algorithm 1. Initially we let $S^- = V \setminus (\overline{U} \cup \{t\})$, $\overline{S^-} = \overline{U} \cup \{t\}$ and $E^- = E(S^-, \overline{S^-})$. We process each subgraph G_i separately. Let W_i denote $V(G_i) \cap \overline{U}$. We want to cover all paths from s to W_i in G_i with arcs of minimum total capacity. We choose a node from W_i , and we examine the path from s to v in G_i , and denote it by $\pi_i(s, v)$. Let $e = uv$ be the first tight arc of the path $\pi_i(s, v)$. Note that all such paths contain at least one tight arc by Proposition 2. We extend the set of cut-arcs E^- with arc e and delete those cut-arcs from E^- that are in the subgraph $\sigma(w)$. We delete the nodes of $\sigma(w)$ from S^- and extend $\overline{S^-}$ with the same nodes. We delete the terminal nodes of $\sigma(w)$ from W_i . This is repeated while W_i is not the empty set. By Proposition 2, this algorithm finds a minimal dominating $s - t$ cut.

Algorithm 1 Find minimal dominating $s - t$ cut w.r.t. U

Input: $U \subseteq V_t$
Output: $[S^-, \overline{S^-}]$ minimal dominating $s - t$ cut w.r.t. U and E^- cut-arcs
 Find a maximum $s - t$ flow in (G, c^U)
 $S^- \leftarrow V \setminus (\overline{U} \cup \{t\}), \overline{S^-} \leftarrow \overline{U} \cup \{t\}, E^- \leftarrow E(S^-, \overline{S^-})$
for $i = 1, \dots, m$ **do**
 $W_i \leftarrow \overline{U} \cap V(G_i)$
 while $W_i \neq \emptyset$ **do**
 Let $v \in W_i$
 $\pi_i(s, v) \leftarrow$ unique path from s to v in G_i
 $e = uv \leftarrow$ first tight arc of $\pi_i(s, v)$
 $W_i \leftarrow W_i \setminus \sigma(w)$
 $S^- \leftarrow (S^- \setminus \sigma(w)) \cup U$
 $\overline{S^-} \leftarrow (\overline{S^-} \cup \sigma(w)) \setminus U$
 $E^- \leftarrow (E^- \setminus E(\sigma(w))) \cup \{e\}$
 end while
end for
return $[S^-, \overline{S^-}], E^-$

For finding the maximal dominating $s - t$ cut w.r.t. U , the procedure is the same, except that we choose the last arc of the path $\pi_i(s, v)$. The corresponding algorithm is depicted in Algorithm 2.

6.2 Facet separation algorithm for polytope P_{\leq}^{emb}

Consider Algorithm 3. The following results ensure that the algorithm always gives the correct answer in polynomial time.

Consider the minimal dominating $s - t$ cut $[S^-, \overline{S^-}]$ w.r.t. U , and suppose that the subgraph of G induced by S^- is not connected. Let S_0, S_1, \dots, S_N denote the node sets of its connected components such that $s \in S_0$. Let $U_0 = S_0 \cap V_t$. See Fig. 3.

Proposition 9 For each $\ell \in [1, N]$ there exists $v \in V_t$ such that $S_\ell = \{v\}$.

Algorithm 2 Find maximal dominating $s - t$ cut w.r.t. U

Input: $U \subseteq V_t$
Output: $[S^+, \overline{S^+}]$ maximal dominating $s - t$ cut w.r.t. U and E^+ cut-arcs

Find a maximum $s - t$ flow in (G, c^U)
 $E^+ \leftarrow \emptyset, S^+ \leftarrow V \setminus \overline{U}, \overline{S^+} \leftarrow \overline{U}$
for $i = 1, \dots, m$ **do**
 $W_i \leftarrow \overline{U} \cap V(G_i)$
 while $W_i \neq \emptyset$ **do**
 Let $v \in W_i$
 $\pi_i(s, v) \leftarrow$ unique path from s to v in G_i
 $e = uv \leftarrow$ last tight arc of $\pi_i(s, v)$
 $W_i \leftarrow W_i \setminus (\sigma(w) \cap W_i)$
 $S^+ \leftarrow (S^+ \setminus \sigma(w)) \cup U$
 $\overline{S^+} \leftarrow (\overline{S^+} \cup \sigma(w)) \setminus U$
 $E^+ \leftarrow E^+ \setminus E(\sigma(w)) \cup \{e\}$
 end while
end for
return $[S^+, \overline{S^+}], E^+$

Algorithm 3 Find violated facet for polytope P_{\leq}^{emb}

Input: $U \subset V_t, (x^*, \lambda^*) \in R_{\leq}$ such that the inequality (14) induced by U is violated by (x^*, λ^*)
Output: U' facet inducing and violated by (x^*, λ^*)

$[S^-, \overline{S^-}] \leftarrow$ minimal dominating $s - t$ cut w.r.t. U found by Algorithm 1
if S^- is not connected **then**
 $S_0 \leftarrow$ vertex set of the connected component of $G(S^-)$ containing s
 $U \leftarrow S_0 \cap V_t$
end if
 $[S^+, \overline{S^+}] \leftarrow$ maximal dominating $s - t$ cut w.r.t. U found by Algorithm 2
if $G(\overline{S^+} \setminus \{t\})$ is not connected **then**
 $T_1, \dots, T_K \leftarrow$ connected components of $\overline{S^+} \setminus \{t\}$
 for $\ell = 1, \dots, K$ **do**
 if $U \setminus \bigcup_{k \in [K] \setminus \{\ell\}} (T_k \cap V_t)$ is violated by (x^*, λ^*) **then**
 $U' \leftarrow U \setminus \bigcup_{k \in [K] \setminus \{\ell\}} (T_k \cap V_t)$
 end if
 end for
 else
 $U' \leftarrow U$
 end if
return inequality (14) for U'

Proof We argue that $S_\ell \subset V_t$ for each $\ell \in [1, N]$, from which the statement follows. If not, then observe that there is no arc uv of the graph G such that $u \in S_0, v \in S_\ell$, since S_ℓ is the node set of a connected component of $G(S^-)$. Hence,

$$c_{x, \lambda} [S^- \setminus (S_\ell \setminus V_t), \overline{S^-} \cup (S_\ell \setminus V_t)] \leq c_{x, \lambda} [S^-, \overline{S^-}].$$

Consequently, $[S^- \setminus (S_\ell \setminus V_t), \overline{S^-} \cup (S_\ell \setminus V_t)]$ is a dominating $s - t$ cut w.r.t. U , while $S^- \setminus (S_\ell \setminus V_t)$ is a proper subset of S^- , a contradiction. \square

For convenience, let x_ℓ denote the capacity of the unique arc in $E(S_\ell, t)$ (Figs. 3 and 4).




Fig. 3 Illustration of S^- , $S^-\setminus\{t\}$, U_0 and S_0, S_1, \dots, S_N node sets




Fig. 4 There is a path (in the undirected sense) unsaturated by g from $U_\ell \setminus U$ to s and from s to U

Proposition 10 *If $(x^*, \lambda^*) \in R_-$ violates the dominating $s - t$ cuts w.r.t. U , then it also violates the dominating $s - t$ cuts w.r.t. U_0 .*

Proof The minimal dominating $s - t$ cut w.r.t. U_0 is $[S_0, S^-\cup(\bigcup_{\ell=1}^N S_\ell)]$. There exists no arc uv such that $u \in S_0$ and $v \in S_\ell$ for any $\ell \in \llbracket N \rrbracket$, since S^- is not connected. Also, there exists no arc uv such that $u \in S_\ell$ and $v \in S^-\setminus\{t\}$ by Proposition 9. The inequality induced by $[S^-, S^-]$ is violated by (x^*, λ^*) , i.e.

$$\sum_{i=1}^m k_i \lambda_i^* < \sum_{j: v_j \in \bar{U}} x_j^*, \tag{29}$$

and the inequality induced by $[S_0, S^-\cup(\bigcup_{\ell=1}^N S_\ell)]$ is

$$\sum_{i=1}^m k_i \lambda_i \geq \sum_{j: v_j \in \bar{U}} x_j + \sum_{\ell=1}^N x_\ell, \tag{30}$$

which is also violated by (x^*, λ^*) . \square

From now on we assume that $G(S^-)$ is connected.

Now suppose that $[S, \bar{S}] \in C_{\min}(U)$ is violated by (x^*, λ^*) , and the subgraph induced by $\bar{S}^+ \setminus \{t\}$ is not connected in the undirected sense, and the node sets of its connected components are T_1, \dots, T_K for some $K > 1$. Let

$$U_\ell = U \cup \left(\bigcup_{k \in \llbracket K \rrbracket \setminus \{\ell\}} (V_t \cap T_k) \right)$$

for all $\ell \in \llbracket K \rrbracket$. We will show that at least one of the sets U_ℓ induces a facet of P_{\leq}^{emb} violated by (x^*, λ^*) .

Proposition 11 *There exists $\ell \in \llbracket K \rrbracket$ such that the dominating $s - t$ cut w.r.t. U_ℓ is violated by (x^*, λ^*) .*

Proof Since (x^*, λ^*) violates the dominating $s - t$ cuts w.r.t. U , we have

$$\sum_{i=1}^m k_i \lambda_i^* < \sum_{j: v_j \in V_t \setminus U} x_j^*. \quad (31)$$

Since there are no arcs between T_r and T_q for any $r \neq q$, the above inequality decomposes to

$$\sum_{i=1}^m \sum_{\ell=1}^K k_i^\ell \lambda_i^* < \sum_{\ell=1}^K \sum_{j: v_j \in T_\ell \cap V_t} x_j^* \quad (32)$$

where k_i^ℓ denotes the portion of k_i that enters T_ℓ , i.e.,

$$k_i^\ell = \sum_{e \in E_i(S^-, T_\ell)} \beta_e.$$

Suppose for a contradiction that for each $\ell \in \llbracket K \rrbracket$ the set U_ℓ does not induce a violated cut, i.e.,

$$\sum_{i=1}^m k_i^\ell \lambda_i^* \geq \sum_{j: v_j \in T_\ell \cap V_t} x_j^*. \quad (33)$$

Summing up these inequalities gives

$$\sum_{i=1}^m \sum_{\ell=1}^K k_i^\ell \lambda_i^* \geq \sum_{\ell=1}^K \sum_{j: v_j \in T_\ell \cap V_t} x_j^*,$$

which contradicts inequality (32), hence, there is at least one index ℓ such that (33) does not hold for ℓ . \square

Theorem 7 *If the subgraph induced by S^- is connected, then for any $\ell \in \llbracket K \rrbracket$, the dominating $s - t$ cut w.r.t. U_ℓ is facet inducing for P_{\leq}^{emb} .*

Proof Let $[Z^+, \overline{Z}^+]$ denote the maximal dominating $s - t$ cut w.r.t. U_ℓ . We first prove that U_ℓ satisfies condition *i*) of Theorem 5. To this end, it suffices to verify that $Z^+ = S^+ \cup (\bigcup_{r \neq \ell} T_r)$, since then $\overline{Z}^+ = T_\ell \cup \{t\}$, and T_ℓ is connected by definition. Recall that $\overline{S}^+ = \bigcup_{r=1}^K T_r \cup \{t\}$ and $S^+ \subset Z^+$ by Proposition 5. We have to prove that we get Z^+ from S^+ by moving all $T_r, r \neq \ell$ to the source side of the cut.

Firstly, we argue that $T_\ell \cap Z^+ = \emptyset$. Suppose for contradiction that $\emptyset \neq T_\ell \cap Z^+$. There is no path from any node of $T_\ell \cap Z^+$ to T_r for any $r \neq \ell$. Let $[S', \overline{S}']$ denote the $s - t$ cut we get as the result of extending S^+ by $Z^+ \cap T_\ell$. Observe that $S' \cap V_t = U$, since $T_\ell \cap Z^+ \cap V_t = \emptyset$. Let $Z^* = Z^+ \setminus T_\ell$. Then $[Z^*, \overline{Z}^*]$ is an $s - t$ cut w.r.t. U_ℓ . Recall the definition of k_i in (10), and define k'_i analogously for $[S', \overline{S}']$. If $[S^+, \overline{S}^+]$ dominates $[S', \overline{S}']$, then there exists i such that $k_i < k'_i$. Observe that

$$k'_i = k_i + \sum_{e \in E_i(Z^+ \cap T_\ell, T_\ell \setminus Z^+)} \beta_e - \sum_{e \in E_i(S^+, Z^+ \cap T_\ell)} \beta_e. \tag{34}$$

Since $k_i < k'_i$ by assumption, we have

$$\sum_{e \in E_i(Z^+ \cap T_\ell, T_\ell \setminus Z^+)} \beta_e > \sum_{e \in E_i(S^+, Z^+ \cap T_\ell)} \beta_e. \tag{35}$$

We define the coefficients k_i^+ and k_i^* for $[Z^+, \overline{Z}^+]$ and $[Z^*, \overline{Z}^*]$ using (10), respectively. We have

$$k_i^* = k_i^+ - \sum_{e \in E_i(Z^+ \cap T_\ell, T_\ell \setminus Z^+)} \beta_e + \sum_{e \in E_i(S^+, Z^+ \cap T_\ell)} \beta_e. \tag{36}$$

Now inequality (35) implies that $k_i^+ > k_i^*$, which means that $[Z^+, \overline{Z}^+]$ is dominated by $[Z^*, \overline{Z}^*]$, a contradiction. Otherwise, if $[S', \overline{S}']$ dominates $[S^+, \overline{S}^+]$, then $[S^+, \overline{S}^+]$ is not a dominating $s - t$ cut w.r.t. U , a contradiction. Finally, if $[S', \overline{S}']$ neither dominates $[S^+, \overline{S}^+]$, nor is dominated by $[S^+, \overline{S}^+]$, then they have the same capacity. But then, $[S^+, \overline{S}^+]$ is not a maximal dominating $s - t$ cut w.r.t. U , since $S^+ \subset (S^+ \cup (Z^+ \cap T_\ell))$.

Finally, we argue that for all $r \neq \ell$ we have $T_r \subseteq Z^+$. Suppose for contradiction that $T_r \setminus Z^+ \neq \emptyset$. By definition, there is no path between T_r and T_ℓ , however, there exists a path from $S^+ \subset Z^+$ to T_r , hence moving $T_r \setminus Z^+$ to Z^+ decreases the capacity of the resulting $s - t$ cut, which contradicts that the $s - t$ cut $[Z^+, \overline{Z}^+]$ is dominating.

Now we prove that condition *ii*) of Theorem 5 holds for Z^- , where $[Z^-, \overline{Z}^-]$ denotes the minimal dominating $s - t$ cut w.r.t. U_ℓ . We claim that there is no cut arc on any path from s to $\bigcup_{r \neq \ell} T_r$ in the $s - t$ cut $[Z^-, \overline{Z}^-]$. Let c^U, c^{U_ℓ} be capacity functions as described by Eq. (11) w.r.t. U and U_ℓ , respectively, and f be maximum flow in the network (G, c^U) . We construct a maximum flow g in (G, c^{U_ℓ}) . First let $g = f$. Let $\pi = s, v_0, \dots, v_n$ be a directed path in subgraph G_i where $v_n \in \bigcup_{r \neq \ell} T_r \cap V_t$. Since $c^{U_\ell}(v_n t) = 0$, we have $g_{v_{n-1} v_n} = 0$. For each arc e of π , let $g_e = f_e - f_{v_{n-1} v_n}$. Repeat this for all paths from s to the terminal nodes in $\bigcup_{r \neq \ell} T_r$, for all subgraphs G_i . The flow g is a feasible flow in (G, c^{U_ℓ}) and it is maximum flow, since we cannot increase the flow value on any paths from s to $T_\ell \cap V_t$, because f and

thus g saturates all cut arcs separating S^+ from T_ℓ . Now $g_e < f_e$ holds for all arcs of all directed paths from s to $\cup_{r \neq \ell} T_r$, hence, none of these arcs are cut arcs in $[Z^-, \overline{Z^-}]$, because those cut-arcs are saturated in every maximum flows in (G, c^{U_ℓ}) , see Fig. 4. Since $S^- \subseteq Z^-$ by Proposition 4, and $T_r, r \neq \ell$, is a subset of Z^- , by the above arguments, it follows that Z^- is connected. \square

6.3 Facet separation algorithm for $P_{\underline{=}}^{\text{emb}}$

The transformation of a set U into a facet inducing set U' is similar to the one described in Sect. 6.2. For simplicity we assume that $\kappa = 1$, i.e. $G \setminus \{s, t\}$ is connected in the undirected sense, otherwise we run the algorithm for the components separately. See Algorithm 4.

Algorithm 4 Facet separation for polytope $P_{\underline{=}}^{\text{emb}}$

Input: $U \subset V_t, (x^*, \lambda^*) \in R_{\underline{=}}$ such that the inequality (14) induced by U is violated by (x^*, λ^*)

Output: U' facet inducing and violated by (x^*, λ^*)

```

 $[S^-, \overline{S^-}] \leftarrow$  minimal dominating cut w.r.t.  $U$ 
if  $S^- \setminus \{s\}$  is not connected then
   $S_1, \dots, S_N \leftarrow$  connected components of  $S^- \setminus \{s\}$ 
  for  $\ell = 1, \dots, N$  do
    if  $S_\ell \cap V_t$  induces an inequality violated by  $(x^*, \lambda^*)$  then
       $U \leftarrow S_\ell \cap V_t$ 
      break
    end if
  end for
end if
 $[S^+, \overline{S^+}] \leftarrow$  maximal dominating cut w.r.t.  $U$ 
if  $\overline{S^+} \setminus \{t\}$  is not connected then
   $T_1, \dots, T_K \leftarrow$  connected components of  $\overline{S^+} \setminus \{t\}$ 
  for  $\ell = 1, \dots, K$  do
    if  $U \setminus \bigcup_{r \in [K] \setminus \{\ell\}} (T_r \cap V_t)$  induces an inequality violated by  $(x^*, \lambda^*)$  then
       $U' = U \setminus \bigcup_{r \in [K] \setminus \{\ell\}} (T_r \cap V_t)$ 
    end if
  end for
else
   $U' = U$ 
end if
return inequality (14) for  $U'$ 

```

The following statements ensure the correctness of the algorithm. First, assume that $G(S^- \setminus \{s\})$ is not connected, and let S_1, \dots, S_N be the node-sets of its connected components. Let $U_\ell = S_\ell \cap V_t$.

Proposition 12 *There exists $\ell \in [N]$ such that the inequality induced by $[S_\ell, \overline{S_\ell}]$ is violated by (x^*, λ^*) .*

Proof Suppose (x^*, λ^*) violates inequality (14) for set U . By Eq. (5), this is equivalent to

$$\sum_{i=1}^m (\alpha_i - k_i) \lambda_i^* > \sum_{j: v_j \in U} x_j^*. \quad (37)$$




Fig. 5 Illustration of the $k_i^0, k_i^1, \dots, k_i^N$ and $\alpha_i^0, \alpha_i^1, \dots, \alpha_i^N$ values

We will decompose the α_i and the k_i the following way. Let

$$\begin{aligned}
 k_i^\ell &= \sum_{e \in E_i(s_\ell, \bar{S}^-)} \beta_e \quad \forall \ell \in \llbracket N \rrbracket, \forall i \in \llbracket m \rrbracket \\
 \alpha_i^\ell &= \sum_{e \in E_i(s, S_\ell)} \beta_e \quad \forall \ell \in \llbracket N \rrbracket, \forall i \in \llbracket m \rrbracket \\
 k_i^0 = \alpha_i^0 &= \sum_{e \in E_i(s, \bar{S}^-)} \beta_e \quad \forall i \in \llbracket m \rrbracket.
 \end{aligned}
 \tag{38}$$

See Fig. 5 for an illustration. Observe that

$$\alpha_i = \alpha_i^0 + \sum_{\ell=1}^N \alpha_i^\ell, \quad k_i = k_i^0 + \sum_{\ell=1}^N k_i^\ell.$$

We can decompose the left hand side of inequality (37) as follows:

$$\begin{aligned}
 \sum_{i=1}^m (\alpha_i - k_i) \lambda_i^* &= \sum_{i=1}^m \left(\alpha_i^0 + \sum_{\ell=1}^N \alpha_i^\ell - k_i^0 - \sum_{\ell=1}^N k_i^\ell \right) \lambda_i^* \\
 &= \sum_{i=1}^m \left(\sum_{\ell=1}^N \alpha_i^\ell - \sum_{\ell=1}^N k_i^\ell \right) \lambda_i^* \\
 &= \sum_{\ell=1}^N \sum_{i=1}^m (\alpha_i^\ell - k_i^\ell) \lambda_i^*,
 \end{aligned}$$

and the right hand side as

$$\sum_{j: v_j \in U} x_j^* = \sum_{\ell=1}^N \sum_{j: v_j \in U_\ell} x_j^*.$$

Therefore, inequality (37) implies that there exists an ℓ such that

$$\sum_{i=1}^m (\alpha_i^\ell - k_i^\ell) \lambda_i^* > \sum_{j: v_j \in U_\ell} x_j^*.$$

Subtracting this from equality (5), we get

$$\sum_{i=1}^m (\alpha_i - \alpha_i^\ell + k_i^\ell) \lambda_i^* < \sum_{j: v_j \in V_i \setminus U_\ell} x_j^*. \quad (39)$$

Observe that (39) means that the inequality induced by the $s-t$ cut $[S_\ell, \overline{S_\ell}]$ is violated by (x^*, λ^*) . \square

Now suppose that $S^- \setminus \{s\}$ is connected, but $\overline{S^+} \setminus \{t\}$ is not, and the node-sets of its connected components are T_1, \dots, T_M . Now let

$$S_\ell^+ = S^+ \cup \left(\bigcup_{r \in \llbracket M \rrbracket \setminus \{\ell\}} T_r \right) \text{ and } U_\ell = V_i \setminus T_\ell.$$

Proposition 13 *There exists $\ell \in \llbracket M \rrbracket$ such that the inequality induced by $[S_\ell^+, \overline{S_\ell^+}]$ is violated by (x^*, λ^*) .*

Proof Recall that the dominating cut w.r.t. U being violated is equivalent to inequality (31). We decompose k_i the following way. Let

$$k_i^\ell = \sum_{e \in E_i(S^+, T_\ell)} \beta_e \quad \forall \ell \in \llbracket M \rrbracket, \forall i \in \llbracket m \rrbracket. \quad (40)$$

Observe that $k_i = \sum_{\ell=1}^M k_i^\ell$. Now we can decompose inequality (31):

$$\sum_{\ell=1}^M \sum_{i=1}^m k_i^\ell \lambda_i^* < \sum_{\ell=1}^M \sum_{j: v_j \in V_i \setminus U_\ell} x_j^*. \quad (41)$$

Therefore, for at least one index ℓ we have

$$\sum_{i=1}^m k_i^\ell \lambda_i^* < \sum_{j: v_j \in V_i \setminus U_\ell} x_j^*, \quad (42)$$

which implies the statement. \square

Theorem 8 *If the subgraph $S^- \setminus \{s\}$ is connected, then for any $\ell \in \llbracket M \rrbracket$ the set U_ℓ is facet inducing.*

Proof We prove that the conditions (i) and (ii) of Theorem 6 hold for U_ℓ for any $\ell \in \llbracket K \rrbracket$. Let $\left[Z^+, \overline{Z^+} \right]$ denote the maximal dominating $s - t$ cut w.r.t. U_ℓ . First we prove that $Z^+ = S^+ \cup \left(\bigcup_{r \neq \ell} T_r \right)$ and consequently $\overline{Z^+} = T_\ell \cup \{t\}$. Note that this implies that U_ℓ satisfies condition (i) of Theorem 6, since T_ℓ is connected by definition.

We define the arc capacities c^U, c^{U_ℓ} , and $s - t$ flows f and g similarly to those in the proof of Theorem 7. Then f carries 0 flow to the nodes in U , and maximum flow to the nodes in $V_i \setminus U$, and likewise, g carries 0 flow to the nodes in U_ℓ and maximum flow to the nodes in $V_i \setminus U_\ell$. First we prove that for all $r \neq \ell$ we have $T_r \subseteq Z^+$. Since there is a cut arc between S^+ and T_r which is saturated by f by construction, there is a directed path π from s to some $v_j \in T_r \cap U_\ell$ with a positive f -flow. Since $U \subset U_\ell$, g does not carry any flow to v_j by construction, $g_e < f_e$ for each arc e of π , and thus g does not saturate any arc on π . This means that there is no cut-arc (for the capacities c^{U_ℓ}) on any path from s to any T_r , $r \neq \ell$, hence $T_r \subset Z^+$ for $r \neq \ell$. Moreover, this shows that $T_r \subseteq Z^-$. \square

Second, we prove that $T_\ell \cap Z^+ = \emptyset$. The arcs that cut T_ℓ from S^+ are saturated by g , hence they cut T_ℓ from Z^+ . Hence, $Z^+ = S^+ \cup \left(\bigcup_{r \neq \ell} T_r \right)$ as claimed.

Let $\left[Z^-, \overline{Z^-} \right]$ denote the minimal dominating $s - t$ cut w.r.t. U_ℓ , and we verify the condition (ii) of Theorem 6. We proceed with the following:

Claim 1 *The subgraph spanned by $S^+ \setminus \{s\}$ is connected in the undirected sense.*

Proof Suppose for contradiction that the subgraph spanned by $S^+ \setminus \{s\}$ has multiple components, S_0, S_1, \dots, S_K . Since the subgraph spanned by $S^- \setminus \{s\}$ is connected by assumption, one of these components contains $S^- \setminus \{s\}$. Let S_0 be that component. Since $S^+ \cap V_i = U = S^- \cap V_i$, no other component contains any terminal nodes. There exists at least one arc that emanates S_k and enters T_r for some $k \in \llbracket K \rrbracket$ and $r \in \llbracket N \rrbracket$, otherwise no terminal node is reachable from the nodes of S_k , which contradicts that from S_k at least one terminal node is reachable. Since S_k contains no terminal nodes or node s , it is a rooted tree with root $w \in V(G_i)$ for some i . If $c_w > \beta_{e_i(w)}$, then the $s - t$ cut $\left[S^+, \overline{S^+} \right]$ is dominated by $s - t$ cut $\left[S^+ \setminus S_k, \overline{S^+} \cup S_k \right]$, contradiction. If $c_w \leq \beta_{e_i(w)}$ then the capacity function is not reduced, since all arcs leaving S_k enter $\overline{S^+}$, contradiction. Hence we have that no such S_k exists, and $S^+ \setminus \{s\}$ is connected. \square

To finish the proof of the theorem, it suffices to prove the following:

Claim 2 *For each $r \neq \ell$, there exists an undirected path connecting a node in $T_r \cap U_\ell$ with a node in U containing no cut arc in $\left[Z^-, \overline{Z^-} \right]$.*

Proof First suppose there exists an arc e between S^- and T_r .

So there is an undirected path π containing e , which connects some node in U to some node in $T_r \cap U_\ell$. No arc of π is saturated by g , since the part of π in S^- contains no saturated arc by Proposition 2 5), the part of π in T_r has zero flow by definition of g , and similarly $g_e = 0$. Since no arc of π is saturated by g , it contains no cut arc.

If the arc e above does not exist, then there is an undirected path between T_r and S^- which contains a node from $S^+ \setminus S^-$. Recall that by Proposition 6, the subgraph spanned by $S^+ \setminus S^-$ decomposes to rooted trees. Let τ denote the rooted tree component of $S^+ \setminus S^-$ that contains a part of the said undirected path, with root node v . The parent node $p_i(v)$ of v is either in




Fig. 6 If $p_i(v) \in S^- \setminus \{s\}$, there is an undirected path unsaturated by g from $U_\ell \setminus U$ to $p_i(v)$ and from $p_i(v)$ to U


$S^- \setminus \{s\}$ or it is s . If $p_i(v) \in S^- \setminus \{s\}$, then there is an undirected path from $T_r \cap U_\ell$ to U with no saturated arc, since any path from s to T_r has no arcs saturated by g as seen in the proof of Theorem 7, hence, the path from $p_i(v)$ to T_r has no saturated arcs, and any path from $p_i(v)$ to U within S^- has no saturated arcs by Proposition 2 5), see Fig. 6.

If $p_i(v) = s$, then there is at least one arc from τ to S^- , since $S^+ \setminus \{s\}$ is connected as shown above. Since there is a path from node v to T_r with arcs unsaturated by flow g , it is enough to show that there exists a path from v to S^- with arcs unsaturated by g . Suppose for contradiction, that all paths from v to S^- contain at least one arc saturated by g , and hence saturated by f , see Fig. 7a. Observe that τ admits a maximal subtree τ' rooted at v such that the flow f saturates no arc in τ' . Then there exists no arc from any node of τ' to some node in S^- by our indirect assumption. Since all arcs leaving τ' are saturated by f , we have

$$\beta_{e_i(v)} = f_{e_i(v)} = \sum_{e \in \delta(\tau')} f_e = \sum_{e \in \delta(\tau')} \beta_e. \tag{43}$$

Hence, all arcs of τ' are saturated by f , therefore $V(\tau') = \{v\}$ (see Fig. 7b), and thus the network G is not reduced (see Proposition 1), a contradiction. Therefore, there must exist at least one path from v to S^- with all arcs unsaturated by flow f , and consequently unsaturated by flow g .

This completes the proof of the theorem. □



(a) If $p_i(v) \in S^- \setminus \{s\}$, there is a path (in the undirected sense) unsaturated by g from $U_\ell \setminus U$ to $p_i(v)$.

(b) The τ and τ' subtrees rooted at v . By the indirect assumption each path from v to S^- has at least one arc saturated by f .

Fig. 7 The wavy lines symbolize paths with no arcs saturated by g , the thick straight lines symbolize paths with at least one saturated arc by g

7 Computational experiments

To assess the computational efficiency of the proposed facet separation procedure (Algorithm 4), we implemented it within a branch-and-cut framework, and we performed computational experiments on a set of problem instances described below. The goal of these experiments is to demonstrate the potential computational advantage of the separation algorithm, and we chose a rather general test problem to do so.

7.1 Test problem

Let $K, m, n \in \mathbb{N}_{>0}$ and $\mathcal{L}_i^k \subseteq 2^{[n]}$ be a laminar family of subsets of $[n]$ for $k \in [K]$ and $i \in [m]$. For all $L \in \mathcal{L}_i^k$ define $b_L \geq 0$, and for all $k \in [K]$ and $i \in [m]$, let $\alpha_i^k > 0$. For each $i \in [m]$, we define the polytope $P(i, k)$ as the convex hull of the non-negative solutions of the system of inequalities described by the laminar family \mathcal{L}_i^k , α_i^k and $\{b_L\}_{L \in \mathcal{L}_i^k}$, as in Theorem 4, i.e.,

$$P(i, k) = \left\{ x \in \mathbb{R}_{\geq 0}^n \mid \sum_{j=1}^n x_j = \alpha_i^k, \sum_{j \in L} x_j \leq b_L \forall L \in \mathcal{L}_i^k \right\}. \tag{44}$$

By Theorem 4, $\text{conv}(\bigcup_{i=1}^m P(i, k)^{emb})$ has a network flow representation.

Let $w_j, \beta_j > 0$ for all $j \in \llbracket n \rrbracket$, while the c_i^k (for $k \in \llbracket K \rrbracket, i \in \llbracket m \rrbracket$) are arbitrary rational numbers. Our goal is to solve the following problem:

$$\begin{aligned} \min \quad & \sum_{j=1}^n w_j y_j + \sum_{k=1}^K \sum_{i=1}^m c_i^k \lambda_i^k \\ \text{s.t.} \quad & \sum_{k=1}^K x_j^k \leq \beta_j + y_j, \quad \forall j \in \llbracket n \rrbracket \\ & (x^k, \lambda^k) \in \bigcup_{i=1}^m P(i, k)^{\text{emb}}, \quad \forall k \in \llbracket K \rrbracket. \end{aligned} \quad (45)$$

In this problem we have to find a set of vectors $x^k \in \mathbb{R}^n, k \in \llbracket K \rrbracket$, such that each x^k is in one of the polytopes $P(i, k), i \in \llbracket m \rrbracket$, the λ^k indicates which one, and the x^k also satisfy a set of linear constraints, with the objective of minimizing the violation of these constraints and the total cost of the chosen alternatives.

We can restate (45) as a mixed integer-linear program:

$$\begin{aligned} \min \quad & \sum_{j=1}^n w_j y_j + \sum_{k=1}^K \sum_{i=1}^m c_i^k \lambda_i^k \\ \text{s.t.} \quad & \sum_{k=1}^K x_j^k \leq \beta_j + y_j, \quad \forall j \in \llbracket n \rrbracket \\ & \sum_{j \in L} x_j^k \leq b_L + (1 - \lambda_i^k) M_k, \quad \forall L \in \mathcal{L}_i^k, i \in \llbracket m \rrbracket, \forall k \in \llbracket K \rrbracket \\ & \sum_{j=1}^n x_j^k = \sum_{i=1}^m \alpha_i^k \lambda_i^k, \quad \forall k \in \llbracket K \rrbracket \\ & \lambda_i^k \in \{0, 1\}, \quad \forall i \in \llbracket m \rrbracket, \forall k \in \llbracket K \rrbracket \\ & x_j^k \geq 0, \quad \forall j \in \llbracket n \rrbracket, \forall k \in \llbracket K \rrbracket. \end{aligned} \quad (46)$$

The constant M_k in (46) equals $\sum_{i=1}^m \alpha_i^k + 1$ for each $k \in \llbracket K \rrbracket$. We will refer to (46) as the *big-M formulation*.

7.2 Methods compared and test environment

We implemented our separation procedure (Algorithm 4) in C++. We used FICO XPRESS v9.4.1 for solving the mixed-integer linear program (46) with branch-and-cut. In our method, called D-cuts, we used presolve with default settings, except for dual reductions, which was turned off to enable cut generation. The built-in cuts of XPRESS were disabled. We separated disjunctive cuts in the root node of the search tree in at most 20 rounds, and then in one round in every node of depth at most 20. A violated cut found by Algorithm 4 was added to the LP relaxation of a node only if the absolute violation was at least 0.1. We compared the performance of our D-cuts method to three other approaches.

The second method, called B&B, in our comparison was the branch-and-bound procedure of XPRESS applied to (46) with presolve turned on (no cut generation at all).

The third approach, called XPRS-cuts, is the default branch-and-cut of the XPRESS solver applied to (46), using presolve and the built-in cuts of the solver.

In the fourth method we applied Balas' reformulation (Theorem 2) to each disjunctive constraint in (45), and solved the resulting MIP with presolve turned on.

Table 1 Results on small and medium size instances

n	D-cuts					B&B				
	Gap	Time	Nodes	Cuts [†]	Root gap	Gap	Time	Nodes	Cuts	Root gap
30	–	2.4s	1.35	109.9	0.0013	–	9.62s	1156.6	–	0.107
40	–	4.7s	1.1	161	0.0004	0.003 (9)	68.5s	7323	–	0.1009
50	–	6.7s	1.4	149.8	0.0001	0.016 (4)	189.6s	14907.8	–	0.1096
60	–	14.6s	1	217.5	0.0002	0.061 (0)	240s	10514.4	–	0.1272
70	–	26.6s	3	425.9	0.0007	0.084 (1)	237.5s	4601.7	–	0.1334

n	XPRS-cuts					Balas' reformulation				
	Gap	Time	Nodes	Cuts [‡]	Root gap	Gap	Time	Nodes	Cuts [‡]	Root gap
30	–	4.28s	2.6	n.a	0.0045	–	2 s	1.05	0	0.0001
40	–	6.7s	4.5	n.a	0.0026	–	5.9s	1.25	0	0.0001
50	–	10.6s	2.5	n.a	0.0014	–	11.9s	1.1	0	0.0001
60	–	26.4s	38.2	n.a	0.0073	–	24.7s	1	0	0.0001
70	–	65.1s	165.3	n.a	0.0031	–	47.1s	1	n.a	0.0001

[†] Average number of D-cuts separated

[‡] Built-in cuts of XPRESS separated

The experiments were performed on a notebook computer with i7-8850H CPU @ 2.60GHz and Windows operating system. The root relaxation was always solved using 8 CPU threads by the barrier solver and 1 thread by the dual simplex method, whereas the subsequent tree-search and cut generation use only one CPU thread. The run-time limit was set to 240 s for the smaller instances and 1200 s for the larger ones.

7.3 Design and evaluation of computational experiments

The test problems were generated as follows. The laminar families defining each polyhedron $P(i, k)$ have depth 3 in all test instances, cf. Sect. 3. We generated 10 random problem instances for each $n \in \{30, 40, 50, 60, 70, 100, 130\}$, while m and K were both set to n in all cases. The laminar sets \mathcal{L}_i^k , the parameters α_i^k , and the right-hand-sides b_L for each $L \in \mathcal{L}_i^k$ were chosen randomly while ensuring that the corresponding network be reduced. Finally, the parameters β_j , and the weights w_j and c_i^k were chosen as follows: β_j is a random number from the interval $[\underline{b}_j/2, 1.5\underline{b}_j]$, where $\underline{b}_j = \sum_{k=1}^K \min\{b_L \mid j \in L, L \in \mathcal{L}_i^k, i \in \llbracket m \rrbracket\}$ is the sum of the smallest non-zero upper bounds on x_j over all the disjunctive constraints. The weights w_j and c_i^k are chosen uniformly at random from the intervals $[0, 10.0]$ and $[-200, +200]$, respectively. The above choices were made after some preliminary tests to make difficult instances.

7.3.1 Results on small and medium size instances

In this section we summarize our computational results on those instances, where $30 \leq n \leq 70$. We ran all four methods on each problem instance in this class. Table 1 shows for each n average values over the 10 instances of the group. For each method and group of instances, we provide the average optimality gap (*gap*), the average running time in seconds (*time*), the average number of search tree nodes (*nodes*), the average number of cuts (*cuts*), and

the average root gap (*root gap*). The optimality gap of a method on a problem instance is computed as $(ub - lb)/ub$, where ub and lb are the best upper and lower bounds obtained by the method, respectively. A dash '-' in the column 'gap' indicates that the optimum was found for all instances in the group. In case of a positive average gap, we provide in parenthesis the number of instances solved optimally among the 10 instances in the group. The content of the column *cuts* depends on the method. For the D-cuts method, it contains the average number of cuts separated, for B&B it is a '-' throughout, since no cuts are generated at all. In the case of the XPRS-cuts method, only the built-in cuts of the solver are used, but we have no data available for their number (n.a.), whereas when using Balas' reformulation, no built-in cuts were generated in most cases, except for one instance for $n = 70$. The *root-gap* is calculated by the formula $(ub - root_lb)/ub$, where ub is the best upper bound, and $root_lb$ is the lower bound after processing the root node including the separation of cuts when it applies.

In all the approaches, the node LPs were presolved by the XPRESS solver, which reduced the computation times significantly. The root LPs were solved with default setting in all cases, which means that the solver applied the dual simplex using one thread, and the parallel barrier solver using 8 threads. After solving the root LP, the D-cuts method strengthened it by at most 20 rounds of cut generation, which led to the optimal solution in most cases. This can be seen by the low average number of search tree nodes. We can observe that as n increases, the average computation time, and the average number of cuts increase as well, but the average number of search tree nodes remains low. In contrast, without cut generation, the B&B method found the optimum only for small values of n . As n increases, the number of instances solved optimally decreases, while the average optimality gap increases. This is due to the run-time limit of 240 s, since as n increases, the size of the MIP formulation increases as well. By the same token, the average solution time increases, and the number of search tree nodes visited decreases. The XPRS-cuts method clearly outperforms the B&B method, but performs worse than D-cuts, especially for larger values of n .

Balas' extended formulation has a stable behavior. Note that the resulting MIP has much more variables and constraints than the big-M formulation. The reason is that the x^k vector variables are copied as many times as the number of alternatives in the disjunctive constraints, cf. (2). In practice, this means that if $x^k \in \mathbb{R}^n$, then there are n copies of x^k , since we have $m = n$ alternatives in every disjunctive constraint in all the problem instances. The number of constraints is multiplied analogously. We also observed that presolve could not reduce the size of the resulting MIPs. However, solving the root LPs using 9 CPU threads helped a lot in reducing the computation times. As it turned out, the barrier solver implementation of XPRESS exploited parallel processing to a great extent. Without this feature, Balas' extended formulation would not be a competitive method for solving the problem at hand. In most cases, the solution of the root LP and presolve found an optimal solution, and the built-in cuts were separated only for one problem instance with $n = 70$.

Based on the results, we can conclude that for $n \geq 40$, the fastest method is D-cuts, and the second place is shared between XPRS-cuts and Balas' extended formulation. Note though that with Balas' extended formulation the optimum was found in the root node in most cases, and it had a more stable behavior than XPRS-cuts. The longer running time with Balas' extended formulation is primarily due to the significantly larger problem sizes the solver must handle with this approach. However, parallel processing also plays a crucial role in efficiently solving the root LPs, particularly when using Balas' extended formulation.

Table 2 Results on large instances

n	D-cuts					D-cuts + XPRS-cuts				
	Gap	Time	Nodes	Cuts [†]	Root gap	Gap	Time	Nodes	Cuts [†]	Root gap
100	–	125.9s	3	605.7	0.0007	–	119.1s	3.2	97.5	0.0010
130	–	358.4s	12.1	915.8	0.0195	–	294.1s	8.3	145.4	0.0179
n	XPRS-cuts					Balas' reformulation				
	Gap	Time	Nodes	Cuts [‡]	Root gap	Gap	Time	Nodes	Cuts [‡]	Root gap
100	–	133.2s	91.8	n.a	0.0119	–	242.5s	1.8	n.a	0.0001
130	–	393.5s	372.1	n.a	0.1238	–	553.2s	1.3	n.a	0.0001

[†] Average number of D-cuts separated

[‡] Built-in cuts of XPRESS separated

7.3.2 Results on large instances

We also tested our methods on instances with $n \in \{100, 130\}$ jobs. We run four methods on each problem instance: D-cuts, XPRS-cuts, Balas' reformulation, and a combination of D-cuts, and XPRS-cuts, which we call "D-cuts + XPRS cuts". In this method, we configured the xpress solver to use built-in cuts, and also our separation procedure for D-cuts. Our separation subroutine was called by the solver after generating built-in cuts. In the root node of the search tree D-cuts cuts were separated in at most 10 rounds, and for one round in all other nodes. We have also tested B&B, but it was always inferior to any other methods, so we do not provide detailed results with that method. We set the run-time limit to 1200 s.

Table 2 presents the results, highlighting that the "D-cuts + XPRS-cuts" method is the fastest on average, followed closely by D-cuts. Both of these methods generate only a small number of search tree nodes. In contrast, XPRS-cuts is slower on average, while Balas' reformulation, though the slowest, generates the fewest search tree nodes.

Figure 8 illustrates the average run time and variance across the 10 instances for $n \in \{100, 130\}$. Notably, D-cuts and Balas' reformulation exhibit the lowest run time variance across both instance groups, while XPRS-cuts shows the highest variance.

We conclude that on the large instances, the best method is "D-cuts + XPRS-cuts", and D-cuts is the second best.

8 Final remarks

In this paper, we have described polynomial time exact separation algorithms for disjunctive constraints with a network-flow representation. We have also identified the disjunctive constraints that can be represented in this manner. Our computational experiments on a set of benchmark problems demonstrate the superiority of our approach. The results indicate that the new cuts can significantly reduce the computational time of a branch-and-cut procedure, outperforming general cutting planes applied to the same formulation.

Balas' reformulation shows a solid performance. In most cases, the optimum of a test problem was found already in the root node without cutting and branching (with the help of presolve and heuristics). In fact, the LP relaxation of Balas' reformulation of (45) is tighter than that of (46) without adding our cutting planes. Nevertheless, when a disjunctive constraint can be represented by a network flow, our modeling approach and separation




Fig. 8 Average run time and variance

procedures constitute a viable alternative to Balas' extended formulation, as our preliminary computational results demonstrate.

A further approach for solving MILPs with disjunctive constraints is Dantzig–Wolfe reformulation with subsequent column generation. For instance, Sadykov and Vanderbeck (2013) applied branch-and-price for solving bin packing problems with conflicts, while Almathkour et al. (2024) proposed branch-and-cut-and-price for the 2-connected subgraph problem with disjunctive constraints.

Our methodology can be applied to any disjunctive constraint with a network-flow representation, facilitated by our general separation procedure that only requires a weighted network as specified in Sect. 2.

In the future, we plan to apply our methodology to specific problems that include network-flow representable disjunctive constraints.

Appendix A Proofs of statements in Sect. 2

Proof of Proposition 1 First we reduce the capacity function such that for each $i \in \llbracket m \rrbracket$, and $v \in V(G_i) \setminus (\{s\} \cup V_i)$:

$$\begin{aligned} \beta_e &\leq \beta_{e_i(v)} \quad \forall e \in \delta(v), \\ \beta_{e_i(v)} &\leq \sum_{e \in \delta(v)} \beta_e. \end{aligned} \tag{A1}$$

Let $c'_{x,\lambda}$ be the resulting capacity function. Then the set of feasible flows in $N = (G, c_{x,\lambda})$ and $N'' = (G, c'_{x,\lambda})$ is the same, see Kis and Horváth (2022). Next we apply the following




Fig. 9 Replacement of $uv \cup \delta(v)$

transformation for any $v \in V(G_i) \setminus (\{s\} \cup V_i)$ such that

$$\beta_{e_i(v)} = \sum_{e \in \delta(v)} \beta_e.$$

Let $u = p_i(v)$ be the parent node of v . We drop the edge uv of G_i and replace each edge $vw \in \delta(v)$ by the edge uw of the same capacity, see Fig. 9. Let $N' = (G', c'_{x,\lambda})$ be the resulting network. It is easy to see that if f is a feasible $s - t$ flow in $N = (G, c_{x,\lambda})$, then there exists a feasible $s - t$ flow f' in N' such that $\sum_{e \in \delta_G(s)} f_e = \sum_{e \in \delta_{G'}(s)} f'_e$, and vice versa, from which the statement follows. \square

Proof of Proposition 2 Properties (1) and (2) follow from $U \subset S^-$ and $\bar{U} \subset \bar{S}^-$. Property (3) holds, because the cut-arcs are tight. Because of property (ii), all paths from s to \bar{U} contains at least one tight arc. Property (4) holds, otherwise if there were a proper subset of E^- which is still an $s - t$ cut, we could decrease the capacity of the cut. For property (5), suppose for the sake of a contradiction that there exist an arc $uv \in E^-$ and a tight arc ww' on the unique path from s to u . Note that since $uv \in E^-$, it is tight, hence, $v \notin U$. Then $[S^- \setminus \sigma(w'), \bar{S}^- \cup \sigma(w')]$ is also an $s - t$ cut w.r.t. U , and $S^- \setminus \sigma(w') \subset S^-$. We claim that the capacity of the cut $[S^- \setminus \sigma(w'), \bar{S}^- \cup \sigma(w')]$ is the same as that of $[S^-, \bar{S}^-]$. By the definition of c^U , all the flow through ww' goes to the nodes in \bar{U} , hence the total capacity of cut arcs in $\sigma(w')$ equals to that of arc ww' . Hence the two $s - t$ cuts have the same capacity. This contradicts that $[S^-, \bar{S}^-]$ is a minimal dominating $s - t$ cut. \square

Proof of Proposition 3 Properties (1) to (4) can be proved analogously to those of Proposition 2. For property (5), suppose for contradiction that there exists an arc $uv \in E^+$ such that $t \neq v$ and each path from v to \bar{U} has a tight arc. Let E' be a subset of these cut-arcs such that no subset of E' covers all paths from v to \bar{U} . By flow conservation, the total capacity of the arcs of E' equals to that of uv . Then S^+ can be extended by the nodes of $\sigma(v)$ from v to the tails of the arcs of E' . The resulting $s - t$ cut has the same capacity as $[S^+, \bar{S}^+]$, and its source side strictly contains S^+ . Hence, $[S^+, \bar{S}^+]$ is not a maximal dominating $s - t$ cut, which is a contradiction. \square

Proof of Proposition 4 Let $c^U, c^{U'}$ be capacity functions as described by Eq. (11) w.r.t. U and U' , respectively, and f be maximum flow in the networks (G, c^U) . We construct a maximum flow g in $(G, c^{U'})$. First let $g = f$. Let $\pi = s, v_0, \dots, v_n$ be a directed path in

G where $v_n \in U' \setminus U$. Since $c^{U'}(v_n t) = 0$, we have $g_{v_{n-1}v_n} = 0$. For each arc e of π , let $g_e = f_e - f_{v_{n-1}v_n}$. Repeat this for all paths from s to the terminal nodes in $U' \setminus U$. Let μ_e denote the total decrease of flow on arc e . Let π' denote a path from s to $\overline{U'}$. For each arc e on the path π' , let $g_e = g_e + \min_{e \in \pi'} \{\beta_e - g_e\}$. Repeat this for all paths from s to $\overline{U'}$. Let v_e denote the total increase of flow on arc e . We argue that for each arc spanned by S^- we have $\mu_e \geq v_e$. If not, then f was not a maximum $s - t$ flow in (G, c^U) .

The flow g is a feasible flow in $(G, c^{U'})$ and it is a maximum $s - t$ flow in $(G, c^{U'})$, since we cannot increase the flow value on any paths from s to $\overline{U'}$.

Observe that no arc e spanned by S^- is saturated by g , since $\mu_e \geq v_e$. \square

Proof of Proposition 5 Let $c^U, c^{U'}$ be capacity functions as described by Eq. (11) w.r.t. U and U' , respectively, and f be maximum flow in the networks (G, c^U) . We construct a maximum flow g in $(G, c^{U'})$. First let $g = f$. Let $\pi = s, v_0, \dots, v_n$ be a directed path in G where $v_n \in U' \setminus U$. Since $c^{U'}(v_n t) = 0$, we have $g_{v_{n-1}v_n} = 0$. For each arc e of π , let $g_e = f_e - f_{v_{n-1}v_n}$. Repeat this for all paths from s to the terminal nodes in $U' \setminus U$. Let μ_e denote the total decrease of flow on arc e . Let π' denote a path from s to $\overline{U'}$. For each arc e on the path π' , let $g_e = g_e + \min_{e \in \pi'} \{\beta_e - g_e\}$. Repeat this for all paths from s to $\overline{U'}$. Let v_e denote the total increase of flow on arc e .

If there is an arc uv spanned by S^+ which is saturated by g , consider the subtree τ_v rooted at v . Let $E' = E(S^+, \overline{S^+}) \cap E(\tau_v)$. By definition, f saturates all arcs in E' . Since $\mu_{uv} \geq v_{uv}$, f saturates arc uv as well, hence, the capacity of uv is equal to the sum of capacities of arcs in E' . The flow g on arc uv can only go through the arcs of E' , since there is no flow from v to any node in U . Therefore, g must saturate all arcs in E' . Hence, by Proposition 3 5), uv is not a cut arc. Consequently, no arc spanned by S^+ is a cut arc in $(G, c^{U'})$, hence, $S^+ \subset Z^+$. \square

Proof of Proposition 6 It follow from the structure of the network that any subgraph spanned by a subset of $V(G) \setminus (V_t \cup \{s, t\})$ decomposes to rooted trees. The node-set $S^+ \setminus S^-$ does not contain terminal nodes, and the nodes s and t , hence, it spans rooted trees. Let τ_v denote the rooted tree with root node $v \in S^+ \setminus S^-$. Suppose $v \in V(G_i)$. Since $v \in S^+$, it is on a directed path from source s to some terminal node in U . Recall the definition of c_v^i in Sect. 1. By definition, $c_v^i(S^+) = \sum_{e \in E_i(V(\tau_v), \overline{S^+})} \beta_e$.

Suppose for contradiction that $c_v^i \neq \beta_{e_i(v)}$. If $c_v^i > \beta_{e_i(v)}$, then the $s - t$ cut $[S^+, \overline{S^+}]$ is dominated by the $s - t$ cut $[S^+ \setminus V(\tau_v), \overline{S^+} \cup V(\tau_v)]$. If $c_v^i < \beta_{e_i(v)}$, then the $s - t$ cut $[S^-, \overline{S^-}]$ is dominated by the $s - t$ cut $[S^+, \overline{S^+}]$. \square

Appendix B Proofs from Sect. 5

Proof of Theorem 6 Necessity: Since $P_{\leq}^{\text{emb}} \subseteq P_{\leq}^{\text{emb}}$, the set of the facets of P_{\leq}^{emb} is a subset of the set of the facets of P_{\leq}^{emb} , therefore, condition (1) is necessary, as proved in Theorem 5. Indirectly, suppose that condition (2) is not satisfied by U , and the subgraph of G induced by $S^- \setminus \{s\}$ has multiple connected components Z_1, \dots, Z_q . Let U_i denote $V(Z_i) \cap V_t$. Recall the inequality (14) induced by U , and the corresponding face F_U defined by Eq. (15), or equivalently, by Eq. (16).

For $\ell \in [q]$, let α_i^ℓ denote the sum of the coefficients of λ_i on those arcs of G_i that enter the component Z_ℓ , α_i^0 is that of those arcs from s to $\overline{S^-}$, k_i^ℓ is the sum of coefficients of λ_i

on those cut arcs that leave the component Z_ℓ , and k_i^0 is that of those cut arcs from s to $\overline{S^-}$. Clearly, $k_i^0 = \alpha_i^0$, and

$$\alpha_i = \sum_{\ell=0}^q \alpha_i^\ell, \quad k_i = \sum_{\ell=0}^q k_i^\ell. \tag{B2}$$

Using this, we can rewrite the left hand side of equality (16):

$$\sum_{i=1}^m (\alpha_i - k_i) \lambda_i = \sum_{i=1}^m \left(\sum_{\ell=1}^q (\alpha_i^\ell - k_i^\ell) \right) \lambda_i \tag{B3}$$

We can observe that $U = \cup_{\ell=1}^q U_\ell$. The right hand side of Eq. (16) can be decomposed as

$$\sum_{j: v_j \in U} x_j = \sum_{\ell=1}^q \sum_{j: v_j \in U_\ell} x_j. \tag{B4}$$

Using Eqs. (B3) and (B4) we obtain an equation equivalent to (16):

$$\sum_{\ell=1}^q \sum_{i=1}^m (\alpha_i^\ell - k_i^\ell) \lambda_i = \sum_{\ell=1}^q \sum_{j: v_j \in U_\ell} x_j \tag{B5}$$

We will show that (B5) implies q equations all valid for F_U , and independent of (16). That is, fix some index $\mu \in \llbracket q \rrbracket$, and consider the $s - t$ cut induced by $[\{s\} \cup Z_\mu, \overline{\{s\} \cup Z_\mu}]$. The corresponding inequality is

$$\sum_{\substack{\ell=0 \\ \ell \neq \mu}}^q \sum_{i=1}^m \alpha_i^\ell \lambda_i + \sum_{i=1}^m k_i^\mu \lambda_i \geq \sum_{j: v_j \in V_t \cup U_\mu} x_j. \tag{B6}$$

This inequality is valid for $P_{=}^{\text{emb}}$. Subtracting it from (5) we obtain

$$\sum_{i=1}^m (\alpha_i^\mu - k_i^\mu) \lambda_i \leq \sum_{j: v_j \in U_\mu} x_j. \tag{B7}$$

Since (B5) is valid for F_U , for each $\mu \in \llbracket q \rrbracket$, the corresponding inequality in (B7) is also valid for F_U . Therefore, (B5) decomposes into $q \geq 2$ linearly independent equations, all valid for F_U . Hence, (16) cannot induce a facet of $P_{=}^{\text{emb}}$.

Sufficiency:

Suppose conditions (1) and (2) of this theorem hold., On the one hand, we will show that that for each pair of terminal nodes $v_j, v_k \in \overline{U}$, there exists $(x, \lambda) \in F_U$ such that $(x', \lambda) \in F_U$, where $x'_j = x_j + \varepsilon, x'_k = x_k - \varepsilon$ for $\varepsilon > 0$ sufficiently small, and $x'_{j'} = x_{j'}$ for all $j' \notin \{j, k\}$. Moreover, for each pair of terminal nodes $v_j, v_k \in U$, there exists $(x, \lambda) \in F_U$ such that $(x', \lambda) \in F_U$, where $x'_j = x_j + \varepsilon, x'_k = x_k - \varepsilon$ for $\varepsilon > 0$ sufficiently small, and $x'_{j'} = x_{j'}$ for all $j' \notin \{j, k\}$. From these two claims it follows that the points of F_U cannot satisfy any other equations, but (4), (5), and (14).

Since $\overline{S^+} \setminus \{t\}$ induce a connected graph G^+ , for any pair of $v_j, v_k \in \overline{U}$, there is a non-directed path in $\overline{S^+} \setminus \{t\}$ connecting them. For $\lambda = (1/m)\mathbf{1}$, we define a feasible $s - t$ flow of value $\sum_{i=1}^m \alpha_i / m$ in the network (G, c) with arc capacities c defined as follows: $c_e = \beta_e / m$ for each arc not adjacent to t , and $c_{v_j, t} = +\infty$. Let f be a maximum $s - t$ flow in this network, which is neither 0, nor saturating on any arc in G^+ , and saturates all the cut arcs

of $[S^+, \overline{S^+}]$. Such f exists: for each node of $V(G^+) \setminus V_t$, the incoming flow is distributed among the out-arcs proportionally to their capacities. Then let x_j be the total flow entering $v_j \in V_t$. Then $(x, \lambda) \in F_U$. Since G^+ is connected, we can pick an undirected path π in G^+ between any pair of nodes $v_j, v_k \in \overline{U}$, and perturb the flow and x as follows. We increase the flow f on all forward arcs on the path from v_j to v_k by $\varepsilon > 0$, and decrease the flow on all the backward arcs by the same amount. We choose $\varepsilon > 0$ sufficiently small such that the new flow f' is feasible, i.e.

$$\varepsilon \leq \min \{ \beta_e - f_e, f_e \mid e \in \pi \}. \quad (\text{B8})$$

Then the corresponding vector x' is as required, only differs from x in j and k , and $(x', \lambda) \in F_U$.

The above reasoning can be applied analogously to U and the subgraph spanned by $S^- \setminus \{s\}$. \square

Funding Open access funding provided by HUN-REN Institute for Computer Science and Control. This research has been supported by the TKP2021-NKTA-01 NRDIO grant on "Research on cooperative production and logistics systems to support a competitive and sustainable economy".

Declarations

Conflict of interest There are no Conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Almalkhour, F., Diarrassouba, I., & Hadhbi, Y. (2024). Extended formulation and branch-and-cut-and-price algorithm for the two connected subgraph problem with disjunctive constraints. *Annals of Operations Research*. <https://doi.org/10.1007/s10479-024-06123-0>
- Balas, E. (1979). Disjunctive programming. In P. L. Hammer, E. L. Johnson, & B. H. Korte (Eds.), *Discrete optimization II. Annals of discrete mathematics* (vol. 5, pp. 3–51). North Holland, Amsterdam. [https://doi.org/10.1016/S0167-5060\(08\)70342-X](https://doi.org/10.1016/S0167-5060(08)70342-X)
- Balas, E. (1985). Disjunctive programming and a hierarchy of relaxations for discrete optimization problems. *SIAM Journal on Algebraic Discrete Methods*, 6(3), 466–486. <https://doi.org/10.1137/0606047>
- Balas, E. (1998). Disjunctive programming: Properties of the convex hull of feasible points. *Discrete Applied Mathematics*, 89(1–3), 3–44.
- Balas, E. (2018). *Disjunctive programming*. Springer. <https://doi.org/10.1007/978-3-030-00148-3>
- Edmonds, J., & Giles, R. (1977). A min-max relation for submodular functions on graphs. In P. L. Hammer, E. L. Johnson, B. H. Korte, G. L. Nemhauser (Eds.), *Studies in integer programming. Annals of discrete mathematics*, vol. 1, (pp. 185–204). North Holland, Amsterdam. [https://doi.org/10.1016/S0167-5060\(08\)70734-9](https://doi.org/10.1016/S0167-5060(08)70734-9)
- Huber, B., Rambau, J., & Santos, F. (2000). The Cayley trick, lifting subdivisions and the Bohne–Dress theorem on zonotopal tilings. *Journal of the European Mathematical Society*, 2(2), 179–198. <https://doi.org/10.1007/s100970050003>
- Huchette, J., & Vielma, J. P. (2019). A combinatorial approach for small and strong formulations of disjunctive constraints. *Mathematics of Operations Research*, 44(3), 793–820. <https://doi.org/10.1287/moor.2018.0946>

- Huchette, J., & Vielma, J. P. (2019). A geometric way to build strong mixed-integer programming formulations. *Operations Research Letters*, 47(6), 601–606. <https://doi.org/10.1016/j.orl.2019.10.003>
- Kis, T., & Horváth, M. (2022). Ideal, non-extended formulations for disjunctive constraints admitting a network representation. *Mathematical Programming*, 194(1–2), 831–869. <https://doi.org/10.1007/s10107-021-01652-z>
- Sadykov, R., & Vanderbeck, F. (2013). Bin packing with conflicts: A generic branch-and-price algorithm. *INFORMS Journal on Computing*, 25(2), 244–255.
- Schrijver, A. (2004). *Combinatorial optimization: Polyhedra and efficiency. Algorithms and combinatorics* (Vol. 24). Springer.
- Vielma, J. P. (2018). Embedding formulations and complexity for unions of polyhedra. *Management Science*, 64(10), 4721–4734. <https://doi.org/10.1287/mnsc.2017.2856>
- Vielma, J. P. (2019). Small and strong formulations for unions of convex sets from the Cayley embedding. *Mathematical Programming*, 177(1–2), 21–53. <https://doi.org/10.1007/s10107-018-1258-4>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.