

A Note on the “Smith is Huq” Condition

Nelson Martins-Ferreira · Tim Van der Linden

Received: 10 December 2009 / Accepted: 16 June 2010 / Published online: 7 July 2010
© Springer Science+Business Media B.V. 2010

Abstract We show that two known conditions which arose naturally in commutator theory and in the theory of internal crossed modules coincide: *every star-multiplicative graph is multiplicative* if and only if *every two effective equivalence relations commute as soon as their normalisations do*. This answers a question asked by George Janelidze.

Keywords Commutator · Internal reflexive graph · Star-multiplication · Groupoid · Protomodular category · Semi-abelian category

Mathematics Subject Classifications (2010) 18D35 · 18G50 · 20J15

N. Martins-Ferreira was supported by IPLeiria/ESTG-CDRSP and Fundação para a Ciência e a Tecnologia (grant number SFRH/BPD/4321/2008).

T. Van der Linden was supported by Centro de Matemática da Universidade de Coimbra and by Fundação para a Ciência e a Tecnologia (grant number SFRH/BPD/38797/2007). He wishes to thank the Instituto Politécnico for its kind hospitality during his stay in Leiria.

N. Martins-Ferreira
Departamento de Matemática, Escola Superior de Tecnologia e Gestão,
Instituto Politécnico de Leiria,
Centre for Rapid and Sustainable Product Development,
Leiria, Portugal
e-mail: nelsonmf@estg.ipleiria.pt

T. Van der Linden (✉)
Centro de Matemática da Universidade de Coimbra,
3001-454 Coimbra, Portugal
e-mail: tvdlinde@vub.ac.be

1 Introduction

The purpose of this work is to prove that for a semi-abelian category, the following conditions are equivalent:

- (SM) every star-multiplicative graph is an internal groupoid;
- (SH) two equivalence relations commute if and only if their normalisations commute.

The first condition comes from the study of internal crossed modules. In a semi-abelian category \mathcal{A} , the internal crossed modules introduced by Janelidze [12] form a category which is equivalent to the category of internal groupoids in \mathcal{A} . To define a crossed module of groups, however, less structure is needed: a reflexive graph equipped with a star-multiplication already determines a crossed module. Nevertheless, there exist examples of semi-abelian categories where this is not true. Thus the question arose under which conditions on \mathcal{A} the star-multiplicative graphs in \mathcal{A} are internal groupoids.

The second condition was first considered by Bourn and Gran in [2]. On one hand, there is the commutator of internal (effective) equivalence relations which was introduced by Smith [20] in the context of Mal'tsev varieties and made categorical by Pedicchio [19]. On the other hand, in the article [11], Huq introduced a commutator for normal subobjects in a context which is roughly equivalent to that of semi-abelian categories. This definition was further studied by several authors, see e.g., [2] and [1]. Since, in any semi-abelian category, there is a bijective correspondence between the normal subobjects of an object and the effective equivalence relations on it, it is natural to ask how the two concepts of commutator correspond to each other. The answer is that commuting equivalence relations induce commuting normal subobjects [2, Proposition 3.2], but in general, the concepts are not equivalent—not even in a variety of Ω -groups, as the counterexample of digroups shows [1]. On the other hand, it was shown in [10] that an equivalence relation R on an object A commutes with the largest equivalence relation ∇_A as soon as the normalisation k of R is *Huq-central*, i.e., as soon as k commutes with the normalisation 1_A of ∇_A . In fact, a result obtained by Gran says that any two equivalence relations of which the normalisations commute and are jointly strongly epic, commute; see [9]. Finally, in a category which is, for instance, pointed and strongly protomodular, any two equivalence relations commute if and only if their normalisations commute [2].

We shall prove that (SH) and (SM) are equivalent conditions. We do this in two steps: in the first section we work towards Theorem 2.3 which essentially states that Condition (SH) may be restricted to a special class of effective equivalence relations: those pairs of effective equivalence relations which are the kernel pairs of the domain and codomain morphisms of a reflexive graph. Under this latter condition Mantovani and Metere studied the relation between Peiffer graphs and groupoids [18, Theorem 6.1]. We follow their intuition in Section 3, where we prove that a reflexive graph carries a star-multiplication if and only if it is a Peiffer graph if and only if the kernels of its domain and codomain morphisms commute (Proposition 3.7). This is enough to obtain our main result, Theorem 3.8, which states that (SM) is equivalent to (SH).

2 The Smith is Huq Condition

We show that for a pointed protomodular category, the following two conditions are equivalent:

- (SH) two effective equivalence relations commute as soon as their normalisations do;
- (SH^o) every reflexive graph of which the kernels of the domain and the codomain morphisms commute is a groupoid.

Condition (SH) is the *Smith is Huq* condition in the title of this section; condition (SH^o) is well-known to hold, for instance, in the case of groups: recall the analysis of crossed modules given in the final chapter of Mac Lane’s [16].

2.1 The Context

In this section we shall work in *pointed protomodular* categories. A category is **pointed** when it has a **zero object**, i.e., an initial object that is also terminal. A pointed category is **Bourn protomodular** [5] when it is finitely complete and the **Split Short Five Lemma** holds: given a commutative diagram

$$\begin{array}{ccccc}
 K[f] & \xrightarrow{\text{Ker } f} & A & \begin{array}{c} \xleftarrow{s} \\ \xrightarrow{f} \end{array} & B \\
 k \downarrow & & \downarrow a & & \downarrow b \\
 K[f'] & \xrightarrow{\text{Ker } f'} & A' & \begin{array}{c} \xleftarrow{s'} \\ \xrightarrow{f'} \end{array} & B',
 \end{array}$$

where $bf = f'a$, $s'b = as$, $fs = 1_B$ and $f's' = 1_{B'}$, the morphisms k and b being isomorphisms implies that a is an isomorphism. (Note that s' is equal to asb^{-1} , so we could avoid mentioning this morphism and the conditions on it.)

Lemma 2.1 *Given a commutative diagram*

$$\begin{array}{ccccc}
 K[f] & \xrightarrow{\text{Ker } f} & A & \begin{array}{c} \xleftarrow{s} \\ \xrightarrow{f} \end{array} & B \\
 k \downarrow & & \downarrow a & & \downarrow b \\
 K[f'] & \xrightarrow{\text{Ker } f'} & A' & \xrightarrow{f'} & B',
 \end{array}$$

such that $fs = 1_B$, the morphism k is an isomorphism if and only if the right hand side commutative square $bf = f'a$ is a pullback.

Given a split epimorphism and its kernel as in

$$K \xrightarrow{k} A \begin{array}{c} \xleftarrow{s} \\ \xrightarrow{f} \end{array} B$$

the morphism k and the section s are jointly strongly epic; hence k and s are jointly epic (see [1, Lemma 3.1.22] or [3, Lemma 2.2]). For instance, such are the product inclusions $\langle 1_X, 0 \rangle: X \rightarrow X \times Y$ and $\langle 0, 1_Y \rangle: Y \rightarrow X \times Y$.

2.2 Commuting Normal Monomorphisms

A coterminial pair of morphisms

$$X \xrightarrow{k} A \xleftarrow{l} Y$$

commutes (in the sense of Huq) [2, 11] when there is a (necessarily unique) morphism φ such that the diagram

$$\begin{array}{ccc}
 & X & \\
 \langle 1_X, 0 \rangle \swarrow & & \searrow k \\
 X \times Y & \xrightarrow{\varphi} & A \\
 \langle 0, 1_Y \rangle \swarrow & & \searrow l \\
 & Y &
 \end{array}$$

is commutative.

We shall only consider the case where k and l are normal monomorphisms (i.e., kernels). We are particularly interested in the situation where they are the kernels of the domain and codomain morphisms of a reflexive graph $C = (C_1, C_0, d, c, e)$:

$$C_1 \begin{array}{c} \xrightarrow{d} \\ \xleftarrow{e} \\ \xrightarrow{c} \end{array} C_0, \quad de = ce = 1_{C_0}$$

and $k = \text{Ker } d: X \rightarrow C_1, l = \text{Ker } c: Y \rightarrow C_1$. Using Lemma 2.1 we may show that when the kernels k and l of the morphisms d and c in a reflexive graph $C = (C_1, C_0, d, c, e)$ commute, their domains are isomorphic.

Lemma 2.2 *Let k and l be induced by a reflexive graph C as above. If k and l commute then the following commutative squares are pullbacks.*

$$\begin{array}{ccc}
 X \times Y & \xrightarrow{\pi_X} & X \\
 \varphi \downarrow & & \downarrow h=ck \\
 C_1 & \xrightarrow{c} & C_0
 \end{array}
 \qquad
 \begin{array}{ccc}
 X \times Y & \xrightarrow{\pi_Y} & Y \\
 \varphi \downarrow & & \downarrow dl \\
 C_1 & \xrightarrow{d} & C_0
 \end{array}$$

This makes X and Y isomorphic in a strong sense: there exist morphisms $i: X \rightarrow Y$ and $j: Y \rightarrow X$ such that

$$ji = 1_X, \quad ij = 1_Y, \quad ckj = dl \quad \text{and} \quad ck = dli.$$

Proof The left hand side diagram commutes because $\langle 1_X, 0 \rangle$ and $\langle 0, 1_Y \rangle$ are jointly epimorphic and moreover $c\varphi\langle 1_X, 0 \rangle = ck = ck\pi_X\langle 1_X, 0 \rangle$ and

$$c\varphi\langle 0, 1_Y \rangle = cl = 0 = ck\pi_X\langle 0, 1_Y \rangle.$$

It is a pullback by Lemma 2.1 since the induced morphism between the kernels of π_X and c is 1_Y . Similarly the right hand side square is a pullback.

The morphism $i: X \rightarrow Y$ is obtained through the universal property of the first pullback as follows. The equality $ceck = ck = h1_X$ gives rise to a morphism $\iota: X \rightarrow X \times Y$ such that $\varphi\iota = eck$ and $\pi_X\iota = 1_X$; considering $X \times Y$ as a product now, this ι is a pair $\langle 1_X, i \rangle: X \rightarrow X \times Y$. Clearly,

$$dli = dl\pi_Y\langle 1_X, i \rangle = d\varphi\langle 1_X, i \rangle = deck = ck.$$

Using the second pullback one obtains a morphism $j: Y \rightarrow X$ that satisfies $\varphi\langle j, 1_Y \rangle = edl$, so that $ckj = dl$.

Now we only have to prove that i and j are mutually inverse. This again follows from the universal properties of the pullbacks. Indeed, the morphisms $\langle j, ij \rangle: Y \rightarrow X \times Y$ and $\langle j, 1_Y \rangle: Y \rightarrow X \times Y$ are both universally induced by the equality $cedl = ckj = hj$, hence they are equal. Likewise, $\langle 1_X, i \rangle$ is equal to $\langle ji, i \rangle$ so that $ji = 1_X$. □

This result may be interpreted as follows: the two (*a priori* non-equivalent) ways a reflexive graph can be normalised—mapping C to either $ck: X \rightarrow C_0$ or $dl: Y \rightarrow C_0$ —induce naturally isomorphic functors from the category of reflexive graphs with commuting kernels to the category of objects over C_0 .

One usually views the elements of C_1 as arrows between the elements of C_0 , so that the morphism $\varphi: X \times Y \rightarrow C_1$ is nothing but a partial composition on C_1 which sends a pair of arrows

$$.\overset{\alpha}{\longleftarrow} 0 \overset{\beta}{\longleftarrow} .$$

to its composite $\varphi(\alpha, \beta)$. The central question studied in this paper is under which conditions such a partial composition extends to a composition on the entire graph. To answer it, we shall need the concept of commuting effective equivalence relations and its connection with commuting normal monomorphisms.

2.3 Commuting Effective Equivalence Relations

Consider a pair of equivalence relations (R, S) on a common object A

$$R \begin{array}{c} \xrightarrow{r_0} \\ \xleftarrow{\Delta_R} \\ \xrightarrow{r_1} \end{array} A \begin{array}{c} \xleftarrow{s_1} \\ \xrightarrow{\Delta_S} \\ \xleftarrow{s_0} \end{array} S,$$

and consider the induced pullback of r_1 and s_0 .

$$\begin{array}{ccc}
 R \times_A S & \xrightarrow{\pi_S} & S \\
 \pi_R \downarrow & \lrcorner & \downarrow s_0 \\
 R & \xrightarrow{r_1} & A
 \end{array} \tag{1}$$

The pair (R, S) **commutes (in the sense of Smith)** [2, 19, 20] when there is a (necessarily unique) morphism θ such that the diagram

$$\begin{array}{ccccc}
 & & R & & \\
 & \swarrow & & \searrow & \\
 (1_R, \Delta_S r_1) & & & & r_0 \\
 & & R \times_A S & \xrightarrow{\theta} & A \\
 & \swarrow & & \searrow & \\
 (\Delta_R s_0, 1_S) & & & & s_1 \\
 & & S & &
 \end{array}$$

is commutative.

We shall only consider the case where R and S are effective equivalence relations (i.e., kernel pairs). It is well-known that when for a span

$$\begin{array}{ccc}
 & C_1 & \\
 d \swarrow & & \searrow c \\
 C_0 & & C'_0
 \end{array} \tag{2}$$

the kernel pairs $R[d]$ and $R[c]$ commute, this means that (d, c) carries an internal pregroupoid structure [15]; briefly, any zigzag

$$\cdot \xleftarrow{\alpha} \cdot \xrightarrow{\beta} \cdot \xleftarrow{\gamma} \cdot$$

in C_1 may be composed to a single arrow $\theta(\alpha, \beta, \gamma)$, in such a way that $\theta(\alpha, \beta, \beta) = \alpha$ and $\theta(\beta, \beta, \gamma) = \gamma$. In particular, a reflexive graph $C = (C_1, C_0, d, c, e)$ is an internal groupoid if and only if $R[d]$ and $R[c]$ commute: then $\theta(\alpha, \beta, \gamma) = \alpha \circ \beta^{-1} \circ \gamma$.

It is also well-known that when a pair (R, S) of (effective) equivalence relations commutes, then so do their normalisations

$$X = K[r_0] \xrightarrow{k=r_1 \text{Ker } r_0} A \xleftarrow{l=s_1 \text{Ker } s_0} K[s_0] = Y :$$

see [2, Proposition 3.2]. In particular, for any internal groupoid C the composition on C restricts in such a way that the kernels of its domain and codomain morphisms commute. The converse is not true: in general, it is not possible to extend the partial composition on a reflexive graph which is given by its commuting kernels to a composition on the entire graph which makes it into a groupoid. This is explained by the following result (inspired by Lemma 2.1 in [14]), together with the fact that

a pair of effective equivalence relations of which the normalisations commute need not commute itself [1].

Theorem 2.3 *For a pointed protomodular category, the following conditions are equivalent:*

- (SH) *two effective equivalence relations commute as soon as their normalisations do;*
- (SH') *every reflexive graph with commuting kernels of the domain and the codomain morphisms is a groupoid.*

Proof It is clear that (SH') is just (SH) in the special case where the effective equivalence relations considered are the kernel pairs of the domain and the codomain morphisms of a reflexive graph. This special case implies the general case. Indeed, let $R = R[d]$ and $S = R[c]$ be the effective equivalence relations induced by a span (2) and assume that the normal monomorphisms $k = \text{Ker } d$ and $l = \text{Ker } c$ commute in the sense of Huq. We have to prove that R and S commute in the sense of Smith, i.e., the span (d, c) is a pregroupoid.

If one thinks of the “elements” of the object C_1 as arrows $d(\alpha) \xrightarrow{\alpha} c(\alpha)$ then R and S consist of pairs

$$\cdot \xleftarrow{\beta} \cdot \xrightarrow{\alpha} \cdot \quad \text{and} \quad \cdot \xrightarrow{\gamma} \cdot \xleftarrow{\delta} \cdot,$$

respectively. Forming the pullback (1) of r_1 and s_0 we obtain a reflexive graph

$$R \times_{C_1} S \begin{array}{c} \xrightarrow{\text{dom}=r_0\pi_R} \\ \xleftarrow{(\Delta_R, \Delta_S)} \\ \xrightarrow{\text{cod}=s_1\pi_S} \end{array} C_1. \tag{3}$$

An element of $R \times_{C_1} S$ is a triple

$$\cdot \xleftarrow{\alpha} \cdot \xrightarrow{\beta} \cdot \xleftarrow{\gamma} \cdot$$

considered as an arrow β with domain $\alpha = \text{dom}(\alpha, \beta, \gamma) = r_0\pi_R(\alpha, \beta, \gamma)$ and codomain $\gamma = \text{cod}(\alpha, \beta, \gamma) = s_1\pi_S(\alpha, \beta, \gamma)$. The kernels dom and cod commute because so do k and l : the needed morphism

$$K[\text{dom}] \times K[\text{cod}] \rightarrow R \times_{C_1} S$$

takes a pair

$$\left(\cdot \xleftarrow{0} \cdot \xrightarrow{\beta} \cdot \xleftarrow{\gamma} \cdot, \cdot \xleftarrow{\delta} \cdot \xrightarrow{\epsilon} \cdot \xleftarrow{0} \cdot \right)$$

in the product $K[\text{dom}] \times K[\text{cod}]$ and maps it to the element

$$\cdot \xleftarrow{\delta} \cdot \xrightarrow{\varphi(\beta, \epsilon)} \cdot \xleftarrow{\gamma} \cdot$$

of $R \times_{C_1} S$. The hypothesis that (SH') holds now implies that this reflexive graph is a groupoid. This, in turn, establishes a pregroupoid structure on the span (d, c) : the required morphism $\theta: R \times_{C_1} S \rightarrow C_1$ is determined by

$$\left(\cdot \xleftarrow{\gamma} \cdot \xrightarrow{\theta(\alpha, \beta, \gamma)} \cdot \xleftarrow{\alpha} \cdot \right) = \left(\cdot \xleftarrow{\beta} \cdot \xrightarrow{\alpha} \cdot \xleftarrow{\alpha} \cdot \right) \circ \left(\cdot \xleftarrow{\gamma} \cdot \xrightarrow{\gamma} \cdot \xleftarrow{\beta} \cdot \right)$$

where the composition takes place in the groupoid (3). Indeed, in this groupoid

$$(\beta, \beta, \beta) \circ (\gamma, \gamma, \beta) = (\gamma, \gamma, \beta)$$

so that $\theta(\beta, \beta, \gamma) = \gamma$. Likewise, $\theta(\alpha, \beta, \beta) = \alpha$. □

Condition (SH) is sometimes called the **Smith is Huq** property. It is known to hold in quite diverse situations: in pointed and *strongly protomodular* categories (by [2]; see also [1] and [8]) and in pointed and *action accessible* categories (as explained in [18]; see also [4]). This condition is also weaker than the *reflected admissibility* condition studied in [17].

Remark 2.4 As explained to us by Tomas Everaert, the condition (SH) may be replaced by its non-effective version

(SH'') two equivalence relations commute as soon as their normalisations do,

using the same proof, even when the category is not Barr exact. Then the kernels should be replaced by normal monomorphisms in the sense of Bourn [6].

3 Star-Multiplication

In this section we show that, in a semi-abelian category, three types of (uniquely determined) structure on a reflexive graph $C = (C_1, C_0, d, c, e)$ coincide: a reflexive graph C is *star-multiplicative* if and only if it is *Peiffer* if and only if the kernels of d and c commute (Proposition 3.7). This allows us to prove Theorem 3.8 which states that a semi-abelian category has the *Smith is Huq* property if and only if every star-multiplicative graph is a groupoid.

3.1 The Context

A category is **semi-abelian** [13] when it is pointed, Bourn protomodular and Barr exact with binary coproducts. **Barr exact** means that every internal equivalence relation is **effective** (i.e., it is a kernel pair) and the category is **regular**: finitely complete with pullback-stable regular epimorphisms and coequalisers of effective equivalence relations. A **homological** category is pointed, regular and protomodular [1].

In a homological category regular epimorphisms (coequalisers), strong epimorphisms and normal epimorphisms (cokernels) coincide, and every morphism $f: A \rightarrow B$ may be factored as a regular epimorphism $A \rightarrow I[f]$ followed by a monomorphism $\text{lm } f: I[f] \rightarrow B$. The monomorphism $\text{lm } f$ is the **image** of f . A morphism f is **proper** when it has a normal image, i.e., $\text{lm } f$ is a normal monomorphism.

In a semi-abelian category, the **direct image** $\text{Im}(pm)$ of a normal monomorphism m along a regular epimorphism p is always a normal monomorphism (condition (SA*6) in [13]).

We need the following strengthening of Lemma 2.1; see [1] or [7, Proposition 7].

Lemma 3.1 *In a homological category, given a commutative diagram*

$$\begin{array}{ccccc}
 K[f] & \xrightarrow{\text{Ker } f} & A & \xrightarrow{f} & B \\
 k \downarrow & & \downarrow a & & \downarrow b \\
 K[f'] & \xrightarrow{\text{Ker } f'} & A' & \xrightarrow{f'} & B'
 \end{array}$$

where f is a regular epimorphism, the morphism k is an isomorphism if and only if the right hand side square $b f = f' a$ is a pullback.

3.2 Star-Multiplicative Graphs

A reflexive graph $C = (C_1, C_0, d, c, e)$ is **star-multiplicative** [12] when there is a (necessarily unique) morphism

$$\zeta : C_1 \times_{C_0} X \rightarrow X$$

such that $\zeta \langle k, 0 \rangle = 1_X$ and $\zeta \langle eck, 1_X \rangle = 1_X$. Here the square

$$\begin{array}{ccc}
 C_1 \times_{C_0} X & \xrightarrow{\pi_1} & X \\
 \pi_0 \downarrow & & \downarrow h=ck \\
 C_1 & \xrightarrow{d} & C_0
 \end{array}$$

is a pullback. A star-multiplication takes a composable pair of arrows

$$\cdot \xleftarrow{\alpha} \cdot \xleftarrow{\beta} 0$$

and sends it to their composite $\zeta(\alpha, \beta)$.

3.3 Peiffer Graphs

A reflexive graph $C = (C_1, C_0, d, c, e)$ is **Peiffer** when there is a (necessarily unique) morphism

$$\omega : X \times X \rightarrow C_1$$

such that $\omega\langle 1_X, 0 \rangle = k$ and $\omega\langle 1_X, 1_X \rangle = eck$. (This definition is not the original one given in [18], but it is equivalent to it in the present context; see [18, Theorem 5.3].) The structure ω sends a composable pair of arrows

$$\cdot \xleftarrow{\alpha} 0 \xrightarrow{\beta} \cdot$$

to the composite $\omega(\alpha, \beta)$ —which should be considered as $\alpha \circ \beta^{-1}$.

In [18] these two structures are shown to be equivalent; we recall the argument.

Proposition 3.2 *A reflexive graph $C = (C_1, C_0, d, c, e)$ in a pointed protomodular category is star-multiplicative if and only if it is Peiffer.*

Proof Given $\zeta: C_1 \times_{C_0} X \rightarrow X$ put $\omega = \pi_0\langle \zeta, \pi_1 \rangle^{-1}$; conversely, given $\omega: X \times X \rightarrow C_1$ put $\zeta = \pi_0\langle \omega, \pi_1 \rangle^{-1}$. Notations are as above. The inverse morphisms exist by the Split Short Five Lemma. □

Now we work towards an equivalence with reflexive graphs of which the kernel of the domain morphism commutes with the kernel of the codomain morphism. In Lemma 3.6 we need the surrounding category to be semi-abelian.

Lemma 3.3 ([18], Theorem 5.3) *Any Peiffer graph C induces commutative squares*

$$\begin{array}{ccc} X \times X & \xrightarrow{\pi_1} & X \\ \omega \downarrow & \text{(i)} & \downarrow h=ck \\ C_1 & \xrightarrow{d} & C_0 \end{array} \quad \text{and} \quad \begin{array}{ccc} X \times X & \xrightarrow{\pi_0} & X \\ \omega \downarrow & \text{(ii)} & \downarrow h=ck \\ C_1 & \xrightarrow{c} & C_0. \end{array}$$

Furthermore, the square (i) is a pullback.

Proof The morphisms $\langle 1_X, 0 \rangle$ and $\langle 1_X, 1_X \rangle$ are jointly epic and

$$\begin{aligned} d\omega\langle 1_X, 0 \rangle &= dk = 0 = h\pi_1\langle 1_X, 0 \rangle, \\ d\omega\langle 1_X, 1_X \rangle &= deck = ck = h\pi_1\langle 1_X, 1_X \rangle, \\ c\omega\langle 1_X, 0 \rangle &= ck = ck\pi_0\langle 1_X, 0 \rangle \end{aligned}$$

and

$$c\omega\langle 1_X, 1_X \rangle = ceck = ck = h\pi_0\langle 1_X, 1_X \rangle$$

so that the two squares commute. Taking kernels horizontally in (i) induces the identity morphism 1_X ; hence the square is a pullback by Lemma 2.1. □

Lemma 3.4 *Let $g: X \times X \rightarrow A$ be a morphism with $g\langle 0, 1_X \rangle = 0$ and write $g_0 = g\langle 1_X, 0 \rangle$. Then $g = g_0\pi_0$, so that $g\langle 1_X, 1_X \rangle = g_0$.*

Proof The morphism g is uniquely determined by the equalities $g\langle 0, 1_X \rangle = 0$ and $g\langle 1_X, 0 \rangle = g_0$. Since also $g_0\pi_0\langle 0, 1_X \rangle = 0$ and $g_0\pi_0\langle 1_X, 0 \rangle = g_0$ we have that $g = g_0\pi_0$. \square

Lemma 3.5 *For any Peiffer graph C , the morphism c is the cokernel of the composite $\omega\langle 0, 1_X \rangle: X \rightarrow C_1$.*

Proof First note that $c\omega\langle 0, 1_X \rangle = 0$ by commutativity of the square (ii) in Lemma 3.3. Consider $f: C_1 \rightarrow A$ with $f\omega\langle 0, 1_X \rangle = 0$; we claim that the morphism $fe: C_0 \rightarrow A$ satisfies $fec = f$. Indeed, by Lemma 3.4 the equalities $f\omega\langle 0, 1_X \rangle = 0$ and $f\omega\langle 1_X, 0 \rangle = fk$ imply $f\omega\langle 1_X, 1_X \rangle = fk$, so that $feck = fk$. Since also $fece = fe$ and k and e are jointly epic we may conclude that $fec = f$. \square

Lemma 3.6 *For any Peiffer graph C in a semi-abelian category the induced commutative square (ii) from Lemma 3.3 is a pullback.*

Proof Taking kernels vertically gives rise to the reflexive graph

$$\begin{array}{ccc}
 & \xrightarrow{\pi'_0} & \\
 K[\omega] & \xleftarrow{\Delta} \xrightarrow{\quad} & K[h]; \\
 & \xrightarrow{\pi'_1} &
 \end{array}$$

Since (i) is a pullback, the morphism π'_1 , and hence also π'_0 , is an isomorphism. It follows by Lemma 3.1 that the top square in the vertical regular epi-mono factorisation

$$\begin{array}{ccccc}
 X & \xrightarrow{\langle 0, 1_X \rangle} & X \times X & \xrightarrow{\pi_0} & X \\
 \vdots & & \downarrow & \lrcorner & \downarrow \\
 X & \xrightarrow{\text{Ker } \bar{c}} & I[\omega] & \xrightarrow{\bar{c}} & I[h] \\
 \vdots & & \downarrow & & \downarrow \\
 Y & \xrightarrow{i} & C_1 & \xrightarrow{c} & C_0 \\
 & & \text{Ker } c & &
 \end{array}$$

(iii)

of (ii) is a pullback. Taking kernels to the left induces morphisms as indicated. We have to show that i is an isomorphism.

Being a composite $h = ck$ of a normal monomorphism with a regular epimorphism, the morphism h is proper, i.e., its image $\text{Im } h$ is a normal monomorphism. Since the square (i) is a pullback, ω is also proper, so that $\text{Im } \omega$ is a normal monomorphism. The morphism $\text{Im } h$ being mono implies that the square (iii) is a pullback. Since both $\text{Im } \omega$ and $\text{Ker } c$ are normal monomorphisms, this implies that the diagonal of (iii)—the morphism $\omega\langle 0, 1_X \rangle$ —is also a normal monomorphism. Lemma 3.5 tells us that c is its cokernel, so that $\omega\langle 0, 1_X \rangle$ is the kernel of c . This means that i is an isomorphism, and the square (ii) is a pullback by Lemma 2.1. \square

Proposition 3.7 *For a reflexive graph $C = (C_1, C_0, d, c, e)$ in a semi-abelian category, the following three conditions are equivalent:*

1. C is star-multiplicative;
2. C is Peiffer;
3. $\text{Ker } d$ and $\text{Ker } c$ commute.

Proof Conditions 1. and 2. are equivalent by Proposition 3.2. If C is Peiffer then $\text{Ker } d$ and $\text{Ker } c$ commute. Indeed, by Lemma 3.6 we can put $\varphi = \omega$ since $\omega\langle 0, 1_X \rangle$ is the kernel l of c . Conversely, if Condition 3. holds then by Lemma 2.2 we have

$$\iota = \langle 1_X, i \rangle: X \rightarrow X \times X$$

such that $\varphi\iota = eck$. Now $\omega = \varphi(1 \times i): X \times X \rightarrow C_1$ is a Peiffer structure on C because $\omega\langle 1_X, 0 \rangle = \varphi(1_X \times i)\langle 1_X, 0 \rangle = \varphi\langle 1_X, 0 \rangle = k$ and $\omega\langle 1_X, 1_X \rangle = \varphi(1_X \times i)\langle 1_X, 1_X \rangle = \varphi\iota = eck$. □

Theorem 3.8 *For a semi-abelian category, the following conditions are equivalent:*

- (SM) every star-multiplicative graph is multiplicative;
- (SH) two (effective) equivalence relations commute if and only if their normalisations commute.

Proof We already explained above that one implication of (SH) always holds by [1, Proposition 2.7.7]. Hence by Theorem 2.3 we may replace the second condition with

- (SH') every reflexive graph with commuting kernels of the domain and the codomain morphisms is a groupoid.

The result now follows from Proposition 3.7 and the fact that in a semi-abelian category, multiplicative graphs (i.e., categories) and groupoids coincide. □

Note that Lemma 3.6 is the only place where we use that the underlying category is semi-abelian rather than pointed protomodular. This suggests an extension of the concept of Peiffer graph to pointed protomodular categories, where the pullback property of square (ii) in Lemma 3.3 becomes an axiom. (Or, equivalently, in the homological case, the morphism $\omega\langle 0, 1_X \rangle$ is demanded to be a normal monomorphism.) The concept of star-multiplicative graph allows a similar modification, where now one asks that the morphism of reflexive graphs

$$\begin{array}{ccc}
 C_1 \times_{C_0} X & \xrightarrow{\pi_0} & C_1 \\
 \pi_1 \downarrow \uparrow \downarrow & \zeta & d \downarrow \uparrow \downarrow c \\
 X & \xrightarrow{h} & C_0
 \end{array}$$

is not just a discrete cofibration (i.e., the square $h\pi_1 = d\pi_0$ is a pullback) but also a discrete fibration ($h\zeta = c\pi_0$ is a pullback). These definitions extend Theorem 3.8 to the pointed protomodular context.

Acknowledgements Thanks to Tomas Everaert, Julia Goedecke and the referee for interesting suggestions and for their comments on the text.

References

1. Borceux, F., Bourn, D.: Mal'cev, protomodular, homological and semi-abelian categories. In: *Mathematics and its Applications*, vol. 566. Kluwer Academic Publishers (2004)
2. Bourn, D., Gran, M.: Centrality and normality in protomodular categories. *Theory Appl. Categ.* **9**(8), 151–165 (2002)
3. Bourn, D., Gran, M.: Regular, protomodular, and abelian categories. In: Pedicchio, M.C., Tholen, W. (eds.) *Categorical Foundations: Special Topics in Order, Topology, Algebra and Sheaf Theory Encyclopedia of Mathematics and its Applications*, vol. 97, pp. 165–211. Cambridge University Press (2004)
4. Bourn, D., Janelidze, G.: Centralizers in action accessible categories. Technical report, L.M.P.A. (2007)
5. Bourn, D.: Normalization equivalence, kernel equivalence and affine categories. In: Carboni, A., Pedicchio, M.C., Rosolini, G. (eds.) *Category Theory, Proceedings Como 1990 Lecture Notes in Mathematics*, vol. 1488, pp. 43–62. Springer (1991)
6. Bourn, D.: Normal subobjects and abelian objects in protomodular categories. *J. Algebra* **228**, 143–164 (2000)
7. Bourn, D.: 3×3 Lemma and protomodularity. *J. Algebra* **236**, 778–795 (2001)
8. Bourn, D.: Commutator theory in strongly protomodular categories. *Theory Appl. Categ.* **13**(2), 27–40 (2004)
9. Everaert, T., Van der Linden, T.: Relative commutator theory in semi-abelian categories. (in preparation, 2010)
10. Gran, M., Van der Linden, T.: On the second cohomology group in semi-abelian categories. *J. Pure Appl. Algebra* **212**, 636–651 (2008)
11. Huq, S.A.: Commutator, nilpotency and solvability in categories. *Quart. Math. J. Oxford* **19**(2), 363–389 (1968)
12. Janelidze, G.: Internal crossed modules. *Georgian Math. J.* **10**(1), 99–114 (2003)
13. Janelidze, G., Márki, L., Tholen, W.: Semi-abelian categories. *J. Pure Appl. Algebra* **168**, 367–386 (2002)
14. Johnstone, P.T.: The ‘closed subgroup theorem’ for localic herds and pregroupoids. *J. Pure Appl. Algebra* **70**, 97–106 (1991)
15. Janelidze, G., Pedicchio, M.C.: Pseudogroupoids and commutators. *Theory Appl. Categ.* **8**(15), 408–456 (2001)
16. Mac Lane, S.: *Categories for the working mathematician*, 2nd edn. Graduate texts in Mathematics, vol. 5. Springer (1998)
17. Martins-Ferreira, N.: Star-multiplicative graphs in pointed protomodular categories. *Theory Appl. Categ.* **23**(9), 170–198 (2010)
18. Mantovani, S., Metere, G.: Internal crossed modules and Peiffer condition. *Theory Appl. Categ.* **23**(6), 113–135 (2010)
19. Pedicchio, M.C.: A categorical approach to commutator theory. *J. Algebra* **177**, 647–657 (1995)
20. Smith, J.D.H.: Mal'cev varieties. *Lecture Notes in Mathematics*, vol. 554. Springer (1976)