

Light intensity and wavelength influence development, reproduction and locomotor activity in the predatory flower bug *Orius sauteri* (Poppius) (Hemiptera: Anthocoridae)

S. Wang · X. L. Tan · J. P. Michaud · F. Zhang ·
X. Guo

Received: 9 February 2013 / Accepted: 15 May 2013 / Published online: 22 May 2013
© International Organization for Biological Control (IOBC) 2013

Abstract Light wavelength and intensity are physical factors that can affect arthropod development and reproduction. The present study examined the development, reproduction and locomotor activity of the predatory flower bug, *Orius sauteri* (Poppius) (Hemiptera: Anthocoridae), under five light intensities (1000, 2000, 3000, 4000 and 5000 lux) and five wavelengths [red (678.5 nm), green (620.0 nm), yellow (581.7 nm), blue (478.1 nm) and white (all wavelengths)] at constant temperature (25 °C) and RH (70 %). The duration of nymphal development was extended at lower light intensities, primarily due to effects on the first three instars. Under white, yellow and green light, *O. sauteri*

completed development in 18.0 days, but blue light extended development by 3.2 days and red light extended it by 7.4 days. Although lower light intensities extended the preoviposition period and reduced fecundity, they improved egg fertility. Both red and blue light negatively affected preoviposition period, fecundity and egg fertility. Whereas adult female mean walking speed over a five min period was reduced at lower light intensities, longer wavelengths (yellow and red) increased it, ostensibly reflecting an avoidance response. The respiration quotient of adult *O. sauteri* females was also elevated under red light conditions. These findings are informative for optimizing *O. sauteri* mass-rearing procedures and maximizing its efficacy as a biological control agent in greenhouse cultures.

Handling Editor: Patrick De Clercq.

S. Wang · F. Zhang · X. Guo
Institute of Plant and Environmental Protection,
Beijing Academy of Agriculture and Forestry Sciences,
Beijing 100097, China
e-mail: septempunctata@yahoo.com.cn

F. Zhang
e-mail: zf6131@163.com

X. Guo
e-mail: gxjbaafs@163.com

X. L. Tan
State Key Laboratory of Crop Stress Biology on the Arid
Areas, Key Laboratory of Crop Pest Management on the
Losses Plateau of Ministry of Agriculture, Northwest
Agriculture and Forestry University, Yangling 712100,
Shaanxi, China
e-mail: tanxiaoling2010@163.com

J. P. Michaud (✉)
Department of Entomology, Kansas State University,
Agricultural Research Center-Hays, 1232 240th Ave,
Hays, KS 67601, USA
e-mail: jpmi@ksu.edu

Keywords Fecundity · Fertility · Locomotion · Mating · Oviposition · Respiration quotient

Introduction

Various aspects of the photic environment, including photoperiod, intensity, wavelength and even polarization, may influence the development, metabolism and reproduction of arthropods. In particular, light intensity and wavelength are known to influence biological and physiological processes in insects (Bishop 1969; Manuwoto and Scriber 1985; Sakai et al. 2002). Predators and parasitoids have received substantial attention in this regard because of the need to understand photic influences on biological control outcomes (e.g., Omkar and Singh 2005; Malaquias et al. 2010) and the suitability of lighting regimes for mass rearing natural enemies for augmentative biological control applications (Chambers 1977). For example, the aphid parasitoid *Aphidius ervi* Haliday (Hymenoptera: Aphidinae) orients preferentially to green light as opposed to blue-green or yellow-green, ostensibly because of the association of aphid hosts with green foliage (Goff and Nault 1984). The predatory stonefly *Eusthenia costalis* Banks (Plecoptera: Eustheniidae) alters its foraging behaviour in response to small differences in wavelength between red and infrared light (Turnbull and Barmuta 2006). Adults of *Harmonia axyridis* (Pallas) (Coleoptera: Coccinellidae) prefer to aggregate on white or highly reflective objects during flights to overwintering sites (Wang et al. 2011). Both photoperiod and light spectrum affect development and reproductive capacity in the aphidophagous coccinellid *Coelophora saucia* (Mulsant) (Coleoptera: Coccinellidae) (Omkar and Pathak 2006). Changes in light intensity have been linked to the induction and termination of diapause in various predatory Hemiptera and Coleoptera (Danks 1987). Diminishing light intensity eventually limits the ability of the bee *Megalopta genalis* Meade-Waldo (Hymenoptera: Halictidae) to find the nest entrance and thus determines the duration of crepuscular foraging behavior (Kelber et al. 2006). The whitefly parasitoids *Encarsia formosa* Gahan and *Eretmocerus eremicus* Rose and Zolnerowich (Hymenoptera: Aphelinidae) parasitized approximately twice as many hosts under conditions of high light intensity and long daylength as under low light intensity and short daylength (Zilahi-balogh et al. 2006).

Flower bugs of the genus *Orius* (Wolff) (Hemiptera: Anthocoridae) are efficient predators of small insect pests, especially thrips, in many agroecosystems (Jacobson 1993; Pfannenstiel and Yeargan 1998; Lattin 1999). Previous work suggests that photic conditions affect many aspects of *Orius* spp. biology and behavior, including development, reproduction and orientation (Askari and Stern 1972; Smith and Stadelbacher 1978; Abenes and Khan 1990). For example, short daylength induces reproductive diapause in *Orius strigicollis* (Poppius) (Hemiptera: Anthocoridae), whereas long daylength tends to increase developmental duration, albeit with an interaction with temperature (Musolin et al. 2004). Similar results have been reported for *O. insidiosus* (Say) (Ruberson et al. 1991). Similarly, photoperiod was shown to influence survival, fecundity and longevity in *O. thyestes* Herring (Carvalho et al. 2006) and *O. niger* Wolff (Bahşi and Tunç 2008). Fewer studies have examined the effect of light intensity on the locomotor activity of *Orius* spp. or their distribution in habitats (Shields 1980; Gossner 2009), and studies of spectral effects (color) have not yet been conducted.

The present study examined the effects of different light intensities and wavelengths on development, reproduction and locomotor activity of the predatory flower bug *Orius sauteri* (Poppius) (Hemiptera: Anthocoridae). This beneficial species is mass-reared for augmentative release in vegetable greenhouses in Asia for suppression of arthropod pests such as thrips and aphids (Nakata 1995; Nagai and Yano 2000). Short daylength induces reproductive diapause in *O. sauteri* (Ito and Nakata 1998a, b) and photic conditions are known to influence other aspects of its development and reproduction in mass-reared cultures (Saulich and Musolin 2009). The primary objective of this study was to generate better understanding of photic effects on aspects of *O. sauteri* developmental and reproductive biology and behavior that may affect either the efficacy of mass-rearing procedures or the insect's performance as a biological control agent following release in augmentation programs.

Materials and methods

Insects

A total of 390 *O. sauteri* adults (213 females and 177 males) were collected from alfalfa (*Medicago sativa*)

fields around Wang Jia-Yuan Biodiversity Research Station (40°14' N, 116°13' E) in Changping County, Beijing, China during the season of April to July, 2011. Species identity was confirmed by Dr. Cai Wanzhi, taxonomist of Hemiptera at Chinese Agricultural University, Beijing, China and these bugs were used to establish a colony in the Natural Enemy Research Laboratory at the Beijing Academy of Agriculture and Forestry Sciences, Haidian District, Beijing. Bugs were reared in aluminum frame cages (45.0 × 50.0 × 60.0 cm) covered with fabric net (100 mesh), beginning with 30 bugs per cage, in a climate-controlled growth chamber (MH-351, Sanyo, Japan) set to 25 ± 1 °C, 70 ± 2 % RH, 16:8 L:D photoperiod, 3,000 lux fluorescent light. The cowpea aphid, *Aphis craccivora* Koch (Hemiptera: Aphididae), reared on broad bean, *Vicia faba* L., was provided as food and kidney bean plants, *Phaseolus vulgaris* L., as an oviposition substrate (Murai et al. 2001). The stock colony was managed as a continuous culture. New plants were introduced every four days and heavily damaged plants removed, their stems clipped and left in cages until bugs enclosed.

Luminosity and wavelength treatments

Experiments were conducted in climate-controlled growth chambers under the same physical conditions as above, unless otherwise specified. For the luminosity treatments, white fluorescent lights were set to one of five light intensities (1000, 2000, 3000, 4000 or 5000 lux). For wavelength treatments, colored LED lamps (LENSER-Illumine™, Shenzhen, China) were used in place of the stock fluorescent lamps and luminosity was set at 3,000 lux in all treatments. The following wavelength treatments were established: 678.5 nm (red), 620.0 nm (yellow), 581.7 nm (green), 478.1 nm (blue) and fluorescent white (control). Actual wavelengths were measured to an accuracy of ±10–15 nm using an LED analyzer (Feasa Enterprises limited, Limerick, Ireland). A thermal probe was used to verify that temperatures experienced by the insects in various containers remained at 25 ± 1 °C across all lighting treatments.

Development assay

Bugs for experiments were obtained by placing kidney bean stems bearing freshly laid eggs of *O. sauteri*

(24 ± 2 h old post-moulting) in clean plastic Petri dishes (10.0 cm diameter), three per dish. As nymphs hatched, they were transferred to smaller Petri dishes (4.5 cm diameter, 1 bug per dish) within 1 h of eclosion. Each dish was provisioned with ca. 40 3rd instar aphids (replaced every 24 h without removing containers from growth chambers) and covered with perforated Parafilm®. Nymphs in experiments were examined every 4 h, using an infrared camera to make night-time observations, the developmental status of each was recorded, and the proportion surviving to the adult stage was tallied. A total of 25 replicates were conducted simultaneously at each light intensity and wavelength treatment.

Reproduction assay

Virgin *O. sauteri* adults (24 ± 2 h old) were paired in plastic Petri dishes (10.0 cm diam) with abundant food (>60 aphids, replaced daily) and assigned to one of the five light intensity or wavelength treatments. Pairs were observed continuously from onset of copulation (maintenance of genital contact for ≥5 min) until male and female separated and then every 4 h thereafter. Males were then removed and each female provided five *P. vulgaris* stems as oviposition substrates. The duration of copula and preoviposition period were determined by direct observation. Bean stems were replaced daily and held under the same physical conditions until eclosion of neonate bugs. Fecundity and egg fertility was followed for five days by examining all oviposition punctures and dividing the number of bugs eclosing by the total number of hatched and unhatched eggs. The test was repeated five times with five pairs in each light intensity for a total of 25 replicates per treatment.

Respiration assay

We reasoned that changes in the respiratory quotient (RQ) should be evident if light conditions affect *O. sauteri* metabolic rate. A respirometry system (Sable Systems International, Las Vegas, NV, USA) was used to measure the RQ of individual newly molted adult female bugs (24 ± 2 h old) under the various conditions of luminosity and wavelength described above. The apparatus consisted of a CO₂ infrared analyzer (Testo 535, Testo, Lenzkirch, Germany), an O₂ gas sensor (InPro 6000G, Mettler-Toledo, Greifensee,

Switzerland), an air pump, a flow meter, a CO₂ purification tube, a drying tube, a glass sample chamber (2.0 cm ID × 10.0 cm), a data collector (Vernier LabPro, Beaverton, OR, USA) and a temperature sensor (TMP-112, Texas Instruments, Dallas, TX, USA). The apparatus was assembled within an aluminum box (65.0 × 40.0 × 40.0 cm) with light sources mounted on the inside and temperature was maintained at 25 ± 1 °C. Air flow was adjusted to 0.1 l min⁻¹ in the apparatus and initial O₂ and CO₂ concentrations were recorded. Once all readings were stable, insects (*n* = 10 adult females per treatment) were placed singly in the sample chamber and measurements made on each for a period of 30 min. The RQ was calculated according to manufacturer's specifications as:

$$\text{RQ} = \frac{(\% \text{CO}_2 \text{ expired} - \% \text{CO}_2 \text{ inspired})}{(\% \text{O}_2 \text{ inspired} - \% \text{O}_2 \text{ expired})}$$

Locomotion assay

A tracksphere locomotion compensator with behavior recorder (Syntech LC-100, Hilversum, Netherlands) was used to monitor differences in locomotor activity under various light intensities and wavelengths. After adjustment of focus, the device was covered with tin foil shade to exclude ambient light and four white LED lamps were placed in the shade. Light intensity was regulated by an auto regulator (light intensity measurer, TPM-Z1, Leaderlab[®], Hangzhou, China). Newly emerged *O. sauteri* females were placed singly on top of the monitoring sphere. The sphere adjusted its speed to maintain the insect on the top and the bug's

movement was recorded continuously for 5 min. The behavior recorder sums all movements of the sphere to yield an estimate of linear distance that is then divided by the time of the observation to yield a walking speed. A total of ten females were observed under each light intensity treatment, and 35 under each wavelength treatment.

Statistical analysis

Data from all experiments were analyzed by one-way ANOVA (SPSS 1998) and means separated by LSD ($\alpha = 0.05$).

Results

The total duration of *O. sauteri* development decreased with increasing light intensity, largely due to faster development in the first, second and third nymphal instars, the fourth and fifth instars remaining largely unaffected (Table 1). Nymphal development was fastest under white, green and yellow light, slower under blue light, and very slow under red light, with significant effects of wavelength on the durations of all nymphal instars (Table 2).

There were significant effects of light intensity on all reproductive parameters. Duration of copula and preoviposition period were both decreased by increasing light intensity, but fecundity was maximal at intermediate light intensity and egg fertility was maximal at low intensity (Table 3). White, green and yellow light yielded the longest durations of

Table 1 Mean duration (days ± SE) of various *O. sauteri* nymphal stages (*n* = 5 replicates of five bugs each per treatment) when reared at 25 ± 1 °C under five different intensities of white fluorescent light

Light intensity (lux)	Nymphal instar					Total
	First	Second	Third	Fourth	Fifth	
1,000	4.3 ± 0.2a	3.7 ± 0.2a	3.5 ± 0.1a	4.1 ± 0.3a	4.2 ± 0.3a	19.7 ± 0.5a
2,000	3.9 ± 0.2ab	3.3 ± 0.2ab	3.3 ± 0.2a	3.8 ± 0.2a	4.1 ± 0.2a	18.3 ± 0.4b
3,000	3.8 ± 0.1b	3.2 ± 0.1b	3.0 ± 0.2a	3.8 ± 0.1a	4.0 ± 0.3a	17.8 ± 0.3b
4,000	3.1 ± 0.1c	2.8 ± 0.1c	2.4 ± 0.1b	3.9 ± 0.2a	4.2 ± 0.1a	16.5 ± 0.2c
5,000	2.9 ± 0.1c	2.6 ± 0.2c	2.3 ± 0.1b	4.1 ± 0.1a	4.2 ± 0.1a	16.2 ± 0.3c
<i>F</i>	19.794	15.557	16.875	0.940	0.555	13.446
<i>df</i>	4,120	4,120	4,120	4,120	4,120	4,120
<i>P</i>	<0.001	<0.001	<0.001	0.443	0.696	<0.001

Values bearing the same letter were not significantly different within columns (LSD test, $\alpha = 0.05$)

Table 2 Mean duration (days \pm SE) of various *O. sauteri* nymphal stages ($n = 5$ replicates of five bugs each per treatment) when reared at 25 ± 1 °C under five different wavelengths of light at 3,000 lux

Wavelength (nm)	Nymphal instar					Total
	First	Second	Third	Fourth	Fifth	
Red (678.5)	5.9 \pm 0.1a	5.1 \pm 0.2a	4.7 \pm 0.2a	4.7 \pm 0.1a	5.1 \pm 0.1a	25.4 \pm 0.4a
Yellow (620.0)	4.1 \pm 0.1c	3.1 \pm 0.1c	3.3 \pm 0.1c	3.6 \pm 0.1bc	4.0 \pm 0.1b	18.0 \pm 0.3c
Green (581.7)	3.9 \pm 0.1c	3.2 \pm 0.2c	3.2 \pm 0.3c	3.5 \pm 0.2c	4.2 \pm 0.1b	18.0 \pm 0.3c
Blue (478.1)	5.2 \pm 0.1b	4.1 \pm 0.1b	3.6 \pm 0.1b	3.9 \pm 0.2b	4.5 \pm 0.2b	21.2 \pm 0.5b
White	3.9 \pm 0.1c	3.2 \pm 0.2c	3.3 \pm 0.1c	3.5 \pm 0.1c	4.1 \pm 0.2b	18.0 \pm 0.2c
<i>F</i>	75.242	70.372	37.664	30.130	13.257	223.605
df	4,120	4,120	4,120	4,120	4,120	4,120
<i>P</i>	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001

Values bearing the same letter were not significantly different within columns (LSD test, $\alpha = 0.05$)

Table 3 Mean reproductive parameters (\pm SE) for *O. sauteri* adult pairs ($n = 5$ replicates of five pairs per treatment) when reared at 25 ± 1 °C under five different intensities of white fluorescent light

Light intensity (lux)	Parameter			
	Duration of copula (min)	Preoviposition period (days)	5-day fecundity (no. eggs)	Fertility (% eggs hatching)
1,000	11.6 \pm 0.4a	11.2 \pm 0.4a	30.2 \pm 1.4b	92.7 \pm 2.0a
2,000	10.4 \pm 0.3a	11.1 \pm 0.6ab	34.1 \pm 1.8b	85.3 \pm 3.2b
3,000	7.6 \pm 0.4b	9.8 \pm 0.3b	40.3 \pm 2.1a	73.8 \pm 2.1c
4,000	6.1 \pm 0.4c	8.4 \pm 0.4c	39.1 \pm 1.9a	61.7 \pm 4.0d
5,000	5.9 \pm 0.3c	7.7 \pm 0.8c	39.3 \pm 1.1a	54.2 \pm 2.5e
<i>F</i>	66.848	18.210	13.725	91.977
df	4,120	4,120	4,120	4,120
<i>P</i>	<0.001	<0.001	<0.001	<0.001

Values bearing the same letter were not significantly different within columns (LSD test, $\alpha = 0.05$)

Table 4 Mean reproductive parameters (\pm SE) for *O. sauteri* adult pairs ($n = 5$ replicates of five bugs per treatment) when reared at 25 ± 1 °C under five different wavelengths of light at 3,000 lux

Wavelength (nm)	Variable			
	Duration of copula (min)	Preoviposition period (days)	5-day fecundity (no. eggs)	Fertility (% eggs hatching)
Red (678.5)	6.2 \pm 0.3b	14.9 \pm 1.1a	37.2 \pm 2.3b	59.1 \pm 2.6b
Yellow (620.0)	7.8 \pm 0.4ab	10.6 \pm 0.6bc	42.5 \pm 1.3a	72.7 \pm 2.5a
Green (581.7)	8.2 \pm 0.5a	9.9 \pm 0.5c	43.1 \pm 1.4a	74.1 \pm 2.4a
Blue (478.1)	6.9 \pm 0.2b	12.4 \pm 0.6b	33.3 \pm 1.7b	63.6 \pm 3.6b
White	7.4 \pm 0.2ab	9.5 \pm 0.3c	41.8 \pm 1.8a	74.5 \pm 2.8a
<i>F</i>	12.687	25.893	13.536	14.718
df	4,120	4,120	4,120	4,120
<i>P</i>	<0.001	<0.001	<0.001	<0.001

Values bearing the same letter were not significantly different within columns (LSD test, $\alpha = 0.05$)




Fig. 1 Mean (\pm SE) respiration quotients of adult female *O. sauteri* measured under different wavelengths of light. Data points bearing the same letters were not significantly different (LSD test, $\alpha = 0.05$)

copula, shortest preoviposition periods, and the highest levels of fecundity and egg fertility (Table 4). Light intensity did not significantly affect the respiration quotient ($F_{4,45} = 1.86$, $P = 0.133$) but wavelength did ($F_{4,45} = 2.97$, $P = 0.029$). Bugs under red light exhibited a significantly higher RQ value than did those under other wavelength treatments, none of which differed significantly (Fig. 1).

The walking speed of *O. sauteri* females averaged over five 1-min intervals increased almost two-fold as light intensity was increased from 1,000 to 5,000 lux (Table 5). Vagility was highest under red light conditions, intermediate under yellow, and lowest under white, green and blue (Table 6).

Table 5 Mean walking speed ($\text{mm sec}^{-1} \pm \text{SE}$) of *O. sauteri* female adults ($n = 10$ per treatment) in 1-min intervals under five different intensities of white fluorescent light

Light intensity (lux)	Time interval					Average
	First	Second	Third	Fourth	Fifth	
1,000	8.8 \pm 0.7b	8.6 \pm 0.9b	7.1 \pm 0.9b	5.7 \pm 0.6d	5.4 \pm 1.1d	7.1 \pm 0.4d
2,000	9.0 \pm 1.4b	8.9 \pm 0.8b	7.7 \pm 0.5b	7.4 \pm 0.3c	7.1 \pm 0.2c	8.0 \pm 0.4cd
3,000	9.2 \pm 0.7b	8.9 \pm 0.6b	8.2 \pm 0.5b	8.0 \pm 0.6c	7.8 \pm 0.9c	8.4 \pm 0.3c
4,000	12.6 \pm 1.3a	11.2 \pm 0.9ab	10.5 \pm 0.7ab	10.2 \pm 0.5b	9.9 \pm 0.7b	10.9 \pm 0.4b
5,000	13.1 \pm 0.9a	13.1 \pm 1.0a	12.2 \pm 1.4a	12.3 \pm 0.7a	12.4 \pm 1.2a	12.6 \pm 0.5a
<i>F</i>	4.12	5.10	6.03	20.52	9.71	34.03
df	4,45	4,45	4,45	4,45	4,45	4,45
<i>P</i>	0.006	0.002	0.001	<0.001	<0.001	<0.001

Values bearing the same letter were not significantly different within columns (LSD test, $\alpha = 0.05$)

Table 6 Mean walking speed ($\text{mm sec}^{-1} \pm \text{SE}$) of *O. sauteri* female adults ($n = 35$ per treatment) during 1-min intervals under five different wavelengths of light

Wavelength (nm)	Time interval					Average
	First	Second	Third	Fourth	Fifth	
Red (678.5)	13.2 \pm 0.7a	11.1 \pm 0.5a	10.1 \pm 0.7a	9.4 \pm 0.2a	7.8 \pm 0.3a	10.3 \pm 0.4a
Yellow (620.0)	11.7 \pm 0.7a	9.7 \pm 0.3b	9.2 \pm 0.4a	9.1 \pm 0.8a	7.4 \pm 0.8a	9.4 \pm 0.8b
Green (581.7)	9.6 \pm 0.7b	9.3 \pm 0.5bc	8.2 \pm 0.3b	7.4 \pm 0.5b	6.7 \pm 0.4a	8.2 \pm 0.6c
Blue (478.1)	8.3 \pm 0.7b	8.3 \pm 0.5c	8.2 \pm 0.4b	7.3 \pm 0.7b	6.4 \pm 0.5a	7.7 \pm 0.9c
White	9.2 \pm 0.4b	9.0 \pm 0.6bc	8.1 \pm 0.4b	7.6 \pm 0.2b	7.0 \pm 0.3a	8.2 \pm 0.6c
<i>F</i>	12.69	6.52	7.10	5.75	0.68	15.18
df	4,45	4,45	4,45	4,45	4,45	4,45
<i>P</i>	<0.001	<0.001	<0.001	<0.001	0.608	<0.001

Values bearing the same letter were not significantly different within columns (LSD test, $\alpha = 0.05$)

Discussion

Although *O. sauteri* was able to develop and reproduce successfully under the full range of photic conditions tested, both light intensity and wavelength exerted significant influences on its biology and behavior. Early instar nymphs of *O. sauteri* were particularly sensitive to light intensity and their development was delayed below 4,000 lux, suggesting a diurnal species adapted to development in brightly lit habitats. The delayed development observed under red and blue light is perhaps suggestive of adaptation to development under light in the yellow–green range, consistent with what would be expected in a natural plant canopy. The equivalent suitability of white light to that of yellow and green suggests that it is not so much the perception of wavelengths longer or shorter than yellow–green that delays development, but rather the absence of wavelengths within this optimal range. Nevertheless, previous studies have demonstrated negative effects of short wavelength light on mating behaviour in *Drosophila melanogaster* (Jacobs 1960; Sakai et al. 2002) and on feeding behaviour in both Diptera (Lunau and Wacht 1997), and Lepidoptera (White et al. 1994), whereas red light has been shown to impede development in coccinellids (Omkar and Singh 2005).

Since developmental rate was almost 20 % faster at the highest light intensity compared to the lowest, cultures of *O. sauteri* should employ light intensity higher than 4,000 lux if the objective is to speed the production of large numbers. Likewise, adult locomotor activity increased with light intensity (Table 5), which might translate into a beneficial effect on foraging behaviour under supplementary lighting in a greenhouse environment. White light was generally most suitable for reproduction, although not significantly superior to yellow or green (Table 4). However, bugs reproducing under red light took longer to lay eggs and had reduced fecundity and fertility. Thus, the greater walking speed of adult females under red light (Table 6) likely reflects an avoidance response to this wavelength of light, an inference that would be consistent with the higher observed RQ (Fig. 1). This might result in lower rates of prey consumption and thus account for the negative effect on reproduction. Both red and blue light were shown to cause erratic reproductive activity in two species of Coccinellidae (Omkar and Singh 2005). Notably, Castrejon and

Rojas (2010) found differences between larvae and mated female adults of *Estigmene acrea* (Drury) (Lepidoptera: Arctiidae) in selective orientation to different wavelengths of light, the former preferring green, the color of food plants, and the latter wavelengths in the range of blue to UV, ostensibly useful as cues for orienting to oviposition sites.

Light intensity had inconsistent effects on different parameters of *O. sauteri* reproduction (Table 3). Notably, Nissenan et al. (2008) found that gravid and virgin female carrot psyllids, *Trioza apicalis* Forster (Hemiptera: Triozidae), responded differently to subtle differences in light intensity. Higher luminosities shortened the duration of copula for *O. sauteri*, accelerated the onset of oviposition, and yielded higher fecundities, but significantly reduced fertility, possibly due in part to the effect on mating time. Thus, the high light intensities that favor rapid bug development and reproduction are associated with reductions in egg fertility. One possible application of this finding in commercial production of *O. sauteri* might be the regular harvest of egg-bearing plant parts so that these can be held under lower light conditions until emergence of nymphs. Alternatively, the cage environment could be modified so that oviposition sites are shaded and experience lower light intensity than the rest of the enclosure. Further experiments could test whether females respond to differences in light intensity when selecting oviposition sites.

Acknowledgments This study was supported by the National Key Basic Research and Development Program (973 program) No. 2013CB127605, the Key Topics in Innovation Engineering Projects, Chinese Academy of Sciences (KSCX2-EW-B-02), the National Basic Research Program of China (2012CB017359) and the Special Fund for Agro-scientific Research in the Public Interest (201303108). The authors are grateful to Prof. Chaodong Zhu for comments which improved the manuscript.

References

- Abenes M, Khan Z (1990) Attractiveness of light color to selected predators of rice pests. *Int Rice Res Newsl* 15:24–25
- Askari A, Stern V (1972) Effect of temperature and photoperiod on *Orius tristicolor* feeding on *Tetranychus pacificus*. *J Econ Entomol* 65:132–135
- Bahşi Ş, Tunç İ (2008) Development, survival and reproduction of *Orius niger* (Hemiptera: Anthocoridae) under different photoperiod and temperature regimes. *Biocontrol Sci Technol* 18:767–778

- Bishop J (1969) Light control of aquatic insect activity and drift. *Ecology* 50:371–380
- Carvalho AR, Bueno VHP, Pedroso EC, Kon LI, Diniz AJF, Silva RJ (2006) Influence of photoperiod on *Orius thyestes* Herring (Hemiptera: Anthocoridae) reproduction and longevity. *Neotrop Entomol* 35:489–492
- Castrejón F, Rojas JC (2010) Behavioral responses of larvae and adults of *Estigmene acrea* (Lepidoptera: Arctiidae) to light of different wavelengths. *Fla Entomol* 93:505–509
- Chambers D (1977) Quality control in mass rearing. *Annu Rev Entomol* 22:289–308
- Danks H (1987) Insect dormancy: an ecological perspective. Biological survey of Canada monograph series 1, Ottawa
- Goff A, Nault L (1984) Response of the pea aphid parasite *Aphidius ervi* Haliday (Hymenoptera: Aphidiidae) to transmitted light. *Environ Entomol* 13:595–598
- Gossner M (2009) Light intensity affects spatial distribution of Heteroptera in deciduous forests. *Eur J Entomol* 106: 241–252
- Ito K, Nakata T (1998a) Diapause and survival in winter in two species of predatory bugs, *Orius sauteri* and *O. minutus*. *Entomol Exp Appl* 89:271–276
- Ito K, Nakata T (1998b) Effect of photoperiod on reproductive diapause in the predatory bugs, *Orius sauteri* (Poppus) and *O. minutus* (Linnaeus) (Heteroptera: Anthocoridae). *Appl Entomol Zool* 33:115–120
- Jacobs M (1960) Influence of light on mating of *Drosophila melanogaster*. *Ecology* 41:182–188
- Jacobson R (1993) Control of *Frankliniella occidentalis* with *Orius majusculus*: experiences during the first full season of commercial use in the U.K. *IOBC/WPRS Bull* 16:81–84
- Kelber A, Warrant E, Pfaff M, Wallén R, Theobald J, Wcislo W, Raguso R (2006) Light intensity limits foraging activity in nocturnal and crepuscular bees. *Behav Ecol* 17:63–72
- Lattin J (1999) Bionomics of the Anthocoridae. *Annu Rev Entomol* 44:207–231
- Lunau K, Wacht S (1997) Innate flower recognition in the hoverfly *Eristalis tenax* L. *Mitteil Deutsch Gesellsch Allgem Angew Entomol* 11:481–484
- Malaquias J, Ramalho F, Fernandes F, Nascimento Júnior J, Correia E, Zanuncio J (2010) Effects of photoperiod on reproduction and longevity of *Podisus nigrispinus* (Heteroptera: Pentatomidae). *Ann Entomol Soc Am* 103: 603–610
- Manuwoto S, Scriber J (1985) Neonate larval survival of European corn borers, *Ostrinia nubilalis*, on high and low dimboia genotypes of maize: Effects of light intensity and degree of insect inbreeding. *Agric Ecosyst Environ* 14:221–236
- Murai T, Narai Y, Sugiura N (2001) Utilization of germinated broad bean seeds as an oviposition substrate in mass rearing of the predatory bug, *Orius sauteri* (Poppus) (Heteroptera: Anthocoridae). *Appl Entomol Zool* 36:489–495
- Musolin D, Tsytulina K, Ito K (2004) Photoperiodic and temperature control of reproductive diapause induction in the predatory bug *Orius strigicollis* (Heteroptera: Anthocoridae) and its implications for biological control. *Biol Control* 31:91–98
- Nagai K, Yano E (2000) Predation by *Orius sauteri* (Poppus) (Heteroptera: Anthocoridae) on *Thrips palmi* Karny (Thysanoptera: Thripidae): Functional response and selective predation. *Appl Entomol Zool* 35:565–574
- Nakata T (1995) Population fluctuations of aphids and their natural enemies on potato in Hokkaido, Japan. *Appl Entomol Zool* 30:129–138
- Nissenan A, Kristoffersen L, Anderbrant O (2008) Physiological state of female and light intensity affect the host-plant selection of carrot psyllid, *Trioza apicalis* (Hemiptera: Triozidae). *Eur J Entomol* 105:227–232
- Omkar, Pathak S (2006) Effects of different photoperiods and wavelengths of light on the life-history traits of an aphidophagous ladybird, *Coelophora saucia* (Mulsant). *J Appl Entomol* 130:45–50
- Omkar MG, Singh K (2005) Effects of different wavelengths of light on the life attributes of two aphidophagous ladybirds (Coleoptera: Coccinellidae). *Eur J Entomol* 102:33–37
- Pfannenstiel RS, Yeargan KV (1998) Association of predatory Hemiptera with selected crops. *Environ Entomol* 27: 232–239
- Ruberson J, Bush L, Kring T (1991) Photoperiodic effect on diapause induction and development in the predator *Orius insidiosus* (Heteroptera: Anthocoridae). *Environ Entomol* 20:786–789
- Sakai T, Isono K, Tomaru M, Fukatami A, Oguma Y (2002) Light wavelength dependency of mating activity in the *Drosophila melanogaster* species subgroup. *Genes Genet Syst* 77:187–195
- Saulich AKh, Musolin DL (2009) Seasonal development and ecology of anthocorids (Heteroptera, Anthocoridae). *Entomol Rev* 89:501–528
- Shields EJ (1980) Locomotory activity of *Orius tristicolor* under various intensities of flickering and non-flickering light. *Ann Entomol Soc Am* 73:74–77
- Smith JW, Stadelbacher EA (1978) Predatory arthropods: seasonal rise and decline of populations in cotton fields in the Mississippi delta. *Environ Entomol* 7:367–371
- SPSS (1998) SPSS 8.0 for windows. SPSS Inc. Prentice Hall, New Jersey, USA
- Turnbull DKR, Barmuta LA (2006) Different light conditions influence the activity of *Eusthenia costalis* Banks (Plecoptera: Eustheniidae). *Int J Freshw Entomol* 28:211–217
- Wang S, Michaud JP, Tan XL, Zhang F, Guo XJ (2011) The aggregation behavior of *Harmonia axyridis* in its native range in Northeast China. *BioControl* 56:193–206
- White RH, Stevenson RD, Bennett RR, Culter DE (1994) Wavelength discrimination and the role of ultraviolet vision in the feeding behavior of hawkmoths. *Biotropica* 26:427–435
- Zilahi-balogh GM, Shipp JL, Cloutier C, Brodeur J (2006) Influence of light intensity, photoperiod, and temperature on the efficacy of two aphelinid parasitoids of the greenhouse whitefly. *Environ Entomol* 35:581–589