

Production of an anti-fungal substance for biological control of *Phytophthora capsici* causing phytophthora blight in red-peppers by *Streptomyces halstedii*

Gil-Jae Joo

Institute of Agricultural Science & Technology, Kyungpook National University, Daegu 702-701, Korea (Fax: +82-53-953-6972; E-mail: gjoo@knu.ac.kr)

Received 4 June 2004; Revisions requested 18 June 2004/26 August 2004; Revisions received 16 August 2004/10 December 2004; Accepted 13 December 2004

Key words: biological control, phytophthora blight, red-pepper, *Streptomyces halstedii*

Abstract

The culture broth of *Streptomyces halstedii* AJ-7 suppressed the growth of *Phytophthora capsici*, which causes phytophthora blight in red-peppers, with less than 1% survival of the pathogen after 12 h of treatment. The low molecular fraction (≤ 10 kDa) of the culture broth retained anti-fungal activity against *P. capsici* after being held at 100 °C for 6 h.

Introduction

Red-peppers (*Capsicum annum* L.) are widely cultivated around the world. Phytophthora blight, which is a serious problem in their cultivation, is caused by *Phytophthora capsici* (Akihiri *et al.* 1992). The pathogen is usually controlled with chemicals but their excessive use has led to environmental pollution. Such chemicals can also kill useful soil insects and beneficial microorganisms in the rhizosphere; they may also enter the food chain (Bartlett *et al.* 2002). Moreover, their efficiency is decreased due to the evolution of resistant pathogens (Rosenberger & Meyer 1981). Over the last 25–30 years, alternative control methods, including the use of biological control microorganisms, have been attempted (Mandee & Baker 1991, Larkin *et al.* 1993).

Actinomycetes have been investigated as broad-spectrum biological control agents for fungal plant pathogens (Pedziwilk 1995, El-Tarabily *et al.* 2000). In the present study, the potential of *Streptomyces halstedii* AJ-7 was evaluated to control *Phytophthora capsici* as a soil-borne pathogen of red-peppers.

Materials and methods

Bacterial strain and culture conditions

Streptomyces halstedii AJ-7 was isolated from soil (unpublished data). It was cultured in a modified Bennet medium (10 g glucose, 2 g peptone, 1 g beef extract, 1 g yeast extract, 10 g colloidal chitin, 0.5 g MgSO₄, 0.5 g K₂HPO₄, 1 l water, pH 7.2) at 28 °C for 5 d. The culture was centrifuged (1000 × g at 4 °C for 20 min) and the supernatant was used for anti-fungal activity.

The phytopathogenic fungi, *Colletotrichum gloeosporioides* KCTC6169 and *Fusarium oxysporium* KCTC16341, were obtained from the Korean Collection for Type Cultures, Daejeon, Korea. *Alternaria alternata*, *Botrytis cinerea*, *Phytophthora capsici*, *Pythium ultimum*, *Rhizopus stolonifer*, and *Stemphylium lycopersici* were from the Laboratory of Phytopathology, Kyungpook National University, Daegu, Korea. Fungi were incubated at 28 °C on a potato/dextrose agar (PDA; Difco, USA).

To obtain spores of *Phytophthora capsici*, mycelia were grown on a potato/dextrose (PD) liquid medium at 28 °C for 7 d and the spores

were collected by filtration and centrifugation (1000 × *g* at 4 °C for 10 min). The spore density was then adjusted to 10⁹ spores per ml with sterile distilled water.

Anti-fungal activity test

Anti-fungal activity was estimated by a growth inhibition assay. The fungal spores (10⁶ spores per ml) were spread on a PDA plate and a paper disk containing a cell-free culture broth of *S. halstedii* AJ-7 (40 µl) was then placed in the center of the plate. After 5 d at 28 °C, the growth of the fungus was measured. The degree of inhibition was divided by the range of clear zone. The anti-fungal activity by the culture broth was classified as no inhibition (–; ≤2 mm), weak inhibition (±; 2–3 mm), moderate inhibition (+; 3–10 mm), strong inhibition (++; 10–20 mm), or very strong inhibition (+++; ≥20 mm).

Fractionation of *S. halstedii* AJ-7 culture broth

The cell-free culture broth of *S. halstedii* AJ-7 was fractionated into a high molecular weight component (high molecular fraction; ≥10 kDa) and a low molecular component (low molecular fraction; ≤10 kDa) by ultrafiltration through an YM10 membrane (Ø43 mm; Amicon Co., MA).

Biological control of phytophthora blight on red-pepper plug seedlings

Red-pepper seeds were soaked in the culture broth, the low molecular fraction, and the high molecular fraction of *S. halstedii* AJ-7 for 2 h. The control seeds were treated with distilled water. After being sown, red-pepper plug seedlings were grown in wet potting soil inoculated with spores of *Phytophthora capsici* (10⁶ spores per g soil), which causes phytophthora blight in red-peppers. Red-pepper plug seedlings were grown as reported previously (Joo *et al.* 2004). During 30 d plant culture, the suppression of phytophthora blight by the culture broth of *S. halstedii* AJ-7 was measured in 105 seedlings.

Results and discussion

Anti-fungal activity of *S. halstedii* AJ-7

The low molecular fraction of *S. halstedii* AJ-7 suppressed the growth of various fungal phytopathogens: soil-borne pathogens (*Phytophthora capsici* and *Pythium ultimum*), foliar plant pathogens (*A. alternata*, *B. cinerea*, *C. gloeosporioides* and *Stemphylium lycopersici*), and post-harvest storage pathogens (*F. oxysporium* and *R. stolonifer*) (Table 1).

Table 1. Anti-fungal activity of *Streptomyces halstedii* AJ-7 against red-pepper pathogens.

Red-pepper pathogens (disease)	Degree of inhibition ^a	
	Low molecular fraction ^b	High molecular fraction ^b
<i>Alternaria alternata</i> (black mold)	++	++
<i>Botrytis cinerea</i> (gray mold)	++	–
<i>Colletotrichum gloeosporioides</i> (anthracnose)	++	++
<i>Fusarium oxysporium</i> (fruit rot)	++	+++
<i>Phytophthora capsici</i> (phytophthora blight)	+++	++
<i>Pythium ultimum</i> (damping-off)	+	±
<i>Rhizopus stolonifer</i> (rhizopus fruit rot)	++	–
<i>Stemphylium lycopersici</i> (leaf spot)	++	+

S. halstedii AJ-7 was cultured in a modified Bennet medium at 28 °C for 5 d. A cell-free culture broth was used for anti-fungal activity.

^aThe degree of inhibition was divided by the range of clear zone. Anti-fungal activity by the cell-free culture broth of *S. halstedii* AJ-7 was classified as no inhibition (–; ≤2 mm), weak inhibition (±; 2–3 mm), moderate inhibition (+; 3–10 mm), strong inhibition (++; 10–20 mm), or very strong inhibition (+++; ≥20 mm).

^bThe culture broth was fractionated into a high molecular weight component (≥10 kDa) and a low molecular component (≤10 kDa) by ultrafiltration through a 10 kDa membrane.

Table 2. Anti-fungal effect of *Streptomyces halstedii* AJ-7 culture broth against *Phytophthora capsici*.

Treated with culture broth (h)	Survival rate of <i>Phytophthora capsici</i> (%)	
	Culture broth	Heated culture broth ^a
0	100	100
2	36 ± 2	56 ± 3
4	24 ± 5	43 ± 4
6	16 ± 4	30 ± 1
8	7 ± 1	26 ± 5
10	4 ± 2	18 ± 3
12	1 ± 3	13 ± 4
14	0 ± 1	10 ± 2
24	0 ± 1	0 ± 1

^aA cell-free culture broth was treated at 80 °C for 30 min. The remaining anti-fungal activity was assayed by the germination inhibitory ratio. Spores of *Phytophthora capsici* (10⁶ spores per ml) were inoculated in a potato/dextrose broth and 20% (v/v) of the culture broth was added. The resultant solution was incubated at 28 °C. Samples were taken at various intervals and spread on a potato/dextrose agar plate. All assays were performed in triplicate.

Growth suppression of *Phytophthora capsici* by culture broth of *S. halstedii* AJ-7

The culture broth of *S. halstedii* AJ-7 suppressed the growth of *Phytophthora capsici* with less than 1% survival of the pathogen after 12 h of treatment (Table 2). The culture broth treated at 80 °C for 30 min also retained anti-fungal activity, suggesting that a thermostable anti-fungal material was present in the culture broth.

Table 3. Thermostability of low molecular fraction of *Streptomyces halstedii* AJ-7.

Heat treated time (h)	Relative activity (%)		
	80 °C	90 °C	100 °C
0	100	100	100
2	100	100	96
4	100	99	86
6	100	95	53
8	100	80	30
10	98	60	24
12	95	56	9

A culture broth of *S. halstedii* AJ-7 was fractionated into a low molecular fraction as described in Table 1. The low molecular fraction was treated at 80, 90, and 100 °C for the indicated times. The remaining anti-fungal activity was estimated by measuring the growth inhibition of the fungus. The fungal spores (10⁶ spores per ml) were spread on a potato/dextrose agar plate and a paper disk containing the low molecular fraction (40 µl) was then placed in the center of the plate. After 5 d incubation at 28 °C, the growth of the fungus was measured. All assays were performed in triplicate.

Thermostability of low molecular fraction of *S. halstedii* AJ-7

After heat treatment at 80 °C for 30 min, the high molecular fraction of *S. halstedii* AJ-7 lost anti-fungal activity (data not shown) but the low molecular fraction retained anti-fungal activity against *Phytophthora capsici* (Table 3). The half-life of the low molecular fraction was 6.3 h at 100 °C. From the above results, it was concluded that a thermostable anti-fungal material was present in the low molecular fraction.



Fig. 1. Abnormal hyphal morphology of *Phytophthora capsici* caused by a culture broth of *Streptomyces halstedii* AJ-7. Spores of *P. capsici* (10⁶ spores per ml) were pre-inoculated for 24 h in a potato/dextrose broth. Twenty % (v/v) of low molecular fractionation or high molecular fractionation of *S. halstedii* AJ-7 was then added to the culture. The culture was incubated at 28 °C for 24 h and the hyphal morphology was observed with a light microscope. (a) Normal mycelia of *P. capsici*; (b) abnormal swelling, curling, and branching of *P. capsici* mycelia grown with the high molecular fraction; (c) abnormal mycelia lysis of *P. capsici* mycelia grown with the low molecular fraction. The scale bar indicates 30 µm.

Table 4. Suppression of phytophthora blight by culture broth of *Streptomyces halstedii* AJ-7.

Growth (d)	Disease incidence (%)			
	Culture broth	Low molecular fraction ^a	High molecular fraction ^a	Not treated
3	0	0	2	15
6	0	0	5	32
9	0	0	13	49
12	3	2	16	70
15	4	3	23	99
18	5	4	38	100
21	10	6	46	100
24	18	9	60	100
27	20	12	68	100
30	28	20	72	100

^aA culture broth of *S. halstedii* AJ-7 was fractionated into a low molecular fraction and a high molecular fraction, as described in Table 1. Red-pepper seeds were soaked in the culture broth, the low molecular fraction, and the high molecular fraction for 2 h. After being sown, red-pepper plug seedlings were grown in wet potting soil inoculated with the spore of the pathogen *Phytophthora capsici* (10^6 spores per g soil). During 30 d plant culture, the suppression of phytophthora blight by the culture broth was measured. The data are the mean of the effect on 105 seedlings.

Abnormal hyphal morphology of Phytophthora capsici caused by culture broth of S. halstedii AJ-7

An abnormal hyphal swelling, degradation, and lysis of mycelia were observed when *Phytophthora capsici* were grown with the high or low molecular fraction of *S. halstedii* AJ-7 (Figure 1). It is well known that chitinases lyse hyphae of some plant pathogenic fungi (De Boer *et al.* 1998, Patil *et al.* 2000). *S. halstedii* AJ-7 also produced extracellular chitinase (data not shown), suggesting that the abnormal hyphal morphology resulted from the chitinase activity produced from *S. halstedii* AJ-7.

Biocontrol of phytophthora blight by culture broth of S. halstedii AJ-7

As shown in Table 4, the phytophthora blight was significantly suppressed by the culture broth and the low molecular fraction of *S. halstedii* AJ-7 gave results that compared with those of the control red-peppers. The high molecular fraction did not significantly suppress the disease. These results indicate that *S. halstedii* AJ-7 is a good candidate to control *Phytophthora capsici*, which causes the phytophthora blight, and that the antifungal activity against the pathogen mainly originated from the low molecular fraction of the culture broth.

Acknowledgement

This work was supported by a grant from the Biogreen 21 Program, the Rural Development Administration, Republic of Korea.

References

- Akihiri O, Takashi A, Makoto S (1992) Production of antifungal antibiotic, iturin in a solid fermentation by *Bacillus subtilis* NB22 using wheat bran as a substrate. *Biotechnol. Lett.* **14**: 817–821.
- Bartlett DW, Clough JM, Godwin JR, Hall AA, Hamer M, Parr-Dobrzanski B (2002) The strobilurin fungicides. *Pest Manage. Sci.* **58**: 649–662.
- De Boer W, Gunnewiek PJAK, Lafeber P, Janse JD, Spit BE, Woldendorp JW (1998) Anti-fungal properties of chitinolytic soil bacteria. *Soil Biol. Biochem.* **30**: 193–203.
- El-Tarabily KA, Soliman MH, Nassar AH, Al-Hassani HA, Sivasithamparam K, McKenna F, Hardy GE (2000) Biological control of *Sclerotinia minor* using a chitinolytic bacterium and actinomycetes. *Plant Pathol.* **49**: 573–583.
- Joo GJ, Kim YM, Lee IJ, Song KS, Rhee IK (2004) Growth promotion of red pepper plug seedlings and the production of gibberellins by *Bacillus cereus*, *Bacillus macroides* and *Bacillus pumilus*. *Biotechnol. Lett.* **26**: 487–491.
- Larkin RP, Hopkins DL, Martin FN (1993) Effect of successive plantings on *Fusarium oxysporum* and other microorganisms in soils suppressive and conducive to fusarium wilt of watermelon. *Phytopathology* **83**: 1097–1105.
- Mandeel Q, Baker R (1991) Mechanisms involved in biological control of fusarium wilt of cucumber with strains of nonpathogenic *Fusarium oxysporum*. *Phytopathology* **81**: 1097–1105.
- Patil RS, Gormade V, Deshpande MV (2000) Chitinolytic enzymes: an exploration. *Enzyme Microb. Technol.* **26**: 473–483.

Pedziwik Z (1995) The numbers and the fungistatic activity of actinomycetes in different soils supplemented with pesticides and organic substances. *Pol. J. Soil Sci.* **28**: 45–52.

Rosenberger DA, Meyer FW (1981) Post-harvest fungicides for apples: development of resistance to benomyl vinclozolin and iprodione. *Plant Dis.* **65**: 1010–1013.