

BRIEF COMMUNICATION

## Localization of cadmium in the root cells of *Allium cepa* by energy dispersive X-ray analysis

D. LIU\*, I. KOTTKE\*\* and D. ADAM\*\*\*

*Department of Biology, College of Chemistry and Life Sciences, Tianjin Normal University, Tianjin 300074, P.R. China\**

*Botanisches Institut\*\* and Institut für Angewandte Physik\*\*\*, Eberhard-Karls Universität Tübingen, D-72076 Tübingen, Germany*

### Abstract

*Allium cepa* L. roots were exposed to 0.1 and 1.0 mM Cd for 6, 24 and 48 h and the localization of Cd in the root tissue was investigated. Scanning electron microscopy (SEM) and energy dispersive X-ray analysis (EDXA) were performed on frozen-dried tissues of roots. No Cd was detected in the roots treated with only 0.1 mM Cd, while after exposure to higher Cd concentration (1.0 mM) Cd was observed in cell wall and in cytoplasm in the epidermis, cortex and vascular tissues in the roots.

*Additional key words:* cell wall, cytoplasm, scanning electron microscopy (SEM).

Cadmium is an important environmental pollutant with high toxicity to animals and plants. Cadmium causes a decrease of mitotic index, inhibits cell division and cell proliferation, and has toxic effects on chromosomal morphology including c-mitosis, anaphase bridges, and chromosome stickiness of *Allium cepa* and *A. sativum* (Liu *et al.* 1992, 2003). Cadmium alters the synthesis of RNA and impedes ribonuclease activity in rice (Shah and Dubey 1995). Although the effects of Cd on plants were widely discussed at biochemistry and cell physiology levels (Rauser 1999, Sanità di Toppi and Gabrielli 1999, Cobbett and Goldsbrough 2002, Hall 2002, Kamnev and Lelie 2000), conflicting results were reported for the storage site of Cd in the cells. Ernst (1975), Lignell *et al.* (1982), Khan *et al.* (1984) and Vázquez *et al.* (1992a) indicated that cell walls are the main site of Cd accumulation. The data obtained by electron energy loss spectroscopy (EELS) by Liu and Kottke (2003, 2004) showed that no Cd was detected in the cell walls. Energy dispersive X-ray analysis (EDXA) following freezing of the plant sample allows the precise localization of many elements in the cell walls (Vázquez *et al.* 1992a), and thus

can provide information on mechanisms of metal toxicity and tolerance (Tsezos *et al.* 1997, Barceló and Poschenrieder 1999, White and Gadd 2000, Küpper *et al.* 2001). The objectives of this investigation were to increase our understanding of Cd accumulation sites at the cellular level in the root of *A. cepa*.

Healthy and equal-sized onion (*Allium cepa* L.) cloves were selected and germinated in plastic containers by dipping the base of the bulb and the emerging root tips in Hoagland's nutrient solution (Stephan and Prochazka 1989). The plants were cultivated in greenhouse at 20 °C with 15-h photoperiod at photosynthetic photon flux density of 150  $\mu\text{mol m}^{-2} \text{s}^{-1}$  and relative humidity of 65 %. When the roots reached a length of 1 cm, 6 seedlings per treatment were selected and exposed to different concentrations of Cd in Hoagland's solution for 6, 24, and 48 h. Cadmium was applied as  $\text{CdCl}_2 \cdot 2 \text{H}_2\text{O}$  at concentration 0.1 mM and 1.0 mM. Hoagland's nutrient solution was used for the control. The solutions were continuously aerated with an aquarium air pump and changed every day.

Received 15 August 2005, accepted 27 April 2006.

*Abbreviation:* EDAX - energy dispersive X-ray microanalysis; EDXA - energy dispersive X-ray analysis; EELS - electron energy loss spectroscopy; SEM - scanning electron microscopy; SIMS - secondary-ion mass-spectrometry.

*Acknowledgements:* The work, undertaken in Botanisches Institut, Spezielle Botanik, Eberhard-Karls Universität Tübingen, Germany, was supported by granting of a DAAD/K.C. Wong Fellowship to D.H. Liu, and by the National Natural Science Foundation of China.

\* Corresponding author; fax: (+86) 022 23358489, e-mail address: [donghua@mail.zlnet.com.cn](mailto:donghua@mail.zlnet.com.cn)

For energy dispersive X-ray analyses (EDXA) ten root samples of 1 cm length were cut from the root tips (as young root tissue including the meristem) and basal parts of root segments (as mature tissue) of each treatment and rapidly frozen in liquid nitrogen and lyophilized. Cross-sections of the roots (about 1 mm) were coated by gold using the sputter/coater (*AGAR Scientific Ltd.*, Stansted, Essex, England). The energy dispersive X-ray

microanalytical studies were carried out using scanning electron microscopy (*Philips XL30*, Eindhoven, The Netherlands) provided with *EDAX DX-4 eDXi* System, version 2.11. EDAX spectra were collected between 0 and 20 keV and X-ray detector equipped with a super ultra thin window. The collection time of spectra was 120 s.

The results indicate that there was a direct correlation in the amounts of Cd in root cells of different regions with




Fig. 1. EDXA spectrum taken from the site of the SEM micrographs, showing Cd localization in root cells of *A. cepa*. *A* - Cd on the surface of epidermis of basal parts of roots (1 mM Cd; 48 h). *B* - Cd in cytoplasm of the epidermis of the basal parts of roots (1 mM Cd; 48 h). *C* - Cd in the third cortical cells (0.1 mM Cd; 48 h). *D* - Cd in cytoplasm of the cortical cells (1 mM Cd; 48 h). *E* - Cd on cell walls of the cortical cells (1 mM Cd; 24 h). *F* - Cd in secondary walls of vessel (1 mM Cd; 24 h). *C* - cortical cell. *E* - epidermis. “+” site of the analysis; x-axis - energy [keV].

increasing Cd concentration in the solution and treatment time.

No Cd could be observed in roots treated with 0.1 mM Cd for 6 and 24 h. Only trace amounts of Cd were detected after exposure of 48 h. In roots treated with 1 mM Cd for 6, 24 and 48 h, Cd was found in the epidermal cell wall of root tips and basal parts of roots. Cadmium ions were localized on the surface of epidermis (Fig. 1A) and in cytoplasm in the epidermis of the basal parts of root (Fig. 1B) after 48 h treatment.

Cd was found in all cortical cells exposed to 1 mM Cd for 6, 24 and 48 h and to 0.1 mM Cd for 48 h, while no Cd was found at 0.1 mM Cd for 6 and 24 h. EDXA spectrum revealed that Cd was accumulated in several layers of cortical cells adjacent to the epidermis after treatment with 0.1 mM Cd for 48 h (Fig. 1C). At 1 mM Cd, Cd ions were present in cytoplasm (Fig. 1D) and in cell walls (Fig. 1E).

The vascular tissues were also storage sites of Cd. Cd was found in the parenchyma, and vessels. Cd appeared in the secondary walls of the vessel (Fig. 1F).

EDXA as an analytical technique is very useful for localization of elements in biological specimens at subcellular level. The present studies clearly indicate that Cd is accumulated in the tissues and the cell walls. The results of this investigation are in line with the findings of other workers that cell walls are the main site of Cd accumulation (Ernst 1975, Lignell *et al.* 1982, Khan *et al.* 1984, Vázquez *et al.* 1992a). However, conflicting results on the subcellular compartmentation of Cd have been reported. The results of our previous electron energy loss spectroscopy (EELS) studies using chemical fixation showed that Cd was localised mainly in the vacuoles and nucleoli of cortical cells of differentiating and mature root tissues in *Allium sativum* treated with 10 mM Cd for 9 d (Liu and Kottke 2003) and in *Allium cepa* exposed to 1 mM and 10 mM Cd for 2 and 3 d (Liu and Kottke 2004), while no Cd was detected in cell walls. Generally speaking, EELS is a good method and can present a better energetic

resolution and a higher sensitivity than X-ray microanalysis (Castaing 1987). EELS is more sensitive than both energy dispersive X-ray microanalysis (EDAX) and secondary-ion mass-spectrometry (SIMS) (Huxham *et al.* 1999). Why was no Cd found on cell walls using EELS in our previous investigations and was Cd detected in cell walls using EDXA in the present study? According to the results of previous studies by other research workers and us, the method of sample preparation can be involved and thought to be very important. In our previous studies, chemical fixed material was used. Due to an increasing permeability of membranes during the chemical procedure, losses and dislocations of mobile electrolytes cannot be excluded (Hall and Gupta 1984). Rauser and Ackerley (1987), using EDAX in conventionally fixed material, could not find Cd on cell walls of maize and *Agrostis* roots. Former works on samples not prepared by freeze-substitution revealed the absence of Cd in root cell walls (Vázquez *et al.* 1991, 1992b). Rauser and Ackerley (1987) indicated that one reason for lack of Cd signals from regions other than the electron-dense granules could be due to loss of Cd ions during the preparation of the tissue for electron microscopy. Cryofixation of the material can overcome some of the problems arising from chemical fixation (Bücking *et al.* 1998). According to Vázquez *et al.* (1992a) X-ray microanalysis on freeze-substituted samples of the Cd hyperaccumulator plant *Thlaspi caerulescens* revealed that high amounts of Cd, Ca and Fe were found in cell walls, whilst in samples fixed with glutaraldehyde in presence of sodium sulphide (Danscher 1981) no significant amounts of Cd were generally detected in these cell walls. Therefore, method of sample preparation is a very important link. Both sensitive and accurate analytical means and perfect sample preparation method are needed in identification of sites of localization of heavy metals at the subcellular level in cell organelles, cytoplasm or cell walls.

## Reference

- Barceló, J., Poschenrieder, C. Structural and ultrastructural changes in heavy metal exposed plants. - In: Prasad, M.N.V., Hagemeyer, J. (ed.): Heavy Metal Stress in Plants. Pp. 183-205. Springer-Verlag, Berlin - Heidelberg 1999.
- Bücking, H., Beckmann, S., Heyser, W., Kottke, I.: Elemental contents in vacuolar granules of ectomycorrhizal fungi measured by EELS and EDXS. A comparison of different methods and preparation techniques. - *Micron* **29**: 53-61, 1998.
- Castaing, R.: Progrès récents en microanalyse des solides. - *Analisis* **15**: 517-527, 1987.
- Cobbett, D., Goldsbrough, P.: Phytochelatins and metallothioneins: roles in heavy metal detoxification and homeostasis. - *Annu. Rev. Plant Biol.* **53**: 159-182, 2002.
- Danscher, G.: Histochemical demonstration of heavy metals. A revised version of the sulphide silver method suitable for both light and electron microscopy. - *Histochemistry* **71**: 1-16, 1981.
- Ernst, W.H.O.: Physiology of heavy metal resistance in plants. - In: Hutchinson, T.C., Epstein, S., Page, A.L., Van Loon, J., Davey, T. (ed.): Proceedings of an International Conference on Heavy Metals in the Environment. Pp. 121-136. CEP Consultants, Edinburgh 1975.
- Hall, J.L.: Cellular mechanisms for heavy metal detoxification and tolerance. - *J. exp. Bot.* **53**: 1-11, 2002.
- Hall, T.A., Gupta, B.L.: The application of EDXS to the biological sciences. - *J. Microscopy* **136**: 193-208, 1984.
- Huxham, I.M., Jarvis, M.C., Shakespeare, L., Dover, C.J., Knox, J.P., Seymour, G.B.: Electron energy loss spectroscopic imaging of calcium and nitrogen in the cell walls of apple fruits. - *Planta* **208**: 438-443, 1999.
- Kamnev, A.A., Van der Lelie, D.: Chemical and biological parameters as tools to evaluate and improve heavy metal phytoremediation. - *Biosci. Rep.* **20**: 239-258, 2000.
- Khan, D.H., Duckett, J.G., Frankland, B., Kirkham, B.: An X-ray microanalytical study of the distribution of cadmium in roots

- of *Zea mays* L. - J. Plant Physiol. **115**: 19-28, 1984.
- Küpper, H., Lombi, E., Zhao, F.J., Wieshammer, G., McGrath, S.P.: Cellular compartmentation of nickel in the hyperaccumulators *Alyssum lesbiacum*, *Alyssum bertolonii* and *Thlaspi goesingense*. - J. exp. Bot. **52**: 2291-2300, 2001.
- Lignell, Å., Roomans, G.M., Pedersén, M.: Localization of absorbed cadmium in *Fucus vesiculosus* L. by X-ray microanalysis. - Z. Pflanzenphysiol. **105**: 103-109, 1982.
- Liu, D.H., Jiang, W.S., Gao, X.Z.: Effects of cadmium on root growth, cell division and nucleoli in root tip cells of garlic. - Biol. Plant. **47**: 79-83, 2003.
- Liu, D.H., Jiang, W.S., Li, M.X.: Effects of Cd<sup>2+</sup> on root growth and cell division of *Allium cepa*. - Acta Sci. Circumstantiae **12**: 439-446, 1992.
- Liu, D.H., Kottke, I.: Subcellular localization of Cd in the root cells of *Allium sativum* by electron energy loss spectroscopy. - J. Biosci. **28**: 471-478, 2003.
- Liu, D. H., Kottke, I.: Subcellular localization of Cd in the root cells of *Allium cepa* by electron energy loss spectroscopy and cytochemistry. - J. Biosci. **29**: 329-335, 2004.
- Rausser, W.E.: Structure and function of metal chelators produced by plants. - Cell Biochem. Biophys. **31**: 19-48, 1999.
- Rausser, W.E., Ackerley, C.A.: Localization of cadmium in granules within differentiating and mature root cells. - Can. J. Bot. **65**: 643-646, 1987.
- Sanità di Toppi, L., Gabbriellini, R.: Response to cadmium in higher plants. - Environ. exp. Bot. **41**: 105-130, 1999.
- Shah, K., Dubey, R.S.: Effect of cadmium on RNA level as well as activity and molecular forms of ribonuclease in growing rice seedlings. - Plant Physiol. Biochem. **33**: 577-584, 1995.
- Stephan, U.W., Prochazka, Z.: Physiological disorders of the nicotianamine-auxotroph tomato mutant *chloronerva* at different levels of iron nutrition. I. Growth characteristics and physiological abnormalities as related to iron and nicotianamine supply. - Acta bot. neerl. **38**: 147-153, 1989.
- Tsezos, M., Remoudaki, E., Angelatou, V.: Biosorption sites of selected metals using electron microscopy. - Biochem. Physiol. **118A**: 481-487, 1997.
- Vázquez, M.D., Barceló, J., Poschenrieder, C.: Location of cadmium in young bush bean roots. - In: Vogeli-Lange, R., Wagner, G.J. (ed.): International Conference on Heavy Metals in the Environment. Pp. 115-118. CEP Consultants, Edinburgh 1991.
- Vázquez, M.D., Barceló, J., Poschenrieder, C., Mácico, J., Hatton, P., Baker, A.J.M., Cope, G.H.: Localization of zinc and cadmium in *Thlaspi caerulescens* (Brassicaceae), a metallophyte that can hyperaccumulate both metals. - J. Plant Physiol. **140**: 350-355, 1992a.
- Vázquez, M.D., Poschenrieder, C., Barceló, J.: Ultrastructural effects and localization of low cadmium concentrations in bean roots. - New Phytol. **120**: 215-226, 1992b.
- White, C., Gadd, G.M.: Copper accumulation by sulfate-reducing bacterial biofilms. - FEMS Microbiol. Lett. **183**: 313-318, 2000.

Schekman, R., Goldstein, L., Rossant, J. (ed.): **Annual Review of Cell and Developmental Biology. Volume 22.** - Annual Reviews Inc., Palo Alto 2006. 795 pp. ISBN 0-8243-3122-2.

The introductory chapter "From nuclear transfer to nuclear reprogramming: the reversal of cell differentiation" is written by John B. Gordon, one of the great developmental biologists of our time. Readers of *Biologia Plantarum* will find especially interesting the reviews: "Cellulose synthesis in higher plants" (C. Somerville) and "*Agrobacterium tumefaciens* and plant cell interactions and activities for interkingdom macromolecular transfer" (C.A. McCullen and A.N. Binns). Other topics covered include: "Voltage induced channel activation", "Mitochondrial dynamics", "Sterol transport and esterification", "Protein modification by UbIs", "Endocytosis and *Drosophila* development", "Tight junctions and cell polarity", "Germ cell migration", "Neural crest stem and progenitor cells", "Tissue architecture regulated development, homeostasis and cancer", "Pancreatic  $\beta$ -cell growth proliferation", "Epidermal stem cell of the skin", "GAG modifications shape of animal development", "TLR recognition and

signaling", "TGN-to-plasma-membrane transport carriers", "Iron-sulfur protein biogenesis", "Unfolded protein response", "Cellular basis of kidney development", "Telomeres: cancer to human aging", "Interferon-inducible GTPase", "Extracellular matrix function", "Cell death in *Drosophila*", "Commissural axon pathfinding", "Evaluation of blood cell development", and "Axonal wiring in the mouse olfactory system".

All reviews, written by the foremost experts in the field, include extensive references to the literature and are written in such a way, as to be valuable both to advanced students and scientist. Annual Review of Cell and Developmental Biology as well as other Annual Reviews are available online ([www.annualreviews.org](http://www.annualreviews.org)). Current individual subscriptions include online access to full-text articles, PDFs, reviews in advance, bibliographies, and other supplementary material in the current volume and the prior 4 years' volumes.

T. GICHNER (*Praha*)