

## Plasma enterolactone and breast cancer risk in the Nurses' Health Study II

Jing Xie · Shelley S. Tworoger · Adrian A. Franke · Kathryn L. Terry ·  
Megan S. Rice · Bernard A. Rosner · Walter C. Willett · Susan E. Hankinson ·  
A. Heather Eliassen

Received: 24 April 2013 / Accepted: 29 May 2013 / Published online: 13 June 2013  
© Springer Science+Business Media New York 2013

**Abstract** Lignans are plant-based phytoestrogens with both estrogenic and anti-estrogenic properties that may be important for breast carcinogenesis. Retrospective studies have observed decreased breast cancer risk associated with high circulating enterolactone concentrations, a biomarker of lignan intake, but results from prospective studies are conflicting. To prospectively examine this association, we measured plasma enterolactone levels in 802 breast cancer cases and 802 matched controls nested among predominantly premenopausal women in the Nurses' Health Study II cohort. We used conditional logistic regression and polytomous logistic regression models, adjusting for known breast cancer risk factors, to calculate relative risks

(RR) and 95 % confidence intervals (CI). Compared to women with enterolactone concentrations  $\leq 4$  nmol/L, the multivariate-adjusted RRs for breast cancer were 1.18 (95 % CI 0.86–1.62), 0.91 (95 % CI 0.66–1.25), and 0.96 (95 % CI 0.70–1.33) for women with enterolactone levels in the second to the fourth quartiles, respectively;  $P_{\text{trend}} = 0.60$ . Results were similar across tumors defined by estrogen and progesterone receptor status. Among premenopausal women with follicular estradiol levels below the median ( $<47$  pg/mL), women in the highest category of enterolactone levels had a 51 % lower breast cancer risk compared to those in the lowest category (95 % CI 0.27–0.91);  $P_{\text{trend}} = 0.02$ . No association was observed among women with high-follicular estradiol levels ( $\geq 47$  pg/mL), (comparable RR = 1.39, 95 % CI 0.73–2.65;  $P_{\text{interaction}} = 0.02$ ). We did not observe an overall association between plasma enterolactone and

**Electronic supplementary material** The online version of this article (doi:10.1007/s10549-013-2586-y) contains supplementary material, which is available to authorized users.

J. Xie · S. S. Tworoger · M. S. Rice · B. A. Rosner ·  
W. C. Willett · S. E. Hankinson · A. H. Eliassen (✉)  
Channing Division of Network Medicine,  
Department of Medicine, Brigham and Women's Hospital and  
Harvard Medical School, 181 Longwood Avenue, Boston,  
MA 02115, USA  
e-mail: nhahe@channing.harvard.edu

J. Xie · S. S. Tworoger · K. L. Terry · M. S. Rice ·  
W. C. Willett · S. E. Hankinson · A. H. Eliassen  
Department of Epidemiology, Harvard School of Public Health,  
Boston, MA, USA

A. A. Franke  
University of Hawaii Cancer Center, Honolulu, HI, USA

K. L. Terry  
Obstetrics and Gynecology Epidemiology Center,  
Department of Obstetrics and Gynecology, Brigham and  
Women's Hospital and Harvard Medical School, Boston,  
MA, USA

B. A. Rosner  
Department of Biostatistics, Harvard School of Public Health,  
Boston, MA, USA

W. C. Willett  
Department of Nutrition, Harvard School of Public Health,  
Boston, MA, USA

S. E. Hankinson  
Division of Biostatistics and Epidemiology,  
School of Public Health and Health Sciences,  
University of Massachusetts, Amherst, MA, USA

breast cancer risk in a large nested case–control study of US women. However, a significant inverse association was observed among premenopausal women with low-follicular estradiol levels, suggesting that enterolactone may be important in a low-estrogen environment. This should be confirmed in future studies.

**Keywords** Enterolactone · Lignan · Breast cancer · Biomarker · Prospective study · Premenopausal

## Introduction

Higher levels of both endogenous estrogens (e.g., circulating estradiol) and exogenous estrogens (e.g., postmenopausal hormone use) have been associated with increased risk of breast cancer among postmenopausal women [1, 2]. Among premenopausal women, a positive association of endogenous estrogen with breast cancer risk is suggestive but not conclusive [3–6]. Enterolactone is a phytoestrogen that may be important in breast carcinogenesis.

Enterolactone is an enterolignan formed by intestinal microbiota from lignan precursors which are contained mainly in vegetables, whole grain products, berries, and flaxseeds [7]. Other factors influencing the enterolactone concentrations in plasma include smoking; obesity; constipation; antibiotics use; intake of fiber, fruits, coffee, or tea; and postmenopausal hormone use [8–11]. Since enterolactone is the major lignan metabolite with the highest concentrations in plasma [12], it is most commonly examined in epidemiologic studies as a biomarker for the dietary lignan exposures.

Enterolactone has an aromatic structure similar to estradiol and is weakly estrogenic, such that it can stimulate the proliferation of breast cancer cells [13, 14]. Conversely, anti-estrogenic properties also have been observed; cell-based studies suggest that enterolactone competitively binds the estrogen receptor (ER) and inhibits estradiol-induced breast cancer cell growth [15–17]. Several mechanisms for enterolactone action in breast cancer development have been proposed. First, it may stimulate the production of sex hormone-binding globulin (SHBG), which binds to estradiol and lowers the estradiol bioactivity [13]. Second, high concentrations of enterolactone may inhibit aromatase, an enzyme that converts androgens to estrogens in fat, muscle, and other tissue, and a primary source of estrogens in postmenopausal women [18–20]. Third, it may act as a strong antioxidant [21]. Finally, it may have preferential activation of ERs [17, 22–24].

Epidemiologic studies that examined enterolactone and breast cancer risk have yielded inconsistent results, as discussed in a recent meta-analysis [25]. Retrospective studies have reported a decreased breast cancer risk with higher

circulating enterolactone concentrations [26–30], but results from prospective studies have been conflicting [31–41]. However, previous studies suggested an inverse association between high enterolactone levels and postmenopausal breast cancer risk [29, 37, 38]. Thus, we prospectively examined the association of plasma enterolactone concentrations and breast cancer risk in a large study (802 cases and 802 matched controls) of predominantly premenopausal women nested within the Nurses' Health Study II (NHSII) cohort for whom we have detailed information on case characteristics [e.g., ER and progesterone receptor (PR) status] as well as dietary intake and circulating estradiol concentrations in the follicular and luteal phases.

## Methods

### Study population

The NHSII is a prospective cohort established in 1989, when 116,430 US female nurses, 25–42 years of age, completed an initial questionnaire. Participants have been followed by biennially mailed questionnaires to update exposure information, lifestyle factors, and ascertain non-fatal incident diseases. Women completed a semi-quantitative food frequency questionnaire (FFQ) every 2–4 years since 1991. Deaths were reported by family or postal authorities. We also searched for names of nonresponders in the National Death Index [42, 43].

Between 1996 and 1999, 29,611 NHS II participants (aged 32–54) provided blood samples and completed a short questionnaire recording their weight, menstrual cycle length, first day of the menstrual cycle, and the time, date, and number of hours since the last meal for each blood sample (details in [44, 45]). Among these, 18,521 premenopausal women, who had not used oral contraceptives, been pregnant, or breastfed within 6 months, provided timed blood samples within the menstrual cycle. Blood samples were drawn on the 3rd–5th day of their menstrual cycle (follicular) and 7–9 days before the anticipated start of their next cycle (luteal). The remaining 11,090 women (e.g., postmenopausal, using hormonal contraception, or unwilling to give timed samples) provided an untimed sample. Women had their blood drawn and shipped overnight on ice to our laboratory, where the blood was processed and separated into plasma, red blood cells, and white blood cells. Samples were stored in liquid nitrogen freezers ( $\leq -130$  °C) after collection.

### Measurement of exposure and laboratory assays

To measure the enterolactone levels, frozen plasma was sent to the Analytical Biochemistry Shared Resource of the

University of Hawaii Cancer Center, Honolulu, HI in 2011 and 2012. Extraction and analysis of enterolactone was performed from freshly thawed samples by liquid–liquid extraction after enzymatic hydrolysis followed by isotope dilution liquid chromatography electrospray ionization tandem mass spectrometry as described in detail previously [46] except for the replacement of diethyl ether by methyl tertiary butyl ether during extraction and the replacement of arylsulfatase (which was discontinued by Boehringer Mannheim) by using 30  $\mu$ L arylsulfatase isolated from *Helix pomatia* at 2,000 U/mL in aqueous solution (Sigma, St. Louis, MO, # S9751).

Laboratory personnel were blinded to case–control and quality control status and matched cases and controls (ordered randomly within each set) were assayed in the same laboratory batch. Samples were sent in two batches and we assayed the luteal sample for women with timed samples. The detection limit of enterolactone was 1 nmol/L. The inter-assay coefficient of variation (CV) from quality control samples was 14.1 % and the intra-assay CV was 5.4 %.

#### Assessment of covariates

Information on covariates was collected from the biennial questionnaires and the questionnaire completed at blood collection. Participants provided information on height, weight at age 18, and age at menarche at baseline. Dietary information was taken from the 1999 FFQ, since it was the questionnaire closest to the time of blood collection. Smoking, current weight, antibiotics use, menopausal status, and postmenopausal hormone use (among postmenopausal women) were ascertained at the time of blood collection. Physical activity, history of benign breast disease, parity, age at first birth, oral contraceptive use, and family history of breast cancer were assessed on the 1997 questionnaire. We measured both follicular and luteal estradiol levels among women providing timed samples previously and details can be found elsewhere [3].

#### Identification of breast cancer cases

We identified incident breast cancer cases by self-report on biennial questionnaires, and from reports of death from family members, the National Death Index, and the US Postal Service. For reported breast cancer cases, we obtained medical records or records from cancer registries for confirmation and to obtain information on invasiveness and ER and PR status. We identified 804 breast cancer cases after blood collection, but before June 2009. Each case was matched to one control on race/ethnicity, age at blood draw, time of day of blood draw and fasting status for follicular and luteal samples (or the one untimed

sample), month/year of blood draw, menopausal status at blood draw and diagnosis, and postmenopausal hormone use at blood draw. For women with timed samples, we additionally matched on the luteal day of the blood draw (date of next period minus date of blood draw). Two cases were excluded because the plasma samples of the matched controls were not available for assay. Therefore, 802 cases of breast cancer and 802 matched controls were included in the analyses. The Institutional Review Board of the Brigham and Women's Hospital approved this analysis. Informed consent was implied by receipt of completed questionnaires and blood samples.

#### Statistical analyses

We did not identify any statistical outliers for enterolactone levels using the generalized extreme Studentized deviate many-outlier detection approach [47]. Nine samples had values below the limit of detection; these were set to be half the limit of detection or 0.5 nmol/L. We calibrated the enterolactone levels in one laboratory batch to the other one,  $R^2 = 0.98$ , and categorized enterolactone levels into quartiles based on the distribution in controls. We used conditional logistic regression models to estimate the relative risks (RR) and 95 % confidence intervals (CI) of breast cancer. The models were conditioned on matching factors and adjusted for covariates including family history of breast cancer (yes, no), history of benign breast disease (yes, no), oral contraceptive use (never, >0–1, >1–5, and >5 years), BMI at age 18 (continuous), weight change since age 18 (continuous), parity and age at first birth (nulliparous, 1–2 children and <25 years, 1–2 children and 25–29 years, 1–2 children and  $\geq 30$  years,  $\geq 3$  children and <25 years, and  $\geq 3$  children and  $\geq 25$  years), age at menarche (continuous), antibiotics use within last month (yes, no), alcohol consumption (continuous), smoking (never, past and <10 years since quitting, past and  $\geq 10$  years since quitting, current), and physical activity (0–3.8, >3.8–11.1, >11.1–26.4, >26.4 MET-h/wk). Tests for trend were conducted by modeling the natural log transformed quartile median concentrations and calculating the Wald statistic.

In stratified analyses by menopausal status at blood draw, menopausal status at diagnosis, median plasma estradiol level in the follicular phase, and median plasma estradiol level in the luteal phase, we used unconditional logistic regression, additionally adjusting for matching factors, since results were essentially the same from multivariate conditional and unconditional logistic regression models. Our previous analyses of plasma estradiol levels and breast cancer risk have been published elsewhere [3, 6]. We used likelihood ratio tests to assess the effect modification by menopausal status and estradiol levels. We also examined if the association varied by invasive versus in situ breast cancer as well as by ER/PR status

using polytomous logistic regression [48]. All  $P$  values were based on two-sided tests and were considered statistically significant if  $\leq 0.05$ . We used SAS 9.3 software (SAS Institute, Cary, NC, USA) and STATA 12.1 software (StataCorp. College Station, TX, USA).

## Results

Our analysis included 802 incident cases of breast cancer and 802 matched controls. There were 553 invasive breast cancer cases and 218 in situ cases, 373 ER+/PR+, 89 ER-/PR-, and 55 ER+/PR- breast cancer cases, and 408 premenopausal cases, 302 postmenopausal cases, and 92 cases with unknown menopausal status. The average time interval between blood draw and time of diagnosis was 6.2 years with a standard deviation of 3.4 years. Compared to those in the lowest quartile of enterolactone, controls in the highest quartile of enterolactone had higher average intakes of vegetables (27.8 vs. 25.4 cups/week), whole grains (10.4 vs. 8.7 cups/week), fiber (6.1 vs. 5.2 g/day), caffeine (241.2 vs. 205.4 mg/day), and tofu (0.5 vs. 0.2 oz/day) (Table 1). Further, those in the highest versus lowest quartile of enterolactone had a lower average BMI (24.5 vs. 27.7 kg/m<sup>2</sup>), and weight change between age 18 and blood draw (21.6 vs. 38.7 lb). Comparing cases to controls, the median levels (10th–90th percentiles) of enterolactone were 11 (1–38) nmol/L among cases and 11 (1–40) nmol/L controls (Supplemental Table 1).

Overall, plasma enterolactone concentrations were not associated with risk of breast cancer (Table 2). Compared to women in the first quartile of enterolactone concentration ( $\leq 4$  nmol/L), the multivariate-adjusted RRs for the second to fourth quartiles were 1.18 (95 % CI 0.86–1.62), 0.91 (95 % CI 0.66–1.25), and 0.96 (95 % CI 0.70–1.33), respectively. No significant linear trend was observed ( $P_{\text{trend}} = 0.60$ ).

No significant associations were observed when we examined invasive and in situ tumors separately. For invasive tumors, the RR comparing the top versus bottom quartile of enterolactone levels was 1.16 (95 % CI 0.83–1.61);  $P_{\text{trend}} = 0.38$  and for in situ tumors, the comparable RR was 0.82 (95 % CI 0.53–1.26);  $P_{\text{trend}} = 0.22$  ( $P_{\text{heterogeneity}} = 0.21$ ). Similarly, no associations were observed for ER+/PR+ tumors (comparable RR = 1.23, 95 % CI 0.84–1.80;  $P_{\text{trend}} = 0.28$ ), ER-/PR- tumors (RR = 1.05, 95 % CI 0.56–1.97;  $P_{\text{trend}} = 0.76$ ), or ER+/PR- tumors (RR = 0.77, 95 % CI 0.32–1.82;  $P_{\text{trend}} = 0.46$ ;  $P_{\text{heterogeneity}} = 0.56$ ). When additionally adjusting for fiber, coffee, tofu, total vegetables, and whole grains individually in the multivariate model, similar results were observed (results not shown).

In stratified analyses, we observed that the association differed by the levels of estradiol (stratified at the median

of 47 pg/mL) during the follicular phase, ( $P_{\text{interaction}} = 0.02$ ) (Table 3). Among women with low follicular estradiol levels ( $< 47$  pg/mL), the multivariate-adjusted RRs were 1.32 (95 % CI 0.75–2.32), 0.64 (95 % CI 0.36–1.15), and 0.49 (95 % CI 0.27–0.91) for the second to fourth quartiles of enterolactone, respectively, compared with the first quartile;  $P_{\text{trend}} = 0.02$ . However, among women with follicular estradiol levels  $\geq 47$  pg/mL, the comparable RRs (95 % CI) were 1.23 (0.67–2.27), 1.03 (0.55–1.93), and 1.39 (0.73–2.65);  $P_{\text{trend}} = 0.44$ . No significant interactions were observed by BMI, fiber intake, alcohol consumption, menopausal status at blood draw, menopausal status at diagnosis, or estradiol levels during the luteal phase ( $P_{\text{interaction}} > 0.15$ , data not shown).

## Discussion

This is the largest prospective study to date examining plasma enterolactone concentrations in relation to breast cancer risk, and it is the first study examining the association stratified by follicular and luteal phase estradiol levels among premenopausal women. The median enterolactone level measured in our study is in accordance with the range of previously published circulating enterolactone concentrations in US and Europe [28–30, 33, 36, 37, 49–55], but lower than that in Asia [56]. We did not observe an association between plasma enterolactone, a biomarker of lignan intake, and breast cancer risk either overall or by hormone receptor status. However, a significant inverse association was observed among premenopausal women with lower estradiol levels during the follicular phase.

Inconsistent results regarding the enterolactone-breast cancer association have been observed between retrospective studies and prospective studies, with case-control studies generally observing an inverse association [26–41]. It is possible that breast cancer or its treatment could result in altered circulating enterolactone levels (e.g., due to changes in dietary intake post-diagnosis, or changes in gut flora as a result of breast cancer treatment), making retrospective case-control study results less reliable. On the other hand, results from ten prospective studies, with a range of sample sizes, menopausal status of participants, and biospecimen types, have been inconsistent [31–40]. Seven reported an overall null association, while two studies reported inverse associations [37, 40], and one reported an increased breast cancer risk for either very low or very high plasma enterolactone concentrations [32]. Two studies have reported inverse associations among either ER- or ER+ subgroups [33, 37]. The variation in results may be due, in part, to relatively small number of cases in six out of the ten studies (range 88–300), but more importantly due to heterogeneity of the study populations.

**Table 1** Selected characteristics of participants across quartiles of enterolactone levels among controls in the nested case–control study of breast cancer

|  | Enterolactone (nmol/L) |                |                  |                  |
|--|------------------------|----------------|------------------|------------------|
|  | Q1                     | Q2             | Q3               | Q4               |
| N  | 209                    | 197            | 196              | 200              |
| Median (10–90 percentile, nmol/L)  |                        |                |                  |                  |
| Enterolactone  | 1.1 (0–4.0)            | 8.0 (5.0–10.9) | 16.0 (12.0–20.0) | 34.0 (23.0–83.5) |
| Enterolactone (premenopausal)  | 1.5 (0–4.0)            | 7.7 (5.0–10.0) | 16.0 (12.0–20.0) | 34.0 (23.3–85.0) |
| Enterolactone (postmenopausal)   | 1.0 (0–4.0)            | 8.0 (6.0–11.0) | 15.5 (12.0–19.0) | 40.0 (24.9–66.0) |
| Mean (SD)  |                        |                |                  |                  |
| Age <sup>a</sup> (years)   | 45.2 (4.4)             | 44.8 (4.2)     | 44.2 (4.8)       | 45.5 (4.2)       |
| BMI (kg/m <sup>2</sup> )   | 27.7 (6.4)             | 26.0 (5.5)     | 25.1 (5.2)       | 24.5 (4.3)       |
| BMI at age 18 (kg/m <sup>2</sup> )   | 21.3 (3.4)             | 21.0 (2.9)     | 21.0 (3.0)       | 20.9 (2.5)       |
| Weight change since age 18 (lb)  | 38.7 (33.9)            | 29.8 (29.2)    | 23.1 (25.8)      | 21.6 (25.3)      |
| Duration of OC use among ever OC users (years)   | 3.8 (3.6)              | 4.4 (4.2)      | 3.9 (4.0)        | 3.4 (3.4)        |
| Parity among parous women  | 2.9 (1.3)              | 3.0 (1.3)      | 3.0 (1.3)        | 2.9 (1.2)        |
| Alcohol intake (g/day)   | 3.6 (7.5)              | 4.2 (8.3)      | 4.9 (8.4)        | 5.4 (8.3)        |
| Physical activity (MET-h/week)   | 15.2 (19.1)            | 21.1 (35.4)    | 19.0 (22.2)      | 19.0 (18.8)      |
| Age at menarche (years)  | 12.5 (1.6)             | 12.6 (1.4)     | 12.5 (1.3)       | 12.4 (1.4)       |
| Age at first birth (years)   | 25.6 (4.7)             | 25.8 (4.4)     | 26.4 (4.5)       | 27.3 (4.8)       |
| Luteal day of blood draw <sup>a, b</sup>   | 7.9 (3.6)              | 7.1 (2.1)      | 7.5 (2.8)        | 7.5 (2.6)        |
| Total calorie intake (kcal/day)  | 1796 (564)             | 1813 (534)     | 1870 (580)       | 1805 (510)       |
| Vegetable intake (cup/week)  | 25.4 (14.4)            | 26.1 (14.5)    | 25.0 (8.7)       | 27.8 (10.2)      |
| Whole grains intake (cup/week)   | 8.7 (4.2)              | 8.4 (4.2)      | 9.3 (3.9)        | 10.4 (5.4)       |
| Fiber (g/day)  | 5.2 (2.1)              | 5.4 (2.2)      | 5.6 (2.2)        | 6.1 (2.5)        |
| Caffeine (mg/day)  | 205 (198)              | 207 (189)      | 240 (195)        | 241 (193)        |
| Tofu (g/day)   | 5.7 (22.3)             | 8.5 (31.2)     | 11.3 (34.0)      | 14.2 (31.2)      |
| Estradiol (follicular, pg/mL)  | 52.5 (44.3)            | 60.4 (43.9)    | 56.4 (37.1)      | 54.7 (36.6)      |
| Estradiol (luteal, pg/mL)  | 138 (107)              | 150 (85)       | 147 (77)         | 152 (74)         |
| Percentage   |                        |                |                  |                  |
| Caucasian <sup>a</sup>   | 96.7                   | 99.5           | 98.5             | 98.5             |
| Family history of breast cancer  | 8.6                    | 10.7           | 11.2             | 10.0             |
| History of benign breast disease   | 14.8                   | 19.3           | 14.3             | 22.5             |
| Antibiotics use in last month  | 17.7                   | 11.2           | 7.7              | 8.0              |
| Current smokers  | 12.4                   | 6.1            | 4.1              | 4.0              |
| Postmenopausal at blood collection <sup>a</sup>  | 18.2                   | 10.7           | 12.2             | 14.5             |
| Postmenopausal hormone use at blood collection among postmenopausal women <sup>a</sup> | 83.6                   | 87.8           | 87.0             | 83.3             |

<sup>a</sup> Matching factors<sup>b</sup> Date of next menstrual cycle minus date of blood collection

Several studies observed an inverse association between enterolactone levels and breast cancer risk among postmenopausal women [29, 37, 38], and the other study that observed an inverse association only included women with palpable cysts and adjusted for cyst type in the analysis [40]. Further, three studies measured enterolactone in urine [31, 36, 38] and eight studies measured enterolactone either in serum or plasma [32–37, 39, 40], although, a strong correlation between enterolactone in urine and blood has been reported ( $r = 0.94$ ,  $P < 0.001$ ) [41]. Also, the studies

varied in the assay types. Seven studies used time-resolved fluoroimmunoassay [31–34, 37, 39, 40], and three studies used mass spectrometry [35, 36, 38].

Our results are consistent with the results from most prospective studies, in that no overall significant association was observed, including by ER status. In contrast with the previous prospective studies, we were able to evaluate the association between enterolactone and breast cancer by premenopausal estradiol levels. With our large sample size and the collection of timed samples within menstrual cycle in premenopausal

**Table 2** Multivariate-adjusted RR (95 % CI) for breast cancer by quartiles of plasma enterolactone

|                                  | Enterolactone (nmol/L) |         |                  |                  |                  | <i>P</i> trend <sup>a</sup> |
|----------------------------------|------------------------|---------|------------------|------------------|------------------|-----------------------------|
|                                  | Cases                  | Q1      | Q2               | Q3               | Q4               |                             |
| All                              |                        |         |                  |                  |                  |                             |
| Cases/controls                   |                        | 199/209 | 227/197          | 179/196          | 197/200          |                             |
| Cut-points <sup>b</sup> (nmol/L) |                        | 0–4     | >4–11            | >11–21           | >21              |                             |
| Simple RR <sup>c</sup>           | 802                    | 1.00    | 1.23 (0.92–1.64) | 0.97 (0.72–1.30) | 1.05 (0.79–1.40) | 0.97                        |
| RR <sup>d</sup>                  | 802                    | 1.00    | 1.18 (0.86–1.62) | 0.91 (0.66–1.25) | 0.96 (0.70–1.33) | 0.60                        |
| By subtype                       |                        |         |                  |                  |                  |                             |
| Invasive <sup>e</sup>            | 553                    | 1.00    | 1.32 (0.95–1.81) | 1.15 (0.83–1.60) | 1.16 (0.83–1.61) | 0.38                        |
| In situ <sup>e</sup>             | 218                    | 1.00    | 1.06 (0.70–1.60) | 0.72 (0.46–1.13) | 0.82 (0.53–1.26) | 0.22                        |
| ER+ PR+ <sup>e</sup>             | 373                    | 1.00    | 1.49 (1.04–2.15) | 1.23 (0.84–1.80) | 1.23 (0.84–1.80) | 0.28                        |
| ER– PR– <sup>e</sup>             | 89                     | 1.00    | 0.97 (0.51–1.84) | 1.18 (0.64–2.19) | 1.05 (0.56–1.97) | 0.76                        |
| ER+ PR– <sup>e</sup>             | 55                     | 1.00    | 1.22 (0.56–2.67) | 0.74 (0.31–1.75) | 0.77 (0.32–1.82) | 0.46                        |

<sup>a</sup> Linear trend test across log transformed median enterolactone levels of each quartile using the Wald test

<sup>b</sup> Cut-points obtained from controls

<sup>c</sup> Conditioning on matching factors

<sup>d</sup> Conditioning on matching factors and adjusting for family history of breast cancer (yes, no), history of benign breast disease (yes, no), OC use (never,  $\leq 1$ ,  $>1$ – $<5$ ,  $>5$  years), BMI at age 18 (continuous), weight change since age 18 (continuous), age at first birth and parity (nulliparous, 1–2 children and  $<25$  years, 1–2 children and 25–29 years, 1–2 children and  $\geq 30$  years,  $\geq 3$  children and  $<25$  years,  $\geq 3$  children and  $\geq 25$  years), age at menarche (continuous), antibiotics use within last month (yes, no), alcohol consumption (continuous), smoking (never, past and  $<10$  years since quitting, past and  $\geq 10$  years since quitting, current), and physical activity (0–3.8,  $>3.8$ –11.1,  $>11.1$ –26.4,  $>26.4$  MET-h/wk)

<sup>e</sup> Polytomous logistic regression adjusting for matching factors and the same covariates as listed above. *P*-heterogeneity were 0.21 and 0.56 for invasive versus in situ and ER+/PR+, ER–/PR–, ER+/PR– tumors

**Table 3** Multivariate-adjusted breast cancer RR (95 % CI) by quartiles of plasma enterolactone from stratified analyses

| Stratified by                          | Enterolactone (nmol/L) |      |                  |                  |                  | <i>P</i> trend <sup>a</sup> |
|--|------------------------|------|------------------|------------------|------------------|-----------------------------|
|  | Cases                  | Q1   | Q2               | Q3               | Q4               |                             |
| Premenopausal at blood draw            | 614                    | 1.00 | 1.21 (0.87–1.69) | 0.84 (0.59–1.19) | 0.85 (0.60–1.21) | 0.26                        |
| Postmenopausal at blood draw           | 110                    | 1.00 | 1.30 (0.50–3.37) | 1.94 (0.82–4.59) | 1.51 (0.63–3.64) | 0.20                        |
| Premenopausal at diagnosis             | 408                    | 1.00 | 1.23 (0.81–1.86) | 0.71 (0.46–1.09) | 0.80 (0.51–1.25) | 0.15                        |
| Postmenopausal at diagnosis            | 302                    | 1.00 | 1.25 (0.76–2.04) | 1.10 (0.66–1.85) | 1.23 (0.75–2.01) | 0.44                        |
| Estradiol $< 47$ pg/mL (follicular)    | 229                    | 1.00 | 1.32 (0.75–2.32) | 0.64 (0.36–1.15) | 0.49 (0.27–0.91) | 0.02                        |
| Estradiol $\geq 47$ pg/mL (follicular) | 220                    | 1.00 | 1.23 (0.67–2.27) | 1.03 (0.55–1.93) | 1.39 (0.73–2.65) | 0.44                        |
| Estradiol $< 132$ pg/mL (luteal)       | 243                    | 1.00 | 1.39 (0.81–2.38) | 1.07 (0.62–1.83) | 1.09 (0.61–1.95) | 0.76                        |
| Estradiol $\geq 132$ pg/mL (luteal)    | 228                    | 1.00 | 0.95 (0.53–1.72) | 0.53 (0.28–1.01) | 0.77 (0.42–1.44) | 0.19                        |

<sup>a</sup> Linear trend test across log transformed median enterolactone levels of each quartile using the Wald test

*P* for interactions were 0.26, 0.26, 0.02, and 0.15 for postmenopausal status at blood draw, postmenopausal status at diagnosis, median estradiol during the follicular phase, and median estradiol during the luteal phase, respectively

women, we observed a differential association by follicular estradiol levels, with an inverse association between enterolactone levels and breast cancer risk among women with low follicular estradiol levels. We did not observe a similar difference by luteal phase estrogens, even though enterolactone levels were measured in the luteal phase. Although no studies have examined whether plasma enterolactone levels differ by phase of the menstrual cycle, one study observed

similar urinary levels in the follicular and luteal phases [57], such that a measure of luteal enterolactone likely represents exposure in the follicular phase. Our data suggest that enterolactone could have a protective effect only in a low estrogen environment. While we did not observe similar effect modification by luteal phase estradiol, the median estradiol level is two to three times higher than that during the follicular phase [3]. We also did not observe an inverse association

among postmenopausal women, who generally have low hormone levels. This may be because we had few postmenopausal women ( $N = 222$ ), and over 85 % of those were using postmenopausal hormones at blood draw; thus, we did not have power to separately examine the postmenopausal women not using hormones. Interestingly, other prospective studies that reported inverse associations support the hypothesis that enterolactone may only be important in a low estrogen environment. First, the participants in the Sonestedt et al., Goodman et al., and Zaineddin et al. [29, 37, 38] studies were mostly postmenopausal women whose endogenous hormone production was lower than that in premenopausal women. Adjusting for postmenopausal hormone use strengthened the risk estimates in the Sonestedt et al. [37] study. Second, the Boccardo et al. study observed that high serum enterolactone concentrations were associated with a reduced breast cancer risk. They also evaluated associations by cyst type, and observed a suggestive inverse association only among type II cysts [40]. However, the Boccardo et al. study had only 383 participants and 18 breast cancer cases, which may have limited power. Experimental studies reported that estradiol concentrations are substantially lower in type II cysts [58]. This evidence suggests that it is necessary to either adjust for or stratify by estrogen environment to reveal a potential protective association between enterolactone and breast cancer risk.

Enterolactone is a phytoestrogen with complex and potentially contradicting properties, as it can act in both an estrogenic and anti-estrogenic capacity [13, 15]. Based on these results, it is possible that a protective effect of enterolactone may be important when the mammary epithelium proliferation is low. Evidence suggests that mammary epithelium proliferation is significantly lower during the follicular phase, compared to that in the luteal phase [59, 60], in part because estrogens which are higher in the luteal phase increase mammary epithelial proliferation and mitotic activity [61]. Another possibility is that enterolactone is important when ER expression is low; some studies suggest that ER expression is lower during the follicular phase [62], though other studies observed constant expression of ER over time [63, 64]. We observed no significant difference by ER status; however, prior prospective studies in postmenopausal women have reported stronger inverse associations for ER—breast cancer [29, 33]. Since our study is the first to report this potential interaction by follicular phase estradiol levels, replication in other large case–control studies nested in prospective cohort studies and more biologic studies exploring enterolactone mechanisms are warranted.

A strength of our study is that we used isotope dilution mass spectrometry to assay for enterolactone, which provided higher specificity and sensitivity than time-resolved fluoroimmunoassay [65]. Although the assay CVs were modest ( $\sim 15\%$ ), this is likely because many women in our

population had relatively low enterolactone concentrations. This measurement error may have modestly attenuated the association. Another strength of our study is the comprehensive collection of dietary and lifestyle information, as well as information on breast cancer subtypes. The prospective design of our study and the large number of incident breast cancer cases are other advantages, although we had limited statistical power to examine case subgroups. A limitation in our study is that we only measured plasma enterolactone concentration at one point in time, while the true exposure of interest is the long-term average level of enterolactone. The reliability study conducted within our cohort has shown that stability of circulating enterolactone was fair over 2–3 years ( $ICC = 0.44$ ; 95 % CI 0.22–0.68) [45]. In addition, any measurement error would be nondifferential misclassification with respect to case–control status, and would have biased our results toward the null.

## Conclusions

Overall, we did not observe an association between a biomarker of lignan intake, plasma enterolactone, and breast cancer risk in a large nested case–control study of primarily premenopausal US women. However, the association varied by follicular phase estradiol levels, with a significant inverse association observed among premenopausal women with low estradiol levels. This suggests that enterolactone may influence breast carcinogenesis only in a low estrogen environment. Further prospective studies with sufficient sample sizes, long follow-up, and accounting for circulating estrogen levels are needed to confirm our findings.

**Acknowledgments** This project was supported by the National Institutes of Health (NIH) R01 CA67272, R01 CA50385, and P30 CA71789. J.X. is supported by scholarships from Harvard University. M.S.R. is supported by the NIH T32 CA09001. The authors thank participants of the Nurses' Health Study II for their longstanding contributions and support to this study. Laurie Custer, University of Hawai'i Cancer Center, is acknowledged for her skillful performance of the LCMS assay. The authors also thank the following state cancer registries for their help: AL, AZ, AR, CA, CO, CT, DE, FL, GA, ID, IL, IN, IA, KY, LA, ME, MD, MA, MI, NE, NH, NJ, NY, NC, ND, OH, OK, OR, PA, RI, SC, TN, TX, VA, WA, and WY.

**Conflict of interest** The authors declare that they have no conflict of interest. The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; and preparation, review, or approval of the manuscript.

## References

1. Key T, Appleby P, Barnes I, Reeves G (2002) Endogenous sex hormones and breast cancer in postmenopausal women: reanalysis of nine prospective studies. *J Natl Cancer Inst* 94:606–616

2. Breast cancer and hormone replacement therapy: collaborative reanalysis of data from 51 epidemiological studies of 52,705 women with breast cancer and 108,411 women without breast cancer (1997). *Lancet* 350 (9084):1047–1059
3. Eliassen AH, Missmer SA, Tworoger SS, Spiegelman D, Barbieri RL, Dowsett M, Hankinson SE (2006) Endogenous steroid hormone concentrations and risk of breast cancer among premenopausal women. *J Natl Cancer Inst* 98(19):1406–1415
4. Kaaks R, Berrino F, Key T, Rinaldi S, Dossus L, Biessy C, Secreto G, Amiano P, Bingham S, Boeing H, de Mesquita HBB, Chang-Claude J, Clavel-Chapelon F, Fournier A, van Gils CH, Gonzalez CA, Gurra AB, Critselis E, Khaw KT, Krogh V, Lahmann PH, Nagel G, Olsen A, Onland-Moret NC, Overvad K, Palli D, Panico S, Peeters P, Quirós JR, Roddam A, Thiebaut A, Tjønneland A, Chirlaque MD, Trichopoulou A, Trichopoulos D, Tumino R, Vineis P, Norat T, Ferrari P, Slimani N, Riboli E (2005) Serum sex steroids in premenopausal women and breast cancer risk within the European prospective investigation into cancer and nutrition (EPIC). *J Natl Cancer Inst* 97(10):755–765
5. Walker K, Bratton DJ, Frost C (2011) Premenopausal endogenous oestrogen levels and breast cancer risk: a meta-analysis. *Br J Cancer* 105(9):1451–1457
6. Fortner R, Eliassen AH, Spiegelman D, Willett W, Barbieri R, Hankinson S (2013) Premenopausal endogenous steroid hormones and breast cancer risk: results from the Nurses' Health Study II. *Breast Cancer Res* 15(2):R19
7. Setchell KDR, Borriello SP, Gordon H, Lawson AM, Harkness R, Morgan DML, Kirk DN, Adlercreutz H, Anderson LC, Axelson M (1981) Lignan formation in man—microbial involvement and possible roles in relation to cancer. *Lancet* 318(8236):4–7
8. Johnsen NF, Hausner H, Olsen A, Tetens I, Christensen J, Knudsen KEB, Overvad K, Tjønneland A (2004) Intake of whole grains and vegetables determines the plasma enterolactone concentration of Danish women. *J Nutr* 134(10):2691–2697
9. Nurmi T, Mursu J, Peñalvo JL, Poulsen HE, Voutilainen S (2010) Dietary intake and urinary excretion of lignans in Finnish men. *Br J Nutr* 103:677–685
10. Adlercreutz H (2007) Lignans and human health. *Crit Rev Clin Lab Sci* 44(5–6):483–525
11. Horner NK, Kristal AR, Prunty J, Skor HE, Potter JD, Lampe JW (2002) Dietary determinants of plasma enterolactone. *Cancer Epidemiol Biomark Prev* 11(1):121–126
12. Rowland IR, Wiseman H, Sanders TAB, Adlercreutz H, Bowey EA (2000) Interindividual variation in metabolism of soy isoflavones and lignans: influence of habitual diet on equol production by the gut microflora. *Nutr Cancer* 36(1):27–32
13. Wang L-Q (2002) Mammalian phytoestrogens: enterodiol and enterolactone. *J Chromatogr B* 777(1–2):289–309
14. Welshons WV, Murphy CS, Koch R, Calaf G, Jordan VC (1987) Stimulation of breast cancer cells in vitro by the environmental estrogen enterolactone and the phytoestrogen equol. *Breast Cancer Res Treat* 10(2):169–175. doi:10.1007/bf01810580
15. Mousavi Y, Adlercreutz H (1992) Enterolactone and estradiol inhibit each other's proliferative effect on MCF-7 breast cancer cells in culture. *J Steroid Biochem Mol Biol* 41(3–8):615–619
16. Bergman Jungeström M, Thompson LU, Dabrosin C (2007) Flaxseed and its lignans inhibit estradiol-induced growth, angiogenesis, and secretion of vascular endothelial growth factor in human breast cancer xenografts in vivo. *Clin Cancer Res* 13(3):1061–1067
17. Martin P, Horwitz K, Ryan D, McGuire W (1978) Phytoestrogen interaction with estrogen receptors in human breast cancer cells. *Endocrinology* 103(5):1860–1867. doi:10.1210/endo-103-5-1860
18. Adlercreutz H, Bannwart C, Wahala K, Makela T, Brunow G, Hase T, Arosemena P, Kellis JJ, Vickery L (1993) Inhibition of human aromatase by mammalian lignans and isoflavonoid phytoestrogens. *J Steroid Biochem Mol Biol* 44(2):147–153
19. Brooks JD, Thompson LU (2005) Mammalian lignans and genistein decrease the activities of aromatase and 17 $\beta$ -hydroxysteroid dehydrogenase in MCF-7 cells. *J Steroid Biochem Mol Biol* 94(5):461–467
20. Siiteri PK (1987) Adipose tissue as a source of hormones. *Am J Clin Nutr* 45(1):277–282
21. Prasad K (2000) Antioxidant activity of secoisolariciresinol diglucoside-derived metabolites, secoisolariciresinol, enterodiol, and enterolactone. *Int J Angiol* 9(4):220–225
22. Carreau C, Flouriot G, Bennetau-Pelissero C, Potier M (2008) Enterodiol and enterolactone, two major diet-derived polyphenol metabolites have different impact on ER $\alpha$  transcriptional activation in human breast cancer cells. *J Steroid Biochem Mol Biol* 110(1–2):176–185
23. Kuiper GG, Lemmen JG, Carlsson B, Corton JC, Safe SH, Van der Saag PT, van der Burg B, Gustafsson J (1998) Interaction of estrogenic chemicals and phytoestrogens with estrogen receptor beta. *Endocrinology* 139:4252–4263
24. Penttinen P, Jaehrling J, Damdimopoulos AE, Inzunza J, Lemmen JG, van der Saag P, Pettersson K, Gauglitz G, Mäkelä S, Pongratz I (2007) Diet-derived polyphenol metabolite enterolactone is a tissue-specific estrogen receptor activator. *Endocrinology* 148(10):4875–4886
25. Buck K, Zaineddin AK, Vrieling A, Linseisen J, Chang-Claude J (2010) Meta-analyses of lignans and enterolignans in relation to breast cancer risk. *Am J Clin Nutr* 92(1):141–153
26. Ingram D, Sanders K, Kolybaba M, Lopez D (1997) Case-control study of phyto-oestrogens and breast cancer. *Lancet* 350(9083):990–994
27. Dai Q, Franke AA, Jin F, Shu X-O, Hebert JR, Custer LJ, Cheng J, Gao Y-T, Zheng W (2002) Urinary excretion of phytoestrogens and risk of breast cancer among Chinese women in Shanghai. *Cancer Epidemiol Biomarkers Prev* 11(9):815–821
28. Pietinen P, Stumpf K, Mannisto S, Kataja V, Uusitupa M, Adlercreutz H (2001) Serum enterolactone and risk of breast cancer: a case-control study in eastern Finland. *Cancer Epidemiol Biomarkers Prev* 10:339–344
29. Zaineddin AK, Vrieling A, Buck K, Becker S, Linseisen J, Fleisch-Janys D, Kaaks R, Chang-Claude J (2012) Serum enterolactone and postmenopausal breast cancer risk by estrogen, progesterone and herceptin 2 receptor status. *Int J Cancer* 130(6):1401–1410
30. Piller R, Chang-Claude J, Linseisen J (2006) Plasma enterolactone and genistein and the risk of premenopausal breast cancer. *Eur J Cancer Prev* 15:225–232
31. den Tonkelaar I, Keinan-Boker L, Van't Veer P, Arts CJM, Adlercreutz H, Thijssen JHH, Peeters PHM (2001) Urinary phytoestrogens and postmenopausal breast cancer risk. *Cancer Epidemiol Biomarkers Prev* 10(3):223–228
32. Hultén K, Winkvist A, Lenner P, Johansson R, Adlercreutz H, Hallmans G (2002) An incident case-referent study on plasma enterolactone and breast cancer risk. *Eur J Nutr* 41(4):168–176
33. Olsen A, Knudsen KEB, Thomsen BL, Loft S, Stripp C, Overvad K, Møller S, Tjønneland A (2004) Plasma enterolactone and breast cancer incidence by estrogen receptor status. *Cancer Epidemiol Biomarkers Prev* 13(12):2084–2089
34. Kilkkinen A, Virtamo J, Vartiainen E, Sankila R, Virtanen MJ, Adlercreutz H, Pietinen P (2004) Serum enterolactone concentration is not associated with breast cancer risk in a nested case-control study. *Int J Cancer* 108(2):277–280
35. Verheus M, van Gils CH, Keinan-Boker L, Grace PB, Bingham SA, Peeters PHM (2007) Plasma phytoestrogens and subsequent breast cancer risk. *J Clin Oncol* 25(6):648–655
36. Ward H, Chapelais G, Kuhnle G, Luben R, Khaw K-T, Bingham S (2008) Breast cancer risk in relation to urinary and serum biomarkers of phytoestrogen exposure in the European



- Prospective into Cancer-Norfolk cohort study. *Breast Cancer Res* 10(2):R32
37. Sonestedt E, Borgquist S, Ericson U, Gullberg B, Olsson H, Adlercreutz H, Landberg G, Wirfält E (2008) Enterolactone is differently associated with estrogen receptor  $\beta$ -negative and positive breast cancer in a Swedish nested case–control study. *Cancer Epidemiol Biomarkers Prev* 17(11):3241–3251
  38. Goodman MT, Shvetsov YB, Wilkens LR, Franke AA, Le Marchand L, Kakazu KK, Nomura AMY, Henderson BE, Kolonel LN (2009) Urinary phytoestrogen excretion and postmenopausal breast cancer risk: the multiethnic cohort study. *Cancer Prev Res* 2(10):887–894
  39. Zeleniuch-Jacquotte A, Adlercreutz H, Shore R, Koenig K, Kato I, Arslan A, Toniolo P (2004) Circulating enterolactone and risk of breast cancer: a prospective study in New York. *Br J Cancer* 91:99–105
  40. Boccardo F, Lunardi G, Guglielmini P, Parodi M, Murialdo R, Schettini G, Rubagotti A (2004) Serum enterolactone levels and the risk of breast cancer in women with palpable cysts. *Eur J Cancer* 40:84–89
  41. Grace PB, Taylor JJ, Low Y-L, Luben RN, Mulligan AA, Botting NP, Dowsett M, Welch AA, Khaw K-T, Wareham NJ, Day NE, Bingham SA (2004) Phytoestrogen concentrations in serum and spot urine as biomarkers for dietary phytoestrogen intake and their relation to breast cancer risk in European prospective investigation of cancer and nutrition-norfolk. *Cancer Epidemiol Biomarkers Prev* 13(5):698–708
  42. Stampfer MJ, Willett WC, Speizer FE, Dysert DC, Lipnick R, Rosner B, Hennekens CH (1984) Test of the National Death Index. *Am J Epidemiol* 119(5):837–839
  43. Rich-Edwards JW, Corsano KA, Stampfer MJ (1994) Test of the National Death Index and Equifax Nationwide Death Search. *Am J Epidemiol* 140(11):1016–1019
  44. Tworoger SS, Sluss P, Hankinson SE (2006) Association between plasma prolactin concentrations and risk of breast cancer among predominately premenopausal women. *Cancer Res* 66(4):2476–2482
  45. Kotsopoulos J, Tworoger SS, Campos H, Chung F-L, Clevenger CV, Franke AA, Mantzoros CS, Ricchiuti V, Willett WC, Hankinson SE, Eliassen AH (2010) Reproducibility of plasma, red blood cell, and urine biomarkers among premenopausal and postmenopausal women from the nurses' health studies. *Cancer Epidemiol Biomarkers Prev* 19(4):938–946
  46. Franke AA, Custer LJ, Wilkens LR, Le Marchand LL, Nomura AM, Goodman MT, Kolonel LN (2002) Liquid chromatographic-photodiode array mass spectrometric analysis of dietary phytoestrogens from human urine and blood. *J Chromatogr B* 777(1–2):45–59
  47. Rosner B (1983) Percentage points for a generalized ESD many-outlier procedure. *Technometrics* 25:165–172
  48. Marshall R, Chisholm E (1985) Hypothesis testing in the polychotomous logistic model with an application to detecting gastrointestinal cancer. *Stat Med* 4(3):337–344
  49. Aarestrup J, Kyrø C, Knudsen KEB, Weiderpass E, Christensen J, Kristensen M, Würtz AML, Johnsen NF, Overvad K, Tjønneland A, Olsen A (2012) Plasma enterolactone and incidence of endometrial cancer in a case-cohort study of Danish women. *Br J Nutr*, FirstView:1–7
  50. Olsen A, Christensen J, Knudsen K, Johnsen N, Overvad K, Tjønneland A (2011) Prediagnostic plasma enterolactone levels and mortality among women with breast cancer. *Breast Cancer Res Treat* 128(3):883–889
  51. Zeleniuch-Jacquotte A, Lundin E, Micheli A, Koenig KL, Lenner P, Muti P, Shore RE, Johansson I, Krogh V, Lukanova A, Stattin P, Afanasyeva Y, Rinaldi S, Arslan AA, Kaaks R, Berrino F, Hallmans G, Toniolo P, Adlercreutz H (2006) Circulating enterolactone and risk of endometrial cancer. *Int J Cancer* 119(10):2376–2381
  52. Stattin P, Adlercreutz H, Tenkanen L, Jellum E, Lumme S, Hallmans G, Harvei S, Teppo L, Stumpf K, Luostarinen T, Lehtinen M, Dillner J, Hakama M (2002) Circulating enterolactone and prostate cancer risk: a Nordic nested case–control study. *Int J Cancer* 99(1):124–129
  53. Vanharanta M, Voutilainen S, Lakka TA, van der Lee M, Adlercreutz H, Salonen JT (1999) Risk of acute coronary events according to serum concentrations of enterolactone: a prospective population-based case–control study. *Lancet* 354(9196):2112–2115
  54. Lin Y, Wolk A, Håkansson N, Peñalvo JL, Lagergren J, Adlercreutz H, Lu Y (2012) Validation of FFQ-based assessment of dietary lignans compared with serum enterolactone in Swedish women. *Br J Nutr*, FirstView:1–8
  55. Travis RC, Spencer EA, Allen NE, Appleby PN, Roddam AW, Overvad K, Johnsen NF, Olsen A, Kaaks R, Linseisen J, Boeing H, Nothlings U, Bueno-de-Mesquita HB, Ros MM, Sacerdote C, Palli D, Tumino R, Berrino F, Trichopoulos A, Dilis V, Trichopoulos D, Chirlaque MD, Ardanaz E, Larranaga N, Gonzalez C, Suarez LR, Sanchez MJ, Bingham S, Khaw KT, Hallmans G, Stattin P, Rinaldi S, Slimani N, Jenab M, Riboli E, Key TJ (2009) Plasma phyto-oestrogens and prostate cancer in the European Prospective Investigation into Cancer and Nutrition. *Br J Cancer* 100(11):1817–1823
  56. Morton M, Arisaka O, Miyake N, Morgan L, Evans B (2002) Phytoestrogen concentrations in serum from Japanese men and women over forty years of age. *J Nutr* 132:3168–3171
  57. Lampe JW, Martini MC, Kurzer MS, Adlercreutz H, Slavin JL (1994) Urinary lignan and isoflavonoid excretion in premenopausal women consuming flaxseed powder. *Am J Clin Nutr* 60(1):122–128
  58. Boccardo F, Torrisi R, Zanardi S, Valenti G, Pensa F, De Franchis V, Barreca A, Ferraro P, Minuto F (1991) EGF in breast cyst fluid: relationships with intracystic androgens, estradiol and progesterone. *Int J Cancer* 47(4):523–526
  59. Navarrete M, Maier C, Falzoni R, Quadros L, Lima G, Baracat E, Nazario A (2005) Assessment of the proliferative, apoptotic and cellular renovation indices of the human mammary epithelium during the follicular and luteal phases of the menstrual cycle. *Breast Cancer Res* 7(3):R306–R313
  60. Ferguson D, Anderson T (1981) Morphological evaluation of cell turnover in relation to the menstrual cycle in the “resting” human breast. *Br J Cancer* 44(2):177–181
  61. Pike MC, Spicer DV, Dahmouch L, Press MF (1993) Estrogens, progestogens, normal breast cell proliferation, and breast cancer risk. *Epidemiol Rev* 15(1):17–30
  62. Vasei M, Azarpira N, Talei A (2006) Status of estrogen and progesterone receptors in various phases of the menstrual cycle in breast cancer. *Arch Iran Med* 9(3):250–253
  63. Karbowski B, Jackisch C, Deppe B, Schneider H (1999) Relationship of estrogen and progesterone receptors to the menstrual cycle and menopausal status at the time of breast surgery. *Zentralbl Gynakol* 121(8):367–374
  64. Khan SA, Yee KA, Kaplan C, Siddiqui JF (2002) Estrogen receptor  $\alpha$  expression in normal human breast epithelium is consistent over time. *Int J Cancer* 102(4):334–337
  65. Grace P, Taylor J, Botting N, Fryatt T, Oldfield M, Al Maharik N, Bingham S (2003) Quantification of isoflavones and lignans in serum using isotope dilution liquid chromatography/tandem mass spectrometry. *Rapid Commun Mass Spectrom* 17:1350–1357