

# Environmental sensing and response genes in cnidaria: the chemical defensome in the sea anemone *Nematostella vectensis*

J. V. Goldstone

Received: 3 April 2008 / Accepted: 3 October 2008 / Published online: 28 October 2008  
© Springer Science + Business Media B.V. 2008

**Abstract** The starlet sea anemone *Nematostella vectensis* has been recently established as a new model system for the study of the evolution of developmental processes, as cnidaria occupy a key evolutionary position at the base of the bilateria. Cnidaria play important roles in estuarine and reef communities, but are exposed to many environmental stressors. Here, I describe the genetic components of a “chemical defensome” in the genome of *N. vectensis* and review cnidarian molecular toxicology. Gene families that defend against chemical stressors and the transcription factors that regulate these genes have been termed a chemical defensome and include the cytochromes P450 and other oxidases, various conjugating enzymes, the ATP-dependent efflux transporters, oxidative detoxification proteins, as well as various transcription factors. These genes account for about 1% (266/27,200) of the predicted genes in the sea anemone genome, similar to the proportion observed in tunicates and humans, but lower than that observed in sea urchins. While there are comparable numbers of stress-response genes, the

stress sensor genes appear to be reduced in *N. vectensis* relative to many model protostomes and deuterostomes. Cnidarian toxicology is understudied, especially given the important ecological roles of many cnidarian species. New genomic resources should stimulate the study of chemical stress sensing and response mechanisms in cnidaria and allow us to further illuminate the evolution of chemical defense gene networks.

**Keywords** Cytochrome P450 · Glutathione transferase · ABC transporter · Aromatic hydrocarbon · Nuclear receptor · Metal · Superoxide dismutase · Oxidative stress

Cnidaria occupy a key basal evolutionary position within Metazoa (Dunn et al. 2008), with recent evidence suggesting that they are early-diverging bilaterians (de Jong et al. 2006; Matus et al. 2006). Cnidaria have important ecological roles as reef structure builders and as predators and prey in planktonic and benthic ecosystems (e.g., Harborne et al. 2006; Sebens 1981). Cnidaria are sensitive to many environmental stressors and have been used as indicators of water quality (Arkhipchuk et al. 2006; Davies and Freeman 1995; Wiger and Stottum 1985). With a better understanding of regulatory processes and development of appropriate endpoints (e.g., Tarrant 2007), cnidaria will become valuable indicators of exposure to disruptive chemicals and other stressors.

---

**Electronic supplementary material** The online version of this article (doi:10.1007/s10565-008-9107-5) contains supplementary material, which is available to authorized users.

---

J. V. Goldstone (✉)  
Biology Department MS #32,  
Woods Hole Oceanographic Institution,  
Woods Hole, MA 02543, USA  
e-mail: jgoldstone@whoi.edu

The starlet sea anemone *Nematostella vectensis* has been recently established as a new model system for the study of the evolution of developmental processes (Darling et al. 2005; Putnam et al. 2007) and may act as a model for the basic molecular biology of anthozoans. The remarkable amenability of this species to laboratory manipulation has already made it a productive system for exploring cnidarian development and the origins of bilateral symmetry (Finnerty and Martindale 1999; Finnerty et al. 2004; Fritzenwanker et al. 2004; Kusserow et al. 2005; Magie et al. 2005; Matus et al. 2006; Torras and Gonzalez-Crespo 2005).

*N. vectensis* is a burrowing estuarine anemone, with populations in the eastern Pacific, northern English Channel, western North Sea, and western Atlantic (Hand and Uhlinger 1994), although it is likely that all but the western Atlantic represent introduced populations (Reitzel et al. 2008a). It can tolerate a remarkably wide ranges of salinities (2–54 ppt), temperatures (–1°C to 28°C), and dissolved oxygen concentrations (Sheader et al. 1997). The facility with which *Nematostella* populations can be investigated within their natural ecological context (Darling et al. 2005) suggests that this model may also be profitably expanded to address important questions in molecular and evolutionary ecology and toxicology. A mechanistic understanding of stress responses is essential to establishing this model system, as with all model systems.

An important question in biology is how cells and organisms maintain homeostasis in a variable environment. The need to deal with physical, chemical, and biological stressors has driven the evolution of an array of gene families and pathways (also known as “environmental genes” (Ponting 2008)) that afford protection from challenges. The immune system is one such protective mechanism, which responds to biotic stressors such as pathogens (Miller et al. 2007). Another set of genes comprises the “chemical defense”, encoding a network of defensive proteins that allows the organism to sense, transform, and eliminate potentially toxic chemicals (Goldstone et al. 2006).

The chemical defense protects against chemically mediated injury by environmental chemicals such as heavy metals, microbial products, and other natural exogenous compounds, as well as anthropogenically derived compounds such as hydrocarbon derivatives and pesticides. These compounds are structurally


diverse, requiring either non-specific enzymatic responses or a broad array of specific enzymatic actions. In addition, the maintenance of cellular homeostasis requires the inactivation and elimination of endogenous signaling molecules, such as eicosanoids, and defense against endogenously generated toxicants such as reactive oxygen species (ROS).

The chemical defense is comprised of several classes of proteins that function coordinately to protect the cell (Fig. 1). These proteins include enzymes that transform chemicals to less toxic and more readily excretable metabolites, efflux transporters that actively eliminate toxicants and transformed products, antioxidant enzymes protecting against externally and internally generated ROS or other radicals, and soluble receptors and ligand-activated transcription factors that act as sensors of toxicants or cellular damage.

Afflux transporter proteins such as the ATP-binding cassette (ABC) transporters can provide the first line of cellular defense (Dean et al. 2001). Once toxicants enter the cytoplasm, however, biotransformation is often required to inactivate or enhance the elimination of toxicants. Biotransformation enzymes include oxidative enzymes such as the cytochromes P450 (CYPs); reductive enzymes such as aldo-keto reductases (AKR), epoxide hydrolase (EH), and NAD (P)H-quinone oxidoreductase (NQO); and conjugative enzymes including glutathione-S-transferases (GST), sulfotransferases (SULT), UDP-glucuronosyl transferases (UGT), and *N*-acetyl transferases (NAT). Biotransformation generally results in detoxification, but oxidation, *N*-acetylation, sulfate, or glutathione conjugation can lead to toxic metabolites in a chemical and cell-specific manner (Gamage et al. 2006; Guengerich et al. 2003; Surh. 1998).

Gene products that protect against injury from chemicals may be especially important in embryos given the complex chemical signaling pathways governing development (Davidson and Erwin. 2006; Hamdoun and Epel 2007), as well as the need to protect the genome of the germ cells (Epel 2003). In adults, some of these proteins also provide protection from environmental factors, such as oxidative stress, that can lead to senescence (Finkel and Holbrook 2000). Many gene products in this network (e.g., CYPs) perform multiple roles, having important endogenous functions (including but not limited to development), as well as functioning in chemical defense.

**Fig. 1** Conceptual organization of the cellular defensesome. Organic and inorganic toxicants are actively exported and also subjected to a variety of biotransformative reactions. Modified from Goldstone et al. (2006)



Here, I show that the major elements of the network of genes and pathways that allow an organism to mount a defense against toxic chemicals appear to be conserved in cnidaria and review relevant aspects of cnidarian molecular toxicology. Almost all of the gene families or superfamilies that are characteristic of the chemical defensive network in deuterostomes (Goldstone et al. 2006) are also represented in the sea anemone (Fig. 1; see also Reitzel et al. 2008b), indicating the presence of this system in the bilaterian ancestor and evolutionary conservation. However, while there is general conformity in the presence of higher order gene groups across taxa, in most cases, gene orthology is more difficult to determine.

## Methods

Different types of evidence are available for the genes discussed in this paper. Predicted genes are derived from the US Department of Energy Joint Genome Institute (JGI) predictions of the whole genome shotgun assembly ([www.jgi.doe.gov](http://www.jgi.doe.gov)). Many of these predicted genes are supported by expression data from an extensive EST collection (Sullivan et al. 2008). Resources are available online at stellabase.

[org.cnidbase.bu.edu](http://org.cnidbase.bu.edu) and [nematostella.org](http://nematostella.org). In this study, defensesome genes were identified by Hidden Markov Model searches (Hmmer v2.3.2; Eddy 1998) of the JGI gene predictions with conserved domains of known defense genes using the PFAM models. Gene homologies were confirmed by reciprocal BLAST of the predicted genes against Genbank. For this study, the JGI “best models” were used without significant refinement. Alignments were constructed using Muscle v3.6b (Edgar 2004) and are available upon request of the author.

The exact nomenclature of many genes presented in this paper is tentative because of the uncertainty in classifying genes to specific subfamilies within the major superfamilies represented in these analyses. I have attempted to follow the nomenclature guidelines for many of the defined gene superfamilies (Hyndman et al. 2003; Jez and Penning 2001; Mackenzie et al. 2005; Nebert and Vasiliou 2004; Nelson et al. 1993; Vasiliou and Nebert 2005; Vasiliou et al. 2006), but due to evolutionary distances, some of the subfamily assignments are tentative. Thus, new genes here are given names, indicating my understanding of the homologous relationships, but that should not be taken as formal assignments. Formal assignments of new gene names are often reserved by specific nomenclature committees (e.g., the Cytochrome

P450 Nomenclature Committee or the Aldo-Keto Reductase Nomenclature Committee). Based on evidence from our previous analysis of the sea urchin genome (Goldstone et al. 2006), gene orthologies may also not be predictive of function.

Phylogenetic trees were constructed by analyzing amino acid sequences using maximum likelihood (RAxML 7.0.3; Stamatakis 2006). Regions of alignment uncertainty were excluded from phylogenetic analysis (Kreil and Ouzounis 2003) by automatic masking using a custom-written script. The WAG-CAT model of amino acid substitution (Whelan and Goldman 2001) with a gamma distribution of substitution rates was used in all likelihood analyses, based on likelihood tests using RAxML.

## Defensome gene families

### Receptors and signal transduction

Homologs of most important stress receptors are present in the sea anemone genome, including the aryl hydrocarbon receptor (AHR), hypoxia-inducible factor 1 (HIF1 $\alpha$ ), and the aryl hydrocarbon nuclear translocator (ARNT), metal transcription factor 1 (MTF1), nuclear factor-kappa B (NFkB), and nuclear factor erythroid-derived 2 related 2 (NRF2), detailed below (Fig. 2). Although some of the known components of the vertebrate and invertebrate xenobiotic receptor pathways are missing (e.g., pregnane X receptor, liver X receptor, farnesoid X receptor [PXR, LXR, and FXR]), receptors that are not clearly orthologous to known xenobiotic sensors may substitute, or there may be an increased xenobiotic receptor promiscuity.


*Aryl hydrocarbon receptor (AHR) and related bHLH-PAS proteins* Basic helix-loop-helix PER/ARNT/SIM (bHLH-PAS) family genes encode proteins involved in critical physiological and developmental signaling, including those that mediate responses to certain environmental pollutants (including polynuclear aromatic hydrocarbons) and low oxygen tension (Kewley et al. 2004). bHLH-PAS genes in chordates that have been shown to be important to physiological responses to environmental pollutants include the aryl hydrocarbon receptor (AHR), HIF1 $\alpha$ , and the ARNT.

The bHLH gene family has previously been examined in *N. vectensis* and other species (Simionato et al. 2007). Simionato et al. identified 68 bHLH genes, several of which also contained a PAS domain, including one or two ARNT genes and zero to two HIF genes (the range depends on uncertainty in the phylogenetic clustering; Simionato et al. 2007), but could not identify an AHR in the *N. vectensis* genome. However, Reitzel et al. (2008b) identify a gene (gi|156394392) as a putative AHR homolog and note that its expression is confirmed through an EST. Both AHR and HIF1 $\alpha$  form heterodimers with ARNT to regulate transcription of downstream targets through the recognition of specific DNA response elements. Transcriptional responses to potential activators of AHR and HIF have not been well studied in sea anemones, and no data are available to determine if these response elements are conserved in *N. vectensis*.

*Oxidative and metal stress-response transcription factors* Oxidative stress-response factors in vertebrates include the CNC-bZIP family [nuclear factor erythroid-derived 2 and related factors (NRFs)], the BTB-bZIP proteins BACH1 and BACH2, and the small Maf proteins (MafF, MafG, and MafK in particular). Maf proteins in vertebrates are heterodimeric partners of NF-E2, NRFs, and Bach proteins (Igarashi and Sun 2006). In addition to their roles as heterodimerization partners for various CNC proteins, small Maf proteins have critical roles in vertebrate stress signaling, oncogenesis, and may also have links to the inflammation response (Blank 2008).

*N. vectensis* has two homologs of the small Maf proteins, an NRE2-like protein homologous to NRF2 and a KEAP1-like protein, which in the absence of oxidative stress in vertebrates, encodes a protein that retains NRF2 in the cytoplasm and enhances its proteasomal degradation (Nguyen et al. 2003). In vertebrates, the NRF2 signaling pathway provides a rapid response to electrophilic or oxidative compounds and has been shown to attenuate carcinogenesis and inflammation (Osburn and Kensler 2008).

Other important oxidative stress-responding transcription factors with homologs in sea anemone include MTF1 and NFkB. MTF1 is well known as a metal-responsive transcription factor (Laity and Andrews 2007), but has also been proposed as a



**Fig. 2** Some of the stress-response transcription factor pathways with homologs in *N. vectensis*. Hypoxia activates both the HIF1 $\alpha$  and MTF1 pathway, metal stress activates MTF1, oxidative stress activates the NRF2 pathway (as well as others, not shown), and organic xenobiotics activate AHR or various NRs. These transcription factors have specific response

elements (REs) in the regulatory regions of responsive genes, including hypoxia RE (HRE; HIF1 $\alpha$ /ARNT), metal RE (MRE; MTF1), antioxidant RE (ARE; NRF2), xenobiotic RE (XRE; AHR/ARNT), and specific NR-REs (e.g., estrogen response elements)

generalized sensor of oxidative stress (Murphy 2004; Murphy et al. 1999, 2005). MTF1 may also interact with HIF1 $\alpha$  and contribute to HIF1 $\alpha$  activation during hypoxia (Murphy et al. 2005).

**Nuclear receptors** Ligand-activated nuclear receptors (NRs) function as chemically activated transcription factors, primarily with endogenous functions but also importantly in xenobiotic sensing. Of greatest interest with regards to the chemical defenses are those related to NRs in the NR1H and NR1I subfamilies, which contain vertebrate FXR, LXR, and PXR, constitutively active receptor (CAR), and the vitamin D receptor (VDR), as well as arthropod EcR (ecdysone receptor). Other NRs involved in xenobiotic response in vertebrates include estrogen receptor (ER; NR3A subfamily); the peroxisome proliferator receptors (PPARs; NR1C subfamily), which have target genes involved in lipid metabolism, energy homeostasis, and cell differentiation; and the retinoid X receptor (RXR; NR2B subfamily), which has many target genes involved in xenobiotic metabolism.

*N. vectensis* and other cnidaria appear to lack many nuclear receptors traditionally studied in response to toxicants (e.g., NR1s, ER; Grasso et al. 2001; Reitzel et al. 2008b). *N. vectensis* appears to have a modest number of NRs (18), none of which are related to the NR1H (PPAR, LXR, FXR) or NR1I (VDR, PXR, CAR) families. However, there are genes related to hepatocyte nuclear factor 4 (HNF4, NR2A) and to RXR, indicating the presence of ancestral NR2 subfamily members in this cnidarian. An RXR gene has been cloned from a cubozoan, *Tripedalia cystophora*, and the protein binds 9-*cis* retinoic acid with high affinity (Kostrouch et al. 1998).

Although there does not appear to be an ER in *N. vectensis*, the existence of a bilaterian ancestral steroid-binding receptor was inferred based on ancestral protein reconstruction (Thornton et al. 2003). Cnidaria appear to be susceptible to signal disruption by exogenous estrogens (reviewed in Tarrant 2005, 2007), although there appear to be differences between coral and hydra sensitivity (Pascoe et al. 2002; Tarrant et al. 2004). Estrogen signaling may

still be important in corals, however, as estrogens have been found in and around spawning corals, and corals have the ability to metabolize estradiol and testosterone (Atkinson and Atkinson 1992; Blomquist et al. 2006; Tarrant et al. 1999, 2003; Twan et al. 2003, 2006).

#### Efflux transporter proteins

Many toxic compounds are pumped against concentration gradients across membranes in an energy-dependent process. This first line of cellular defense, against amphipathic or slightly lipophilic compounds in particular, is mediated by efflux proteins known as ABC or multidrug efflux transporters, including the *p*-glycoproteins (PGP/ABCB), mitoxantrone resistance protein (MXR/ABCG2), and multidrug resistance proteins (MRP/ABCC; Dean et al. 2001). Efflux transporters function to export both unmodified substrates and substrates modified by other defense enzymes (Deeley et al. 2006). In embryos, efflux transporters may provide the primary defense against exogenous toxicants but also play important roles in developmental programs by establishing morphogen gradients (Hamdoun and Epel 2007).

In chordates, the ABC transporters are organized into eight subfamilies designated ABC *A* through *H* (Annilo et al. 2006). A subset of these families includes proteins known to export toxicants: the ABCB, ABCC, and ABCG transporters. These proteins are commonly called multidrug resistance transporters after their ability to pump out multiple therapeutic drugs, a major obstacle to the efficacy of the treatment of several pathogens (Dean et al. 2005).

Genome searches revealed that sea anemones have 64 ABC genes organized into six subfamilies, including the three multidrug transporter subfamilies

(ABC B, C, and G; Table 1, Supplemental Table S1). There is considerable variation in the total number of ABC genes within eukaryotic genomes, but the relative proportions have tended to stay constant (Annilo et al. 2006; Goldstone et al. 2006). The ABC genes clustered in the ABCA (five genes), ABCD (six genes), and ABCF (four genes) families either do not have known function or do not have known roles in detoxification and will not be considered further here.

*N. vectensis* has seven ABCB genes, including two related to the ABCB1 (pgp) proteins. The pgp transporters are well known as multidrug resistance proteins involved in the efflux of toxic compounds. Additional genes related to known xenobiotic transporters include six ABCC4 (MRP4)-like genes, found in a sea anemone-specific cluster; six other ABCC-like genes including one ABCC5 (MRP5)-like sequence; and 24 other genes that cluster in a large anemone-specific clade within the ABCC family. Finally, analyses conducted in this study identified six ABCG sequences, including one sequence that clusters closely with the vertebrate ABCG2s. In vertebrates, ABCG2 proteins exhibit broad substrate specificity among xenobiotic compounds and play critical roles in the clearance of certain drugs (Allikmets et al. 1998; Kusuhara and Sugiyama 2007; Miyake et al. 1999).

Other potentially important anemone transporters include the organic anion polypeptides (OATP; solute carrier family 21, SLC21) and organic anion and cation transporters (OAT and OCT; solute carrier family 22, SLC22). Both SLC21 and SLC22 are part of the major facilitator superfamily. OAT substrates examined in vertebrates include estrone sulfate, urate, prostaglandins, heavy metals such as mercury and cadmium, and the herbicide 2,4-dichlorophenoxyacetic acid (Eraly et

**Table 1** Gene counts of xenobiotic transporter genes

Superfamily	Gene family	Human	Urchin	Anemone
ABC superfamily	ABCB	11	12	7
	ABCC	12	30	36
	ABCG	5	9	6
	Other	20	14	16
	Total	48	65	65
Major facilitator	SLC21A	11	30	17
	SLC22	5	46	62

Data for human were taken from Dean and Annilo (2005) and for urchin from Goldstone et al. (2006)

al. 2004; Kimura et al. 2002; Sweet 2005). OATPs have partially overlapping substrate specificities for steroid conjugates, bile salts, anionic oligopeptides, and anionic xenobiotics including toxins and drugs (Hagenbuch and Meier 2003; Jacobsson et al. 2007). The anemone genome contains 17 OATP genes and 62 SLC22 (OCT and OAT) genes. However, orthology among non-vertebrate SLC families is difficult to assign, precluding any hypotheses regarding substrate specificity.

#### Oxidative or reductive biotransformation

*Cytochromes P450* Oxidative modification of chemicals to more hydrophilic products is often the initial step leading to excretion. In bilaterians, this is carried out by CYP and flavoprotein monooxygenase (FMO) enzymes, especially members of the CYP1, CYP2, CYP3, CYP6, CYP9, and CYP4 families. Toxicant oxidation can, however, also lead to increased toxicity; for example, oxidation of benzo[*a*]pyrene by CYP1A leads to hepatotoxicity (Uno et al. 2001).

The sea anemone genome contains 82 CYP genes, which are in general not classifiable into established CYP families due to the low (<40%) identity with other known CYPs. A large-scale reclassification of metazoan CYPs taking into account recent genomic data will be required to formally name the *N. vectensis* CYPs (Nelson, personal communication). However, broader classification into the CYP clan framework (Nelson 1998, 2006) is possible: Clan 2, containing CYP families 1, 2, 17, 18, 21, 33, 34, and 35; Clan 3, containing primarily CYPs 3, 5, 6, 9, 28, 309, 310, and 317; Clan 4, containing CYPs 4, 311, 313, 316, and 318; and the mitochondrial clan, CYPs 11, 12, 24, 27, 44, 49, 302, 314, and 315.

Sea anemone CYPs are principally part of Clan 2 and Clan 3, with 39 and 20 genes in these two Clans, respectively, while there are only three Clan 4 genes (Table 2, Fig. 3, Supplemental Figure S1, Supplemental Table S2). Many of the CYPs in these clans are involved in detoxification of exogenous and endogenous compounds (Lewis et al. 2004), although the functional information is primarily from vertebrates and insects. The anemone Clan 2 CYPs are clustered more closely to the vertebrate CYP17s (and thus the important xenobiotic-detoxifying CYP1s, including aryl hydrocarbon hydroxylases) than to the vertebrate CYP2s. However, it is not clear that CYP1

genes exist outside the deuterostomes (Goldstone et al. 2007), and these sea anemone CYPs cannot be considered early CYP1-like genes. The Clan 3 genes are likewise less closely related to the vertebrate CYP3 or insect CYP6 detoxification genes than to other members of Clan 3, in this case, the CYP5-like genes. CYP5 genes have unusual functionality in that they catalyze a rearrangement of a prostaglandin endoperoxide (Hecker and Ullrich 1989), rather than a substrate oxidation. More generally, the sea anemone Clan 3 CYPs may oxidize prostaglandins, which are potent chemical defenses in marine systems (Paul and Puglisi 2004; Paul et al. 2006).

*N. vectensis* does not have a CYP19 (aromatase), despite the fact that (low) aromatase activity has been demonstrated in a scleractinian coral, *Euphylia ancora* (Twan et al. 2003). CYP19 is not present in most invertebrates with a sequenced genome (Goldstone, Nelson, and Stegeman, unpublished data), although it is present in amphioxus (Castro et al. 2005; Mizuta and Kubokawa 2007). It is very possible that a different CYP, not CYP19, possesses aromatase activity.

*Other redox enzymes* Other proteins that oxidize or reduce toxicants include the FMO (Ziegler 2002), AKR (Jin and Penning 2007), aldehyde dehydrogenases (ALDH), NQO, and epoxide hydrolase (EPHX). In contrast to the CYPs, much less is known about many of the substrates of these enzymes, even in humans (Krueger and Williams 2005; Penning and Drury 2007).

In addition to the 82 CYPs, analyses conducted in this paper identified genes for six FMO enzymes. Although both enzyme families are primarily monooxygenases and have some overlapping substrate specificities (Krueger and Williams 2005; Ziegler 2002), FMOs are generally thought to oxidize soft nucleophiles, while CYPs often catalyze C–H abstraction (Cashman 2005). FMOs are less stable enzymes than CYPs, which has contributed to the relative lack of functional knowledge. The sea anemone FMO enzymes are quite distinct from the known human FMOs, and, as with the anemone CYPs, specific functions cannot be easily guessed at.

The sea anemone genome has at least 12 AKR genes, 21 ALDH, and one EPHX gene. These numbers are comparable to deuterostome gene inventories. In vertebrates, EPHX contributes to the toxicity

**Table 2** Gene counts of biotransformative genes in the humans, sea urchin, and sea anemone genomes

Classification	Gene	Human	Urchin	Anemone
Oxidative	CYP Clan 2	21	85	39
	CYP Clan 3	5	13	20
	CYP Clan 4	12	10	3
	FMO	6	16	6
	ALDH	19	20	21
Conjugative	GST	21	38	18
	MGST	3	12	5
	SULT	13	73	22
	UGT	13 <sup>a</sup>	50	9
	NAT	2	1	0
Reductive	AKR-like	8	10	12
	EPHX	2	5	1
	NOQ1	2	0	0


<sup>a</sup>Not including multiple first exon expression in UGT1

of benzo[*a*]pyrene by converting the benzo[*a*]pyrene epoxides produced by CYP1s to benzo[*a*]pyrene dihydrodiols (Shimada 2006), which eventually can be oxidized to redox-cycling benzo[*a*]pyrene quinones by AKR (Palackal et al. 2001; Penning et al. 1999).

One of the most important ALDH reactions in vertebrate development is the irreversible oxidation of retinal to retinoic acid (Lee et al. 1991); retinoids play very important roles in vertebrate patterning and are also likely important in cnidarian development

(Bouzaiene et al. 2007; Johnson and Chun 1989; Kostrouch et al. 1998; Muller 1984). ALDH enzymes may also help maintain the cellular redox balance via ROS scavenging and the production of reducing equivalents as NADPH or NADH.

NQO enzymes catalyze the two-electron reduction of quinones to hydroquinones, reducing the formation of semiquinones and the potential for reactive oxygen generation (Vasiliou et al. 2006). Similar to sea urchins (Goldstone et al. 2006), analyses conducted for this paper revealed that sea anemones do not have NQO-like genes. This is in line with the observed lack of NQO genes in the worm, fly, sea squirt, or plants (Vasiliou et al. 2006).



**Fig. 3** Distribution of genes in CYP Clans in human, tunicate, sea urchin, and sea anemone. Data for the tunicate and sea urchin assignments were taken from Goldstone et al. (2006)

### Conjugative biotransformation

Sea anemones possess relatively few proteins with direct homology to xenobiotic-conjugating enzymes, particularly in comparison to the purple sea urchin (Table 2; Goldstone et al. 2006). Sea anemones have genes for 23 GST, including five microsomal GSTs (MAPEG), nine UGT genes, and 22 SULT genes. No NAT genes were found. These numbers are far lower than the large diversification of these gene families seen in sea urchins, but comparable to the numbers observed in mammalian genomes (the human genome contains 13 SULT, 13 UGT, and 21 GST genes; Gamage et al. 2006; Mackenzie et al. 2005; Nebert and Vasiliou, 2004).

Cytosolic GSTs are soluble proteins that catalyze the transfer of glutathione to an electrophilic substrate



(Hayes et al. 2005). Microsomal or membrane GSTs (MAPEG) form an evolutionarily distinct class of enzymes that exhibit both glutathione transferase and lipid peroxidase activity (Bresell et al. 2005), thus detoxifying both xenobiotic compounds and ameliorating oxidative stress. The majority (15) of the 18 sea anemone GSTs are readily classifiable, including three mu-class, three omega-class, six sigma-class, one theta-class, one fungal-type, and one zeta-class. The three remaining GSTs appear homologous to the xenobiotic-metabolizing alpha/pi GSTs. This search also found a sequence homologous to the translation elongation factor 1g, which contains a GST domain but does not have glutathione transferase activity. The *N. vectensis* genome also codes for a total of five MAPEG sequences, including one homologous to vertebrate MAPEG1, one sequence homologous to MAPEG3, and three sequences homologous to prostaglandin E synthase (PTGS). PTGS enzymes are MAPEG superfamily members important to eicosanoid synthesis and involved in the vertebrate inflammation response (Jakobsson et al. 1999). Prostaglandins in corals are very important in chemical defense (reviewed in Paul and Puglisi 2004; Paul et al. 2006) and have been extensively studied in gorgonians. A prostaglandin synthase with 50% identity to mammalian PTGS has been cloned from an Arctic soft coral (Koljak et al. 2001).

SULT and UGT enzymes catalyze the conjugation of sulfuryl groups donated by 3'-phosphoadenosine-5'-phosphosulfate or UDP-glucuronide, respectively, to a wide variety of substrates, including both xenobiotics and endogenous products (Bock and Kohle 2004; Gamage et al. 2006; Runge-Morris and Kocarek 2005). Cytosolic (soluble) SULTs are responsible for the metabolism of xenobiotic and small endogenous substrates (SULT1 and SULT2), while membrane-bound SULTs are involved in endogenous peptide, lipid, and aminosugar sulfonation (Gamage et al. 2006). I found 22 SULT genes in the sea anemone genome, all of which are more closely related to the SULT genes involved in energy metabolism rather than those SULT genes known from vertebrate studies to participate in detoxification reactions. The anemone genes are divided among the SULT3A family (eight genes), SULT3B (two genes), SULT4 (four genes), and carbohydrate keratan/chondroitin SULTs (eight genes). Chondroitin sulfation has been demonstrated in the nematocysts of *Hydra magnipapillata*

(Yamada et al. 2007), and it is possible that the *N. vectensis* genes are involved in similar functions.

The sea anemone UGT genes are likewise not closely related to the UGT families with known xenobiotic-metabolizing or detoxification roles. UGT1 genes in mammals consist of one gene with as many as 14 different first exons, complicating the assignment of UGT homology (Mackenzie et al. 2005). Based on our previous analysis of the large number of distinct genes in the sea urchin, exon duplication like that observed in the mammalian UGT families is not the only method of UGT diversification. However, the nine anemone UGTs are not classifiable to any of the known vertebrate UGT families, and thus, no function can even be hinted at. This finding is not unique to anemones, as other marine genomes contain what appear to be lineage-specific gene family expansions that are not readily assignable to known UGT functional classes (J. Goldstone, unpublished data).

#### Antioxidant proteins

ROS, including superoxide, hydrogen peroxide, and hydroxyl radicals, are derived from a variety of cellular processes, including leakage from mitochondrial respiration. Reactive oxygen can also be produced by exposure to toxicants and to ultraviolet radiation. ROS contribute to diseases and pathologies generally deriving from altered gene expression or damage to biomolecules, including proteins, lipids, and DNA (Halliwell and Gutteridge 1999; Lesser 2006). General antioxidant defensive genes include superoxide dismutase (SOD), catalases (CAT), and peroxidases, including glutathione peroxidase (GPX), peroxiredoxin, and thioredoxins (TXNs).

The sea anemone genome has a total of six SOD genes: three Cu/Zn SOD genes, one Mn SOD gene, two Fe SODs, as well as an SOD copper chaperone homolog (which contains an SOD domain but has no dismutase activity). Both EST and cDNA libraries support the expression of all six SOD forms under normal conditions (A. Reitzel, A. Tarrant, J. Goldstone, unpublished data). In addition, there is one CAT, 12 glutathione peroxidase genes, and six other heme peroxidase genes. This abundance of antioxidant defense genes is complemented by a complete glutathione system (glutathione reductase and 4 gamma-glutamyl transferases), as well as TXN and thioredoxin reductase (TXNRD).

## Metal detoxification

Heavy metals are important aquatic pollutants resulting from sewage, urban and agricultural runoff, and antifouling paint. Bioconcentration of heavy metals can lead to tissue concentrations that are ten- to 10,000-fold higher than environmental levels, resulting in a variety of toxic effects. Four phytochelatin synthase (PCS) homologs are present in the *N. vectensis* genome and expressed under normal conditions (A. Reitzel, A. Tarrant, J. Goldstone, unpublished data). Phytochelatins are metal-binding peptides composed primarily of glutathione groups that are important metal detoxifying genes in plants and fungi. Until PCS was discovered in the nematode *Caenorhabditis elegans* (Clemens et al. 2001), it was believed that phytochelatins were present only in plants and fungi. Now, it is clear that many other lineages contain PCS homologs (Clemens 2006), including the sea urchin (Goldstone et al. 2006). Currently sequenced vertebrate genomes do not contain a gene homologous to PCS, nor do insect genomes, suggesting that phytochelatin synthesis ability was lost independently in some protostome and deuterostome lineages.

No metallothionein (MT) genes, neither plant- nor fungi- nor metazoan-related, were found in the sea anemone genome despite extensive searching, perhaps because of the presence of the alternative metal-complexing phytochelatin system. The absence of MT genes is apparently due to gene loss, as MT proteins are important metal detoxification proteins in plants (Cobbett and Goldsbrough 2002), mollusks (Amiard et al. 2006), sea urchins (Nemer et al. 1985), and vertebrates, and are also present in sponges (Berthet et al. 2005; Philp 1999).

Active efflux of toxic metals is another important route to detoxification. Both OAT and ABC efflux proteins (see above) export metals (Leslie et al. 2001; Sweet 2005), and the anemone contains genes homologous to the transporters (within both the OAT and ABC families) known to facilitate metal export.

## Heat shock proteins

Heat shock proteins (HSP) have been implicated in the response to various toxicants, including cadmium, arsenic, and free radicals (Feder and Hofmann 1999). The induction of HSP mRNA and protein by heat

shock factor 1 appears to be part of generalized cellular stress response, and HSPs may not only act as chaperones but also assist in refolding of partially denatured proteins (Kim et al. 2006). Sea anemones have several families of heat shock proteins, including HSP 90 and 70 and small alpha crystalline HSPs (HSP20s). The largest family of heat shock proteins is the HSP20 family, containing at least 18 genes. Sea anemones also have at least four HSP90 genes and nine HSP70s. Various coral and anemone HSP60, HSP70, and HSP90 proteins and cDNA sequences have been shown to be strongly induced not only by heat or cold shock (Choresch et al. 2007, 2004, 2001; Hashimoto et al. 2004; Robbart et al. 2004; Rossi and Snyder 2001; Rossi et al. 2006; Sharp et al. 1997, 1994; Snyder and Ross 2004) but also by PCB118 (Wiens et al. 2000).


## Discussion

The chemical defensesome is an integrated network of chemical sensing and response proteins that function as an organized defense against toxic chemicals, both endo- and exogenous (Goldstone et al. 2006). Elucidation of the chemical stress-response repertoire of *N. vectensis* provides a framework for studies on a number of cnidarian-specific questions as well as on broader evolutionary questions. Characterization of these stress-response genes in *N. vectensis* facilitates the use of this and other anemones as sentinel species for changing environmental stressors. *N. vectensis* is a hardy species, tolerating extremes of temperatures unknown to other members of the family Edwardsiidae, which are restricted to temperate and polar coastal seas with less dramatic temperature and salinity variations (Daly 2002). Identification of molecular responses to chemical stress will help us to develop markers that will allow *N. vectensis* and other anemones to act as sentinels of environmental contamination.

The major components of this defensive gene network are conserved in the sea anemone genome (Fig. 4), indicating that they must have origins prior to the cnidarian–bilaterian split. Interphyla comparison of the components and linkages within the chemical defensesome will help us understand the early evolution of the chemical stress response. Despite the fact that the individual genes within the defensesome network may vary across organisms, this network may be comprised of evolutionarily conserved mod-

ules, which are retained across evolution. Comparing the susceptibility of sea anemone embryos with that of deuterostome and protosome embryos, for a range of chemicals, could lead to fundamental insights into how these defensome “kernels” function to protect embryos from the myriad chemical challenges that could derail development. Predictions of defensome interactions (e.g., the roles of nuclear receptors in simultaneously modulating multiple parts of the defensome) are testable using microarray analysis of gene expression in combination with gene knock-down and protein overexpression.

**Signaling network** Ligand-activated transcription factors form a significant component of the defensome, integrating the stress response and potentially activating many different pathways simultaneously. The evolutionary history of both the bZIP (Amoutzias et al. 2007) and bHLH-PAS (Simionato et al. 2007) receptor superfamilies is complicated, but most major clades of these receptors are present in cnidaria.



**Fig. 4** Gene count comparisons for various classes of defensome genes. The area of each circle is proportional to the total number of genes classified into the defensome. Receptors include bHLH-ZIP, NR, and CNC receptors. Transporters are ABC and OAT transporters; oxidative and reductive modification genes include CYP, FMO, ALDH, EH, and AKR; conjugative genes are GST, MAPEG, UGT, and SULT; antioxidant genes are SOD, CAT, PXR, and GPX. Other genes include PCS, HSP20, HSP70, and HSP90

Although some of the known components of the vertebrate and invertebrate xenobiotic receptor pathways are missing, homologs of most important stress receptors are present, including AHR, ARNT, HIF1 $\alpha$ , MTF, and NRF2. The deuterostome xenobiotic-responsive NR1H and NR1I subfamilies are missing, however, and it is not currently known whether other NRs are functioning as xenobiotic receptors. Cnidaria have been shown to have a retinoid response, including a functional homolog of the RXR (Bouzaïene et al. 2007; Johnson and Chun 1989; Kostrouch et al. 1998; Muller 1984).

**Gene family diversification** Many defense gene families have undergone diversification and expansion in marine invertebrates in comparison to vertebrate genomes. In general, analysis of entire genomes is required to determine that a specific family or superfamily has undergone diversification, and thus current examples of such events are scattered. Class or order level diversification may be presumed, based on the model organisms with sequenced genomes, but caution should be exercised when extrapolating. Although there is general conformity in the presence of higher order gene groups, in many cases, gene orthology is more difficult to determine.

For example, the sea anemone contains 82 CYP genes and those related to CYP gene families 1–4 constitute a large proportion (76%) of the total, suggesting evolutionary pressure to maintain broad functionality in these important defense gene families. Multiple gene duplications in the toxicologically important CYP families appear to have taken place in many different lineages, leading to taxon-specific gene clades that are related to known CYP families yet distinct enough to preclude definitive assignment of names based on current CYP nomenclature guidelines. The extensive birth–death process of CYP diversification is not solely represented by invertebrates—within the vertebrates, there is significant evidence for extensive gene duplication and loss within xenobiotic-metabolizing CYP families (Thomas 2007).

In *N. vectensis*, particular examples of family diversification relative to known sequences are the CYPs and the ABC transporters. Other defensome gene families do not appear significantly expanded, nor do they have genes distributed into completely novel subfamilies. This observation is interesting in

light of the fact that *N. vectensis* lives in the challenging environment of a temperate estuary and ranges from subtropical to subarctic estuaries (Hand and Uhlinger 1994).

**Symbiosis** An important consideration for the study of cnidarian chemical defense genes is that many species are host to photosynthetic endosymbionts (zooxanthellae or zoochorellae). There are unique aspects of both normal physiology and toxicological responses that are related to the presence of endosymbionts. Notably, photosynthesis produces oxygen, and surrounding host tissues require additional protection against ROS to withstand hyperoxygenation, such as additional superoxide dismutase genes. Indeed, Allemand and coworkers have characterized multiple SOD forms in the Mediterranean sea anemone *Anemone viridis*. They found up to seven SOD activity bands in various tissues and detected several forms of CuZnSOD, MnSOD, and FeSOD in the various compartments (Richier et al. 2003). Two of the CuZnSODs were cloned and encode both extra- and intracellular CuZnSODs with different putative transcription binding sites (Plantivaux et al. 2004).

Although *N. vectensis* is an apparently asymbiotic anemone, the genome has genes for six different SODs, all of which are expressed under normal conditions. This abundance of SOD genes may be a general pattern, particularly in anthozoans. Interestingly, greater diversity in SOD activities was found in a symbiotic anemone species (*A. viridis*) than in an asymbiotic species (*Actinia schmidtii*), and the asymbiotic anemone experienced significantly greater oxidative protein damage upon exposure to hyperoxia (Richier et al. 2005). Thus, the presence of photosynthetic endosymbionts and the concomitant possibility of hyperoxia may have driven the evolution of multiple additional SOD forms in cnidaria. Several different catalase forms were also characterized in *A. viridis*, with tissue-specific distributions and activities (Merle et al. 2007). Inhibition of the host anemone catalase led to symbiont expulsion, suggesting an active response to increased oxidative stress.

A second effect of algal symbiosis is the sensitivity of symbiotic cnidarian species to herbicidal contamination (Jones 2005; Jones and Kerswell 2003). Notably, there is increasing distribution of herbicides such as the s-triazine Irgarol 1051 that have been

incorporated into marine antifouling paints along with copper (Carbery et al. 2006; Gardinali et al. 2004), and there is significant runoff of herbicide-containing waste from agricultural regions. Irgarol 1051 is a photosystem II binding agent that inhibits photosynthetic electron transport, resulting in a shortage of NADPH and the formation of singlet oxygen (Fufezan et al. 2002). Acute exposure of coral to Irgarol 1051 resulted in induction of SOD and MXR (ABCG) proteins and decreases of GPX, CAT, and certain CYP proteins (Downs and Downs 2007). While other herbicides have been investigated (e.g., diuron; Harrington et al. 2005; Negri et al. 2005; Raberg et al. 2003), there have been relatively few investigations of the molecular mechanisms of herbicide toxicity in cnidaria, and it is generally thought that damage is primarily a result of the disruption of host–algal symbiosis (Jones 2005).

**Reactive oxygen and UV** ROS production can also be an important consequence of UV exposure (Lesser 2006; Mopper and Kieber 2000). While UV responses have been studied in a number of coral species, many coral physiological responses to UV appear to be related to the physiological responses of their algal symbionts (Baruch et al. 2005; Torres et al. 2007; Verde and McCloskey 2002). UV has been shown to interfere with pattern formation in regenerating hydra and promote budding of intact hydra, possibly in response to tissue damage (Ghaskadbi et al. 2005; Znidaric et al. 1992). Both exogenous hydrogen peroxide and UV treatments have been shown to increase DNA strand breaks in cnidaria, demonstrating the potential for genotoxic ROS effects (Baruch et al. 2005; Mitchelmore and Hyatt 2004).

A very important protective mechanism in corals, as well as in diverse other marine organisms, is the accumulation of sunscreens compounds known as mycosporine-like amino acids (MAAs; Shick and Dunlap 2002). MAAs may facilitate larval survival (Wellington and Fitt 2003), as well as adult UV tolerance (Ferrier-Pages et al. 2007; Torres et al. 2007), and may also contribute to antioxidant capacity (Dunlap and Yamamoto 1995; Yakovleva et al. 2004). In contrast to other animals, *N. vectensis* apparently possesses the shikimic acid pathway thought to be necessary for MAA production (Starcevic et al. 2008), presumably obtained via lateral gene transfer from bacteria. Many cnidaria have been thought to

accumulate MAAs from their symbionts (Shick and Dunlap 2002) or from their diet, as is the case for sea urchins (Carroll and Shick 1996). Given the presence of the MAA biosynthetic pathway in the *N. vectensis* genome and the clustering of sea anemone MAA complement by anemone phylogenetic distribution rather than endosymbiont identity, presence, or other environmental factors (Shick et al. 2002), it is likely that the ability of cnidaria to biosynthesize MAAs is not restricted to *N. vectensis*.

In contrast to tropical corals and many littoral anemones, *N. vectensis* is a burrowing anemone, and it is possible that *N. vectensis* adult may be able to avoid UV damage despite living in shallow ponds. However, larval *N. vectensis* may require more protection from ROS than adults, leading to the apparent diversification of ROS defenses. Examination of the UV-stress response of *N. vectensis* will aid in understanding the different roles that antioxidant enzymes and suncreening compounds may play in protecting sea anemones. Comparisons of *N. vectensis*, an apparently asymbiotic anemone, with symbiotic anemones (e.g., *A. viridis*) or symbiotic reef-building corals could elucidate the protective mechanisms required by symbiotic cnidaria and shed light on the role of antioxidant enzymes in thermotolerance and bleaching (Downs et al. 2002; Merle et al. 2007; Richier et al. 2005; Richier et al. 2003).

**Molecular toxicology** Many toxicological studies of cnidaria involve metals, particularly copper, cadmium, and zinc. In particular, the acute and structural effects of copper, cadmium, and zinc have been investigated in various hydrozoan species, including both freshwater and marine hydra (Holdway et al. 2001; Karntanut and Pascoe 2000, 2002, 2005) and a variety of anthozoa, including scleractinian corals (Mitchelmore et al. 2007). Other biological responses to heavy metals in cnidaria include coral bleaching (Jones 1997) and effects on coral metabolism (Alutoin et al. 2001; Nystrom et al. 2001), larval mortality, and inhibition of reproduction, including settlement, motility, and fertilization of larvae (Negri and Heyward 2000, 2001; Reichelt-Brushett and Harrison 2000, 2005). Few studies have examined molecular biomarkers or molecular mechanisms of metal stress (Mitchelmore et al. 2002; Morgan et al. 2001).

The availability of the *N. vectensis* genome will make many mechanistic studies possible and should

spur the development of metal stress biomarkers in various species. In particular, the presence of multiple genes for phytochelatin synthase provides obvious markers for metal stress, despite the lack of metallothionein genes.

As with metal contamination, there are few studies of either molecular markers or mechanisms of exposure to organic contaminants other than herbicides in cnidaria (Rougee et al. 2006). A number of biochemical studies of cnidarian CYP biochemistry have been carried out, however. CYP carbon monoxide difference spectra have been observed in six different species of sea anemone (Heffernan and Winston 1998, 2000; Sole and Livingstone 2005) and three different species of scleractinian corals (*Favia fragum*, *Siderastrea sidea*, and *Montastraea faveolata*; Garcia et al. 2005; Gassman and Kennedy 1992; Ramos and Garcia 2007). Furthermore, benzo[*a*]pyrene hydroxylase activities have been observed in sea anemones, likely due to the action of CYP mixed-function oxygenases (Heffernan et al. 1996; Winston et al. 1998). The presence of inducible (versus constituent) CYP content in corals has also been demonstrated in coral due to benzo[*a*]pyrene or fuel oil exposure (Ramos and Garcia 2007; Rougee et al. 2006). The same PAH exposures induced components of the reactive oxygen defense systems, including CAT, SOD, and GST. Finally, an important molecular marker of genotoxic damage, the Comet assay of DNA damage, has been assessed in cnidaria in response to benzo[*a*]pyrene exposures (Mitchelmore and Hyatt 2004). Benzo[*a*]pyrene was found to increase DNA strand breaks in the temperate anemone *Anthopleura elegantissima*, suggesting also that cnidaria, like vertebrates, are capable of bioactivating benzo[*a*]pyrene to genotoxic metabolites, likely by CYPs.

More subtle effects of organic contamination might include in particular disruption of endogenous signaling pathways by exogenous hormones or hormone mimetics. Recent research indicates that cnidaria are susceptible to this sort of signal disruption, although the precise mechanisms are unknown (Fukuhori et al. 2005; Pachura-Bouchet et al. 2006; Pascoe et al. 2002; Tarrant 2005, 2007; Tarrant et al. 2004). In particular, Tarrant et al (2004) observed that spawning and growth rates were reduced in corals exposed to exogenous steroidal estrogens. As noted above, cnidaria do not possess a homolog of the vertebrate estrogen receptor, although there may be other nuclear

receptors that function as a steroid receptors (Reitzel et al. 2008b; Tarrant 2007). Cnidaria are steroid-rich organisms (Withers et al. 1982), but the roles these steroids play in normal physiology are not clear (Tarrant 2005). Steroids and secosteroids from gorgonian and soft corals have been show to have antimicrobial and antifouling activity (Qi et al. 2008; Sica and Musumeci 2004), suggesting that many of these compounds may be produced for chemical defense. CYP enzymes often participate in steroid synthesis and modification, and the diversity of *N. vectensis* CYPs may relate to the diversity of cnidarian steroids, although the steroid content of *N. vectensis* has not been investigated.

*N. vectensis* is a physical stress-tolerant organism, tolerating a wide range of environmental conditions (Sheader et al. 1997). With this robustness to physical stress, *N. vectensis* is in a prime position to act as a sentinel species toward chemical stress in estuaries. *N. vectensis* is also an excellent laboratory model, with simple maintenance needs, and the generation of clonal stocks by forced regeneration allows great scope for genetic manipulation. Furthermore, this sea anemone is an excellent model for the study of embryonic development (Matus et al. 2006). With the description of these chemical defense genes, we can study the evolution of cellular defense during embryonic development. Development inherently is a robust process; which parts of the process are more susceptible to disruption and from which stressors are not clear (Hamdoun and Epel 2007). The description of this defense gene set will allow us to examine the evolution of generalized cellular stress responses in bilaterian embryos and to understand how these stress responses function in adult cnidaria.

**Acknowledgments** Funding was provided by the WHOI Ocean Life Institute and NIH R01-ES015912. This manuscript benefited from discussions with Ann Tarrant, Adam Reitzel, and Cécile Sabourault and from the comments of several anonymous reviewers.

## References

- Allikmets R, Schriml LM, Hutchinson A, Romano-Spica V, Dean M. A human placenta-specific ATP-binding cassette gene (ABCP) on chromosome 4q22 that is involved in multidrug resistance. *Cancer Res* 1998;58(23):5337–9.
- Alutain S, Boberg J, Nystrom M, Tedengren M. Effects of the multiple stressors copper and reduced salinity on the metabolism of the hermatypic coral *Porites lutea*. *Mar Environ Res* 2001;52(3):289–99. doi:10.1016/S0141-1136(01)00105-2.
- Amiard JC, Amiard-Triquet C, Barka S, Pellerin J, Rainbow PS. Metallothioneins in aquatic invertebrates: their role in metal detoxification and their use as biomarkers. *Aquat Toxicol* 2006;76(2):160–202. doi:10.1016/j.aquatox.2005.08.015.
- Amoutzias GD, Veron AS, Weiner J 3rd, Robinson-Rechavi M, Bornberg-Bauer E, Oliver SG, et al. One billion years of bZIP transcription factor evolution: conservation and change in dimerization and DNA-binding site specificity. *Mol Biol Evol* 2007;24(3):827–35. doi:10.1093/molbev/msl211.
- Annilo T, Chen ZQ, Shulenin S, Costantino J, Thomas L, Lou H, et al. Evolution of the vertebrate ABC gene family: analysis of gene birth and death. *Genomics* 2006;88:1–11.
- Arkipchuk VV, Blaise C, Malinovskaya MV. Use of hydra for chronic toxicity assessment of waters intended for human consumption. *Environ Pollut* 2006;142(2):200–11. doi:10.1016/j.envpol.2005.10.012.
- Atkinson S, Atkinson MJ. Detection of estradiol-17-beta during a mass coral spawn. *Coral Reefs* 1992;11(1):33–5. doi:10.1007/BF00291932.
- Baruch R, Avishai N, Rabinowitz C. UV incites diverse levels of DNA breaks in different cellular compartments of a branching coral species. *J Exp Biol* 2005;208(5):843–8. doi:10.1242/jeb.01496.
- Berthet B, Mouneyrac C, Perez T, Amiard-Triquet C. Metallothionein concentration in sponges (*Spongia officinalis*) as a biomarker of metal contamination. *Comp Biochem Physiol C Toxicol Pharmacol* 2005;141(3):306–13. doi:10.1016/j.cca.2005.07.008.
- Blank V. Small Maf proteins in mammalian gene control: mere dimerization partners or dynamic transcriptional regulators? *J Mol Biol* 2008;376(4):913–25. doi:10.1016/j.jmb.2007.11.074.
- Blomquist CH, Lima PH, Tarrant AM, Atkinson MJ, Atkinson S. 17Beta-hydroxysteroid dehydrogenase (17beta-HSD) in scleractinian corals and zooxanthellae. *Comp Biochem Physiol B Biochem Mol Biol* 2006;143(4):397–403. doi:10.1016/j.cbpb.2005.12.017.
- Bock KW, Kohle C. Coordinate regulation of drug metabolism by xenobiotic nuclear receptors: UGTs acting together with CYPs and glucuronide transporters. *Drug Metab Rev* 2004;36(3–4):595–615. doi:10.1081/DMR-200033455.
- Bouzaïene M, Angers A, Anctil M. Immunohistochemical localization of a retinoic acid-like receptor in nerve cells of two colonial anthozoans (Cnidaria). *Tissue Cell* 2007;39(2):123–30. doi:10.1016/j.tice.2007.02.001.
- Bresell A, Weinander R, Lundqvist G, Raza H, Shimoji M, Sun TH, et al. Bioinformatic and enzymatic characterization of the MAPEG superfamily. *FEBS J* 2005;272(7):1688–703. doi:10.1111/j.1742-4658.2005.04596.x.
- Carbery K, Owen R, Frickers T, Otero E, Readman J. Contamination of Caribbean coastal waters by the anti-fouling herbicide Irgarol 1051. *Mar Pollut Bull* 2006;52(6):635–44. doi:10.1016/j.marpolbul.2005.10.013.
- Carroll AK, Shick JM. Dietary accumulation of UV-absorbing mycosporine-like amino acids (MAAs) by the green sea urchin (*Strongylocentrotus droebachiensis*). *Mar Biol (Berl)* 1996;124(4):561–9. doi:10.1007/BF00351037.

- Cashman JR. Some distinctions between flavin-containing and cytochrome P450 monooxygenases. *Biochem Biophys Res Commun* 2005;338(1):599–604. doi:10.1016/j.bbrc.2005.08.009.
- Castro LF, Santos MM, Reis-Henriques MA. The genomic environment around the Aromatase gene: evolutionary insights. *BMC Evol Biol* 2005;5:43. doi:10.1186/1471-2148-5-43.
- Choresch O, Azem A, Loya Y. Over-expression of highly conserved mitochondrial 70-kDa heat-shock protein in the sea anemone *Anemonia viridis*. *J Therm Biol* 2007;32(7–8):367–73. doi:10.1016/j.jtherbio.2007.04.006.
- Choresch O, Loya Y, Muller WE, Wiedenmann J, Azem A. The mitochondrial 60-kDa heat shock protein in marine invertebrates: biochemical purification and molecular characterization. *Cell Stress Chaperones* 2004;9(1):38–48.
- Choresch O, Ron E, Loya Y. The 60-kDa heat shock protein (HSP60) of the sea anemone *Anemonia viridis*: a potential early warning system for environmental changes. *Mar Biotechnol* 2001;3(5):501–8. doi:10.1007/s10126-001-0007-4.
- Clemens S. Evolution and function of phytochelatin synthases. *J Plant Physiol* 2006;163(3):319–32. doi:10.1016/j.jplph.2005.11.010.
- Clemens S, Schroeder JI, Degenkolb T. *Caenorhabditis elegans* expresses a functional phytochelatin synthase. *Eur J Biochem* 2001;268(13):3640–3. doi:10.1046/j.1432-1327.2001.02293.x.
- Cobbett C, Goldsbrough P. Phytochelatins and metallothioneins: roles in heavy metal detoxification and homeostasis. *Annu Rev Plant Biol* 2002;53:159–82. doi:10.1146/annurev.arplant.53.100301.135154.
- Daly M. A systematic revision of Edwardsiidae (Cnidaria, Anthozoa). *Invertebr Biol* 2002;121(3):212–25.
- Darling JA, Reitzel AR, Burton PM, Mazza ME, Ryan JF, Sullivan JC, et al. Rising starlet: the starlet sea anemone, *Nematostella vectensis*. *Bioessays* 2005;27(2):211–21. doi:10.1002/bies.20181.
- Davidson EH, Erwin DH. Gene regulatory networks and the evolution of animal body plans. *Science* 2006;311(5762):796–800. doi:10.1126/science.1113832.
- Davies WJ, Freeman SJ. The *Hydra attenuata* assay. *Methods Mol Biol* 1995;43:321–6. Clifton, NJ.
- de Jong DM, Hislop NR, Hayward DC, Reece-Hoyes JS, Pontynen PC, Ball EE, et al. Components of both major axial patterning systems of the Bilateria are differentially expressed along the primary axis of a ‘radiate’ animal, the anthozoan cnidarian *Acropora millepora*. *Dev Biol* 2006;298(2):632–43. doi:10.1016/j.ydbio.2006.07.034.
- Dean M, Annilo T. Evolution of the ATP-binding cassette (ABC) transporter superfamily in vertebrates. *Annu Rev Genomics Hum Genet* 2005;6:123–42. doi:10.1146/annurev.genom.6.080604.162122.
- Dean M, Hamon Y, Chimini G. The human ATP-binding cassette (ABC) transporter superfamily. *J Lipid Res* 2001;42(7):1007–17.
- Dean M, Fojo T, Bates S. Tumour stem cells and drug resistance. *Nat Rev Cancer* 2005;5(4):275–84. doi:10.1038/nrc1590.
- Deeley RG, Westlake C, Cole SP. Transmembrane transport of endo- and xenobiotics by mammalian ATP-binding cassette multidrug resistance proteins. *Physiol Rev* 2006;86(3):849–99. doi:10.1152/physrev.00035.2005.
- Downs C, Downs A. Preliminary examination of short-term cellular toxicological responses of the coral *Madracis mirabilis* to acute Irgarol 1051 exposure. *Arch Environ Contam Toxicol* 2007;52(1):47–57. doi:10.1007/s00244-005-0213-6.
- Downs CA, Fauth JE, Halas JC, Dustan P, Bemiss J, Woodley CM. Oxidative stress and seasonal coral bleaching. *Free Radic Biol Med* 2002;33(4):533–43. doi:10.1016/S0891-5849(02)00907-3.
- Dunlap WC, Yamamoto Y. Small-molecule antioxidants in marine organisms: antioxidant activity of mycosporine-glycine. *Comp Biochem Physiol B* 1995;112(1):105–14. doi:10.1016/0305-0491(95)00086-N.
- Dunn CW, Hejnal A, Matus DQ, Pang K, Browne WE, Smith SA, et al. Broad phylogenomic sampling improves resolution of the animal tree of life. *Nature* 2008;452(7188):745–9. doi:10.1038/nature06614.
- Eddy SR. Profile hidden Markov models. *Bioinformatics* 1998;14:755–63. doi:10.1093/bioinformatics/14.9.755.
- Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 2004;32(5):1792–7. doi:10.1093/nar/gkh340.
- Epel D. Protection of DNA during early development: adaptations and evolutionary consequences. *Evol Dev* 2003;5(1):83–8. doi:10.1046/j.1525-142X.2003.03013.x.
- Eraly SA, Monte JC, Nigam SK. Novel slc22 transporter homologs in fly, worm, and human clarify the phylogeny of organic anion and cation transporters. *Physiol Genomics* 2004;18(1):12–24. doi:10.1152/physiolgenomics.00014.2004.
- Feder ME, Hofmann GE. Heat-shock proteins, molecular chaperones, and the stress response: evolutionary and ecological physiology. *Annu Rev Physiol* 1999; 61: 243–82.
- Ferrier-Pages C, Richard C, Forcioli D, Allemand D, Pichon M, Shick JM. Effects of temperature and UV radiation increases on the photosynthetic efficiency in four scleractinian coral species. *Biol Bull* 2007;213(1):76–87.
- Finkel T, Holbrook NJ. Oxidants, oxidative stress and the biology of ageing. *Nature* 2000;408(6809):239–47. doi:10.1038/35041687.
- Finnerty JR, Martindale MQ. Ancient origins of axial patterning genes: Hox genes and ParaHox genes in the Cnidaria. *Evol Dev* 1999;1(1):16–23. doi:10.1046/j.1525-142x.1999.99010.x.
- Finnerty JR, Pang K, Burton P, Paulson D, Martindale MQ. Origins of bilateral symmetry: Hox and dpp expression in a sea anemone. *Science* 2004;304(5675):1335–7. doi:10.1126/science.1091946.
- Fritzenwanker JH, Saina M, Technau U. Analysis of forkhead and snail expression reveals epithelial–mesenchymal transitions during embryonic and larval development of *Nematostella vectensis*. *Dev Biol* 2004;275(2):389–402. doi:10.1016/j.ydbio.2004.08.014.
- Fufezan C, Rutherford AW, Krieger-Liszak A. Singlet oxygen production in herbicide-treated photosystem II. *FEBS Lett* 2002;532(3):407–10. doi:10.1016/S0014-5793(02)03724-9.
- Fukuhori N, Kitano M, Kimura H. Toxic effects of bisphenol A on sexual and asexual reproduction in *Hydra oligactis*. *Arch Environ Contam Toxicol* 2005;48(4):495–500. doi:10.1007/s00244-004-0032-1.
- Gamage N, Barnett A, Hempel N, Duggleby RG, Windmill KF, Martin JL, et al. Human sulfotransferases and their role in

- chemical metabolism. *Toxicol Sci* 2006;90(1):5–22. doi:10.1093/toxsci/kfj061.
- Garcia E, Ramos R, Bastidas C. Presence of cytochrome P450 in the Caribbean corals *Siderastrea siderea* and *Motastrea faveolata*. *Cinenc* 2005;31:1–7.
- Gardinali PR, Plasencia MD, Maxey C. Occurrence and transport of Irgarol 1051 and its major metabolite in coastal waters from South Florida. *Mar Pollut Bull* 2004;49(11–12):1072–83. doi:10.1016/j.marpolbul.2004.08.003.
- Gassman NJ, Kennedy CJ. Cytochrome P450 content and xenobiotic metabolizing enzyme activities in the scleractinian coral, *Favia fragum* (ESPER). *Bull Mar Sci* 1992;50:320–30.
- Ghaskadbi SS, Shetye L, Chiplonkar S, Ghaskadbi S. Ultraviolet irradiation initiates ectopic foot formation in regenerating hydra and promotes budding. *J Biosci* 2005;30(2):177–82. doi:10.1007/BF02703697.
- Goldstone J, Hamdoun AM, Cole B, Ashby MH, Scally M, Dean M, et al. The chemical defenseome: environmental sensing and response genes in the *Strongylocentrotus purpuratus* genome. *Dev Biol* 2006;300(1):366–84. doi:10.1016/j.ydbio.2006.08.066.
- Goldstone JV, Goldstone HM, Morrison AM, Tarrant A, Kern SE, Woodin BR, et al. Cytochrome P450 1 genes in early deuterostomes (tunicates and sea urchins) and vertebrates (chicken and frog): origin and diversification of the CYP1 gene family. *Mol Biol Evol* 2007;24(12):2619–31. doi:10.1093/molbev/msm200.
- Grasso LC, Hayward DC, Trueman JW, Hardie KM, Janssens PA, Ball EE. The evolution of nuclear receptors: evidence from the coral *Acropora*. *Mol Phylogenet Evol* 2001;21(1):93–102. doi:10.1006/mpev.2001.0994.
- Guengerich FP, McCormick WA, Wheeler JB. Analysis of the kinetic mechanism of haloalkane conjugation by mammalian theta-class glutathione transferases. *Chem Res Toxicol* 2003;16(11):1493–9. doi:10.1021/tx034157r.
- Hagenbuch B, Meier PJ. The superfamily of organic anion transporting polypeptides. *Biochim Biophys Acta* 2003;1609(1):1–18. doi:10.1016/S0005-2736(02)00633-8.
- Halliwell B, Gutteridge JMC. *Free radicals in biology and medicine*. 3rd ed. Oxford: Oxford University Press; 1999.
- Hamdoun A, Epel D. Embryo stability and vulnerability in an always changing world. *Proc Natl Acad Sci U S A* 2007;104(6):1745–50. doi:10.1073/pnas.0610108104.
- Hand C, Uhlinger K. The unique, widely distributed sea anemone, *Nematostella vectensis* Stephenson: a review, new facts, and questions. *Estuaries* 1994;17:501–8. doi:10.2307/1352679.
- Harborne AR, Mumby PJ, Micheli F, Perry CT, Dahlgren CP, Holmes KE, et al. The functional value of Caribbean coral reef, seagrass and mangrove habitats to ecosystem processes. *Adv Mar Biol* 2006;50:57–189. doi:10.1016/S0065-2881(05)50002-6.
- Harrington L, Fabricius K, Eaglesham G, Negri A. Synergistic effects of diuron and sedimentation on photosynthesis and survival of crustose coralline algae. *Mar Pollut Bull* 2005;51(1–4):415–27. doi:10.1016/j.marpolbul.2004.10.042.
- Hashimoto K, Shibuno T, Murayama-Kayano E, Tanaka H, Kayano T. Isolation and characterization of stress-responsive genes from the scleractinian coral *Pocillopora damicornis*. *Coral Reefs* 2004;23(4):485–91.
- Hayes JD, Flanagan JU, Jowsey IR. Glutathione transferases. *Annu Rev Pharmacol Toxicol* 2005;45:51–88. doi:10.1146/annurev.pharmtox.45.120403.095857.
- Hecker M, Ullrich V. On the mechanism of prostacyclin and thromboxane A2 biosynthesis. *J Biol Chem* 1989;264(1):141–50.
- Heffernan LM, Winston GW. Spectral analysis and catalytic activities of the microsomal mixed-function oxidase system of the sea anemone (phylum: Cnidaria). *Comp Biochem Physiol* 1998;121(1–3):371–83.
- Heffernan LM, Winston GW. Distribution of microsomal CO-binding chromophores and EROD activity in sea anemone tissues. *Mar Environ Res* 2000;50(1–5):23–7.
- Heffernan LM, Zinn RR, Mayeaux MH, Winston GW. Benzo[a]pyrene metabolism in the sea anemone *Bunodosoma cavernata*. *FASEB J* 1996;10(6):2868.
- Holdway DA, Lok K, Semaan M. The acute and chronic toxicity of cadmium and zinc to two hydra species. *Environ Toxicol* 2001;16(6):557–65. doi:10.1002/tox.10017.
- Hyndman D, Bauman DR, Heredia VV, Penning TM. The aldo-keto reductase superfamily homepage. *Chem Biol Interact* 2003;143–144:621–31. doi:10.1016/S0009-2797(02)00193-X.
- Igarashi K, Sun J. The heme-Bach1 pathway in the regulation of oxidative stress response and erythroid differentiation. *Antioxid Redox Signal* 2006;8(1–2):107–18. doi:10.1089/ars.2006.8.107.
- Jacobsson JA, Haitina T, Lindblom J, Fredriksson R. Identification of six putative human transporters with structural similarity to the drug transporter SLC22 family. *Genomics* 2007;90(5):595–609. doi:10.1016/j.ygeno.2007.03.017.
- Jacobsson PJ, Thoren S, Morgenstern R, Samuelsson B. Identification of human prostaglandin E synthase: a microsomal, glutathione-dependent, inducible enzyme, constituting a potential novel drug target. *Proc Natl Acad Sci U S A* 1999;96(13):7220–5. doi:10.1073/pnas.96.13.7220.
- Jez JM, Penning TM. The aldo-keto reductase (AKR) superfamily: an update. *Chem Biol Interact* 2001;130–132(1–3):499–525. doi:10.1016/S0009-2797(00)00295-7.
- Jin Y, Penning TM. Aldo-keto reductases and bioactivation/detoxication. *Annu Rev Pharmacol Toxicol* 2007;47:263–92. doi:10.1146/annurev.pharmtox.47.120505.105337.
- Johnson EM, Chun YH. In vitro differential developmental toxicity of vitamin A congeners. *Teratology* 1989;39(4):349–61. doi:10.1002/tera.1420390407.
- Jones R. The ecotoxicological effects of Photosystem II herbicides on corals. *Mar Pollut Bull* 2005;51(5–7):495–506. doi:10.1016/j.marpolbul.2005.06.027.
- Jones RJ. Zooxanthellae loss as a bioassay for assessing stress in corals. *Mar Ecol Prog Ser* 1997;149(1–3):163–71. doi:10.3354/meps149163.
- Jones RJ, Kerswell AP. Phytotoxicity of photosystem II (PSII) herbicides to coral. *Mar Ecol Prog Ser* 2003;261:149–59. doi:10.3354/meps261149.
- Karntanut W, Pascoe D. A comparison of methods for measuring acute toxicity to *Hydra vulgaris*. *Chemosphere* 2000;41(10):1543–8. doi:10.1016/S0045-6535(00)00068-0.
- Karntanut W, Pascoe D. The toxicity of copper, cadmium and zinc to four different *Hydra* (Cnidaria: Hydrozoa). *Chemosphere* 2002;47(10):1059–64. doi:10.1016/S0045-6535(02)00050-4.
- Karntanut W, Pascoe D. Effects of removing symbiotic green algae on the response of *Hydra viridissima* (Pallas 1776)



- to metals. *Ecotoxicol Environ Saf* 2005;60(3):301–5. doi:10.1016/j.ecoenv.2004.04.001.
- Kewley RJ, Whitelaw ML, Chapman-Smith A. The mammalian basic helix-loop-helix/PAS family of transcriptional regulators. *Int J Biochem Cell Biol* 2004;36(2):189–204. doi:10.1016/S1357-2725(03)00211-5.
- Kim HP, Morse D, Choi AM. Heat-shock proteins: new keys to the development of cytoprotective therapies. *Expert Opin Ther Targets* 2006;10(5):759–69. doi:10.1517/14728222.10.5.759.
- Kimura H, Takeda M, Narikawa S, Enomoto A, Ichida K, Endou H. Human organic anion transporters and human organic cation transporters mediate renal transport of prostaglandins. *J Pharmacol Exp Ther* 2002;301(1):293–8. doi:10.1124/jpet.301.1.293.
- Koljak R, Jarving I, Kurg R, Boeglin WE, Varvas K, Valmsen K, et al. The basis of prostaglandin synthesis in coral: molecular cloning and expression of a cyclooxygenase from the Arctic soft coral *Gersemia fruticosa*. *J Biol Chem* 2001;276(10):7033–40. doi:10.1074/jbc.M009803200.
- Kostrouch Z, Kostrouchova M, Love W, Jannini E, Piatigorsky J, Rall JE. Retinoic acid X receptor in the diploblast, *Tripedalia cystophora*. *Proc Natl Acad Sci U S A* 1998;95(23):13442–7. doi:10.1073/pnas.95.23.13442.
- Kreil DP, Ouzounis CA. Comparison of sequence masking algorithms and the detection of biased protein sequence regions. *Bioinformatics* 2003;19(13):1672–81. doi:10.1093/bioinformatics/btg212.
- Krueger SK, Williams DE. Mammalian flavin-containing monooxygenases: structure/function, genetic polymorphisms and role in drug metabolism. *Pharmacol Ther* 2005;106(3):357–87. doi:10.1016/j.pharmthera.2005.01.001.
- Kusserow A, Pang K, Sturm C, Hrouda M, Lentfer J, Schmidt HA, et al. Unexpected complexity of the Wnt gene family in a sea anemone. *Nature* 2005;433(7022):156–60. doi:10.1038/nature03158.
- Kusuhara H, Sugiyama Y. ATP-binding cassette, subfamily G (ABCG family). *Pflugers Arch* 2007;453(5):735–44. doi:10.1007/s00424-006-0134-x.
- Laity JH, Andrews GK. Understanding the mechanisms of zinc-sensing by metal-response element binding transcription factor-1 (MTF-1). *Arch Biochem Biophys* 2007;463(2):201–10. doi:10.1016/j.abb.2007.03.019.
- Lee MO, Manthey CL, Sladek NE. Identification of mouse liver aldehyde dehydrogenases that catalyze the oxidation of retinaldehyde to retinoic acid. *Biochem Pharmacol* 1991;42(6):1279–85. doi:10.1016/0006-2952(91)90266-8.
- Leslie EM, Deeley RG, Cole SP. Toxicological relevance of the multidrug resistance protein 1, MRP1 (ABCC1) and related transporters. *Toxicology* 2001;167(1):3–23. doi:10.1016/S0300-483X(01)00454-1.
- Lesser MP. Oxidative stress in marine environments: biochemistry and physiological ecology. *Annu Rev Physiol* 2006;68:253–78. doi:10.1146/annurev.physiol.68.040104.110001.
- Lewis DF, Lake BG, Dickins M. Substrates of human cytochromes P450 from families CYP1 and CYP2: analysis of enzyme selectivity and metabolism. *Drug Metabol Drug Interact* 2004;20(3):111–42.
- Mackenzie PI, Walter Bock K, Burchell B, Guillemette C, Ikushiro S, Iyanagi T, et al. Nomenclature update for the mammalian UDP glycosyltransferase (UGT) gene superfamily. *Pharmacogenet Genomics* 2005;15(10):677–85. doi:10.1097/01.fpc.0000173483.13689.56.
- Magie CR, Pang K, Martindale MQ. Genomic inventory and expression of Sox and Fox genes in the cnidarian *Nematostella vectensis*. *Dev Genes Evol* 2005;215(12):618–30. doi:10.1007/s00427-005-0022-y.
- Matus DQ, Pang K, Marlow H, Dunn CW, Thomsen GH, Martindale MQ. Molecular evidence for deep evolutionary roots of bilaterality in animal development. *Proc Natl Acad Sci U S A* 2006;103(30):11195–200. doi:10.1073/pnas.0601257103.
- Merle PL, Sabourault C, Richier S, Allemand D, Furla P. Catalase characterization and implication in bleaching of a symbiotic sea anemone. *Free Radic Biol Med* 2007;42(2):236–46. doi:10.1016/j.freeradbiomed.2006.10.038.
- Miller DJ, Hemmrich G, Ball EE, Hayward DC, Khalturin K, Funayama N, et al. The innate immune repertoire in cnidaria—ancestral complexity and stochastic gene loss. *Genome Biol* 2007;8(4):R59. doi:10.1186/gb-2007-8-4-r59.
- Mitchelmore CL, Hyatt S. Assessing DNA damage in cnidarians using the Comet assay. *Mar Environ Res* 2004;58(2–5):707–11. doi:10.1016/j.marenvres.2004.03.019.
- Mitchelmore CL, Schwarz JA, Weis VM. Development of symbiosis-specific genes as biomarkers for the early detection of cnidarian-algal symbiosis breakdown. *Mar Environ Res* 2002;54(3–5):345–9. doi:10.1016/S0141-1136(02)00201-5.
- Mitchelmore CL, Verde EA, Weis VM. Uptake and partitioning of copper and cadmium in the coral *Pocillopora damicornis*. *Aquat Toxicol* 2007;85(1):48–56. doi:10.1016/j.aquatox.2007.07.015.
- Miyake K, Mickley L, Litman T, Zhan Z, Robey R, Cristensen B, et al. Molecular cloning of cDNAs which are highly overexpressed in mitoxantrone-resistant cells: demonstration of homology to ABC transport genes. *Cancer Res* 1999;59(1):8–13.
- Mizuta T, Kubokawa K. Presence of sex steroids and cytochrome P450 genes in amphioxus. *Endocrinology* 2007;148(8):3554–65. doi:10.1210/en.2007-0109.
- Mopper K, Kieber DJ. Marine photochemistry and its impacts on carbon cycling. In: De Mora S, Demers S, Vernet M, editors. *The effects of UV radiation in the marine environment*. Cambridge, UK: Cambridge University Press; 2000. p. 101–30.
- Morgan MB, Vogelien DL, Snell TW. Assessing coral stress responses using molecular biomarkers of gene transcription. *Environ Toxicol Chem* 2001;20(3):537–43. doi:10.1897/1551-5028(2001)020<0537:ACSRUM>2.0.CO;2.
- Muller WA. Retinoids and pattern formation in a hydroid. *J Embryol Exp Morphol* 1984;81:253–71.
- Murphy BJ. Regulation of malignant progression by the hypoxia-sensitive transcription factors HIF-1 $\alpha$  and MTF-1. *Comp Biochem Physiol B Biochem Mol Biol* 2004;139(3):495–507. doi:10.1016/j.cbpc.2004.04.009.
- Murphy BJ, Andrews GK, Bittel D, Discher DJ, McCue J, Green CJ, et al. Activation of metallothionein gene expression by hypoxia involves metal response elements and metal transcription factor-1. *Cancer Res* 1999;59(6):1315–22.

- Murphy BJ, Sato BG, Dalton TP, Laderoute KR. The metal-responsive transcription factor-1 contributes to HIF-1 activation during hypoxic stress. *Biochem Biophys Res Commun* 2005;337(3):860–7. doi:10.1016/j.bbrc.2005.09.124.
- Nebert DW, Vasiliou V. Analysis of the glutathione S-transferase (GST) gene family. *Hum Genomics* 2004;1(6):460–4.
- Negri A, Vollhardt C, Humphrey C, Heyward A, Jones R, Eaglesham G, et al. Effects of the herbicide diuron on the early life history stages of coral. *Mar Pollut Bull* 2005;51(1–4):370–83. doi:10.1016/j.marpolbul.2004.10.053.
- Negri AP, Heyward AJ. Inhibition of fertilization and larval metamorphosis of the coral *Acropora millepora* (Ehrenberg, 1834) by petroleum products. *Mar Pollut Bull* 2000;41(7–12):420–7.
- Negri AP, Heyward AJ. Inhibition of coral fertilisation and larval metamorphosis by tributyltin and copper. *Mar Environ Res* 2001;51(1):17–27. doi:10.1016/S0141-1136(00)00029-5.
- Nelson DR. Metazoan cytochrome P450 evolution. *Comp Biochem Physiol* 1998;121(1–3):15–22.
- Nelson DR. Cytochrome P450 nomenclature, 2004. *Methods Mol Biol* 2006;320:1–10. Clifton, NJ.
- Nelson DR, Kamataki T, Waxman DJ, Guengerich FP, Estabrook RW, Feyereisen R, et al. The P450 superfamily: update on new sequences, gene mapping, accession numbers, early trivial names of enzymes, and nomenclature. *DNA Cell Biol* 1993;12(1):1–51.
- Nemer M, Wilkinson DG, Travaglini EC, Sternberg EJ, Butt TR. Sea urchin metallothionein sequence: key to an evolutionary diversity. *Proc Natl Acad Sci U S A* 1985;82(15):4992–4. doi:10.1073/pnas.82.15.4992.
- Nguyen T, Sherratt PJ, Huang HC, Yang CS, Pickett CB. Increased protein stability as a mechanism that enhances Nrf2-mediated transcriptional activation of the antioxidant response element. Degradation of Nrf2 by the 26 S proteasome. *J Biol Chem* 2003;278(7):4536–41. doi:10.1074/jbc.M207293200.
- Nystrom M, Nordemar I, Tedengren M. Simultaneous and sequential stress from increased temperature and copper on the metabolism of the hermatypic coral *Porites cylindrica*. *Mar Biol (Berl)* 2001;138(6):1225–31. doi:10.1007/s002270100549.
- Osburn WO, Kensler TW. Nrf2 signaling: an adaptive response pathway for protection against environmental toxic insults. *Mutat Res* 2008;659(1–2):31–9. doi:10.1016/j.mrrev.2007.11.006.
- Pachura-Bouchet S, Blaise C, Vasseur P. Toxicity of non-ylphenol on the cnidarian *Hydra attenuata* and environmental risk assessment. *Environ Toxicol* 2006;21(4):388–94. doi:10.1002/tox.20201.
- Palackal NT, Burczynski ME, Harvey RG, Penning TM. Metabolic activation of polycyclic aromatic hydrocarbon trans-dihydrodiols by ubiquitously expressed aldehyde reductase (AKR1A1). *Chem Biol Interact* 2001;130–132(1–3):815–24. doi:10.1016/S0009-2797(00)00237-4.
- Pascoe D, Carroll K, Karntanut W, Watts MM. Toxicity of 17 $\alpha$ -ethinylestradiol and bisphenol A to the freshwater Cnidarian *Hydra vulgaris*. *Arch Environ Contam Toxicol* 2002;43(1):56–63. doi:10.1007/s00244-001-0016-3.
- Paul VJ, Puglisi MP. Chemical mediation of interactions among marine organisms. *Nat Prod Rep* 2004;21(1):189–209. doi:10.1039/b302334f.
- Paul VJ, Puglisi MP, Ritson-Williams R. Marine chemical ecology. *Nat Prod Rep* 2006;23(2):153–80. doi:10.1039/b404735b.
- Penning TM, Burczynski ME, Hung CF, McCoull KD, Palackal NT, Tsuruda LS. Dihydrodiol dehydrogenases and polycyclic aromatic hydrocarbon activation: generation of reactive and redox active o-quinones. *Chem Res Toxicol* 1999;12(1):1–18. doi:10.1021/tx980143n.
- Penning TM, Drury JE. Human aldo-keto reductases: function, gene regulation, and single nucleotide polymorphisms. *Arch Biochem Biophys* 2007;464(2):241–50. doi:10.1016/j.abb.2007.04.024.
- Philp RB. Cadmium content of the marine sponge *Microciona prolifera*, other sponges, water and sediment from the eastern Florida panhandle: possible effects on *Microciona* cell aggregation and potential roles of low pH and low salinity. *Comp Biochem Physiol* 1999;124(1):41–9. doi:10.1016/S0305-0491(99)00095-4.
- Plantivaux A, Furla P, Zoccola D, Garelo G, Forcioli D, Richier S, et al. Molecular characterization of two CuZn-superoxide dismutases in a sea anemone. *Free Radic Biol Med* 2004;37(8):1170–81. doi:10.1016/j.freeradbio.2004.06.043.
- Ponting CP. The functional repertoires of metazoan genomes. *Nat Rev* 2008;9(9):689–98.
- Putnam NH, Srivastava M, Hellsten U, Dirks B, Chapman J, Salamov A, et al. Sea anemone genome reveals ancestral eumetazoan gene repertoire and genomic organization. *Science* 2007;317(5834):86–94. doi:10.1126/science.1139158.
- Qi SH, Zhang S, Yang LH, Qian PY. Antifouling and antibacterial compounds from the gorgonians *Subergorgia suberosa* and *Scripearia gracillis*. *Nat Prod Res* 2008;22(2):154–66. doi:10.1080/14786410701642441.
- Raberg S, Nystrom M, Eros M, Plantman P. Impact of the herbicides 2,4-D and diuron on the metabolism of the coral *Porites cylindrica*. *Mar Environ Res* 2003;56(4):503–14. doi:10.1016/S0141-1136(03)00039-4.
- Ramos R, Garcia E. Induction of mixed-function oxygenase system and antioxidant enzymes in the coral *Montastraea faveolata* on acute exposure to benzo(a)pyrene. *Comp Biochem Physiol C Toxicol Pharmacol* 2007;144(4):348–55. doi:10.1016/j.cbpc.2006.11.006.
- Reichelt-Brushett AJ, Harrison PL. The effect of copper on the settlement success of larvae from the scleractinian coral *Acropora tenuis*. *Mar Pollut Bull* 2000;41(7–12):385–91.
- Reichelt-Brushett AJ, Harrison PL. The effect of selected trace metals on the fertilization success of several scleractinian coral species. *Coral Reefs* 2005;24(4):524–34. doi:10.1007/s00338-005-0013-5.
- Reitzel AM, Darling JA, Sullivan JC, Finnerty JR. Global population genetic structure of the starlet anemone *Nematostella vectensis*: multiple introductions and implications for conservation policy. *Biol Invasions*. 2008a (in press).
- Reitzel AM, Sullivan JC, Traylor-Knowles N, Finnerty JR. Genomic survey of candidate stress-response genes in the estuarine anemone *Nematostella vectensis*. *Biol Bull* 2008b;214(3):233–54.
- Richier S, Furla P, Plantivaux A, Merle PL, Allemand D. Symbiosis-induced adaptation to oxidative stress. *J Exp Biol* 2005;208(Pt 2):277–85. doi:10.1242/jeb.01368.
- Richier S, Merle PL, Furla P, Pigozzi D, Sola F, Allemand D. Characterization of superoxide dismutases in anoxia- and

- hyperoxia-tolerant symbiotic cnidarians. *Biochim Biophys Acta* 2003;1621(1):84–91.
- Robbart ML, Peckol P, Scordilis SP, Curran HA, Brown-Saracino J. Population recovery and differential heat shock protein expression for the corals *Agaricia agaricites* and *A tenuifolia* in Belize. *Mar Ecol Prog Ser* 2004;283:151–60. doi:10.3354/meps283151.
- Rossi S, Snyder MJ. Competition for space among sessile marine invertebrates: changes in HSP70 expression in two Pacific cnidarians. *Biol Bull* 2001;201(3):385–93. doi:10.2307/1543616.
- Rossi S, Snyder MJ, Gili JM. Protein, carbohydrate, lipid concentrations and HSP70-HSP 90 (stress protein) expression over an annual cycle: useful tools to detect feeding constraints in a benthic suspension feeder. *Helgol Mar Res* 2006;60(1):7–17. doi:10.1007/s10152-005-0009-0.
- Rougee L, Downs CA, Richmond RH, Ostrander GK. Alteration of normal cellular profiles in the scleractinian coral (*Pocillopora damicornis*) following laboratory exposure to fuel oil. *Environ Toxicol Chem* 2006;25(12):3181–7. doi:10.1897/05-510R2.1.
- Runge-Morris M, Kocarek TA. Regulation of sulfotransferases by xenobiotic receptors. *Curr Drug Metab* 2005;6(4):299–307. doi:10.2174/1389200054633871.
- Sebens KP. Recruitment in a sea anemone population: juvenile substrate becomes adult prey. *Science* 1981;213(4509):785–7. doi:10.1126/science.213.4509.785.
- Sharp VA, Brown BE, Miller D. Heat shock protein (hsp 70) expression in the tropical reef coral *Goniopora djiboutiensis*. *J Therm Biol* 1997;22(1):11–9. doi:10.1016/S0306-4565(96)00029-0.
- Sharp VA, Miller D, Bythell JC, Brown BE. Expression of low-molecular-weight Hsp-70 related polypeptides from the symbiotic sea-anemone *Anemonia-viridis* Forskall in response to heat-shock. *J Exp Mar Biol Ecol* 1994;179(2):179–93. doi:10.1016/0022-0981(94)90113-9.
- Shearer M, Suwailem A, Rowe G. The anemone, *Nematostella vectensis*, in Britain: considerations for conservation management. *Aquat Conserv: Mar Freshwat Ecosyst* 1997;7:13–25. doi:10.1002/(SICI)1099-0755(199703)7:1<13::AID-AQC210>3.0.CO;2-Y.
- Shick JM, Dunlap WC. Mycosporine-like amino acids and related Gadusols: biosynthesis, accumulation, and UV-protective functions in aquatic organisms. *Annu Rev Physiol* 2002;64:223–62. doi:10.1146/annurev.physiol.64.081501.155802.
- Shick JM, Dunlap WC, Pearse JS, Pearse VB. Mycosporine-like amino acid content in four species of sea anemones in the genus *Anthopleura* reflects phylogenetic but not environmental or symbiotic relationships. *Biol Bull* 2002;203(3):315–30. doi:10.2307/1543574.
- Shimada T. Xenobiotic-metabolizing enzymes involved in activation and detoxification of carcinogenic polycyclic aromatic hydrocarbons. *Drug Metab Pharmacokinet* 2006;21(4):257–76. doi:10.2133/dmpk.21.257.
- Sica D, Musumeci D. Secosteroids of marine origin. *Steroids* 2004;69(11–12):743–56. doi:10.1016/j.steroids.2004.09.001.
- Simionato E, Ledent V, Richards G, Thomas-Chollier M, Kerner P, Coornaert D, et al. Origin and diversification of the basic helix-loop-helix gene family in metazoans: insights from comparative genomics. *BMC Evol Biol* 2007;7:33. doi:10.1186/1471-2148-7-33.
- Snyder MJ, Ross S. Stress protein (HSP70 family) expression in intertidal benthic organisms: the example of *Anthopleura elegantissima* (Cnidaria: Anthozoa). *Sci Mar* 2004;68:155–62.
- Sole M, Livingstone DR. Components of the cytochrome P450-dependent monooxygenase system and ‘NADPH-independent benzo[a]pyrene hydroxylase’ activity in a wide range of marine invertebrate species. *Comp Biochem Physiol C Toxicol Pharmacol* 2005;141(1):20–31. doi:10.1016/j.cca.2005.04.008.
- Stamatidakis A. RAXML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 2006;22(21):2688–90. doi:10.1093/bioinformatics/btl446.
- Starcevic A, Akthar S, Dunlap WC, Shick JM, Hranueli D, Cullum J, et al. Enzymes of the shikimic acid pathway encoded in the genome of a basal metazoan, *Nematostella vectensis*, have microbial origins. *Proc Natl Acad Sci U S A* 2008;105:2533–37.
- Sullivan JC, Reitzel AM, Finnerty JR. Upgrades to StellaBase facilitate medical and genetic studies on the starlet sea anemone, *Nematostella vectensis*. *Nucleic Acids Res* 2008;36:D607–11, (Database issue) doi:10.1093/nar/gkm941.
- Surh YJ. Bioactivation of benzylic and allylic alcohols via sulfo-conjugation. *Chem Biol Interact* 1998;109(1–3):221–35. doi:10.1016/S0009-2797(97)00134-8.
- Sweet DH. Organic anion transporter (Slc22a) family members as mediators of toxicity. *Toxicol Appl Pharmacol* 2005;204(3):198–215. doi:10.1016/j.taap.2004.10.016.
- Tarrant AM. Endocrine-like signaling in cnidarians: current understanding and implications for ecophysiology. *Integr Comp Biol* 2005;45:201–14. doi:10.1093/icb/45.1.201.
- Tarrant AM. Hormonal signaling in cnidarians: do we understand the pathways well enough to know whether they are being disrupted? *Ecotoxicology* 2007;16(1):5–13. doi:10.1007/s10646-006-0121-1.
- Tarrant AM, Atkinson MJ, Atkinson S. Effects of steroidal estrogens on coral growth and reproduction. *Mar Ecol Prog Ser* 2004;269:121–9. doi:10.3354/meps269121.
- Tarrant AM, Atkinson S, Atkinson MJ. Estrone and estradiol-17 beta concentration in tissue of the scleractinian coral, *Montipora verrucosa*. *Comp Biochem Physiol A Mol Integr Physiol* 1999;122(1):85–92. doi:10.1016/S1095-6433(98)10155-1.
- Tarrant AM, Blomquist CH, Lima PH, Atkinson MJ, Atkinson S. Metabolism of estrogens and androgens by scleractinian corals. *Comp Biochem Physiol B Biochem Mol Biol* 2003;136(3):473–85. doi:10.1016/S1096-4959(03)00253-7.
- Thomas JH. Rapid birth–death evolution specific to xenobiotic cytochrome P450 genes in vertebrates. *PLoS Genet* 2007;3(5):e67. doi:10.1371/journal.pgen.0030067.
- Thornton JW, Need E, Crews D. Resurrecting the ancestral steroid receptor: ancient origin of estrogen signaling. *Science* 2003;301(5640):1714–7. doi:10.1126/science.1086185.
- Torrás R, Gonzalez-Crespo S. Posterior expression of nanos orthologs during embryonic and larval development of the

- anthozoan *Nematostella vectensis*. *Int J Dev Biol* 2005;49(7):895–9. doi:10.1387/ijdb.051980rt.
- Torres JL, Armstrong RA, Corredor JE, Gillbes F. Physiological responses of *Acropora cervicornis* to increased solar irradiance. *Photochem Photobiol* 2007;83(4):839–50.
- Twan WH, Hwang JS, Chang CF. Sex steroids in scleractinian coral, *Euphyllia ancora*: implication in mass spawning. *Biol Reprod* 2003;68(6):2255–60. doi:10.1095/biolreprod.102.012450.
- Twan WH, Hwang JS, Lee YH, Wu HF, Tung YH, Chang CF. Hormones and reproduction in scleractinian corals. *Comp Biochem Physiol A Mol Integr Physiol* 2006;144(3):247–53. doi:10.1016/j.cbpa.2006.01.011.
- Uno S, Dalton TP, Shertzer HG, Genter MB, Warshawsky D, Talaska G, et al. Benzo[a]pyrene-induced toxicity: paradoxical protection in *Cyp1a1*( $-/-$ ) knockout mice having increased hepatic BaP-DNA adduct levels. *Biochem Biophys Res Commun* 2001;289(5):1049–56. doi:10.1006/bbrc.2001.6110.
- Vasiliou V, Nebert DW. Analysis and update of the human aldehyde dehydrogenase (ALDH) gene family. *Hum Genomics* 2005;2(2):138–43.
- Vasiliou V, Ross D, Nebert DW. Update of the NAD(P)H: Quinone oxidoreductase (NQO) gene family. *Hum Genomics* 2006;2(5):329–35.
- Verde EA, McCloskey LR. A comparative analysis of the photobiology of zooxanthellae and zoochlorellae symbiotic with the temperate clonal anemone *Anthopleura elegantissima* (Brandt)—II. Effect of light intensity. *Mar Biol (Berl)* 2002;141(2):225–39. doi:10.1007/s00227-002-0824-7.
- Wellington GM, Fitt WK. Influence of UV radiation on the survival of larvae from broadcast-spawning reef corals. *Mar Biol (Berl)* 2003;143(6):1185–92. doi:10.1007/s00227-003-1150-4.
- Whelan S, Goldman N. A general empirical model of protein evolution derived from multiple protein families using a maximum-likelihood approach. *Mol Biol Evol* 2001;18(5):691–9.
- Wiens M, Ammar MS, Nawar AH, Koziol C, Hassanein HMA, Eisinger M, et al. Induction of heat-shock (stress) protein gene expression by selected natural and anthropogenic disturbances in the octocoral *Dendronephthya klunzingeri*. *J Exp Mar Biol Ecol* 2000;245(2):265–76. doi:10.1016/S0022-0981(99)00167-7.
- Wiger R, Stottum A. In vitro testing for developmental toxicity using the *Hydra attenuata* assay. *NIPH Ann* 1985;8(2):43–7.
- Winston GW, Mayeaux MH, Heffernan LM. Benzo[a]pyrene metabolism by the intertidal sea anemone, *Bunodosoma cavernata*. *Mar Environ Res* 1998;45(1):89–100. doi:10.1016/S0141-1136(97)00026-3.
- Withers NW, Kokke WC, Fenical W, Djerassi C. Sterol patterns of cultured zooxanthellae isolated from marine invertebrates: synthesis of gorgosterol and 23-desmethylgorgosterol by aposymbiotic algae. *Proc Natl Acad Sci U S A* 1982;79(12):3764–8. doi:10.1073/pnas.79.12.3764.
- Yakovleva I, Bhagooli R, Takemura A, Hidaka M. Differential susceptibility to oxidative stress of two scleractinian corals: antioxidant functioning of mycosporine-glycine. *Comp Biochem Physiol B Biochem Mol Biol* 2004;139(4):721–30. doi:10.1016/j.cbpc.2004.08.016.
- Yamada S, Morimoto H, Fujisawa T, Sugahara K. Glycosaminoglycans in *Hydra magnipapillata* (Hydrozoa, Cnidaria): demonstration of chondroitin in the developing nematocyst, the sting organelle, and structural characterization of glycosaminoglycans. *Glycobiology* 2007;17(8):886–94. doi:10.1093/glycob/cwm051.
- Ziegler DM. An overview of the mechanism, substrate specificities, and structure of FMOs. *Drug Metab Rev* 2002;34(3):503–11. doi:10.1081/DMR-120005650.
- Znidaric D, Liu A, Kalafatic M. Regeneration and reproduction of irradiated green hydra (Cnidaria). *Period Biol* 1992;94(2):99–104.