

Erratum to: Molecular dynamics simulation study of xyloglucan adsorption on cellulose surfaces: effects of surface hydrophobicity and side-chain variation

Zhen Zhao · Vincent H. Crespi ·
James D. Kubicki · Daniel J. Cosgrove ·
Linghao Zhong

Published online: 19 October 2013
© Springer Science+Business Media Dordrecht 2013

Erratum to: Cellulose DOI 10.1007/s10570-013-0041-1

In the original publication, the labels for the Miller indices of (110), (1–10) and (200) for the cellulose microfibril model in Fig. 2 are incorrect. In the text, figures and tables, the index “(1–10)” should be replaced with “(110)”, “(110)” should be replaced

with “(1–10)” and “(200)” should be replaced with “(100)”. We are grateful to Hervé Bizot for pointing out the error. The corrected Fig. 2 is provided in this Erratum.

The online version of the original article can be found under doi:[10.1007/s10570-013-0041-1](https://doi.org/10.1007/s10570-013-0041-1).

Z. Zhao (✉) · J. D. Kubicki
Department of Geosciences, Pennsylvania State
University, University Park, PA 16802, USA
e-mail: zxz13@psu.edu

Z. Zhao · V. H. Crespi · J. D. Kubicki ·
D. J. Cosgrove · L. Zhong
Center for Lignocellulose Structure and Formation,
Pennsylvania State University, University Park,
PA 16802, USA

V. H. Crespi
Departments of Physics, Chemistry, and Materials
Science and Engineering, Pennsylvania State University,
University Park, PA 16802, USA

D. J. Cosgrove
Department of Biology, Pennsylvania State University,
University Park, PA 16802, USA

L. Zhong
Department of Chemistry, Pennsylvania State University,
Mont Alto, PA 17237, USA




Fig. 2 Initial and final (25 ns) structures of $(XXXG)_3$ adsorbed onto cellulose I- β microfibril (1-10), (100), (010), (100) and (110) surfaces. The cellulose microfibril has 25 glucan chains,

each 20 glucose residues long. Same *color code* as Fig. 1 is used for $(XXXG)_3$. (Color figure online)