

Warning SINEs: *Alu* elements, evolution of the human brain, and the spectrum of neurological disease

Peter A. Larsen · Kelsie E. Hunnicutt ·
Roxanne J. Larsen · Anne D. Yoder ·
Ann M. Saunders

Received: 6 December 2017 / Revised: 14 January 2018 / Accepted: 15 January 2018 / Published online: 19 February 2018
© The Author(s) 2018. This article is an open access publication

Abstract *Alu* elements are a highly successful family of primate-specific retrotransposons that have fundamentally shaped primate evolution, including the evolution of our own species. *Alus* play critical roles in the formation of neurological networks and the epigenetic regulation of biochemical processes throughout the central nervous system (CNS), and thus are hypothesized to have contributed to the origin of human cognition. Despite the benefits that *Alus* provide, deleterious *Alu* activity is associated with a number of neurological and neurodegenerative disorders. In particular,

neurological networks are potentially vulnerable to the epigenetic dysregulation of *Alu* elements operating across the suite of nuclear-encoded mitochondrial genes that are critical for both mitochondrial and CNS function. Here, we highlight the beneficial neurological aspects of *Alu* elements as well as their potential to cause disease by disrupting key cellular processes across the CNS. We identify at least 37 neurological and neurodegenerative disorders wherein deleterious *Alu* activity has been implicated as a contributing factor for the manifestation of disease, and for many of these disorders, this activity is operating on genes that are essential for proper mitochondrial function. We conclude that the epigenetic dysregulation of *Alu* elements can ultimately disrupt mitochondrial homeostasis within the CNS. This mechanism is a plausible source for the incipient neuronal stress that is consistently observed across a spectrum of sporadic neurological and neurodegenerative disorders.

Responsible Editor: Beth Sullivan.

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s10577-018-9573-4>) contains supplementary material, which is available to authorized users.

P. A. Larsen · K. E. Hunnicutt · A. D. Yoder
Department of Biology, Duke University, Durham, NC 27708,
USA

P. A. Larsen · A. D. Yoder
Duke Lemur Center, Duke University, Durham, NC 27708, USA

P. A. Larsen (✉)
Department of Biology, Duke University, 130 Science Drive, Box
90338, Durham, NC 27708, USA
e-mail: peter.larsen@duke.edu

R. J. Larsen
Duke University School of Medicine, Duke University, Durham,
NC 27710, USA

A. M. Saunders
Zinfandel Pharmaceuticals Inc, Chapel Hill, NC 27709, USA

Keywords A-to-I editing · Alzheimer's disease · Brain connectome · Epigenetics · Mitochondria · Mosaic brain · Parkinson's disease

Abbreviations

A-to-I	Adenosine-to-inosine
AD	Alzheimer's disease
ADAR	Adenosine deaminase acting on RNA
ALS	Amyotrophic lateral sclerosis
AMPA	α -Amino-3-hydroxy-5methyl-4-isoxazole propionate
APP	Amyloid precursor protein

circRNAs	Circular RNAs
CNS	Central nervous system
FLAM	Free left <i>Alu</i> monomer
LINE	Long interspersed element
L1	Long interspersed element-1
LTR	Long-terminal repeat
mRNA	Messenger RNA
PD	Parkinson's disease
pre-mRNA	Precursor messenger RNA
SEDs	Super-enhancer domains
SINE	Short interspersed element
TADs	Topologically associating domains
TOMM	Translocase of outer mitochondrial membrane

Introduction

Retrotransposons are mobile genetic elements that utilize an RNA intermediate to copy and paste themselves throughout the genome. There are two primary groups of retrotransposons, those having long-terminal repeats (LTRs) and those without (non-LTR) (Cordaux and Batzer 2009). In the human genome, non-LTR retrotransposons consist of long interspersed elements (LINEs) and short interspersed elements (SINEs), and these collectively account for a remarkable ~33% of total genome sequence (Cordaux and Batzer 2009). *Alu* elements are primate-specific SINEs that are approximately 300 nucleotides in length and are abundant in the human genome, with over 1.3 million elements accounting for at least 11% of overall DNA sequence (Deininger et al. 2003; Hancks and Kazazian 2016). Although once considered to be useless “junk DNA,” the prevalence, diversity, and non-random distribution of *Alu* elements across primate genomes are suggestive of a functional advantage. Indeed, a large body of evidence documents that *Alu* elements have directly influenced primate evolution by facilitating genome innovation through novel gene formation, elevated transcriptional diversity, long non-coding RNA and microRNA evolution (including circular RNAs), transcriptional regulation, and creation of novel response elements (Vansant and Reynolds 1995; Norris et al. 1995; Britten 1997; Lev-Maor et al. 2003; Polak and Domany 2006; Laperriere et al. 2007; Lin et al. 2008, 2016; Lehnert et al. 2009; Cordaux and Batzer 2009; Shen et al. 2011;

Jeck et al. 2013; Töhönen et al. 2015; Luco 2016; Chen and Yang 2017). Moreover, *Alus* fundamentally alter the three-dimensional architecture and spatial organization of primate genomes by defining the boundaries of chromatin interaction domains (i.e., topologically associating domains (TADs); Dixon et al. 2012). Genome architecture has a direct influence on biological function, and the observation that *Alus* are enriched within both TADs and super-enhancer domains (SEDs) supports the hypothesis that *Alus* directly influence a wide range of critically important processes within primates across multiple levels, from overall genome stability to tissue-specific gene regulation (Huda et al. 2009; Dixon et al. 2012; Soibam 2017; Glinsky 2018). In light of the functional benefits that *Alus* provide primates, it is interesting to note that *Alu* retrotransposition events occurred at an estimated 15-fold higher rate in the human, chimpanzee, and bonobo lineage (as compared to other great apes) and a 2.2-fold higher rate in humans when compared to chimpanzee and bonobo (Hedges et al. 2004; Prüfer et al. 2012; Hormozdiari et al. 2013). These evolutionary patterns indicate that positive selection is acting to maintain *Alu* elements in primate genomes, especially within humans (Mattick and Mehler 2008; Tsigiris and Rigoutsos 2009).

One of the most fascinating and biologically important aspects of *Alu* elements is that they serve an important role in the formation and function of the brain connectome (Oliver and Greene 2011; Li and Church 2013; Smalheiser 2014; Sakurai et al. 2014; Prendergast et al. 2014; Linker et al. 2017; Bitar and Barry, 2017). Many lines of evidence connect *Alu* elements with neurogenesis and critical neuronal biochemical processes, including somatic retrotransposition in developing neurons (in parallel to L1 retrotransposition; Baillie et al. 2011; Kurnosov et al. 2015), formation of regulatory circRNAs that are enriched in the central nervous system (CNS) and concentrated at synapses (Jeck et al. 2013; Rybak-Wolf et al. 2015; Chen and Schuman 2016; Floris et al. 2017), regulation of genes that are essential for proper neuron function (e.g., *ACE*, *SMN1*, *SMN2*, *SLC6A4*; Wu et al. 2013; Ottesen et al. 2017; Schneider et al. 2017), and elevated adenosine-to-inosine (A-to-I) RNA editing in the brain (Mehler and Mattick 2007; Kurnosov et al. 2015; Behm and Öhman 2016). In particular, epigenetic A-to-I editing plays a significant role in mediating neuronal gene expression pathways (Tariq and Jantsch 2012) with *Alus* serving as the primary target for RNA editing in primates (Picardi

et al. 2015; Behm and Öhman 2016). Beyond RNA editing mechanisms, human neuronal gene pathways are regulated by non-coding RNAs originating from *Alu* elements (e.g., BC200 and NDM29) and specific *Alu* subfamilies contain retinoic acid response elements which help to regulate neural patterning, differentiation, and axon outgrowth (Vansant and Reynolds 1995; Laperriere et al. 2007; Maden 2007; Castelnovo et al. 2010; Smalheiser 2014). Moreover, recent discoveries indicate that *Alu* elements underlie the formation of a vast number of human-specific circRNAs that are hypothesized to play important roles in neurological gene expression pathways (Jeck et al. 2013; Rybak-Wolf et al. 2015; Chen and Schuman 2016; Dong et al. 2017). There is a deep connection between *Alus* and the formation and function of primate neurological networks, and this has led to the hypothesis that *Alu* elements were essential for development of the transcriptional diversity and regulation required for the genesis of human cognitive function (Mattick and Mehler 2008; Oliver and Greene 2011; Li and Church 2013; Sakurai et al. 2014).

Despite the functional benefits that *Alus* have provided primate genomes, *Alu* elements can disrupt gene expression and function through many pathways (Fig. 1; Deininger and Batzer 1999; Deininger 2011; Tarallo et al. 2012; Ade et al. 2013; Elbarbary et al. 2016; Varizhuk et al. 2016). For this reason, the genome tightly regulates *Alus* using both DNA methylation and histone (H3K9 methylation) modification in order to control their expression and *de novo* retrotransposition (Varshney et al. 2015; Elbarbary et al. 2016; Mita and Boeke 2016), and there is mounting evidence indicating that the loss of these epigenetic control mechanisms (due to aging, cellular senescence, environmental factors, and stress) contributes to many forms of cancer, diabetes, osteoporosis, and several mental and neurodegenerative disorders (Szapkowski et al. 2009; Belancio et al. 2010; Muotri et al. 2010; Jintaridh et al. 2013; Dannlowski et al. 2014; Erwin et al. 2014; Bundo et al. 2014; Sun et al. 2014; Goodier 2016; Neven et al. 2016; Bedrosian et al. 2016; Shpyleva et al. 2017; Thongsroy et al. 2017). With respect to deleterious *Alu* pathways and neurological disease, there are at least 37 mental and neurodegenerative disorders wherein *Alu* elements are hypothesized to disrupt key cellular processes, thereby resulting in or contributing to the diseased state (Table 1).

Given the tight connection between *Alu* elements and the formation and function of the nervous system, it is likely that the dysregulation of *Alu* elements contributes




Fig. 1 Select mechanisms whereby *Alu* elements can alter gene expression and function (also see Elbarbary et al. 2016). **a** Sequence homology and orientation of *Alu* elements contribute to the formation of distinct secondary structures in both DNA and RNA. DNA *Alu* G-quadruplex structures can alter transcription kinetics (Varizhuk et al. 2016), and pre-mRNA *Alu* binding forms stem-loop structures that are the primary site for A-to-I editing (see Fig. 2). **b** Recombination of intra-gene *Alu* elements resulting in exon deletion. **c** Exonification of intronic *Alus* contributing to the production of alternative mRNAs. **d** Environmental or traumatic stress cascades resulting in increased expression of *Alu* RNAs that contribute to inflammation (Li and Schmid 2001; Tarallo et al. 2012; Hunter et al. 2015; Lapp et al. 2016), the disruption of global gene transcription through Pol II binding (Mariner et al. 2008), and an increase of H3K9 histone methylation that alters *Alu* epigenetic pathways (Varshney et al. 2015; Lapp and Hunter 2016; Larsen et al. 2017)

to many sporadic or idiopathic neurological disorders observed across the global human population (Larsen et al. 2017). Here, we highlight both the beneficial neurological aspects of *Alu* elements as well as their potential to cause neurological disease. We focus on a novel hypothesis that identifies a potential epigenetic vulnerability to neurological networks that has likely escaped purifying selection. The *Alu* neurodegeneration

Table 1 Genes associated with neurological and neurodegenerative disorders wherein deleterious *Alu* activity has been documented experimentally or is hypothesized to disrupt gene function

Gene name	Disorder	<i>Alu</i> mechanism of disruption	Reference
ABCD1	Adrenoleukodystrophy	Deletion events	Kutsche et al. (2002)
ACAT1 (T2)	Mitochondrial acetoacetyl-CoA thiolase deficiency	Deletion event	Zhang et al. (2006)
ACE	Alzheimer's disease	Insertion events	Wu et al. (2013)
ADAR2	Glioma	Exonization	Li et al. (2015)
ALDH7A1	Pyridoxine-dependent epilepsy	Recombination	Mefford et al. (2015)
ALMS1	Alström syndrome	Insertion event	Taşkesen et al. (2012)
APOB	Hypobetalipoproteinemia	Recombination	Huang et al. (1989)
ATP5J	Alzheimer's disease	Duplication	Antonell et al. (2012)
ATP7A	Menkes disease	Insertion event	Gu et al. (2007); Bhattacharjee et al. (2016)
ATP7B	Wilson's disease	Alternative splicing	Mameli et al. (2015)
C9orf72	ALS, FTL	Loss of epigenetic control, elevated <i>Alu</i> transcripts	Prudencio et al. (2017)
CHD7	CHARGE syndrome	Deletion	Udaka et al. (2007)
CLN3	Batten disease	Deletion	Lerner et al. (1995)
COL4A5	Alport syndrome	Deletion and exonization	Nozu et al. (2014)
DICER1 ^a	Age-related macular degeneration	<i>Alu</i> RNA build-up with reduced DICER1 activity	Kaneko et al. (2011), Kim et al. (2014)
FXN	Friedreich's ataxia	<i>Alu</i> repeat expansion, alternative splicing events	Pandolfo (2006)
GK	Glycerol kinase deficiency	Insertion event	Zhang et al. (2000)
GLA	Fabry disease	Deletion event	Dobrovolny et al. (2011)
HPRT	Lesch-Nyhan disease	Recombination	Brooks et al. (2001)
HMBS	Acute intermittent porphyria	Insertion event	Mustajoki et al. (1999)
LPL	Lipoprotein lipase deficiency	Complex deletion-insertion	Okubo et al. (2007)
MFN2 (CMT2a)	Charcot-Marie-Tooth type 2A	Copy number variants	Pehlivan et al. (2016)
MPO	Alzheimer's disease	<i>Alu</i> hormone response variant; estrogen dysregulation	Reynolds et al. (1999)
NDUFS2	Leigh syndrome	Exonization	Larsen et al. (2017)
NF1	Neurofibromatosis type I	Deletion and chimeric gene fusion	Wimmer et al. (2011), Ferrari et al. (2017)
NFIX	Marshall-Smith syndrome	Deletions	Schanze et al. (2014)
OPA1	Autosomal dominant optic atrophy	Alternative splicing events	Gallus et al. (2010)
PARK2	Parkinson's disease	Recombination	Morais et al. (2016)
PARK7 (DJ-1)	Parkinson's disease	Deletion	Bonifati et al. (2003)
PDHA1	Pyruvate dehydrogenase deficiency	Exonization	Larsen et al. (2017)
PIGL	CHIME syndrome	Deletion	Johnson et al. (2017)
PMM2	Congenital disorders of glycosylation type Ia	Complex deletion	Schollen et al. (2007)
POMT1	Walker Warburg syndrome	Insertion	Bouchet et al. (2007)
PSEN1	Alzheimer's disease	Deletion	Le Guennec et al. (2017)
PXMP2 (PMP22)	Charcot-Marie-Tooth type 2A	<i>Alu-Alu</i> -mediated rearrangement	Choi et al. (2011), Gu et al. (2015)

Table 1 (continued)

Gene name	Disorder	<i>Alu</i> mechanism of disruption	Reference
RP2 (NUDT19)	X-linked retinitis pigmentosa	<i>Alu</i> -L1 recombination	Schwahn et al. (1998), Jiang et al. (2017)
SLC6A4	Depression, reduced hippocampal volume	Altered promoter methylation	Dannowski et al. (2014)
SLC25AC	Intellectual disability	Deletion	Vandewalle et al. (2013)
SLC30A6	Alzheimer's disease, dementia, ALS	Gene fusion event	Boone et al. (2014)
SMN1	Spinal muscular atrophy	Exonization, deletion events, circularization	Ottesen et al. (2017)
SOD2	Hyperglycemia	Repressed expression	Wang et al. (2016)
SOX10	Waardenburg syndrome type 4	Deletion	Bondurand et al. (2012)
SPAST	Autosomal-dominant spastic paraplegia 4	Deletions, CNVs, gene fusion events	Boone et al. (2014)
SPG7	Hereditary spastic paraplegia	Deletion, recombination	Arnoldi et al. (2008), López et al. (2015)
SPG11	Hereditary spastic paraplegia	Deletion	Conceição et al. (2012)
STAU1	Myotonic dystrophy type 1	Alternative splicing regulation	Bondy-Chorney et al. (2016)
TDP-43	ALS, frontotemporal lobar degeneration	Transposable element dysregulation	Li et al. (2012)
TOMM40	Late-onset Alzheimer's disease	<i>Alu</i> repeat expansion, putative alternative splicing events	Larsen et al. (2017)
TRIM37	Mulibrey nanism	Deletion events	Jobic et al. (2017)

Gene names in bold identify genes essential for mitochondrial function and transport and/or are associated with mitochondrial abnormalities (sensu Dawson et al. 1995; Calvo et al. 2015; Zempel and Mandelkow 2015; Bhattacharjee et al. 2016; Chong-Chong et al. 2016; Checler et al. 2017; Johnson et al. 2017; Wang et al. 2013). For additional *Alu*-associated diseases, see Hancks and Kazazian (2016) and Payer et al. (2017)

^a Genes where mutations result in dysregulation of *Alu* elements

hypothesis (sensu Larsen et al. 2017) posits that the epigenetic dysregulation of *Alu* elements ultimately serves to disrupt mitochondrial homeostasis in neurological networks, thereby setting the stage for increased neuronal stress and neurodegeneration. Given this hypothesis, it is noteworthy that many of the *Alu*-disrupted genes associated with neurological disorders are related to mitochondrial function and trafficking, including nuclear-encoded mitochondrial genes (i.e., mitonuclear) which help to regulate oxidative stress and metabolic processes in the CNS (Table 1). Mitochondrial dysfunction is implicated across the spectrum of neurological and neurodegenerative disorders that are observed in humans, and this pattern is suggestive of a genetic vulnerability that has evolved in humans. Considering this, we begin by reviewing the integral role that *Alu* elements have played in human evolution through brain-specific epigenetic A-to-I RNA editing pathways and neurological network formation. Although these *Alu*-related processes are hypothesized to have contributed

to the origin of human cognition, they are likely accompanied by age or stress-related epigenetic vulnerabilities to the CNS, with mitochondrial pathways being especially sensitive.

Alu elements, A-to-I editing, and evolution of the human brain

Alu elements are non-randomly distributed throughout the genome. They occur most frequently within introns and are enriched within genes involved in metabolic, mitochondrial, cellular transport, and binding pathways (Grover et al. 2003; de Andrade et al. 2011; Larsen et al. 2017). *Alu* nucleotide sequences and lengths (~300 bp) are generally conserved (Batzer and Deininger 2002), and it is this seemingly simple aspect of *Alu* biology that is of monumental biological importance. When inserted within a gene at opposite orientations and at close proximity, *Alus* bind upon themselves post-transcriptionally, resulting in the formation of a duplex stem-loop




Fig. 2 Intronic *Alu* elements located in close proximity (a) can bind to each other within pre-mRNAs (b) thereby producing a stable stem-loop secondary structure that is the primary substrate for A-to-I editing in primates. ADAR proteins bind to pre-mRNA *Alu* structures (c) and convert adenosine residues to inosine. If occurring in coding regions, the translation machinery then interprets the inosine residues as guanosine and this can contribute to amino acid changes and alternative protein conformations (d). The model is also applicable to inverted *Alus* within 3' untranslated regions, wherein promiscuous editing by ADAR can result in nuclear retention of mRNAs (Chen and Carmichael 2008 and 2009)

structure that is stabilized by the *Alu* nucleotide sequence and length (Fig. 2; Athanasiadis et al. 2004). These *Alu*-based secondary structures fundamentally alter the shape of pre-mRNA molecules and serve as the primary binding site for ADAR proteins, which bind to the double-stranded pre-mRNA duplex and edit adenosine (A) residues to inosine (I) thereby recoding pre-mRNAs (Fig. 2). When operating in coding regions (either directly or indirectly), the translation machinery interprets the resulting I residues as guanosine (G) and this mechanism accounts, in part, for the incredible

diversity observed in the human proteome that is not encoded within the original DNA sequence (Nishikura 2016). However, the vast majority of A-to-I editing operating on *Alu* elements occurs within pre-mRNA introns and 3' UTRs, and this can directly influence gene regulation and function in a surprising number of ways, including the creation of novel splice donor and acceptor sites that result in *Alu* exonization and alternative gene splicing (Nishikura 2016), recoding of exons immediately adjacent to *Alus* (Daniel et al. 2014), disruption of RNAi pathways (Chen and Carmichael 2008), production of novel micro-RNA regulatory sites (Borchert et al. 2009), and increased nuclear retention of promiscuously edited mRNAs (Chen and Carmichael 2008 and 2009).

Although A-to-I editing plays an essential role in generating transcriptional diversity across eukaryotes, *Alu* elements provide primate-specific RNA editing opportunities. An example of this is found when comparing the rodent-specific SINE B1 family to *Alu*. Both B1 and *Alu* SINE families originated from 7SL RNA (Ullu and Tschudi 1984; Vassetzky et al. 2003), yet rodent-specific B1 elements are approximately half the length (~140 bp) of primate *Alus* (~300 bp) and have greater levels of intranucleotide variation. When considering the functional mechanics of A-to-I editing (Fig. 2), the shorter lengths and more variable rodent B1 elements result in the formation of shorter and less-stable double-stranded stem-loop structures in rodent pre-mRNAs. Thus, the molecular properties that separate rodent B1 from primate *Alu* translate to key functional genomic differences that have influenced evolutionary processes within each lineage (Eisenberg et al. 2005; Neeman et al. 2006; Picardi et al. 2015; Tan et al. 2017).

A-to-I editing associated with *Alu* elements is perhaps one of the most functionally important yet underappreciated aspect of *Alu* biology. Approximately 90% of A-to-I editing within primate gene networks centers on *Alu* elements, and this has fundamentally shaped primate evolution, including the evolution of our own species where A-to-I editing is estimated to occur at over 100 million sites in the human transcriptome (Bazak et al. 2014). Moreover, recent data supports a connection between A-to-I editing and neurological network formation, with elevated editing levels occurring throughout neurogenesis (Behm and Öhman 2016). Genes encoding for key neurological proteins involved in neurotransmission, neurogenesis, gliogenesis, and synaptogenesis are subject to enhanced A-to-I editing,

and thus, a number of studies have hypothesized a strong link between A-to-I RNA editing pathways and brain development and function (Schmauss and Howe 2002; Mehler and Mattick 2007; Tan et al. 2009; Sakurai et al. 2014; Liscovitch et al. 2014; Behm and Öhman 2016; Hwang et al. 2016; Picardi et al. 2017b). A recent analysis of A-to-I editing in over 8500 human samples identified tissue-specific editing patterns with elevated editing levels in the brain, including unique patterns in the cerebellum (Tan et al. 2017). The vast majority of these neurologic A-to-I editing events are operating on *Alu* elements, and, when combined with human-specific *Alu* evolution (Hedges et al. 2004; Cordaux and Batzer 2009; Prüfer et al. 2012; Hormozdiari et al. 2013), this observation serves as the foundation for the hypothesis that *Alu* elements and *Alu*-related pathways contributed to the evolution of enhanced human cognitive abilities (Mattick and Mehler 2008; Li and Church 2013).

Given the relationship between *Alu*-centric A-to-I editing and the formation and function of the CNS, it is important to expand upon the neuro-specific functions of ADAR proteins. Three ADAR proteins are identified (ADAR1, ADAR2, and ADAR3), and these proteins have distinct tissue-specific expression patterns (Picardi et al. 2015; Tan et al. 2017). ADAR1 and ADAR2 co-opt to regulate neuronal activity by editing key neurotransmitter receptors and ion channels in the CNS (Hood and Emeson 2011). Interestingly, both ADAR2 and ADAR3 have unique brain-specific expression patterns with ADAR2 being highly expressed in the brain and ADAR3 exclusively expressed in the brain (Mehler and Mattick 2007). Until recently, the functional role of the brain-specific ADAR3 protein was largely unknown; however, Oakes et al. (2017) discovered that ADAR3 competes with ADAR2 to regulate glutamate receptor subunit B (*GRIA2*) A-to-I editing. The *GRIA2* protein forms a critical subunit of α -amino-3-hydroxy-5methyl-4-isoxazole propionate (AMPA) receptors, which regulate synaptic calcium and are involved with synaptic plasticity, memory, and learning (Wright and Vissel 2012). Remarkably, A-to-I editing of a specific adenosine nucleotide within *GRIA2* results in an amino acid change that alters the GluR-2 protein conformation, thus disrupting calcium permeability of the AMPA receptor and potentially contributing to epilepsy, amyotrophic lateral sclerosis (ALS), and schizophrenia (see Oakes et al. 2017). In light of Oakes et al. (2017), the brain-specific expression pattern of ADAR3 indicates that this

protein helps to offset A-to-I editing by ADAR2, perhaps serving to mediate enhanced RNA editing processes throughout the CNS.

Considering the essential role that A-to-I editing processes play in the CNS, the dysregulation of these processes can have a profound impact on the stability of neurological networks (Mehler and Mattick 2007; Rice et al. 2012; Hwang et al. 2016). With respect to ADAR proteins, mutations within ADAR1 have been linked to Aicardi-Goutières syndrome (characterized by severe brain dysfunction; Rice et al. 2012) and *Alu*-related alternative splicing events of ADAR2 are linked to glioma (Li et al. 2015). Disruption of ADAR1 editing increases production of unedited RNAs which interact with MAV proteins in the outer mitochondrial membrane, ultimately serving to activate inflammatory response pathways (Bajad et al. 2017; Gallo et al. 2017) and perhaps providing a mechanism for inflammatory diseases of the CNS (Hofer and Campbell 2016). ADAR2 knockout mice display epileptic seizures and neuronal death caused by an influx of calcium owing to the disruption of *GRIA2* editing (see above). The interference of A-to-I editing processes associated with the *KCNA1* gene (encoding a protein essential for potassium regulation and neuron excitability) is hypothesized to underlie episodic ataxia type-1 disorder, a disease of the CNS characterized by seizures, stress-induced ataxia, and myokymia (Ferrick-Kiddie et al. 2017). Moreover, a reduction of A-to-I editing has been observed within hippocampal tissues of Alzheimer's brains versus healthy controls (Khmermesh et al. 2016). From a broader perspective, the disruption of A-to-I editing processes across the CNS has been linked to a wide variety of mental and neurodegenerative disorders including major depression and suicide, epilepsy, schizophrenia, Alzheimer's disease (AD), and ALS (Gurevich et al. 2002; Kawahara et al. 2004; Kwak and Kawahara 2005; Maas et al. 2006; Kubota-Sakashita et al. 2014; Khmermesh et al. 2016; Weissmann et al. 2016; Gal-Mark et al. 2017).

Alu elements, neurogenesis, and the human brain connectome

There is a strong connection between *Alu* A-to-I editing and the development and function of the brain; therefore, it is impossible to disentangle *Alus* from the formation and function of neurologic networks (Mehler and Mattick 2007; Tan et al. 2009; Behm and Öhman

2016). It is estimated that the human brain is comprised of over 100 billion neurons that are organized into functional hubs or parcels collectively forming the brain connectome (Van Den Heuvel and Sporns 2013). Beyond major structures of the brain (e.g., cerebellum, frontal cortex, hippocampus, etc.), the existence of connectome parcels shared across unrelated individuals is indicative of an evolutionary conserved process underlying neurological network formation and operating throughout neurogenesis. For example, a recent study mapped the cerebral cortex using multi-modal magnetic resonance imaging and identified 180 connectome parcels that were largely shared across 210 healthy adults (Glasser et al. 2016). Understanding the molecular processes that contribute to the formation of the human brain connectome is essential for understanding the origin of human-specific neurological disorders and diseases observed across the global distribution of our species. This is especially true for neurodegenerative conditions that are hypothesized to originate in functional network hubs and progress along neuronal network connections (e.g., AD; Seeley 2017; Cope et al. 2018).

A growing body of evidence indicates that retrotransposons (including both LINES and SINES) are active throughout neurogenesis and contribute to mosaic neuron genomes that ultimately form the human brain connectome (Muotri et al. 2005; Erwin et al. 2014; Kurnosov et al. 2015; Evrony 2016; Paquola et al. 2016; Linker et al. 2017). Although somatic L1 retrotransposition events within developing neurons have received much attention, it is noteworthy that *Alu* retrotransposition occurs in parallel with L1 throughout neurogenesis (Baillie et al. 2011; Kurnosov et al. 2015), thus providing primate-specific aspects of neurologic network formation. Furthermore, there is evidence that unites *Alu* elements with retinoic acid regulation (Vansant and Reynolds 1995; Laperriere et al. 2007), which is essential for neuronal patterning and differentiation throughout neurogenesis and is a potential regulator of neuron regeneration (Maden 2007). Retinoic acid is vital for the establishment, maintenance, and repair of neuronal networks, and given the presence of retinoic acid response elements in *Alu* elements, it is possible that *Alu* activity during neurogenesis is connected to retinoic acid signaling processes.

Considering the *Alu* regulatory pathways discussed above, it is of great interest to note that retrotransposition of *Alu* elements is hypothesized to occur at elevated




Fig. 3 Deleterious *Alu* activity operating on mitonuclear genes can disrupt mitochondrial function in the CNS and contribute to a number of diseased phenotypes (see Table 1). The type and severity of associated neurological and neurodegenerative disorders depends on the deleterious *Alu* mechanism of action, the mitonuclear gene pathways involved, the time or developmental stage of induction, level or severity of traumatic stress, and tissue specificity (see Larsen et al. 2017). If operating across the suite of mitonuclear genes through epigenetic pathways, the mechanism helps to explain the origin of incipient mitochondrial stress and CNS connectome destabilization that is observed across the spectrum of neurological and neurodegenerative disorders

levels within the dentate gyrus of the hippocampus, the putative site of adult neurogenesis (Kurnosov et al. 2015). Moreover, A-to-I editing levels steadily increase as neural progenitor cells develop into adult neurons (Behm and Öhman 2016). These data indicate that at least two retrotransposon-centric processes (somatic retrotransposition of both LINES and SINES and enhanced A-to-I editing operating primarily on *Alu* elements) are major contributors to neurogenesis, perhaps

servicing to establish the neuronal and biochemical diversity that underlies the ~100 billion neuron brain connectome. Remarkably, emerging data suggests that a third *Alu*-centric process is associated with the formation and function of neurological networks, this being the production of circRNAs that are enriched in the brain and concentrated at synaptic junctions (Jeck et al. 2013; Rybak-Wolf et al. 2015; Chen and Schuman 2016). Identifying vulnerabilities to each of these retrotransposon-centric processes will likely contribute to the identification of novel mechanisms underlying mental disorders and neurologic disease and could lead to novel therapeutic interventions.

Pathways to incipient neuronal stress and neurological disease

The disruption of *Alu*-centric epigenetic RNA editing processes is implicated across the entire spectrum of neurologic disorders (see above). In light of this observation, it is interesting to note that another, seemingly unrelated, feature of many neurological disorders is mitochondrial dysfunction (Lin and Beal 2006; Rugarli and Langer 2012; Gottschalk et al. 2014; Petschner et al. 2017). However, we have previously shown that mitonuclear genes are enriched with *Alu* elements when compared to random (Larsen et al. 2017), which is consistent with earlier observations regarding the non-random insertion of *Alu* elements into genes associated with transcriptionally active regions of the genome (Grover et al. 2003; de Andrade et al. 2011). Thus, it is likely that *Alu*-mediated gene regulatory processes are actively influencing mitonuclear gene expression, regulation, and protein function through the pathways discussed above and reviewed in Chen and Yang (2017). Knowing this, the dysregulation of epigenetic *Alu* regulatory pathways is a plausible source for mitochondrial stress and dysfunction, with the CNS being particularly vulnerable (Larsen et al. 2017). Such a mechanism could contribute to the initial activation of complex mitochondrial stress pathways and incipient neuronal stress associated with sporadic neurologic disorders (e.g., inflammation, immune response, mitophagy, etc.). Importantly, these processes would precede macroscopic pathologies such as protein aggregation and neuronal atrophy observed in neurodegenerative diseases (Swerdlow et al. 2010; Larsen et al. 2017; Swerdlow 2017).

The *Alu* neurodegeneration hypothesis (*sensu* Larsen et al. 2017) proposes a “double-edged sword,” whereby the beneficial *Alu*-related processes that underlie neuron diversity and function also have the potential to disrupt mitochondrial homeostasis across neurological networks through deleterious cascade events that are facilitated by eroding tissue-specific *Alu* epigenetic control mechanisms. The stability of the brain’s connectome and the entire CNS depends on healthy mitochondrial populations within neurons, astrocytes, microglia, and supporting cells (Cai et al. 2011; Viader et al. 2011; Schwarz 2013; Jackson and Robinson 2017). Mitochondria play critical roles for a wide range of essential neuronal processes including glucose and lipid metabolism, metal ion biosynthesis, cellular trafficking along axons, neurotransmitter relay across synapses, and synaptic calcium homeostasis (Schwarz 2013; Harbauer et al. 2014). Therefore, molecular mechanisms that are known to disrupt gene expression and protein folding of genes that are essential for mitochondrial function can ultimately disrupt neurological function.

Interference of mitochondrial dynamics across the CNS is consistently hypothesized to occur during the earliest stages of mental, neurological, and neurodegenerative disorders ranging from depression, epilepsy, and schizophrenia to ALS, AD, and Parkinson’s disease (PD; Lu 2009; Rezin et al. 2009; Kim et al. 2010; Coskun et al. 2012; Martin 2012; Gottschalk et al. 2014; Larsen et al. 2017; Flippo and Strack 2017; Petschner et al. 2017). Collectively, these disorders are estimated to impact approximately 250 million people globally, accounting for at least 10.2% of the global disease burden (GBD 2015 Neurologic Disorders Collaborator Group 2017). The occurrence of sporadic forms of human-specific neurologic disorders (e.g., non-familial schizophrenia, ALS, late-onset AD, PD, etc.) across the entire distribution of our species is suggestive of a common yet complex genetic mechanism that evolved in primates and is amplified in humans (Larsen et al. 2017). Considering this, we expand on the mitocentric view of idiopathic neurologic disease manifestation by reviewing the evidence that unites primate-specific *Alu* activity with incipient neurologic mitochondrial dysfunction.

Eukaryotic mitochondria are hypothesized to have originated from an endosymbiotic alphaproteobacterium which, over expansive evolutionary time, evolved in parallel with host genomes into the mitochondrial organelles that we observe today (Roger et al. 2017). The

human mitochondrial genome encodes only 13 proteins, yet it is estimated that human mitochondria depend on approximately ~2000 genes encoded within the nuclear genome for their functionality (Calvo et al. 2015; Johnson et al. 2017). These mitonuclear genes are thus subject to deleterious *Alu* activity, and *Alu*-related deleterious events have been linked to many neurologic and neurodegenerative disorders, including epilepsy, Wilson's disease, Leigh syndrome, PD, ALS, and AD (Table 1 and references therein; Fig. 3). When considering the incipient mitochondrial dysfunction observed across the spectrum of neurological neurodegenerative disorders, it is possible that tissue-specific epigenetic dysregulation of *Alu* elements within the CNS can ultimately manifest into distinct disease phenotypes (Larsen et al. 2017).

Several interesting patterns emerge when examining the key neurologic processes that are disrupted through deleterious *Alu* activity (Table 1). For example, mitochondria play an essential role in maintaining intracellular metal ion homeostasis (e.g., iron, copper, and zinc), the disruption of which can result in the increased production of free radicals that damage mitochondria and contribute to the increased production of reduced oxygen species (ROS; Rossi et al. 2004; Madsen and Gitlin 2007). The brain is especially sensitive to ROS production, and iron, copper, and zinc-related oxidative stress has been linked to many neurodegenerative disorders including AD, PD, and Wilson's disease (Rossi et al. 2004; Madsen and Gitlin 2007). It is notable that deleterious *Alu* activity has been identified in several genes that are essential to maintaining proper iron and copper homeostasis, including *FXN*, *ATP7A*, *ATP7B*, *HMBS*, *NDUFS2*, *SLC30A6*, and *PARK7 (DJ-1)* (Table 1; Gu et al. 2007; Kaler 2011; Girotto et al. 2014). Knowing this, it is possible that either global or tissue-specific dysregulation of *Alu* elements within mitonuclear genes can alter mitochondrial metal ion processing pathways thereby contributing to increased ROS production leading to neurologic stress.

A second interesting pattern with respect to deleterious neurologic *Alu* activity concerns metabolic pathways. The efficient processing of glucose and lipids across the CNS is critical for the stability and function of neurons, and the disruption of mitochondrial-mediated metabolic pathways has been linked to many neurologic disorders including AD and peripheral neuropathies (Viader et al. 2013; De La Monte and Tong 2014). Deleterious *Alu* activity occurs in genes that are

critical for glucose and lipid metabolism, including *ABCD1*, *ACAT1*, *ALMS1*, *APOB*, *GK*, *GLA*, *HPRT*, *LPL*, *PDHA1*, *PMM2*, *PSEN1*, *SOD2*, and *SPAST* (Table 1). Several of these genes encode for mitochondrial-related proteins that have been implicated in metabolic diseases that directly, or indirectly, contribute to neurological dysfunction. The connection between *Alu* elements and metabolic pathways is consistent with the observation that *Alu* elements preferentially insert into metabolic genes, and this has led to the hypothesis that *Alus* regulate the expression of genes related to type 1 diabetes (Grover et al. 2003; Mirza et al. 2014; Kaur and Pociot 2015). Moreover, *Alu* RNAs act to suppress the expression of both endothelial nitric oxide synthase (eNOS) and superoxide dismutase 2 (SOD2) during hyperglycemic conditions (Wang et al. 2016), suggesting a regulatory role of *Alu* elements during oxidative stress and strengthening the link between *Alu* element activity and diabetes.

There is growing evidence linking sporadic AD with dysfunctional metabolic pathways, leading some to consider AD as a "type 3 diabetes" wherein glycolysis and lipid homeostasis are altered (Steen et al. 2005; De La Monte et al. 2006; De La Monte and Wands 2008; De La Monte and Tong 2014; De Felice and Lourenco 2015; Mittal et al. 2016). The most well-documented risk factor for AD is a variant within *APOE* (*APOE* ϵ 4), a gene which encodes for a glycoprotein that mediates cholesterol and lipid transport (Saunders et al. 1993; Strittmatter et al. 1993; Mahley and Rall 2000). The *APOE* ϵ 4 allele is strongly associated with earlier onset of AD, and it is hypothesized that this is a result of the disruption of cholesterol processing and subsequent accumulation of amyloid precursor proteins (APP; i.e., the amyloid cascade hypothesis). Although the "amyloid cascade hypothesis" has dominated Alzheimer's research for decades (Hardy and Higgins 1992; Selkoe 2000; McKhann et al. 2011), the failure of multiple drug trials targeting amyloid pathways has led many in the Alzheimer's research community to search for alternative hypotheses that can help explain the origin of neurodegenerative disease as well as novel molecular pathways with therapeutic potential (Herrup 2015).

It is of particular interest then to note that a second genetic risk factor for AD, *TOMM40*, is located immediately adjacent to *APOE* on human chromosome 19, and the two genes are in tight linkage disequilibrium (Lyll et al. 2014; Roses et al. 2016a). *TOMM40* encodes for a beta-barrel protein that ultimately forms a

central pore in the outer mitochondrial membrane (Shiota et al. 2015), and much like *APOE*, genetic variants of *TOMM40* are linked to cognitive impairment and neurodegenerative disease (Roses 2010; Gottschalk et al. 2014; Greenbaum et al. 2014; Payton et al. 2016; Roses et al. 2016b; Arpawong et al. 2017). The most well known of these *TOMM40* variants is the rs10524523 (rs523) homopolymer repeat, a variable stretch of deoxythymidine (T) located within *TOMM40* intron 6 (Roses 2010). The rs523 poly-T varies in length from approximately 12 to 46 nucleotides, and the longer variants are statistically associated with thinning of the hippocampus (independent of the *APOE* ϵ 4 allele; Burggren et al. 2017) and earlier onset of AD (Lutz et al. 2010; Roses et al. 2010). Interestingly, rs523 is embedded within tandemly repeated *Alu* elements and originated from an *Alu* insertion event (Payton et al. 2016; Larsen et al. 2017). At least 149 *Alu* A-to-I editing events are identified within *TOMM40*, the majority of which are associated with *Alu* elements surrounding the rs523 repeat and intron 9 (Picardi et al. 2017a).

There is a potentially important link that unties *APOE* APP processing with the functional mechanics of pre-protein transport through the TOMM pore. It is possible that conformational changes of the Tom40 protein, potentially originating from *Alu*-mediated events (see above, reviewed in Elbarbary et al. 2016; Chen and Yang 2017; Larsen et al. 2017), can ultimately serve to restrict the passage of lipids across the outer mitochondrial membrane (Larsen et al. 2017). When combined with altered APP processing, this process could account for the initial site of intracellular protein accumulation that is hypothesized to precede extracellular plaque formation during very early stages of AD (Skovronsky et al. 1998; D'Andrea et al. 2001; Takahashi et al. 2002). Consistent with this hypothesis is the direct observation of APP accumulation at the TOMM pore (Devi et al. 2006) as well as functional tolerance of Tom40 conformational changes by mitochondria (Mager et al. 2011; Kuszak et al. 2015). Importantly, this mechanism could help to explain the common patterns of protein accumulation (e.g., amyloid plaques and alpha-synuclein Lewy bodies) observed across the spectrum of neurodegenerative disease, including ALS, AD, and PD (Ross and Poirier 2004; Gottschalk et al. 2014; Larsen et al. 2017). An age or stress-related component to Tom40 conformational changes comes with the epigenetic dysregulation

of *Alu* elements associated with the aging process or traumatic stress (see Larsen et al. 2017). Whether or not these processes are directly mediated by deleterious *Alu* events remains to be tested; however, it is notable that *Alu* exons and *Alu* somatic retrotransposition events have been identified in several TOM genes that are required for the stability of the translocase of the outer mitochondrial membrane and pre-protein import, including *TOMM5*, *TOMM7*, *TOMM22*, *TOMM40*, and *TOMM40L* (Baillie et al. 2011; de Andrade et al. 2011; Lin et al. 2016).

With respect to *Alu* elements, mitochondrial dysfunction, and the broader pathological scope of AD and other neurodegenerative diseases, there is evidence suggesting that *Alu*-derived peptides interact with tau proteins, perhaps serving a regulatory role for tau phosphorylation (Hoenicka et al. 2002). Tau is a microtubule-associated protein that functions to stabilize axonal microtubules and to transport mitochondria along axons, and taupathies (including tau hyperphosphorylation) are a characteristic feature of several neurodegenerative diseases including AD, progressive supranuclear palsy, corticobasal degeneration, and Pick's disease (Ittner and Götz 2011; Khanna et al. 2016). The *MAPT* gene encodes for tau, and alternative splicing events of *MAPT* result in multiple tau isoforms (Reddy 2011). Approximately 86 *Alu* elements (including FLAMs) are distributed throughout *MAPT* introns, and A-to-I editing is occurring at 315 *Alu*-related sites with elevated levels at the 3' end of *MAPT* (REDIportal database; Picardi et al. 2017a). When considering the potential for *Alu* structural variants of *MAPT* (including DNA and pre-mRNA secondary structures) and evidence of *Alu* RNAs interacting with tau proteins (Hoenicka et al. 2002), we recommend additional studies aimed at elucidating the regulatory impacts that *Alu* elements might have on *MAPT* gene expression and tau phosphorylation.

A-to-I editing and the potential for mitochondrial stress

Although several neurological disorders are hypothesized to be the result of disruptive A-to-I editing processes across the CNS (see above), it is presently unknown whether or not these processes are actively influencing mitochondrial function. What evidence is there indicating that post-transcriptional modification of mitonuclear genes can alter gene expression or function? Are there particular neurological or neurodegenerative disorders that are associated with mitonuclear

genes that have elevated levels of A-to-I editing? To provide insights into these questions, we searched the REDportal A-to-I editing database (Picardi et al. 2017a) for mitonuclear genes where (1) A-to-I editing has been identified within *Alu* elements in coding regions and (2) A-to-I editing has contributed to non-synonymous amino acid changes. We identified 57 mitonuclear genes with A-to-I editing occurring within putative *Alu* exons, and in 52 of these genes, the post-transcriptional modification resulted in non-synonymous amino acid changes (Supplementary Table 1). Many of these genes are involved with essential neuronal processes including calcium binding and transport, zinc transport, apoptosis regulation, voltage-gated ion channels, and mitochondrial elongation with notable examples including *ADSL*, *BAX*, *CASP2*, *COQ2*, *DFFB*, *FBXO18*, *LYRM4*, *PACRG*, and *SLC30A6* (Supplementary Table 1).

From a broader perspective, we identified enhanced A-to-I editing across 134 mitonuclear genes that are associated with a spectrum of neurologic and neurodegenerative disorders ranging from depression, tobacco use disorder, and bipolar disorder to ALS, Leigh syndrome, PD, and AD (Supplementary Table 2). In light of these patterns, we hypothesize that system-wide or tissue-specific epigenetic dysregulation of *Alu* A-to-I editing within the CNS can serve to disrupt key mitochondrial biochemical processes, thus potentially contributing to incipient mitochondrial and neuronal stress (Fig. 3).

Conclusions

Enhanced somatic retrotransposon throughout neurogenesis contributes to the mosaic brain; however, such activity likely contributes to mosaic pathways leading to disease (Erwin et al. 2014). Elucidating these pathways might ultimately provide insight into the sporadic nature of idiopathic diseases that are impacting the global human population. The disruption of *Alu*-mediated pathways that underlie gene regulation is a plausible mechanism for the origin of complex human-specific neurologic and neurodegenerative disorders. Although many of these disorders have similar phenotypes (e.g., mitochondrial dysfunction), it is possible that these phenotypes arise from deleterious activity operating across tissue-specific gene networks. If correlated with eroding or fluctuating epigenetic control

mechanisms of retrotransposons that are associated with aging, cellular senescence, and/or cellular stress (Belancio et al. 2010; Pal and Tyler 2016; Schneider et al. 2017), then such mechanisms might largely escape purifying selection and would be difficult to detect using traditional methods (e.g., genome-wide association studies). It is important to note that the *Alu*-centric mechanisms discussed herein collectively provide a unified framework for multiple hypotheses that have been put forth regarding the origin of neurodegenerative disease including inflammation, oxidative stress, metabolic dysfunction, and accumulation of protein bodies (see above).

Alu elements have played a pivotal role in the evolution of the human epigenome (Prendergast et al. 2014), and both hypermethylation and hypomethylation of *Alu* elements have been correlated with a number of age-related disorders including Alzheimer's disease, multiple sclerosis, osteoporosis, and many forms of cancer (Bollati et al. 2009; Jintaridith and Mutirangura 2010; Belancio et al. 2010; Jintaridith et al. 2013; Neven et al. 2016). In light of these patterns, as well as the newly discovered regulatory roles of *Alu* elements (Polak and Domany 2006; Chen and Carmichael 2008; Chen and Yang 2017), we recommend additional research that focuses on the epigenetic interplay between *Alu* elements and mitochondrial gene networks in the central nervous system.

Acknowledgements We thank A.D. Brown, W.K. Gottschalk, and M. Mihovilovic for helpful discussion. C. Hohoff, B. Ciapa, and an anonymous reviewer kindly reviewed the manuscript and provided helpful comments. This is Duke Lemur Center publication number 1389. We dedicate this work to the memory of Allen Roses, who believed there was no such thing as “junk DNA”.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Ade C, Roy-Engel AM, Deininger PL (2013) *Alu* elements: an intrinsic source of human genome instability. *Curr Opin Virol*. 3:639–645

- Antonell A, Gelpi E, Sánchez-Valle R et al (2012) Breakpoint sequence analysis of an A β PP locus duplication associated with autosomal dominant Alzheimer's disease and severe cerebral amyloid angiopathy. *J Alzheimers Dis* 28:303–3088
- Arnoldi A, Tonelli A, Crippa F et al (2008) A clinical, genetic, and biochemical characterization of SPG7 mutations in a large cohort of patients with hereditary spastic paraplegia. *Hum Mutat* 29:522–531
- Arpawong TE, Pendleton N, Mekli K et al (2017) Genetic variants specific to aging-related verbal memory: insights from GWASs in a population-based cohort. *PLoS One* 12:e0182448
- Athanasiadis A, Rich A, Maas S (2004) Widespread A-to-I RNA editing of Alu-containing mRNAs in the human transcriptome. *PLoS Biol* 2:e391
- Baillie JK, Barnett MW, Upton KR et al (2011) Somatic retrotransposition alters the genetic landscape of the human brain. *Nature* 479:534–537
- Bajad P, Jantsch MF, Keegan L, O'Connell M (2017) A to I editing in disease is not fake news. *RNA Biol* 14:1223–1231
- Batzler MA, Deininger PL (2002) Alu repeats and human genomic diversity. *Nat Rev Genet* 3:370–379
- Bazak L, Haviv A, Barak M et al (2014) A-to-I RNA editing occurs at over a hundred million genomic sites, located in a majority of human genes. *Genome Res* 24:365–376
- Bedrosian TA, Linker S, Gage FH (2016) Environment-driven somatic mosaicism in brain disorders. *Genome Med* 8:58
- Behm M, Öhman M (2016) RNA editing: a contributor to neuronal dynamics in the mammalian brain. *Trends Genet* 32:165–175
- Belancio VP, Roy-Engel AM, Deininger PL (2010) All y'all need to know 'bout retroelements in cancer. *Semin Cancer Biol* 20:200–210
- Bhattacharjee A, Yang H, Duffy M et al (2016) The activity of menkes disease protein ATP7A is essential for redox balance in mitochondria. *J Biol Chem* 291:16644–16658
- Bitar M, Barry G (2017) Multiple innovations in genetic and epigenetic mechanisms cooperate to underpin human brain evolution. *Mol Biol Evol* msx303. <https://doi.org/10.1093/molbev/msx303>
- Bollati V, Schwartz J, Wright R et al (2009) Decline in genomic DNA methylation through aging in a cohort of elderly subjects. *Mech Ageing Dev* 130:234–239
- Bondurand N, Fouquet V, Baral V et al (2012) Alu-mediated deletion of SOX10 regulatory elements in Waardenburg syndrome type 4. *Eur J Hum Genet* 20:990–994. <https://doi.org/10.1038/ejhg.2012.29>
- Bondy-Chomey E, Crawford Parks TE, Ravel-Chapuis A et al (2016) *Staufen1* regulates multiple alternative splicing events either positively or negatively in DMI indicating its role as a disease modifier. *PLoS Genet* 12:e1005827. <https://doi.org/10.1371/journal.pgen.1005827>
- Bonifati V, Rizzu P, van Baren MJ et al (2003) Mutations in the DJ-1 gene associated with autosomal recessive early-onset parkinsonism. *Science* 299:256–259. <https://doi.org/10.1126/science.1077209>
- Boone PM, Yuan B, Campbell IM et al (2014) The Alu-rich genomic architecture of SPAST predisposes to diverse and functionally distinct disease-associated CNV alleles. *Am J Hum Genet* 95:143–161
- Borchert GM, Gilmore BL, Spengler RM et al (2009) Adenosine deamination in human transcripts generates novel microRNA binding sites. *Hum Mol Genet* 18:4801–4807. <https://doi.org/10.1093/hmg/ddp443>
- Bouchet C, Vuillaumier-Barrot S, Gonzales M et al (2007) Detection of an Alu insertion in the POMT1 gene from three French Walker Warburg syndrome families. *Mol Genet Metab* 90:93–96. <https://doi.org/10.1016/j.ymgme.2006.09.005>
- Britten RJ (1997) Mobile elements inserted in the distant past have taken on important functions. *Gene* 205:177–182
- Brooks EM, Branda RF, Nicklas JA, O'Neill JP (2001) Molecular description of three macro-deletions and an Alu-Alu recombination-mediated duplication in the HPRT gene in four patients with Lesch-Nyhan disease. *Mutat Res* 476:43–54
- Bundo M, Toyoshima M, Okada Y et al (2014) Increased L1 retrotransposition in the neuronal genome in schizophrenia. *Neuron* 81:306–313. <https://doi.org/10.1016/j.neuron.2013.10.053>
- Burggren AC, Mahmood Z, Harrison TM et al (2017) Hippocampal thinning linked to longer TOMM40 poly-T variant lengths in the absence of the APOE ϵ 4 variant. *Alzheimer's Dement* 13:739–748. <https://doi.org/10.1016/j.jalz.2016.12.009>
- Cai Q, Davis ML, Sheng Z-H (2011) Regulation of axonal mitochondrial transport and its impact on synaptic transmission. *Neurosci Res* 70:9–15. <https://doi.org/10.1016/j.neures.2011.02.005>
- Calvo SE, Clauser KR, Mootha VK (2015) MitoCarta2.0: an updated inventory of mammalian mitochondrial proteins. *Nucleic Acids Res* 44:D1251–D1257. <https://doi.org/10.1093/nar/gkv1003>
- Castelnuovo M, Massone S, Tasso R et al (2010) An Alu-like RNA promotes cell differentiation and reduces malignancy of human neuroblastoma cells. *FASEB J* 24:4033–4046. <https://doi.org/10.1096/fj.10-157032>
- Checler F, Goiran T, da Costa AC (2017) Presenilins at the crossroad of a functional interplay between PARK/PARKIN and PINK1 to control mitophagy: implications for neurodegenerative diseases. *Autophagy* 15:1–2. <https://doi.org/10.1080/15548627.2017.1363950>
- Chen L-L, Carmichael GG (2008) Gene regulation by SINES and inosines: biological consequences of A-to-I editing of Alu element inverted repeats. *Cell Cycle* 7:3294–3301. <https://doi.org/10.4161/cc.7.21.6927>
- Chen L-L, Carmichael GG (2009) Altered nuclear retention of mRNAs containing inverted repeats in human embryonic stem cells: functional role of a nuclear noncoding RNA. *Molecular Cell* 35:467–478
- Chen W, Schuman E (2016) Circular RNAs in brain and other tissues: a functional enigma. *Trends Neurosci* 39:597–604. <https://doi.org/10.1016/j.tins.2016.06.006>
- Chen L-L, Yang L (2017) ALU alternative regulation for gene expression. *Trends Cell Biol* 27:480–490. <https://doi.org/10.1016/j.tcb.2017.01.002>
- Choi B-O, Kim NK, Park SW et al (2011) Inheritance of Charcot-Marie-Tooth disease 1A with rare nonrecurrent genomic rearrangement. *Neurogenetics* 12:51–58. <https://doi.org/10.1007/s10048-010-0272-3>
- Chong-Chong X, Denton KR, Wang Z et al (2016) Abnormal mitochondrial transport and morphology as early

- pathological changes in human models of spinal muscular atrophy. *Dis Model Mech* 9:39–49
- Conceição Pereira M, Loureiro JL, Pinto-Basto J et al (2012) Alu elements mediate large SPG11 gene rearrangements: further spatacsin mutations. *Genet Med* 14:143–151. <https://doi.org/10.1038/gim.2011.7>
- Cope TE, Rittman T, Borchert RJ, et al. (2018) Tau burden and the functional connectome in Alzheimer's disease and progressive supranuclear palsy. *Brain*, awx347, [doi.org:10.1093/brain/awx347](https://doi.org/10.1093/brain/awx347)
- Cordaux R, Batzer MA (2009) The impact of retrotransposons on human genome evolution. *Nat Rev Genet* 10:691–703. <https://doi.org/10.1038/nrg2640>
- Coskun P, Wyrembak J, Schriener SE et al (2012) A mitochondrial etiology of Alzheimer and Parkinson disease. *Biochim Biophys Acta (BBA)-General Subj* 1820:553–564
- D'Andrea MR, Nagele RG, Wang HY et al (2001) Evidence that neurones accumulating amyloid can undergo lysis to form amyloid plaques in Alzheimer's disease. *Histopathology* 38: 120–134. <https://doi.org/10.1046/j.1365-2559.2001.01082.x>
- Daniel C, Silberberg G, Behm M, Öhman M (2014) Alu elements shape the primate transcriptome by cis-regulation of RNA editing. *Genome Biol* 15:R28. <https://doi.org/10.1186/gb-2014-15-2-r28>
- Dannlowski U, Kugel H, Redlich R et al (2014) Serotonin transporter gene methylation is associated with hippocampal gray matter volume. *Hum Brain Mapp* 35:5356–5367. <https://doi.org/10.1002/hbm.22555>
- Dawson G, Kilkus J, Siakotos AN et al (1995) Mitochondrial abnormalities in CLN2 and CLN3 forms batten disease. *Mol Chem. Neuro* 29:227–235
- De Andrade A, Wang M, Bonaldo MF et al (2011) Genetic and epigenetic variations contributed by Alu retrotransposition. *BMC Genomics* 12:617
- De Felice FG, Lourenco MV (2015) Brain metabolic stress and neuroinflammation at the basis of cognitive impairment in Alzheimer's disease. *Front Aging Neurosci* 7:94. <https://doi.org/10.3389/fnagi.2015.00094>
- De La Monte SM, Tong M (2014) Brain metabolic dysfunction at the core of Alzheimer's disease. *Biochem. Pharmacol.* 88: 548–559
- De La Monte SM, Wands JR (2008) Alzheimer's disease is type 3 diabetes—evidence reviewed. *J Diabetes Sci Technol* 2: 1101–1113. <https://doi.org/10.1177/193229680800200619>
- De La Monte SM, Tong M, Lester-Coll N et al (2006) Therapeutic rescue of neurodegeneration in experimental type 3 diabetes: relevance to Alzheimer's disease. *J Alzheimers Dis* 10:89–109. <https://doi.org/10.2165/11597760-000000000-00000>
- Deininger P (2011) Alu elements: know the SINEs. *Genome Biol* 12:1
- Deininger PL, Batzer MA (1999) Alu repeats and human disease. *Mol Genet Metab* 67:183–193
- Deininger PL, Moran JV, Batzer MA, Kazazian HH (2003) Mobile elements and mammalian genome evolution. *Curr Opin Genet Dev* 13:651–658
- Devi L, Prabhu BM, Galati DF et al (2006) Accumulation of amyloid precursor protein in the mitochondrial import channels of human Alzheimer's disease brain is associated with mitochondrial dysfunction. *J Neurosci* 26:9057–9068
- Dixon JR, Selvaraj S, Yue F et al (2012) Topological domains in mammalian genomes identified by analysis of chromatin interactions. *Nature* 485:376–380
- Dobrovolsky R, Nazarenko I, Kim J et al (2011) Detection of large gene rearrangements in X-linked genes by dosage analysis: identification of novel α -galactosidase A (GLA) deletions causing Fabry disease. *Hum Mutat* 32:688–695. <https://doi.org/10.1002/humu.21474>
- Dong R, Ma XK, Chen LL, Yang L (2017) Increased complexity of circRNA expression during species evolution. *RNA Biol* 14:1064–1074
- Eisenberg E, Nemzer S, Kinar Y et al (2005) Is abundant A-to-I RNA editing primate-specific? *Trends Genet* 21:77–81. <https://doi.org/10.1016/j.tig.2004.12.005>
- Elbarbary RA, Lucas BA, Maquat LE (2016) Retrotransposons as regulators of gene expression. *Science* 351:aac7247
- Erwin JA, Marchetto MC, Gage FH (2014) Mobile DNA elements in the generation of diversity and complexity in the brain. *Nat Rev Neurosci* 15:497–506
- Evrny GD (2016) One brain, many genomes. *Science* 354:557–558
- Ferrari L, Scuvera G, Tucci A et al (2017) Identification of an atypical microdeletion generating the RNF135-SUZ12 chimeric gene and causing a position effect in an NF1 patient with overgrowth. *Hum Genet* 136:1329–1339. <https://doi.org/10.1007/s00439-017-1832-5>
- Ferrick-Kiddie EA, Rosenthal JJC, Ayers GD, Emeson RB (2017) Mutations underlying Episodic Ataxia type-1 antagonize Kv1.1 RNA editing. *Sci Rep* 7:41095. <https://doi.org/10.1038/srep41095>
- Flippo KH, Strack S (2017) An emerging role for mitochondrial dynamics in schizophrenia. *Schizophr. Res.* 187:26–32
- Floris G, Zhang L, Follesa P, Sun T (2017) Regulatory role of circular RNAs and neurological disorders. *Mol Neurobiol* 54: 156–165
- Gallo A, Vukic D, Michalík D et al (2017) ADAR RNA editing in human disease; more to it than meets the I. *Hum Genet* 136: 1265–1278. <https://doi.org/10.1007/s00439-017-1837-0>
- Gallus GN, Cardaioli E, Rufa A et al (2010) Alu-element insertion in an OPA1 intron sequence associated with autosomal dominant optic atrophy. *Mol Vis* 10:178–183
- Gal-Mark N, Shallev L, Sweetat S et al (2017) Abnormalities in A-to-I RNA editing patterns in CNS injuries correlate with dynamic changes in cell type composition. *Sci Rep* 7: 43421. <https://doi.org/10.1038/srep43421>
- GBD 2015 Neurological Disorders Collaborator Group 2015 NDC (2017) Global, regional, and national burden of neurological disorders during 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet Neurol* 16: 877–897. [https://doi.org/10.1016/S1474-4422\(17\)30299-5](https://doi.org/10.1016/S1474-4422(17)30299-5)
- Giroto S, Cendron L, Bisaglia M et al (2014) DJ-1 is a copper chaperone acting on SOD1 activation. *J Biol Chem* 289: 10887–10899. <https://doi.org/10.1074/jbc.M113.535112>
- Glasser MF, Coalson TS, Robinson EC et al (2016) A multi-modal parcellation of human cerebral cortex. *Nature* 536:171–178. <https://doi.org/10.1038/nature18933>
- Glinsky GV (2018) Contribution of transposable elements and distal enhancers to evolution of human-specific features of interphase chromatin architecture in embryonic stem cells. *Chromosome Res* <https://doi.org/10.1007/s10577-018-9571-6>

- Goodier JL (2016) Restricting retrotransposons: a review. *Mob DNA* 7:16. <https://doi.org/10.1186/s13100-016-0070-z>
- Gottschalk WK, Lutz MW, He YT et al (2014) The broad impact of TOM40 on neurodegenerative diseases in aging. *J Parkinsons Dis Alzheimers Dis* 1
- Greenbaum L, Springer RR, Lutz MW et al (2014) The TOMM40 poly-T rs10524523 variant is associated with cognitive performance among non-demented elderly with type 2 diabetes. *Eur Neuropsychopharmacol* 24:1492–1499
- Grover D, Majumder PP, Rao CB, et al (2003) Nonrandom distribution of *Alu* elements in genes of various functional categories: insight from analysis of human chromosomes 21 and 22. *Mol Biol Evol* 20:1420–1424. doi: <https://doi.org/10.1093/molbev/msg153>
- Gu Y, Kodama H, Watanabe S et al (2007) The first reported case of Menkes disease caused by an *Alu* insertion mutation. *Brain Dev* 29:105–108. <https://doi.org/10.1016/j.braindev.2006.05.012>
- Gu S, Yuan B, Campbell IM et al (2015) *Alu*-mediated diverse and complex pathogenic copy-number variants within human chromosome 17 at p13.3. *Hum Mol Genet* 24:4061–4077. <https://doi.org/10.1093/hmg/ddv146>
- Gurevich I, Tamir H, Arango V et al (2002) Altered editing of serotonin 2C receptor pre-mRNA in the prefrontal cortex of depressed suicide victims. *Neuron* 34:349–356. [https://doi.org/10.1016/S0896-6273\(02\)00660-8](https://doi.org/10.1016/S0896-6273(02)00660-8)
- Hancks DC, Kazazian HH (2016) Roles for retrotransposon insertions in human disease. *Mob DNA* 7(1)
- Harbauer AB, Zahedi RP, Sickmann A et al (2014) The protein import machinery of mitochondria—a regulatory hub in metabolism, stress, and disease. *Cell Metab* 19:357–372
- Hardy J, Higgins G (1992) Alzheimer's disease: the amyloid cascade hypothesis. *Science* (80-) 256:184–185. <https://doi.org/10.1126/science.1566067>
- Hedges DJ, Callinan PA, Cordaux R et al (2004) Differential *Alu* mobilization and polymorphism among the human and chimpanzee lineages. *Genome Res* 14:1068–1075
- Herrup K (2015) The case for rejecting the amyloid cascade hypothesis. *Nat Neurosci* 18:794–799. <https://doi.org/10.1038/nn.4017>
- Hoenicka J, Arrasate M, de Yébenes JG, Avila J (2002) A two-hybrid screening of human Tau protein: interactions with *Alu*-derived domain. *Neuroreport* 13:343–349
- Hofer MJ, Campbell IL (2016) Immunoinflammatory diseases of the central nervous system—the tale of two cytokines. *Br J Pharmacol* 173:716–728. <https://doi.org/10.1111/bph.13175>
- Hood JL, Emeson RB (2011) Editing of neurotransmitter receptor and ion channel RNAs in the nervous system. Springer, Berlin, Heidelberg, pp 61–90
- Hormozdiari F, Konkel MK, Prado-Martinez J et al (2013) Rates and patterns of great ape retrotransposition. *Proc Natl Acad Sci USA* 110:13457–13462. <https://doi.org/10.1073/pnas.1310914110>
- Huang LS, Ripps ME, Korman SH et al (1989) Hypobetalipoproteinemia due to an apolipoprotein B gene exon 21 deletion derived by *Alu*-*Alu* recombination. *J Biol Chem* 264:11394–11400
- Huda A, Mariño-Ramírez L, Landsman D, Jordan IK (2009) Repetitive DNA elements, nucleosome binding and human gene expression. *Gene* 436:12–22
- Hunter R, Gagnidze K, McEwen B, Pfaff D (2015) Stress and the dynamic genome: steroids, epigenetics, and the transposome. *Proc Natl Acad Sci USA* 112:6828–6833
- Hwang T, Park C-K, Leung AKL et al (2016) Dynamic regulation of RNA editing in human brain development and disease. *Nat Neurosci* 19:1093–1099. <https://doi.org/10.1038/nn.4337>
- Ittner LM, Götze J (2011) Amyloid- β and tau—a toxic pas de deux in Alzheimer's disease. *Nat Rev Neurosci* 12:65–72. <https://doi.org/10.1038/nrn2967>
- Jackson JG, Robinson MB (2017) Regulation of mitochondrial dynamics in astrocytes: Mechanisms, consequences, and unknowns. *Glia*. <https://doi.org/10.1002/glia.23252>
- Jeck WR, Sorrentino JA, Wang K et al (2013) Circular RNAs are abundant, conserved, and associated with *ALU* repeats. *RNA* 19:141–157. <https://doi.org/10.1261/ma.035667.112>
- Jiang J, Wu X, Shen D et al (2017) Analysis of RP2 and RPGR mutations in five X-linked Chinese families with retinitis pigmentosa. *Sci Rep* 7:44465. <https://doi.org/10.1038/srep44465>
- Jintaridith P, Mutirangura A (2010) Distinctive patterns of age-dependent hypomethylation in interspersed repetitive sequences. *Physiol Genomics* 41:194–200
- Jintaridith P, Tungtrongchitr R, Preutthipan S, Mutirangura A (2013) Hypomethylation of *Alu* elements in postmenopausal women with osteoporosis. *PLoS One* 8: e70386. <https://doi.org/10.1371/journal.pone.0070386>
- Jobic F, Morin G, Vincent-Delorme C et al (2017) New intragenic rearrangements in non-Finnish mulibrey nanism. *Am J Med Genet Part A* 173:2782–2788. <https://doi.org/10.1002/ajmg.a.38381>
- Johnson SC, Gonzalez B, Zhang Q et al (2017) Network analysis of mitonuclear GWAS reveals functional networks and tissue expression profiles of disease-associated genes. *Hum Genet* 136:55–65. <https://doi.org/10.1007/s00439-016-1736-9>
- Kaler SG (2011) ATP7A-related copper transport diseases—emerging concepts and future trends. *Nat Rev Neurol* 7:15–29. <https://doi.org/10.1038/nrneurol.2010.180>
- Kaneko H, Dridi S, Tarallo V et al (2011) DICER1 deficit induces *Alu* RNA toxicity in age-related macular degeneration. *Nature* 471:325–330. <https://doi.org/10.1038/nature09830>
- Kaur S, Pociot F (2015) *Alu* elements as novel regulators of gene expression in type 1 diabetes susceptibility genes? *Genes (Basel)* 6:577–591. <https://doi.org/10.3390/genes6030577>
- Kawahara Y, Ito K, Sun H et al (2004) Glutamate receptors: RNA editing and death of motor neurons. *Nature* 427:801. <https://doi.org/10.1038/427801a>
- Khanna MR, Kovalevich J, Lee VMY et al (2016) Therapeutic strategies for the treatment of tauopathies: Hopes and challenges. *Alzheimer's Dement*. 12:1051–1065
- Khermesh K, D'Erchia AM, Barak M et al (2016) Reduced levels of protein recoding by A-to-I RNA editing in Alzheimer's disease. *RNA* 22:290–302. <https://doi.org/10.1261/rna.054627.115>
- Kim J, Moody JP, Edgerly CK et al (2010) Mitochondrial loss, dysfunction and altered dynamics in Huntington's disease. *Hum Mol Genet* 19:3919–3935
- Kim Y, Tarallo V, Kerur N et al (2014) DICER1/*Alu* RNA dysmetabolism induces caspase-8-mediated cell death in age-related macular degeneration. *Proc Natl Acad Sci U S A* 111:16082–16087. <https://doi.org/10.1073/pnas.1403814111>

- Knight Johnson A, Schaefer GB, Lee J et al (2017) Alu-mediated deletion of PIGL in a Patient with CHIME syndrome. *Am J Med Genet Part A* 173:1378–1382. <https://doi.org/10.1002/ajmg.a.38181>
- Kubota-Sakashita M, Iwamoto K, Bundo M, Kato T (2014) A role of ADAR2 and RNA editing of glutamate receptors in mood disorders and schizophrenia. *Mol Brain* 7:5. <https://doi.org/10.1186/1756-6606-7-5>
- Kurnosov AA, Ustyugova SV, Nazarov VI et al (2015) The evidence for increased L1 activity in the site of human adult brain neurogenesis. *PLoS One* 10:e0117854. <https://doi.org/10.1371/journal.pone.0117854>
- Kuszak AJ, Jacobs D, Gurnev PA et al (2015) Evidence of distinct channel conformations and substrate binding affinities for the mitochondrial outer membrane protein translocase pore Tom40. *J Biol Chem* 290:26204–26217. <https://doi.org/10.1074/jbc.M115.642173>
- Kutsche K, Ressler B, Katzera H et al (2002) Characterization of breakpoint sequences of five rearrangements in LICAM and ABCD1 (ALD) genes. *Hum Mutat* 19:526–535
- Kwak S, Kawahara Y (2005) Deficient RNA editing of GluR2 and neuronal death in amyotrophic lateral sclerosis. *J Mol Med (Berl)* 83:110–120. <https://doi.org/10.1007/s00109-004-0599-z>
- Laperriere D, Wang T-T, White JH, Mader S (2007) Widespread Alu repeat-driven expansion of consensus DR2 retinoic acid response elements during primate evolution. *BMC Genomics* 8:23. <https://doi.org/10.1186/1471-2164-8-23>
- Lapp HE, Hunter RG (2016) The dynamic genome: transposons and environmental adaptation in the nervous system. *Epigenomics* 8:237–249
- Larsen PA, Lutz MW, Hunnicutt KE et al (2017) The Alu neurodegeneration hypothesis: a primate-specific mechanism for neuronal transcription noise, mitochondrial dysfunction, and manifestation of neurodegenerative disease. *Alzheimer's Dement*. 13:828–838
- Le Guennec K, Veugelen S, Quenez O et al (2017) Deletion of exons 9 and 10 of the presenilin 1 gene in a patient with early-onset Alzheimer disease generates longer amyloid seeds. *Neurobiol Dis* 104:97–103. <https://doi.org/10.1016/j.nbd.2017.04.020>
- Lehnert S, Van Loo P, Thilakarathne PJ et al (2009) Evidence for co-evolution between human microRNAs and Alu-repeats. <https://doi.org/10.1371/journal.pone.0004456>
- Lerner T, Boustany RMN, Anderson JW et al (1995) Isolation of a novel gene underlying Batten disease, CLN3. *Cell* 82:949–957
- Lev-Maor G, Sorek R, Shomron N, Ast G (2003) The birth of an alternatively spliced exon: 3' splice-site selection in Alu exons. *Science* 300:1288–1291
- Li JB, Church GM (2013) Deciphering the functions and regulation of brain-enriched A-to-I RNA editing. *Nat Neurosci* 16:1518–1522. <https://doi.org/10.1038/nn.3539>
- Li TH, Schmid CW (2001) Differential stress induction of individual Alu loci: implications for transcription and retrotransposition. *Gene* 276:135–141
- Li W, Jin Y, Prazak L et al (2012) Transposable elements in TDP-43-mediated neurodegenerative disorders. *PLOS One*: e44099, 1–e44010
- Li Z, Tian Y, Tian N et al (2015) Aberrant alternative splicing pattern of ADAR2 downregulates adenosine-to-inosine editing in glioma. *Oncol Rep* 33:2845–2852. <https://doi.org/10.3892/or.2015.3907>
- Lin MT, Beal MF (2006) Mitochondrial dysfunction and oxidative stress in neurodegenerative diseases. *Nature* 443:787–795
- Lin L, Shen S, Tye A et al (2008) Diverse splicing patterns of exonized Alu elements in human tissues. *PLoS Genet* 4:e1000225. <https://doi.org/10.1371/journal.pgen.1000225>
- Lin L, Jiang P, Park JW et al (2016) The contribution of Alu exons to the human proteome. *Genome Biol* 17:15
- Linker SB, Marchetto MC, Narvaiza I et al (2017) Examining non-LTR retrotransposons in the context of the evolving primate brain. *BMC Biol* 15:68. <https://doi.org/10.1186/s12915-017-0409-z>
- Liscovitch N, Bazak L, Levanon EY, Chechik G (2014) Positive correlation between ADAR expression and its targets suggests a complex regulation mediated by RNA editing in the human brain. *RNA Biol* 11:1447–1456. <https://doi.org/10.4161/15476286.2014.992286>
- López E, Casanovas C, Giménez J et al (2015) Characterization of Alu and recombination-associated motifs mediating a large homozygous SPG7 gene rearrangement causing hereditary spastic paraplegia. *Neurogenetics* 16:97–105. <https://doi.org/10.1007/s10048-014-0429-6>
- Lu B (2009) Mitochondrial dynamics and neurodegeneration. *Curr Neurol Neurosci Rep* 9:212
- Luco RF (2016) Retrotransposons jump into alternative-splicing regulation via a long noncoding RNA. *Nat Struct Mol Biol* 23:952–954. <https://doi.org/10.1038/nsmb.3318>
- Lutz MW, Crenshaw DG, Saunders AM, Roses AD (2010) Genetic variation at a single locus and age of onset for Alzheimer's disease. *Alzheimer's Dement* 6:125–131
- Lyall DM, Harris SE, Bastin ME et al (2014) Alzheimer's disease susceptibility genes APOE and TOMM40, and brain white matter integrity in the Lothian Birth Cohort 1936. *Neurobiol Aging*. <https://doi.org/10.1016/j.neurobiolaging.2014.01.006>
- Maas S, Kawahara Y, Tamburro KM, Nishikura K (2006) A-to-I RNA editing and human disease. *RNA Biol* 3:1–9
- Maden M (2007) Retinoic acid in the development, regeneration and maintenance of the nervous system. *Nat Rev Neurosci* 8:755–765. <https://doi.org/10.1038/nrn2212>
- Madsen E, Gitlin JD (2007) Copper and iron disorders of the brain. *Annu Rev Neurosci* 30:317–337. <https://doi.org/10.1146/annurev.neuro.30.051606.094232>
- Mager F, Gessmann D, Nussberger S, Zeth K (2011) Functional refolding and characterization of two Tom40 isoforms from human mitochondria. *J Membr Biol* 242:11–21
- Mahley RW, Rall SC (2000) Apolipoprotein E: far more than a lipid transport protein. *Annu Rev Genomics Hum Genet* 1:507–537. <https://doi.org/10.1146/annurev.genom.1.1.507>
- Mameli E, Lepori MB, Chiappe F et al (2015) Wilson's disease caused by alternative splicing and Alu exonization due to a homozygous 3039-bp deletion spanning from intron 1 to exon 2 of the ATP7B gene. 569:276–279
- Mariner PD, Walters RD, Espinoza CA et al (2008) Human Alu RNA is a modular transacting repressor of mRNA transcription during heat shock. *Mol Cell* 29:499–509
- Martin LJ (2012) Biology of mitochondria in neurodegenerative diseases. *Prog Mol Biol Transl Sci* 107:355–415. <https://doi.org/10.1016/B978-0-12-385883-2.00005-9>

- Mattick JS, Mehler MF (2008) RNA editing, DNA recoding and the evolution of human cognition. *Trends Neurosci* 31:227–233. <https://doi.org/10.1016/j.tins.2008.02.003>
- McKhann GM, Knopman DS, Chertkow H et al (2011) The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging–Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's Dement*. 7:263–269
- Mefford HC, Zemel M, Geraghty E et al (2015) Intragenic deletions of *ALDH7A1* in pyridoxine-dependent epilepsy caused by Alu–Alu recombination. *Neurology* 85:756–762. <https://doi.org/10.1212/WNL.0000000000001883>
- Mehler MF, Mattick JS (2007) Noncoding RNAs and RNA editing in brain development, functional diversification, and neurological disease. *Physiol Rev* 87:799–823
- Mirza AH, Kaur S, Brorsson CA, Pociot F (2014) Effects of GWAS-associated genetic variants on lncRNAs within IBD and T1D candidate loci. *PLoS One*. <https://doi.org/10.1371/journal.pone.0105723>
- Mita P, Boeke JD (2016) How retrotransposons shape genome regulation. *Curr Opin Genet Dev* 37:90–100. <https://doi.org/10.1016/j.gde.2016.01.001>
- Mittal K, Mani RJ, Katara DP (2016) Type 3 diabetes: cross talk between differentially regulated proteins of type 2 diabetes mellitus and Alzheimer's disease. *Sci Rep* 6:25589. <https://doi.org/10.1038/srep25589>
- Morais S, Bastos-Ferreira R, Sequeiros J, Alonso I (2016) Genomic mechanisms underlying *PARK2* large deletions identified in a cohort of patients with PD. *Neurol Genet* 2:e73. <https://doi.org/10.1212/NXG.0000000000000073>
- Muotri AR, Chu VT, Marchetto MCN et al (2005) Somatic mosaicism in neuronal precursor cells mediated by L1 retrotransposition. *Nature* 435:903–910
- Muotri AR, Marchetto MCN, Coufal NG et al (2010) L1 retrotransposition in neurons is modulated by MeCP2. *Nature* 468:443–446. <https://doi.org/10.1038/nature09544>
- Mustajoki S, Ahola H, Mustajoki P, Kauppinen R (1999) Insertion of Alu element responsible for acute intermittent porphyria. *Hum Mutat* 13:431–438
- Neeman Y, Levanon EY, Jantsch MF, Eisenberg E (2006) RNA editing level in the mouse is determined by the genomic repeat repertoire. *RNA* 12:1802–1809. <https://doi.org/10.1261/rna.165106>
- Neven KY, Piola M, Angelici L et al (2016) Repetitive element hypermethylation in multiple sclerosis patients. *BMC Genet* 17:84. <https://doi.org/10.1186/s12863-016-0395-0>
- Nishikura K (2016) A-to-I editing of coding and non-coding RNAs by ADARs. *Nat Rev Mol Cell Biol* 17:83–96. <https://doi.org/10.1038/nrm.2015.4>
- Norris J, Fan D, Aleman C et al (1995) Identification of a new subclass of Alu DNA repeats which can function as estrogen receptor-dependent transcriptional enhancers. *J Biol Chem* 270:22777–22782. <https://doi.org/10.1074/JBC.270.39.22777>
- Nozu K, Iijima K, Ohtsuka Y et al (2014) Alport syndrome caused by a *COL4A5* deletion and exonization of an adjacent AluY. *Mol Genet genomic Med* 2:451–453
- Oakes E, Anderson A, Cohen-Gadol A, Hundley HA (2017) Adenosine deaminase that acts on RNA 3 (ADAR3) binding to glutamate receptor subunit B pre-mRNA inhibits RNA editing in glioblastoma. *J Biol Chem* 292:4326–4335. <https://doi.org/10.1074/jbc.M117.779868>
- Okubo M, Horinishi A, Saito M et al (2007) A novel complex deletion–insertion mutation mediated by Alu repetitive elements leads to lipoprotein lipase deficiency. *Mol Genet Metab* 92:229–233. <https://doi.org/10.1016/J.YMGME.2007.06.018>
- Oliver KR, Greene WK (2011) Mobile DNA and the TE-Thrust hypothesis: supporting evidence from the primates. *Mob DNA* 2:8. <https://doi.org/10.1186/1759-8753-2-8>
- Ottesen EW, Seo J, Singh NN, Singh RN (2017) A multilayered control of the human survival motor neuron gene expression by Alu elements. *Front Microbiol* 8:2252. <https://doi.org/10.3389/fmicb.2017.02252>
- Pal S, Tyler JK (2016) Epigenetics and aging. *Sci Adv* 2:e1600584–e1600584. <https://doi.org/10.1126/sciadv.1600584>
- Pandolfo M (2006) Friedreich ataxia: detection of GAA repeat expansions and frataxin point mutations. In: *Congenital Heart Disease*. Humana Press, New Jersey, pp 197–216
- Paquola ACM, Erwin JA, Gage FH (2016) Insights into the role of somatic mosaicism in the brain. *Curr Opin Syst Biol* 1:90–94
- Payer LM, Steranka JP, Yang WR et al (2017) Structural variants caused by Alu insertions are associated with risks for many human diseases. *Proc Natl Acad Sci USA* 114:E3984–E3992. <https://doi.org/10.1073/pnas.1704117114>
- Payton A, Sindrewicz P, Pessoa V et al (2016) A TOMM40 poly-T variant modulates gene expression and is associated with vocabulary ability and decline in non-pathological aging. *Neurobiol Aging* 39:217 e1–7
- Pehlivan D, Beck CR, Okamoto Y et al (2016) The role of combined SNV and CNV burden in patients with distal symmetric polyneuropathy. *Genet Med* 18:443–451. <https://doi.org/10.1038/gim.2015.124>
- Petschner P, Gonda X, Baksa D et al (2017) Genes linking mitochondrial function, cognitive impairment and depression are associated with endophenotypes serving precision medicine. *Neuroscience*. <https://doi.org/10.1016/J.NEUROSCIENCE.2017.09.049>
- Picardi E, Manzari C, Mastropasqua F et al (2015) Profiling RNA editing in human tissues: towards the inosinome Atlas. *Sci Rep* 5:14941. <https://doi.org/10.1038/srep14941>
- Picardi E, D'Erchia AM, Lo Giudice C, Pesole G (2017a) REDiportal: a comprehensive database of A-to-I RNA editing events in humans. *Nucleic Acids Res* 45:D750–D757. <https://doi.org/10.1093/nar/gkw767>
- Picardi E, Horner DS, Pesole G (2017b) Single-cell transcriptomics reveals specific RNA editing signatures in the human brain. *RNA* 23:860–865. <https://doi.org/10.1261/rna.058271.116>
- Polak P, Domany E (2006) Alu elements contain many binding sites for transcription factors and may play a role in regulation of developmental processes. *BMC Genomics* 7:133. <https://doi.org/10.1186/1471-2164-7-133>
- Prendergast JGD, Chambers EV, Semple CAM (2014) Sequence-level mechanisms of human epigenome evolution. *Genome Biol Evol* 6:1758–1771. <https://doi.org/10.1093/gbe/evu142>
- Prudencio M, Gonzales PK, Cook CN et al (2017) Repetitive element transcripts are elevated in the brain of C9orf72 ALS/FTLD patients. *Hum Mol Genet* 26:3421–3431. <https://doi.org/10.1093/hmg/ddx233>

- Prüfer K, Munch K, Hellmann I, et al (2012) The bonobo genome compared with the chimpanzee and human genomes *Nature* 486:nature11128. doi: <https://doi.org/10.1038/nature11128>
- Reddy PH (2011) Abnormal tau, mitochondrial dysfunction, impaired axonal transport of mitochondria, and synaptic deprivation in Alzheimer's disease. *Brain Res.* 1415:136–148
- Reynolds WF, Rhees J, Maciejewski D et al (1999) Myeloperoxidase polymorphism is associated with gender specific risk for Alzheimer's disease. *Exp Neurol* 155:31–41. <https://doi.org/10.1006/exnr.1998.6977>
- Rezin GT, Amboni G, Zugno AI et al (2009) Mitochondrial dysfunction and psychiatric disorders. *Neurochem Res* 34: 1021–1029. <https://doi.org/10.1007/s11064-008-9865-8>
- Rice GI, Kasher PR, Forte GMA et al (2012) Mutations in ADAR1 cause Aicardi-Goutières syndrome associated with a type I interferon signature. *Nat Genet* 44:1243–1248. <https://doi.org/10.1038/ng.2414>
- Roger AJ, Muñoz-Gómez SA, Kamikawa R (2017) The origin and diversification of mitochondria. *Curr Biol* 27:R1177–R1192. <https://doi.org/10.1016/j.cub.2017.09.015>
- Roses AD (2010) An inherited variable poly-T repeat genotype in TOMM40 in Alzheimer disease. *Arch Neurol* 67:536–541
- Roses AD, Lutz MW, Amrine-Madsen H et al (2010) A TOMM40 variable-length polymorphism predicts the age of late-onset Alzheimer's disease. *Pharmacogenomics J* 10:375–384
- Roses AD, Sundseth S, Saunders A et al (2016a) Understanding the genetics of APOE and TOMM40 and role of mitochondrial structure and function in clinical pharmacology of Alzheimer's disease. *Alzheimers Dement* 12:687–694. <https://doi.org/10.1016/j.jalz.2016.03.015>
- Roses AD, Akkari PA, Chiba-Falek O et al (2016b) Structural variants can be more informative for disease diagnostics, prognostics and translation than current SNP mapping and exon sequencing. *Expert Opin Drug Metab Toxicol* 12:135–147. <https://doi.org/10.1517/17425255.2016.1133586>
- Ross CA, Poirier MA (2004) Protein aggregation and neurodegenerative disease. *Nat Med*:S10–S17
- Rossi L, Lombardo MF, Ciriolo MR, Rotilio G (2004) Mitochondrial dysfunction in neurodegenerative diseases associated with copper imbalance. *Neurochem. Res.* 29:493–504
- Rugarli EI, Langer T (2012) Mitochondrial quality control: a matter of life and death for neurons. *EMBO J* 31:1336–1349
- Rybak-Wolf A, Stottmeister C, Glazar P et al (2015) Circular RNAs in the mammalian brain are highly abundant, conserved, and dynamically expressed. *Mol Cell* 58:870–885
- Sakurai M, Ueda H, Yano T et al (2014) A biochemical landscape of A-to-I RNA editing in the human brain transcriptome. *Genome Res* 24:522–534. <https://doi.org/10.1101/gr.162537.113>
- Saunders AM, Schmeider K, Breitner JC et al (1993) Apolipoprotein E epsilon 4 allele distributions in late-onset Alzheimer's disease and in other amyloid-forming diseases. *Lancet (London, England)* 342:710–711. [https://doi.org/10.1016/0140-6736\(93\)91709-U](https://doi.org/10.1016/0140-6736(93)91709-U)
- Schanze D, Neubauer D, Cormier-Daire V et al (2014) Deletions in the 3' part of the *NFIX* gene including a recurrent Alu-mediated deletion of exon 6 and 7 account for previously unexplained cases of Marshall-Smith syndrome. *Hum Mutat* 35:1092–1100. <https://doi.org/10.1002/humu.22603>
- Schmauss C, Howe JR (2002) RNA editing of neurotransmitter receptors in the mammalian brain. *Sci STKE* 2002:pe26. <https://doi.org/10.1126/stke.2002.133.pe26>
- Schneider I, Kugel H, Redlich R et al (2017) Association of serotonin transporter gene *AluJb* methylation with major depression, amygdala responsiveness, 5-HTTLPR/rs25531 polymorphism, and stress. *Neuropsychopharmacology online early*:1–9. <https://doi.org/10.1038/npp.2017.273>
- Schollen E, Keldermans L, Foulquier F et al (2007) Characterization of two unusual truncating *PMM2* mutations in two CDG-Ia patients. *Mol Genet Metab* 90:408–413. <https://doi.org/10.1016/j.ymgme.2007.01.003>
- Schwahn U, Lenzner S, Dong J et al (1998) Positional cloning of the gene for X-linked retinitis pigmentosa 2. *Nat Genet* 19: 327–332. <https://doi.org/10.1038/1214>
- Schwarz TL (2013) Mitochondrial trafficking in neurons. *Cold Spring Harb Perspect Biol* 5:a011304
- Seeley WW (2017) Mapping neurodegenerative disease onset and progression. *Cold Spring Harb Perspect Biol* 9:a023622. <https://doi.org/10.1101/cshperspect.a023622>
- Selkoe DJ (2000) Toward a comprehensive theory for Alzheimer's disease. Hypothesis: Alzheimer's disease is caused by the cerebral accumulation and cytotoxicity of amyloid beta-protein. *Ann N Y Acad Sci* 924:17–25. <https://doi.org/10.1111/j.1749-6632.2000.tb05554.x>
- Shen S, Lin L, Cai JJ et al (2011) Widespread establishment and regulatory impact of *Alu* exons in human genes. *Proc Natl Acad Sci* 108:2837–2842
- Shiota T, Imai K, Qiu J et al (2015) Molecular architecture of the active mitochondrial protein gate. *Science* 349:1544–1548
- Shpyleva S, Melnyk S, Pavliv O et al (2017) Overexpression of LINE-1 retrotransposons in autism brain. *Mol Neurobiol*:1–10. <https://doi.org/10.1007/s12035-017-0421-x>
- Skovronsky DM, Doms RW, Lee VM (1998) Detection of a novel intraneuronal pool of insoluble amyloid beta protein that accumulates with time in culture. *J Cell Biol* 141:1031–1039
- Smalheiser NR (2014) The RNA-centred view of the synapse: non-coding RNAs and synaptic plasticity. *Philos Trans R Soc B Biol Sci* 369:20130504–20130504. <https://doi.org/10.1098/rstb.2013.0504>
- Soibam B (2017) Super-lncRNAs: identification of lncRNAs that target super-enhancers via RNA:DNA:DNA triplex formation. *RNA* 23:1729–1742
- Steen E, Terry BM, Rivera EJ et al (2005) Impaired insulin and insulin-like growth factor expression and signaling mechanisms in Alzheimer's disease—is this type 3 diabetes? *J Alzheimers Dis* 7:63–80
- Strittmatter WJ, Saunders AM, Schmechel D et al (1993) Apolipoprotein E: high-avidity binding to beta-amyloid and increased frequency of type 4 allele in late-onset familial Alzheimer disease. *Proc Natl Acad Sci* 90:1977–1981. <https://doi.org/10.1073/pnas.90.5.1977>
- Sun Z, Wu Y, Ordog T et al (2014) Aberrant signature methylation by DNMT1 hot spot mutation in hereditary sensory and autonomic neuropathy 1E. *Epigenetics* 9:1184–1193. <https://doi.org/10.4161/epi.29676>
- Swerdlow RH (2017) Mitochondria and mitochondrial cascades in Alzheimer's disease. *J Alzheimer's Dis Preprint*:1–14. <https://doi.org/10.3233/JAD-170585>

- Swerdlow RH, Burns JM, Khan SM (2010) The Alzheimer's disease mitochondrial cascade hypothesis. *J Alzheimer's Dis* 20:265
- Szpakowski S, Sun X, Lage JM et al (2009) Loss of epigenetic silencing in tumors preferentially affects primate-specific retroelements. *Gene* 448:151–167. <https://doi.org/10.1016/j.gene.2009.08.006>
- Takahashi RH, Milner TA, Li F et al (2002) Intraneuronal Alzheimer β 242 accumulates in multivesicular bodies and is associated with synaptic pathology. *Am J Pathol* 161: 1869–1879
- Tan B, Huang H, Lam R, Soong T (2009) Dynamic regulation of RNA editing of ion channels and receptors in the mammalian nervous system. *Mol Brain* 2:13. <https://doi.org/10.1186/1756-6606-2-13>
- Tan MH, Li Q, Shanmugam R et al (2017) Dynamic landscape and regulation of RNA editing in mammals. *Nature* 550:249–254. <https://doi.org/10.1038/nature24041>
- Tarallo V, Hirano Y, Gelfand BD et al (2012) DICER1 loss and *Alu* RNA induce age-related macular degeneration via the NLRP3 inflammasome and MyD88. *Cell* 149:847–859. <https://doi.org/10.1016/j.cell.2012.03.036>
- Tariq A, Jantsch MF (2012) Transcript diversification in the nervous system: A to I RNA editing in CNS function and disease development. *Front Neurosci* 1–13. <https://doi.org/10.3389/fnins.2012.00099>
- Taşkesen M, Collin GB, Evsikov AV et al (2012) Novel *Alu* retrotransposon insertion leading to Alström syndrome. *Hum Genet* 131:407–413. <https://doi.org/10.1007/s00439-011-1083-9>
- Thongsroy J, Patchesung M, Mutirangura A (2017) The association between *Alu* hypomethylation and severity of type 2 diabetes mellitus. *Clin Epigenetics* 9:93. <https://doi.org/10.1186/s13148-017-0395-6>
- Töhönen V, Katayama S, Vesterlund L et al (2015) Novel PRD-like homeodomain transcription factors and retrotransposon elements in early human development. *Nat Commun* 6:8207. <https://doi.org/10.1038/ncomms9207>
- Tsirigos A, Rigoutsos I (2009) *Alu* and B1 repeats have been selectively retained in the upstream and intronic regions of genes of specific functional classes. *PLOS Comput Biol* 5: e1000610. <https://doi.org/10.1371/JOURNAL.PCBI.1000610>
- Udaka T, Okamoto N, Aramaki M et al (2007) An *Alu* retrotransposition-mediated deletion of CHD7 in a patient with CHARGE syndrome. *Am J Med Genet Part A* 143: 721–726. <https://doi.org/10.1002/ajmg.a.31441>
- Ullu E, Tschudi C (1984) *Alu* sequences are processed 7SL RNA genes. *Nature* 312:171–172. <https://doi.org/10.1038/312171a0>
- Van Den Heuvel MP, Sporns O (2013) Network hubs in the human brain. *Trends Cogn Sci* 17:683–696. <https://doi.org/10.1016/j.tics.2013.09.012>
- Vandewalle J, Bauters M, Van Esch H et al (2013) The mitochondrial solute carrier SLC25A5 at Xq24 is a novel candidate gene for non-syndromic intellectual disability. *Hum Genet* 132:1177–1185
- Vansant G, Reynolds WF (1995) The consensus sequence of a major *Alu* subfamily contains a functional retinoic acid response element. *Proc Natl Acad Sci U S A* 92:8229–8233
- Varizhuk AM, Sekridova AV, Tankevich MV et al (2016) Conformational polymorphism of G-rich fragments of DNA *Alu*-repeats. II. the putative role of G-quadruplex structures in genomic rearrangements. *Biomeditsinskaya Khimiya* 62:630–637. <https://doi.org/10.18097/pbmc20166206630>
- Varshney D, Vavrova-Anderson J, Oler AJ, et al (2015) SINE transcription by RNA polymerase III is suppressed by histone methylation but not by DNA methylation.
- Vassetzky NS, Ten OA, Kramerov DA (2003) B1 and related SINEs in mammalian genomes. *Gene* 319:149–160
- Viader A, Golden JP, Baloh RH et al (2011) Schwann cell mitochondrial metabolism supports long-term axonal survival and peripheral nerve function. *J Neurosci* 31:10128–10140. <https://doi.org/10.1523/JNEUROSCI.0884-11.2011>
- Viader A, Sasaki Y, Kim S et al (2013) Aberrant Schwann cell lipid metabolism linked to mitochondrial deficits leads to axon degeneration and neuropathy. *Neuron* 77:886–898. <https://doi.org/10.1016/j.neuron.2013.01.012>
- Wang W, Li L, Lin W et al (2013) The ALS disease-associated mutant TDP-43 impairs mitochondrial dynamics and function in motor neurons. *Hum Mol Genet* 22:4706–4719
- Wang W, Wang WH, Azadzi KM et al (2016) *Alu* RNA accumulation in hyperglycemia augments oxidative stress and impairs eNOS and SOD2 expression in endothelial cells. *Mol Cell Endocrinol* 426:91–100
- Weissmann D, van der Laan S, Underwood MD et al (2016) Region-specific alterations of A-to-I RNA editing of serotonin 2c receptor in the cortex of suicides with major depression. *Transl Psychiatry* 6:e878. <https://doi.org/10.1038/tp.2016.121>
- Wimmer K, Callens T, Wernstedt A, Messiaen L (2011) The NF1 gene contains hotspots for L1 endonuclease-dependent De Novo insertion. *PLoS Genet*. <https://doi.org/10.1371/journal.pgen.1002371>
- Wright A, Vissel B (2012) The essential role of AMPA receptor GluR2 subunit RNA editing in the normal and diseased brain. *Front Mol Neurosci* 5:34. <https://doi.org/10.3389/fnfmol.2012.00034>
- Wu S-J, Hsieh T-J, Kuo M-C, et al (2013) Functional regulation of *Alu* element of human angiotensin-converting enzyme gene in neuron cells. *Neurobiol Aging* 34:1921.e1–1921921.e7. doi: <https://doi.org/10.1016/j.neurobiolaging.2013.01.003>
- Zempel H, Mandelkow E (2015) Tau missorting and spastin-induced microtubule disruption in neurodegeneration: Alzheimer disease and hereditary spastic paraplegia. *Mol Neurodegener* 10:68. <https://doi.org/10.1186/s13024-015-0064-1>
- Zhang Y-H, Dipple KM, Vilain E et al (2000) *AluY* insertion (IVS4-52ins316alu) in the glycerol kinase gene from an individual with benign glycerol kinase deficiency. *Hum Mutat* 15:316–323. [https://doi.org/10.1002/\(SICI\)1098-1004\(200004\)15:4<316::AID-HUMU3>3.0.CO;2-9](https://doi.org/10.1002/(SICI)1098-1004(200004)15:4<316::AID-HUMU3>3.0.CO;2-9)
- Zhang G, Fukao T, Sakurai S et al (2006) Identification of *Alu*-mediated, large deletion-spanning exons 2–4 in a patient with mitochondrial acetoacetyl-CoA thiolase deficiency. *Mol Genet Metab* 89:222–226. <https://doi.org/10.1016/j.ymgme.2006.06.010>