

# Enhancing carbon sequestration in soil with coal combustion products: a technology for minimising carbon footprints in coal-power generation and agriculture

Isa A. M. Yunusa<sup>1</sup> · Graeme Blair<sup>1</sup> · Ayalsew Zerihun<sup>2</sup> ·  
Shenjiao Yang<sup>1,3</sup> · Susan C. Wilson<sup>1</sup> · Iain M. Young<sup>1</sup>

Received: 22 May 2014 / Accepted: 15 March 2015 / Published online: 18 April 2015  
© Springer Science+Business Media Dordrecht 2015

**Abstract** Coal-fired power generation and agriculture account for more than half of global greenhouse gas emissions, but the coal fly ash (CFA) produced in the former can be a resource for reducing emissions from agriculture to minimise environmental footprints in both industries. Our aim in this study was to test how acidic and alkaline CFA addition could minimise loss of C and N from acidic soil, with or without added manure. We determined composition and structural characteristics of acidic and alkaline CFA for their capacity to adsorb organic carbon, but observed poor adsorption because of low concentrations of cenospheres and unburnt carbon as the primary adsorbents in the ash. Addition of CFA had no impact on the loss of carbon or nitrogen from unmanured soil in which concentrations of these nutrients were low. Loss of carbon from manured soil was reduced by 36 % with alkaline ashes and by 3-fold with acidic ashes; while loss of N was 30–50 % lower with acidic ashes, but 28 % higher with alkaline ashes, compared with no ash treatment. The increases in C sparing with CFA addition were achieved not by direct C absorption but by restraining microbial population and respiration, and potentially emissions. Alkaline CFA increased soil pH and if used to substitute just 10 % of lime for ameliorating soil acidity would reduce CO<sub>2</sub> emission associated with the mining of the lime and its eventual dissolution in soil by ~2.66 Tg or 2.8 % of Australia's annual agricultural emissions. High concentrations of oxides of phosphorus, silicon, titanium and clay particles in acidic ashes, and oxides of cations in alkaline ashes, were associated with potential for promoting C storage and acidity amelioration in soil.

---

✉ Isa A. M. Yunusa  
isa.yunusa@une.edu.au

<sup>1</sup> School of Environmental & Rural Sciences, University of New England, Armidale, NSW 2351, Australia

<sup>2</sup> Department of Environment & Agriculture, Curtin University, Bentley, WA 6102, Australia

<sup>3</sup> Farmland Irrigation Research Institute, Chinese Academy of Agricultural Sciences, Xinxiang, Henan 453003, People's Republic of China

## 1 Introduction

Power generation and agriculture both generate substantial environmental footprints that pose significant risk to the long-term viability of these two industrial sectors. Greenhouse gas (GHG) emission is a major component of the environmental footprint for which the two industries account for ~80 % and ~7 %, respectively, of the total global GHG emissions in 2011 (UNFCCC 2014). The majority of the emission in the energy sector arises from burning of fossil fuel. Coal contributes about 40 % of total power generation (IAEA 2012) but accounts for 56 % of the emission from the sector (ASN Bank/Ecofys 2013). A substantial part of the emissions from the agriculture sector emanate from soils which are responsible for up to 54 % of agricultural emissions in the USA, 18 % in Australia, 25 % in China, and 34 % in Brazil (UNFCCC 2014). Most of this is associated with the use of fertilisers and soil amendments (Garnaut 2011).

Between 5 and 15 % of coal fed into boiler remains unburnt forming solid residues that are collectively termed Coal Combustion Products (CCPs). The majority of these residue (>85 %) comprises of fine particles (<0.2 mm) referred to as coal fly ash (Heidrich et al. 2013) of which only about half of the 780 Mg global production is effectively used. The unused CFA stockpiles grow by 370 Mg annually, and along with gaseous emissions, has to be sustainably managed as part of the *Clean Coal Technology* (Seshadri et al. 2013). Potential use of coal fly ash (CFA) to ameliorate soil constraints on agricultural productivity, along with potential risks, has been extensively reviewed (Seshadri et al. 2013; Shaheen et al. 2014; Yunusa et al. 2012). Protocols for logistical, economic, monitoring and regulatory imperatives for routine agricultural application of CFA have been proposed (Yunusa et al. 2012). However, with the exception of a recent study (Masto et al. 2014), little attention has been given to potential application of CCPs for mitigating GHG emissions from agriculture through promotion of C and N storage in soil proposed earlier (Amonette et al. 2004; Palumbo et al. 2004).

Furthermore, the C in limestone and dolomite used to ameliorate soil acidity on farms is inevitably released as CO<sub>2</sub> to the atmosphere (IPCC 2006). This emission accounts for about 9 Tg CO<sub>2</sub> yearly in the United States of America (USEPA 2013). Liming contributes up to 7 % of total greenhouse gas (GHG) emission from a typical wheat farm in Australia (Brock et al. 2012) and would account for about 30 % of the total agricultural emissions by 2015 in Germany (Benndorf 2013). These farm emissions could be reduced by substituting liming products with CFA that contains non-oxidisable C and which reportedly increased pH of treated acidic soils by up to 2 units (Manoharan et al. 2010; Yunusa et al. 2013). Soil N is transformed into potentially volatile and/or soluble forms and easily lost from the soil during mineralisation of organic matter, and should be considered in strategies for abating farm emissions.

The CFA rich in lime CaO ( $\geq 3.0$  %) can directly adsorb CO<sub>2</sub> on to their particles through carbonation. Up to 7 % w/w of CO<sub>2</sub> was absorbed through this process under natural conditions over a 20- year period, but the process is enhanced at high temperatures and pressure (60–90 °C; ~4.0 MPa) in the laboratory (Muriithi et al. 2013). Particulate organic C can also be absorbed onto ash particles, the unburnt C and/or cenospheres; the latter are ash particles that burst under high temperatures forming hollow spheres with large surface areas (Amonette et al. 2004; Palumbo et al. 2004). However, CFAs contain low amounts of cenospheres (<4% w/w) and unburnt C (Ngu et al. 2007) so that large amounts of ash would be required to achieve significant absorption of particulate C. Hence both carbonation and adsorption of organic C onto particles in CFA are therefore likely to have negligible short-term capacity for soil C sequestration.

Increased C storage in soil with CFA amendment is more likely to be indirect through alterations of processes and properties in the soil, such as aggregation, pH, salinity, and, where the ash contains significant amounts of unburnt C, the C/N (Palumbo et al. 2004; Amonette et al. 2004). Increases in pH due to ash addition reportedly enhanced humification in analytical organic monomers (Amonette et al. 2004), which along with reduced mineralisation stabilised peroxidase and tryrosinase activities and enhanced humification in ash-amended soil (Masto et al. 2014). Improved soil aggregation with ash amendment (Yunusa et al. 2011) should protect organic C from microbial decomposers (Beare et al 1994; Jastrow et al. 2007; Rabbi et al., 2013; Young and Ritz 2000).

In the current study, we determined the potential of acidic and alkaline CFA for enhancing storage of C and N in acidic soil with or without incorporation of sheep manure. Our objectives were to (1) quantify cenosphere and unburnt C contents of the CFA and their capacity to adsorb particulate organic C, (2) determine depletion of C and N from an acidic soil, (3) identify chemical properties that influence loss of C and N from ash-amended soil, and (4) identify knowledge-gaps in developing CFA technology for enhancing C and N storage in soils.

## 2 Materials and methods

Five CFAs were obtained from power stations in Western Australia, South Australia, Victoria, New South Wales and Queensland, in Australia. Three of the ashes were alkaline and the other two acidic; henceforth denoted as acidic1, acidic2, alkaline1, alkaline2 and alkaline3, and all five had low salinity except alkaline3 (Table 1) and used in the following experiments.

### 2.1 Ash characterisation and adsorption of organic carbon

Cenospheres were harvested by dispersing ash samples in distilled water in 1:20 (w/v) ash/water at room temperature, and then followed by a 24-hour sink-float separation. Chemical composition of the cenosphere was determined in a two-step procedure of fusion and analysis prescribed in the Australian Standard AS 1038.14.1–2003 and reported by Yip et al. (2010). A scanning electron microscope (SEM) was used to determine morphological characteristics of cenospheres, while the unburnt C was quantified using thermogravimetric analysis (TGA) following a procedure reported (Ngu et al. 2007).

Adsorption of organic carbon by the CFA cenospheres was determined in a batch adsorption reactor system involving shaking 0.35 g of cenosphere in 200 ml of synthetic water containing methylene blue (10 ppm). The dye concentration in the solutions was determined before and after adsorption using a Spectrophotometer (Spectronic 20 Genesys<sup>TM</sup>, Thermo Scientific, Melbourne, Australia) to measure absorbance at  $\lambda_{\max}$  of 665 nm for methylene blue (Wang et al. 2008).

### 2.2 Soil incubation experiment

An acidic sandy loam soil (Yellow Chromosol) equivalent to a Haplic Xerosol (FAO 1974) was collected from the UNE research farm, Armidale (30° 29' 16" S, 151° 38' 29" E); it belongs to the order Alfisol in the USDA classification (Soil Survey Staff 2012) and had been under lightly fertilized mixture of native pastures. It was amended

**Table 1** Selected chemical properties and elemental composition of the five coal fly ashes, the soil and manure used in this study

Properties	Acidic1	Acidic2	Alkaline1	Alkaline2	Alkaline3	Soil <sup>1</sup>	Manure
<i>Basic chemical properties and particle size distribution</i>							
pH	3.3	4.0	10.7	10.8	9.0	6.3	7.40
EC (dS/m)	1.10	0.14	0.70	0.54	19.0	0.08	10.03
Total C	0.11	0.68	0.04	2.10	0.45	1.03	34.04
Total N (%)	0.001	0.007	0.003	0.018	0.012	0.056	3.39
C/N	110	97	13	117	38	18	10
$\delta^{13}\text{C}$ (‰)	-22.84	-25.38	-24.77	-24.22	-24.32	-18.69	-23.02
Clay (%)	8.9	4.2	5.4	2.4	3.7	60.9	na
Silt (%)	78.2	68.0	66.3	51.9	45.5	23.2	na
Sand (%)	13.1	27.8	28.3	45.9	50.8	15.9	na
<i>Chemical composition (%)</i>							
SiO <sub>2</sub>	50.1	69	61.3	63.6	8.18	nd	nd
Al <sub>2</sub> O <sub>3</sub>	26.5	24	28.1	25.9	5.34	nd	nd
Fe <sub>2</sub> O <sub>3</sub>	13.5	1.7	2.42	2.83	33.1	nd	nd
K <sub>2</sub> O	0.64	0.44	3.01	3.58	0.24	nd	nd
Na <sub>2</sub> O	0.29	0.01	1.16	0.94	5.92	nd	nd
CaO	1.84	0.12	1.64	1.36	8.68	nd	nd
MgO	1.14	0.22	0.60	0.76	13.76	nd	nd
TiO <sub>2</sub>	1.15	1.7	0.87	0.79	0.26	nd	nd
Mn <sub>3</sub> O <sub>4</sub>	0.09	0.01	0.05	0.05	0.80	nd	nd
P <sub>2</sub> O <sub>5</sub>	1.43	0.1	0.24	0.09	0.08	nd	nd
SO <sub>3</sub>	0.37	0.2	0.52	nd	14	nd	nd
BaO	0.32	0.1	0.06	0.06	nd	nd	nd
SrO	nd	0.01	0.04	0.04	nd	nd	nd
Total	97.37	97.61	100.00	100.00	90.36	na	na

<sup>1</sup> determined on fresh soil soon after collection; *na* not applicable, *nd* not determined

with sheep manure and/or CFA for this study. The sheep manure was collected from a local shearing shed. The soil, manure and CFA were air dried, and the soil and manure were ground to pass a 2 mm sieve, and mixed in plastic containers to form the following factorial treatments:

- i. Ash amendment: 85 % soil (153 g) + 15 % ash (27 g) + 0% manure
- ii. Manure amendment: 85 % soil (153 g) + 15 % manure (27 g) + 0 % ash
- iii. Ash & manure amendment: 70 % soil (126 g) + 15 % ash (27 g) + 15 % manure (27 g)
- iv. Control: 100 % soil (180 g) + 0 % ash + 0 % manure

The soil or mixtures were blended uniformly and then weighed at 180 g into 0.5 L clear plastic containers (80 mm x 110 mm x 90 mm) with each treatment prepared in 4 replicates. The containers were arranged randomly on benches in a glasshouse maintained at 22 °C and humidity at ~50 %. The treatments were gradually wetted to bring their weights to predetermined field capacity and weight recorded. The treatments were covered with a black

shade-cloth to minimise evaporation, and brought back to their prescribed weights with addition of water once a week for 3 months.

### 2.2.1 Measurements

Carbon and nitrogen content and their stable isotopes. After three months, half of the sample in each jar was taken by cutting cleanly vertically through the soil; the remaining half was stored in a fridge to be used for microbial analysis. The samples were air-dried and sieved to pass 2.0 mm screen, and used to determine salinity as electrical conductivity (EC) and pH using a portable bench-top pH-EC meter (LabCHEM Model 901-CP, TPS Pty Ltd, Australia) on filtrate of air dried sample mixed into deionised water (1:5). Both total C (TC) and total N (TN) were measured on 5 g samples in a high temperature furnace (LECO®, St Joseph, MI, USA), while  $\delta^{13}\text{C}$  was determined using a mass spectrometer (ANCA-GSL IRMS, Sercon Ltd., Cheshire, UK). The  $\delta^{13}\text{C}$  isotope composition of the samples was computed as (IAEA 2001).

$$\delta^{13}\text{C} (\text{‰}) = \left[ \frac{R_{\text{sample}}}{R_{\text{standard}}} - 1 \right] 10^3 \quad (1)$$

in which  $R$  represents the mass ratio of  $^{13}\text{C}/^{12}\text{C}$  for the sample or 0.0112372 for the standard. The  $\delta^{13}\text{C}$  for the mixtures at the start of incubation was calculated thus:

$$\delta^{13}\text{C}_{\text{mixture at day 0}} = f_{\text{soil}} \delta^{13}\text{C}_{\text{soil}} + f_{\text{manure}} \delta^{13}\text{C}_{\text{manure}} + f_{\text{ash}} \delta^{13}\text{C}_{\text{ash}} \quad (2)$$

where  $f$  represents fractions of C for either the soil, manure or ash in the mixture. The TC and TN lost during the incubation period were estimated from their quantities ( $Q$ ) at the start (day 0) and after the 3-month incubation ( $Q_3$ ):

$$Q_{\text{lost}} = Q_0 - Q_3 \quad (3)$$

The quantities of TC and TN at the start of the incubation were estimated from the masses of soil, CFA and manure in the mixtures, and the respective concentrations of the two nutrients:

$$Q_0 = \sum_{i=1}^{n=3} (m_i \times c_i) \quad (4)$$

where  $m_i$  are the masses of the component in the substrates,  $i$ , (soil, CFA and manure), and  $c_i$  are their respective TC or TN concentrations;  $n$  is the number of components (1–3) in the mixture. The  $Q_3$  values were obtained from the product of the mass of the incubation substrate and its measured TC or TN values.

**Microbial biomass and respiration** These were measured on the stored half of the samples using microplate-based respiration technique (MicroResp™, Craigiebukler, Aberdeen, UK) as per Campbell et al. (2003). Basal respiration rate (BSR) was monitored over three days after which 0.22 g of glucose was stirred into the soil and monitored for a further 24 hours to determined substrate-induced respiration and the dependent microbial biomass.

## 2.3 Data analyses

Data on cenosphere and unburnt carbon in the ashes were evaluated in a one-way analysis of variance. All data from the incubation experiment were subjected to two-way analysis of variance (ANOVA); statistical significance was determined when  $p \leq 0.05$ . Two sets of principal component analyses (PCA) were undertaken to examine (i) interrelationships amongst TC and TN, EC, pH and microbial biomass and BSR at the completion of the incubation period, and (ii) whether TC and TN post-incubation could be associated with specific properties of the CFA. Both ANOVA and PCA were carried out using IBM SPSS Statistics, V22.0 (IBM Corp, Armonk, NY).

## 3 Results

### 3.1 Physico-chemical properties of the ashes and their cenosphere yields

The five ashes consisted mostly of silt- and sand-size particles, whose mineral composition was dominated by  $\text{SiO}_2$  and  $\text{Al}_2\text{O}_3$  (Table 1). The  $\text{SiO}_2$  and  $\text{Al}_2\text{O}_3$  accounted for  $>70\%$  in four ashes, but  $<20\%$  in alkaline3 in which  $\text{Fe}_2\text{O}_3$  dominated (30 %). Other oxides, especially  $\text{CaO}$ ,  $\text{Na}_2\text{O}_5$ ,  $\text{MgO}$ ,  $\text{Mn}_3\text{O}_4$ ,  $\text{SO}_3$ , were generally higher in alkaline3 than in the other ashes.

Only alkaline1 and alkaline2 yielded any cenospheres with an average of 0.5 % w/w. The cenospheres had smooth round and spherical shapes (Fig. 1) with  $\sim 30 - 150 \mu\text{m}$  diameter. Close-up views of individual cenospheres reveals macro-size ( $>10 \mu\text{m}$ ) pores and finer spherical ash particles adhered onto the surface. The amount of unburnt C in the cenospheres was generally small,  $\sim 1.8\%$  w/w in alkaline1 and  $\sim 0.2\%$  in alkaline2.


### 3.2 Carbon adsorption onto cenospheres and unburnt carbon in ash

The adsorption of methylene blue onto the cenospheres of alkaline1 and alkaline2 showed a two-phase trend (Fig. 2a): an initial exponential phase in the first 10 mins, followed by a slow linear phase in the following 4 hours. On the whole, the adsorption capacity of cenosphere after 4 hours was 0.7 mg methylene blue per gram of cenospheres. Over the same duration, the amounts of methylene blue adsorbed by alkaline2 ash directly (2.0 mg/g) or by its unburnt carbon ( $\sim 21 \text{ mg/g}$ ) were larger than adsorption by its cenospheres alone (Fig. 2b).

### 3.3 Soil incubation

#### 3.3.1 Total carbon and nitrogen concentrations in soil

Concentrations of TC and TN at the start of the incubation were larger by factors of 4 and 9, respectively, in the manured soil compared with unmanured soil (Table 2). Addition of CFA to unmanured soil did not alter TC (Fig. 3a) or TN (Fig. 3b) after incubation; however, TC was higher in the manured soil supplied with ash compared with no ash. Post-incubation TN was higher with addition of the acidic CFAs than with the alkaline CFAs. The proportions of TC and TN lost during incubation from the manured soil were generally lower with CFA than without, and also differed with ash type (Table 2). The acidic ashes were more effective than the alkaline ashes in arresting losses



**Fig. 1** Scanning electron microscope images of collection of cenospheres for (a) alkaline1 and (b) alkaline 3, and of enlarged cenospheres for (c) alkaline1 and (d) alkaline3


of C (28 % vs 43 % or 2.65 g vs 4.37 g) and N (13 % vs 39% or 0.11 g vs 0.36 g). With exception of alkaline2 that reduced N loss only marginally, addition of alkaline1 or alkaline3 increased N loss by 28 and 36 %, or 0.37 g and 0.45 g, respectively, compared with 0.29 g when no ash was added. It was not possible to partition the losses in TC and TN amongst the three sources.

The amounts of TC measured in both manured and unmanured soils post-incubation as percentage of their pre-incubation concentrations averaged 57 %; whereas proportionally more TN was lost from the manured (30 %) compared with unmanured soil (2%). The losses of both nutrients from the soils during incubation were coupled:

$$N_{\text{loss}} = 1.57C_{\text{loss}} - 29.63; r^2 = 0.94, n-2 = 18 \quad (5)$$

### 3.3.2 Microbial respiration and biomass

The BSR in unmanured soil was not affected by addition of acidic CFA, but increased with alkaline CFA (Fig. 3c). However, BSR was at least halved in manured soil with addition of any of the five ashes. Microbial biomass in unmanured soil was unaffected with acidic2 and alkaline1, and significantly with alkaline3 (Fig. 3d). Addition of any of the ashes significantly reduced microbial biomass in the manured soil; acidic1 was




**Fig. 2** Adsorption of methyl blue by cenospheres of (a) alkaline1 and alkaline2 ashes, and by (b) the alkaline2 or its unburnt carbon. The equations for the *fitted lines* are given in each graph

the least detrimental to microbial biomass. Both BSR and microbial biomass were 4- and 5- times, respectively, larger in the manured soil compared with unmanured soil.

**Table 2** Calculated total C and total N at the start of incubation of soil amended with 0 or 15 % sheep manure, and proportions lost during the 3 month incubation

Fly ash treatment	Initial total C (%)		C loss (%)	Initial total N (%)		N loss (%)
	0 %	15 %		0 %	15 %	
Control	0.907	4.730	63.6	0.049	0.434	33.4
Acidic1	0.982	4.797	22.8	0.050	0.435	9.6
Acidic2	0.899	4.723	33.1	0.049	0.435	16.4
Alkaline1	1.167	4.960	43.7	0.051	0.436	42.6
Alkaline2	0.952	4.770	39.0	0.050	0.436	30.3
Alkaline3	1.027	5.333	47.4	0.056	0.491	45.3
<i>Mean</i>	<i>0.989</i>	<i>4.940</i>	<i>41.6</i>	<i>0.051</i>	<i>0.444</i>	<i>29.6</i>





**Fig. 3** Impact of acidic or alkaline coal fly ash additions on (a) total carbon (TC), (b) total N (TN), (c) basal microbial respiration rate (BSR), and (d) microbial biomass in soil supplied with 0 or 15 % w/w of sheep manure after three-month incubation. The mean ( $\pm$  standard errors) values for 0 and 15 % manure additions are given

### 3.3.3 Basic soil chemistry after incubation

Changes in the soil chemical variables at the end of the incubation are presented in Table 3. With exception of acidic1 and alkaline3, the pH was lower with ash additions in manured soil than in the unmanured soil. The electrical conductivity (EC) in unmanured soil was increased only with addition of alkaline1 or alkaline3. Addition of any ash to the manured soil either had no impact on EC or reduced it, the exception being alkaline3 that increased it significantly. The mean pH was similar for manured and unmanured soil, while the manured soil was more saline, had more depleted  $\delta^{13}\text{C}$ , but higher C/N, than the unmanured soil.

Addition of alkaline3 enriched  $\delta^{13}\text{C}$  in unmanured soil, while addition of other ashes depleted  $\delta^{13}\text{C}$  in unmanured and manured soil (Table 3). Relative to starting conditions, CFA additions depleted  $\delta^{13}\text{C}$  in unmanured soil during incubation, the exception was alkaline3 that enriched  $\delta^{13}\text{C}$ . There was general  $\delta^{13}\text{C}$  enrichment in the manured soil during incubation due to CFA addition, the exception being alkaline2 that caused a slight depletion.


**Table 3** Impacts of ash additions on some chemical properties of soil amended with 0 or 15 % sheep manure measured in October after 3-month incubation

Fly ash treatment	pH		C/N		Electrical conductivity (dS/m)		$\delta^{13}\text{C}$ (‰) initial <sup>1</sup>		$\delta^{13}\text{C}$ (‰) final	
	0 %	15 %	0 %	15 %	0 %	15 %	0 %	15 %	0 %	15 %
Control	6.96d	7.41c	9.08d	5.97e	0.11c	1.88b	-18.69	-22.30	-19.07b	-20.91c
Acidic1	6.91d	7.41c	10.47c	9.30d	0.11c	1.75b	-18.76	-22.30	-19.27b	-21.38d
Acidic2	6.88d	6.74d	10.45c	8.86d	0.10c	1.95b	-19.30	-22.35	-19.76b	-21.20d
Alkaline1	7.77c	7.13c	9.94b	10.64c	0.18b	1.19c	-18.73	-22.30	-19.28b	-21.49d
Alkaline2	8.23b	7.65b	13.37a	10.17c	0.11c	0.98d	-19.99	-22.39	-20.72c	-22.42e
Alkaline3	9.38a	9.43a	11.82b	10.54c	1.94a	2.63a	-19.04	-22.32	-18.65a	-21.79e
Mean	7.69A	7.66A	10.85A	9.24B	0.43B	1.73B	-19.08	-22.33	-19.46A	-21.53B

<sup>1</sup> calculated with eq. 2; means for the same variable followed by different letter(s) are statistically different at  $p \leq 0.05$  for the respective variables – the small letters compare ash effects within the same response variables and capital letters compare the mean effects of manure amendment for the same response variables

3.3.4 Associations between CFA properties and post incubation total C and TN

Principal component analysis (PCA) showed that most of the total variance (82 %) was accounted by the first two principal axes of variations (Fig. 4a). The first axis (62.8 %) had high negative loadings on TC, TN, microbial biomass and basal respiration, all of which were tightly coupled, and high positive loading on  $\delta^{13}\text{C}$ . There was also a clear separation between manured (suffix \_15) and unmanured (suffix \_0) soil along the first principal component (PC1) axis, showing manured soil as having high levels of TC, TN, microbial biomass and basal respiration, and depleted  $^{13}\text{C}$ . The PC2 (19.5 % of variation) was dominated by pH and EC of



**Fig. 4** Biplots from principal component analysis showing vector loadings for relationships amongst (a) coal fly ash characteristics and TC (total C) and TN (total N) measured, and (b) measured soil variables, after 3 month incubation The ash treatments are expressed as either acidic (Acid1, Acid2) or alkaline (Alk1, Alk2, Alk3) applied to either unmanured (\_0) or manured (\_15) soil; other notations are BSR (basal respiration rate), MB (microbial biomass), and deltaC ( $\delta^{13}\text{C}$ )

the incubation substrate. It showed an inverse correlation between TC, TN and microbial biomass, and also, with BSR and with pH. Four major clusters could be discerned consisting of (i) alkaline1 and alkaline2 with 0 % manure, (ii) acidic ashes with 0 % manure, (iii) alkaline1 and alkaline2 with 0 % manure, and (iv) alkaline3 with either manure treatment.

### 3.3.5 Associations between TC, TN, microbial dynamics and chemical attributes

The PCA extracted 78 % of the total variance in the CFA composition and their correlations with post-incubation TC and TN levels (Fig. 4b). The PC1 accounted for 53 % of the total variance with high positive loadings on CaO, Mn<sub>3</sub>O<sub>4</sub>, MgO, Fe<sub>2</sub>O<sub>3</sub>, SO<sub>3</sub>, Na<sub>2</sub>O and associated EC, all of which had high negative correlations with TiO<sub>2</sub>, P<sub>2</sub>O<sub>5</sub> and SiO<sub>2</sub>. The PC2 extracted a further 25 % of the total variance, with high positive loadings on pH and negative loadings on clay content, suggesting a strong inverse association between the two. Alkaline1 was the most benign of the ashes in terms of direct impact of its properties on TC and TN.

## 4 Discussion

### 4.1 Cenosphere and unburnt carbon yields and their potential for carbon adsorption

The ashes used here generally contained low amounts of cenospheres, and the average yield of 0.5% w/w by the two alkaline ashes was within the range (0.2–3.8%) for a number of CFA in Australia (Ngu et al. 2007). Ngu et al. (2007) reported cenosphere yields of 3.8% w/w for acidic2 contrary to zero yields in the current study. This could be due to variability in the physico-chemical properties of different batches of CFA arising from the operating conditions of the boilers (Raask 1985), the source coal and other factors, including decomposition of carbonates, fusion of silicates and evaporation of pore water (Ngu et al. 2007). For instance, while the 69 % w/w of silicates for acidic2 (Table 1) was comparable to the 65 % reported earlier, the aluminate content of 24 % in the current study was well below the 35 % reported for this ash previously (Ngu et al. 2007). Alkaline3 had low concentrations of silicates and aluminates, suggesting an inherent low potential for cenosphere yield. Similarly low cenosphere production would be expected from acidic1 given its low silicate content. However, both the cenospheres and the unburnt C were poor adsorbents of methylene blue (Fig. 2). The unburnt C had superior C adsorption because its porous structure and associated surface area were preserved, while these were destroyed during melting of the cenospheres (Wang et al. 2005). Thus adsorption by unburnt carbon could be up to 9-fold greater than by CFA particles (Wang et al. 2008), consistent with the findings here (Fig. 2b).

The bulk CFA as a whole therefore has low capacity for physical adsorption of organic carbon, with absorption being only 7–10 mg C/g for Australian CFAs (Wang et al. 2008). The current study, demonstrates that, even accounting for adsorption by the unburnt carbon, CFA is unlikely to have a major impact on soil carbon sequestration since the total C contents of these ashes are generally low (<5.5 % even in acidic1, Table 1). The capacity of unburnt carbon to absorb particulate organic carbon was ≤1 % w/w for the ashes used in this study and demonstrated the poor capacity for direct adsorption of organic carbon by CFA. Therefore, any real benefit for soil C sequestration due to ash addition would accrue only from changes induced in the physico-chemical properties of ash/soil admixtures.

## 4.2 Changes in total carbon and total nitrogen during incubation

Ash addition altered the properties of the media during incubation and explained the differences in the TC and TN observed afterwards. Addition of ash clearly reduced loss of TC, which along with TN and microbial variables, were strongly and inversely correlated with  $\delta^{13}\text{C}$  (Fig. 4a). This was because the manure was the dominant source of C in the mixture and its loss enriched  $\delta^{13}\text{C}$  towards the soil value of -18.69. The general  $\delta^{13}\text{C}$  enrichment during incubation of manured soil was smaller with ash addition that also increased the C/N, compared with no ash addition (Table 3). Maintenance of high C/N with ash addition would have restrained decomposition and subsequent loss of C (Drinkwater et al. 1998). Minimal loss of C (<40 %) and N (<31 %) were observed with addition of CFA that kept the incubate pH within 6.4–7.6 range (Table 2), well below pH of 9.4 for alkaline3 that lost the most C and N. This was consistent with positive association found between acidic pH conditions and enhanced retention of C and N in soil (Bååth and Anderson 2003; Conyers et al. 2012).

The links between TN, TC and microbial variables, and also pH and the EC, are revealed along the PC-2 axis (Fig. 4a). Both TC and TN were coupled with treatment cluster consisting of acidic ashes and alkaline1 and alkaline2 in the third quadrant of PCA. These clusters could also be partly due to salinity of the ashes, in addition to the pH, that stressed the microbial populations and impaired mineralisation of the organic matter (Rietz and Haynes 2003). The clustering of ashes reflected their chemical composition (Fig. 4b) such that alkaline1 and alkaline 2 were closely associated with  $\text{K}_2\text{O}$ , while alkaline3 was associated with a suite of alkali metals ( $\text{Na}_2\text{O}$ ,  $\text{MgO}$ ,  $\text{CaO}$ ,  $\text{Mn}_3\text{O}_4$  and  $\text{Fe}_2\text{O}_3$ ). A third group consisting of the two acidic ashes was closely associated with  $\text{P}_2\text{O}_5$  and  $\text{SiO}_2$ ,  $\text{TiO}_2$  and clay particles. These two groups of oxides can then be used for identifying ashes with potential for C conservation in soil.

Losses of the two nutrients were tightly coupled (eq. 4), although on average about 30 % TN was lost compared with 42 % of TC (Table 2) possibly due to mineralisation of N lagging behind that of C, with the former reportedly commencing only after the microbes become C-starved (Weintraub and Schimel 2003). Differences in the labile fractions of the two nutrients in the manure, similarity in C/N notwithstanding, could affect their mineralisation (Blair et al. 2005), because mineralisation of C is dependent more on pH, while that of N is more on C/N (Tian et al. 2013).

The reductions of 40–70 % in the loss of C with both ash types and N with acidic ashes demonstrated the capacity of CFA to mitigate GHG emission. Although we were unable to directly quantify the fluxes, negligible methane emissions from manure-rich arable soils, such as that used here, have been reported (Hütsch 1998). Despite the neutral impacts of CFA in unmanured soils, the capacity of alkaline ashes in raising soil pH (Table 3) as found in earlier studies (Manoharan et al. 2012; Yunusa et al. 2012) means these ashes can substitute lime for treating soil acidity and thereby avoid associated emissions. For example, applying just 2 Mg/ha of lime to the 1.7 M ha of agricultural land limed annually in Australia (ABS 2014), mining the required 3.4 Tg lime would generate ~2.66 Tg  $\text{CO}_2$  using the lime characteristics ( $\text{CaO}$ ,  $\text{MgO}$ , water content) in Approach1 of the Greenhouse Gas Protocol (WRI 2008). This emission would be avoided as would the expected annual emission of 0.02 Tg  $\text{CO}_2$ , i.e. 0.059 Mg C/Mg lime (West and McBride 2005), from eventual dissolution of the soil applied lime. This 2.66 Tg emission abatement is equivalent to 2.8 % of the ~98 Tg annual emission

from the Australian agricultural sector. Acidic ash additions would also significantly mitigate N emissions from the manure treated soil by as much as two-thirds (Table 2). Aerobic decomposition of soil applied manure disposes of at least half of the N as  $\text{NH}_3$  or  $\text{N}_2\text{O}$  within the first year (Dorahy et al. 2010). Therefore, applying the manure with 3.39 % N *w/w* (Table 1) at 10 Mg/ha over 100 ha would add 3.4 Mg N to the soil and generate 1.7 Mg of gaseous N emissions; this amount would, however, be cut by between 50 and 67% with the acidic ashes. This potential emission mitigation would be even larger in countries such as the United Kingdom where up to 42 Mg/ha of manures are prescribed routine farm applications (DEFRA 2009). Application of an acidic CFA should cut N emission from such practice in the first year to <10 Mg/ha compared with the expected 21 Mg/ha.

In conclusion, use of CFA for promoting storage of carbon and nitrogen in cropping soils is a viable technology for reducing GHG emissions in agriculture and environmental foot prints in this and the coal-power generation by minimising the CFA disposed in landfill. In selecting ashes for mitigating the loss of these nutrients from soil, acidic ashes are preferred over alkaline ashes that were effective in minimising only C loss. High concentrations of  $\text{P}_2\text{O}_5$  and  $\text{SiO}_2$ ,  $\text{TiO}_2$  and clay particles in acidic ashes, and alkali/alkaline-earth metals ( $\text{Na}_2\text{O}$ ,  $\text{MgO}$ ,  $\text{CaO}$ ,  $\text{Mn}_3\text{O}_4$  and  $\text{Fe}_2\text{O}_3$ ) and sulphur in alkaline ashes, provided initial selection criteria for identifying CFAs for soil carbon sequestration. Knowledge-gaps to be addressed in developing this technology further include: (1) characterisation of the enzymatic processes in organic C mineralisation in ash amended soil, (2) major forms of N loss from the system, (3) whether similar C sparing is possible with plant residues, and (4) optimum rates and mode of ash additions.

**Acknowledgments** Cenosphere analyses and adsorption experiments were undertaken by Prof Hongwei Hu, Dr Xiangpeng Gao, Prof Shaobin Wang and their teams at Curtin University; Prof Hu also commented on the manuscript. We acknowledge the assistance of Leanne Leslie, Michael Faint, Jan Carruthers, Gabrielle Ray and Ewan Paterson, and methodology advice by Drs S.M. Rabbi, Rebecca Hailing and V. Manoharan, and Assoc Prof Brian Wilson. We thank the reviewers for their helpful comments, and the power stations and the Ash Development Association of Australia (ADAA) for their support. This study was supported with UNE Seed Grant, and evolved from an earlier project funded by ADAA and the Australian Research Council (LP0455110).

## References

- Amonette JE, Kim J, Russell CK (2004) Enhancement of soil carbon sequestration: a catalytic approach. *Am Chem Soc Div Fuel Chem* 49:366–367
- Australian Bureau of Statistics (2014). Soil management in 2011-12: National and State/Territory. [www.abs.gov.au/ausstats/abs@.nsf/Lookup/4630.0main+features112011-12](http://www.abs.gov.au/ausstats/abs@.nsf/Lookup/4630.0main+features112011-12). (Accessed: February 17, 2015)
- ASN Bank/Ecofys (2013) World GHG Emissions Flow Chart (2010) <http://www.ecofys.com/files/files/asn-ecofys-2013-world-ghg-emissions-flow-chart-2010.pdf>. (Accessed: March 29, 2015)
- Bååth E, Anderson TH (2003) Comparison of soil fungal/bacterial ratios in a pH gradient using physiological and PLFA-based techniques. *Soil Biol Biochem* 35:955–963
- Benndorf R (2013) Germany 2050 - A Greenhouse Gas-Neutral Country. <http://www.umweltbundesamt.de/en/publikationen/germany-2050-a-greenhouse-gas-neutral-country>. (Accessed: January 20, 2015)
- Blair N, Faulkner RD, Till AR, Sanchez P (2005) Decomposition of  $^{13}\text{C}$  and  $^{15}\text{N}$  labelled plant residue materials in two different soil types and its impact on soil carbon nitrogen aggregate stability and aggregate formation. *Soil Res* 43:873–886
- Brock P, Madden P, Schwenke G, Herridge D (2012) Greenhouse gas emissions profile for 1 tonne of wheat produced in Central Zone (East) New South Wales: a life cycle assessment approach. *Crop Pasture Sci* 63: 319–329

- Campbell CD, Chapman SJ, Cameron CM, Davidson MS, Potts JM (2003) A rapid microtiter plate method to measure carbon dioxide evolved from carbon substrate amendments so as to determine the physiological profiles of soil microbial communities by using whole soil. *Appl Environ Microbiol* 69:3593–3599
- Conyers M, Newton P, Condon J, Poile G, Mele P, Ash G (2012) Three long-term trials end with a quasi-equilibrium between soil C N and pH: an implication for C sequestration. *Soil Res* 50:527–535
- DEFRA (2009). Nitrate Vulnerable Zones (NVZs)—Guidance for farmers. Summary of the guidance for farmers in NVZs (Leaflet 1 - PB12736a). Department for Environment Food and Rural Affairs, UK. <http://adlib.eversite.co.uk/adlib/defra/content.aspx?doc=251225&id=251489>. (Accessed: February 25, 2015)
- Dorahy C, Dougherty W, Chan Y, Waters D (2010) Using recycled organics and manures in grain cropping systems. PrimeFact 1008. [http://www.dpi.nsw.gov.au/\\_data/assets/pdf\\_file/0005/341762/Using-recycled-organics-and-manures-in-grain-cropping-systems.pdf](http://www.dpi.nsw.gov.au/_data/assets/pdf_file/0005/341762/Using-recycled-organics-and-manures-in-grain-cropping-systems.pdf). (Accessed: February 24, 2015)
- Drinkwater LE, Wagoner P, Sarrantonio M (1998) Legume-based cropping systems have reduced carbon and nitrogen losses. *Nature* 396:262–265
- FAO (1974) Key to the FAO Soil Units <http://www.fao.org/soils-portal/soil-survey/soil-classification/fao-legend/key-to-the-fao-soil-units/en/> (Accessed: September 23, 2013)
- Garnaut R (2011) Australia's emissions in a global context Chapter 7. [http://www.garnautreview.org.au/pdf/Garnaut\\_Chapter7.pdf](http://www.garnautreview.org.au/pdf/Garnaut_Chapter7.pdf) (Accessed: February 4 2014)
- Heidrich C, Feuerborn HJ, Weir A (2013) Coal Combustion Products: a Global Perspective. <http://www.flyash.info/2013/1/71-Heidrich-Plenary-2013.pdf> (Accessed: February 4 2014)
- Hütsch BW (1998) Methane oxidation in arable soil as inhibited by ammonium, nitrite, and organic manure with respect to soil pH. *Biol Fert Soils* 28:27–35
- IAEA (2001) Use of Isotope and Radiation Methods in Soil and Water Management and Crop. International Atomic Energy Agency Nutrition Training Course Series 14. <http://www.naweb.iaea.org/nafa/swmn/public/TCS14-forward-contents.pdf>. (Accessed: January 20, 2015)
- IAEA (2012) Coal's share of global energy mix to continue rising with coal closing in on oil as world's top energy source by 2017. <http://www.iaea.org/newsroomandevents/pressreleases/2012/december/name,34441,en.html> (Accessed: January 20, 2015)
- IPCC (2006) N<sub>2</sub>O emissions from managed soils and CO<sub>2</sub> emissions from lime and urea application IPCC Guidelines for National Greenhouse Gas Inventories. [http://www.ipcc-nggip.iges.or.jp/public/2006gl/pdf/4\\_Volume4/V4\\_11\\_Ch11\\_N2O&CO2.pdf](http://www.ipcc-nggip.iges.or.jp/public/2006gl/pdf/4_Volume4/V4_11_Ch11_N2O&CO2.pdf) (Accessed: February 4, 2014)
- Jastrow JD, Amonette JE, Bailey VL (2007) Mechanisms controlling soil carbon turnover and their potential application for enhancing carbon sequestration. *Clim Chang* 80:5–23
- Manoharan V, Yunusa IAM, Loganathan P, Lawrie R, Skilbeck CG, Burchett MD, Murray BR, Eamus D (2010) Assessments of Class F coal fly ashes for amelioration of soil acidity and their influence on growth and uptake of Mo and Se by canola. *Fuel* 89:3498–3504
- Masto RE, Sengupta T, George J, Ram LC, Sunar KK, Selvi VA, Sinha AK (2014) The Impact of Fly Ash Amendment on Soil Carbon. *Energ Sour Part A: Recov Util Environ Effect* 36:554–562
- Muriithi GN, Petrik LF, Fatoba O, Gitari WM, Doucet FJ, Nel J, Nyale SM, Chuks PE (2013) Comparison of CO<sub>2</sub> capture by *ex-situ* accelerated carbonation and in *in-situ* naturally weathered coal fly ash. *J Environ Manag* 127:212–220
- Ngu LN, Wu H, Zhang DK (2007) Characterization of ash cenospheres in fly ash from Australian power stations. *Energ Fuels* 21:3437–3445
- Palumbo AV, Mccarthy JF, Amonette JE, Fisher LS, Wullschleger SD, Daniels WL (2004) Prospects for enhancing carbon sequestration and reclamation of degraded lands with fossil-fuel combustion by-products. *Adv Environ Res* 8:425–438
- Raask E (1985) Mineral impurities in coal combustion: behavior problems and remedial measures. Taylor & Francis, Washington, DC, 484p
- Rietz DN, Haynes RJ (2003) Effects of irrigation-induced salinity and sodicity on soil microbial activity. *Soil Biol Biochem* 35:845–854
- Seshadri B, Bolan NS, Naidu R, Wang H, Sajwan K (2013) Clean coal technology combustion products: properties agricultural and environmental applications and risk management. *Adv Agron* 119:310–370
- Shaheen SM, Hooda PS, Tsadilas CD (2014) Opportunities and challenges in the use of coal fly ash for soil improvements—a review. *J Environ Manag* 145:249–267
- Soil Survey Staff (2012) Soil Taxonomy <http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/survey/class/taxonomy/> (Accessed: February 24, 2014)
- Tian Y, Takamashi K, Toda H, Haibara K, Ding F (2013) pH and substrate regulation of nitrogen and carbon dynamics in forest soils in a karst region of the upper Yangtze River basin China. *J For Res* 18:228–237
- UNFCCC (2014) Greenhouse gas inventory data—detailed data by party. United Nations framework convention for climate change. <http://unfccc.int/di/DetailedByParty.do>. (Accessed: February 27 2014)

- US Environment Protection Agency (2013) Inventory of US greenhouse gas emissions and sinks: 1990–2011. <http://www.epa.gov/climatechange/Downloads/ghgemissions/US-GHG-Inventory-2013-Main-Text.pdf>. (Accessed: February 26, 2014)
- Wang S, Li L, Wu H, Zhu ZH (2005) Unburned carbon as a low-cost adsorbent for treatment of methylene blue-containing wastewater. *J Colloid Interface Sci* 292:336–343
- Wang S, Ma Q, Zhu ZH (2008) Characteristics of coal fly ash and adsorption application. *Fuel* 87:3469–3473
- Weintraub MN, Schimel JP (2003) Interactions between carbon and nitrogen mineralization and soil organic matter chemistry in arctic tundra soils. *Ecosystems* 6:129–143
- West TO, McBride AC (2005) The contribution of agricultural lime to carbon dioxide emissions in the United States: dissolution, transport, and net emissions. *Agric Ecosyst Environ* 108:145–154
- World Resource Institute (2008) Greenhouse Gas Protocol: CO<sub>2</sub> emissions from the production of lime. <http://www.ghgprotocol.org/calculation-tools/lime-sector>. (accessed: February 14, 2015)
- Yip K, Tian F, Hayashi JI, Wu H (2010) Effect of alkali and alkaline earth metallic species on biochar reactivity and syngas compositions during steam gasification. *Energ Fuels* 24:173–181
- Young IM, Ritz K (2000) Tillage habitat space and function of soil microbes. *Soil Tillage Res* 53:201–213
- Yunusa IAM, Manoharan V, Odeh IO, Shrestha S, Skilbeck CG, Eamus D (2011) Structural and hydrological alterations of soil due to addition of coal fly ash. *J Soils Sediments* 11:423–431
- Yunusa IAM, Loganathan P, Nissanka SP, Manoharan V, Burchett MD, Skilbeck CG, Eamus D (2012) Application of coal fly ash in agriculture: a strategic perspective. *Crit Rev Environ Sci Technol* 42:559–600
- Yunusa IAM, Manoharan V, Harris R, Lawrie R, Pal Y, Quito JT, Bell R, Eamus D (2013) Differential growth and yield by canola (*Brassica napus* L) and wheat (*Triticum aestivum* L) arising from alterations in chemical properties of sandy soils due to additions of fly ash. *J Sci Food Agric* 93:995–1002