


## Natural hazards in Australia: heatwaves

S. E. Perkins-Kirkpatrick<sup>1</sup>  · C. J. White<sup>2,3</sup> ·  
L. V. Alexander<sup>1</sup> · D. Argüeso<sup>1</sup> · G. Boschat<sup>4</sup> ·  
T. Cowan<sup>5,6</sup> · J. P. Evans<sup>1</sup> · M. Ekström<sup>7</sup> ·  
E. C. J. Oliver<sup>8,9</sup> · A. Phatak<sup>10</sup> · A. Purich<sup>1,5</sup>

Received: 20 September 2015 / Accepted: 4 March 2016 / Published online: 11 April 2016  
© Springer Science+Business Media Dordrecht 2016

**Abstract** As part of a special issue on natural hazards, this paper reviews the current state of scientific knowledge of Australian heatwaves. Over recent years, progress has been made in understanding both the causes of and changes to heatwaves. Relationships between atmospheric heatwaves and large-scale and synoptic variability have been identified, with increasing trends in heatwave intensity, frequency and duration projected to continue throughout the 21st century. However, more research is required to further our understanding of the dynamical interactions of atmospheric

---

This article is part of a Special Issue on “The effect of historical and future climate changes on natural hazards in Australia” edited by Seth Westra, Chris White and Anthony Kiem.

---

✉ S. E. Perkins-Kirkpatrick  
Sarah.Kirkpatrick@unsw.edu.au

<sup>1</sup> Climate Change Research Centre & ARC Centre of Excellence for Climate System Science, UNSW, Sydney, Australia

<sup>2</sup> School of Engineering and ICT, University of Tasmania, Hobart, Australia

<sup>3</sup> Antarctic Climate and Ecosystems Cooperative Research Centre, University of Tasmania, Hobart, Australia

<sup>4</sup> School of Earth Sciences and ARC Centre of Excellence for Climate System Science, Melbourne University, Melbourne, Australia

<sup>5</sup> CSIRO Oceans and Atmosphere, Aspendale, Australia

<sup>6</sup> School of GeoSciences, The University of Edinburgh, Edinburgh, UK

<sup>7</sup> CSIRO Land and Water, Canberra, Australia

<sup>8</sup> Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Australia

<sup>9</sup> ARC Centre of Excellence for Climate System Science, University of Tasmania, Hobart, Australia

<sup>10</sup> Department of Mathematics and Statistics, Curtin University, Bentley, Australia

heatwaves, particularly with the land surface. Research into marine heatwaves is still in its infancy, with little known about driving mechanisms, and observed and future changes. In order to address these knowledge gaps, recommendations include: focusing on a comprehensive assessment of atmospheric heatwave dynamics; understanding links with droughts; working towards a unified measurement framework; and investigating observed and future trends in marine heatwaves. Such work requires comprehensive and long-term collaboration activities. However, benefits will extend to the international community, thus addressing global grand challenges surrounding these extreme events.

## 1 Introduction

Heatwaves are natural hazards that have substantial impacts on human health, the economy and environment, occurring in the atmosphere and the ocean. They are Australia's most deadly natural hazard, causing 55 % of all natural disaster related deaths (Coates et al. 2014) and burden the Australian workforce by ~US\$6.2 billion every year (Zander et al. 2015). The January 2009 Victorian heatwave killed over 370 people (Alexander and Tebaldi 2012) with an insured loss of US\$1.3 billion (Munich Re 2009). Heatwaves are also a key contributor to bushfires.

Heatwaves also occur in the marine environment; for example, the 2011 Western Australia marine heatwave (Pearce and Feng 2013) had substantial impacts on marine biodiversity (Wernberg et al. 2013). Extreme heat events also impact agriculture and aquaculture industries, respectively harming grain harvest yields (Barlow et al. 2013) and reducing livestock in salmon farming.

Despite their importance, research into atmospheric heatwaves in Australia is generally lagging behind the global effort. Recent studies over Europe have demonstrated how the land interacts with synoptic systems (e.g. Fischer et al. 2012; Quesada et al. 2012) to influencing heatwave variability. Moreover, several studies have indicated that anthropogenic forcing has contributed to specific European events (e.g. Stott et al. 2004), while others indicate increases in the frequency of future heatwaves under greenhouse conditions (Orlowsky and Seneviratne 2012). However, a unified approach in understanding and characterizing atmospheric heatwaves in Australia is currently missing. This is lacking despite improved understanding of relationships between heatwaves and large-scale modes of climate variability (Parker et al. 2014a; Perkins et al. 2015) dominant synoptic patterns (Pezza et al. 2012) and increases in heatwave frequency since the 1950s (Indian Ocean Climate Initiative 2012; Perkins and Alexander 2013). In the case of marine heatwaves, only a handful of studies focus on the dynamics and impacts of specific events (Oliver et al. 2014a; Benthuisen et al. 2014), with a measurement framework only recently proposed (Hobday et al. 2016).

This paper reviews the scientific literature on the measurement, causes, observed trends and future projections of both atmospheric and marine heatwaves across Australia. We conclude with principal findings and provide key recommendations on future research priorities. While atmospheric heatwaves also occur during cooler seasons. The focus here is limited to austral summer heatwaves when the scale of impacts are generally larger.

## 2 Understanding heatwaves

### 2.1 Measuring atmospheric heatwaves


Atmospheric heatwaves are often classified as prolonged periods of excessive heat (Perkins and Alexander 2013), although no universal definition exists. They can be measured using different characteristics – such as intensity, frequency, duration, timing and spatial extent – and calculated using daily maximum, minimum, or average temperatures (e.g. Furrer et al. 2010; Fischer and Schär 2010; Russo et al. 2014). In most Australian studies, a relative threshold (or percentile) is used to determine excessive heat, where prolonged periods of heat last for at least 3 days (Tryhorn and Risbey 2006; Alexander and Arblaster 2009; Pezza et al. 2012). A relative threshold is useful since what is considered extreme in one location and/or time of year may not be extreme under other circumstances (Perkins and Alexander 2013).

The Australian Bureau of Meteorology has devised the Excess Heat Factor (EHF; Nairn and Fawcett 2013) index that takes account how hot a three-day period is compared to the previous month, as well as the climatological 95th percentile. Furthermore, a multi-characteristic framework has been developed (Perkins and Alexander 2013), employing five metrics of heatwave intensity, frequency and duration (see Fischer and Schär 2010) for three different definitions. This approach allows for a more consistent analysis, whilst providing useful information to a range of impacts communities, and is similar to the “hot-spell” (periods of extreme heat similar to heatwaves) approach of Furrer et al. (2010), where heatwave characteristics are modelled as a function of covariates, for example, time.

### 2.2 Large-scale mechanisms of atmospheric heatwaves

Recently, there have been Australian and international advances in understanding drivers and mechanisms of atmospheric heatwaves (e.g. Loikith and Broccoli 2012; Horton et al. 2015; Krueger et al. 2015; Grotjahn et al. 2015). Figure 1 explains how physical mechanisms underpinning atmospheric heatwaves over various timescales may interact in the lead-up to a heatwave. Several studies have examined the relationship between modes of large-scale climate variability and land surface temperatures across Australia (Nicholls et al. 1996; Jones and Trewin 2000; Arblaster and Alexander 2012). While the El Niño-Southern Oscillation (ENSO) is regarded as the primary large-scale driver of inter-annual variations of Australian rainfall (Risbey et al. 2009), its role on temperature extremes is more varied (e.g. Arblaster and Alexander 2012; Min et al. 2013). Significantly more heatwave days, and longer and more intense events are observed over northern and eastern Australia during El Niño compared to La Niña (Perkins et al. 2015), yet different relationships occur in the southeast (Trewin 2009; Parker et al. 2014a; Boschat et al. 2015). White et al. (2013a) found that the Indian Ocean Dipole has a positive relationship over southern Australia for austral winter and spring.

Southeastern Australia heatwaves are associated with the Madden Julian Oscillation (MJO) phases 3–6 during austral summer (Parker et al. 2014a), yet during spring, MJO phases 2–3 are more influential (Marshall et al. 2013). Over most of Australia, the likelihood of extreme temperature increases during negative phases of the Southern Annular Mode (Marshall et al. 2013), but relationships with summertime heatwaves are less clear (Perkins et al. 2015). Large-scale teleconnections to sea surface temperature (SST) and atmospheric conditions are also suggested (e.g. Pezza et al. 2012).



**Fig. 1** Heatwave schematic illustrating the various physical processes contributing to heatwaves, the interactions and feedbacks existing between them, and timescales on which they operate. Note that not all processes need to be present for a heatwave to occur (Fischer et al. 2007; Miralles et al. 2014). Coloured shading indicates the severity of a heatwave (red being more severe), the arrow thickness on the x-axis indicates the temporal length of key mechanisms, and the arrow on the y-axis indicates how the mechanisms on their various timescales may amplify heatwave severity. For example, A particular phase of climate modes may increase the likelihood of heatwaves, which then become more severe once other, shorter time-scale mechanisms (e.g. dry soil and high-pressure systems) also occur

### 2.3 Atmospheric heatwave meteorology and land surface influences

The most important weather system for Australian heatwaves is the persistent anticyclone, positioned adjacent to the affected area (Marshall et al. 2013) and largely associated with planetary-scale Rossby waves (Pezza et al. 2012; Parker et al. 2014b). Anticyclonic high-pressure systems bring warm air from the interior of the continent to the heatwave affected area, sustaining conditions for a number of days (Steffen et al. 2014). For southeastern Australia, anticyclone systems are generally centred over the Tasman Sea in line with the subtropical ridge (Hudson et al. 2011; Marshall et al. 2013). Parker et al. (2014b) found an association with propagating and overturning Rossby waves, dynamically influencing the development of southeast heatwaves. Across the southwest, anticyclone systems are typically centred over the Great Australian Bight (Pezza et al. 2012). Other influential features include intra-seasonal drivers of variability (Marshall et al. 2013; White et al. 2013a), rainfall deficits (Nicholls 2004), the Australian monsoon and tropical cyclones (Parker et al. 2013). Mechanisms of extreme-heat build-up include advection from lower latitudes, large-scale subsidence transporting higher potential temperature air from upper levels, or development and replacement of the diurnal mixed layer (McBride et al. 2009).

International studies have shown that the land surface provides important feedbacks (including the albedo, surface roughness and soil moisture; e.g. Miralles et al. 2012, 2014) that can exacerbate or dampen heatwave intensity (Seneviratne et al. 2006). Kala et al. (2015)

demonstrated the impact of soil moisture on the meteorology of the 2009 Black Saturday heatwave, highlighting the significant contribution desiccated soil can have on extreme heat events. Such studies are important, as a better understanding of the strength of these feedback mechanisms may allow for improved land cover management, potentially reducing heatwave severity (Davin et al. 2014). This may be particularly important in urban environments where the urban heat island effect can compound extreme temperature (Argüeso et al. 2014, 2015).

## 2.4 Measuring marine heatwaves

Marine heatwaves are often measured by the magnitude of ocean temperature anomalies above the monthly seasonal climatology (e.g. Pearce and Feng 2013). Temperature anomalies for specific events have been reported on weekly, daily and finer time scales, using satellite measurements and data loggers (e.g. Olita et al. 2007; Mills et al. 2013). Other studies use more sophisticated metrics including a period (~ 3 to 5 days) where ocean temperatures were at least 3–5 °C above average (Sorte et al. 2010), thermal stress anomalies (Selig et al. 2010) or degree-heating weeks (Gleeson and Strong 1995). Extreme ocean temperatures have been examined using the frequency of days above the 95th percentile (Lima and Wethey 2012) and extreme value theory (Oliver et al. 2014a, b). A standardized definition has been recently constructed, based on consecutive exceedances of the calendar day 90th percentile of temperature for at least five consecutive days (Hobday et al. 2016). From this definition, a set of metrics are computed that measure marine heatwave intensity, duration, cumulative intensity and rate of onset/decline.

## 2.5 Large-scale mechanisms of marine heatwaves

Large-scale mechanisms of marine heatwaves are less well understood. ENSO is known to play a role in driving temperature events such as the unprecedented 2011 “Ningaloo Niño” (Pearce and Feng 2013), whereby La Niña conditions drove a stronger than average Leeuwin Current southward along the coast of Western Australia (Kataoka et al. 2014). Off the southeast coast, mesoscale eddies from instabilities in the East Australian Current drive marine heatwaves along the continental shelf (Oliver et al. 2014a). In regions such as coastal South Australia (e.g. Kämpf et al. 2004) and New South Wales (NSW) (e.g. Roughan and Middleton 2004), local winds drive temperature variations due to upwelling and downwelling processes. Globally, high atmospheric temperatures and low winds commonly drive marine heatwaves and this relationship can be expected to hold around Australia (e.g. Olita et al. 2007; Pearce and Feng 2013).

## 3 Observed changes

### 3.1 Observed changes and attribution of atmospheric heatwaves


The continentally averaged Australian mean temperature has increased by 0.9 °C since 1950, which is slightly higher than the combined ocean-land global average of 0.85 °C (Bureau of Meteorology 2012), though it is worth noting that globally averaged land-only temperatures have warmed twice as fast as the combined average. There have been various assessments of Australian extreme temperature trends (e.g. Tryhorn and Risbey 2006; Alexander and

Arblaster 2009; Trewin and Vermont 2010; Pezza et al. 2012; Donat et al. 2013), however heatwave characteristics – and the metrics used to define them – can vary markedly, limiting consistent comparisons.

Heatwave characteristics have increased across many Australian regions since the mid 20th century (Alexander and Arblaster 2009; Donat et al. 2013; Perkins and Alexander 2013). Over 1971–2008, the hottest day in a heatwave increased faster than average heatwave intensity, with a measurable increase in the duration and frequency of heatwaves (Perkins and Alexander 2013). Similar patterns are found when extending the analysis to 1950–2013 (Steffen et al. 2014; see Fig. 2). Throughout southwest Western Australia the frequency and intensity of hot spells (periods of relative extreme heat that can occur throughout the year) increased over 1958–2010, but their duration decreased slightly (Indian Ocean Climate Initiative 2012). Over the same period, inland areas of northwest Western Australia experienced increases in intensity, frequency, and duration, but along coastal areas, intensity decreased. While emerging studies are explaining the dynamic/thermodynamic components of changes in Northern Hemisphere extreme temperatures (e.g. Horton et al. 2015), similar studies have not been undertaken.

Classically, studies analysing the role of human influence on observed extreme temperature events are based on monthly/seasonal anomalies for large spatial domains (e.g. Stott et al. 2004). In Australia, the intensity of the 2012/2013 summer was five times more likely to occur in a climate under the influence of anthropogenic greenhouse gases, compared to a climate without these influences (Lewis and Karoly 2013). Moreover, Australia's hottest spring on record (2013) would not have occurred without anthropogenic influence (Lewis and Karoly 2014; Knutson et al. 2014). Although attribution studies are specific to the event and domain analysed, there is evidence that a relationship exists between larger-scale, longer-term extreme temperature anomalies, and those over smaller spatial and temporal scales (Angéil et al. 2014). Hence, the studies of Lewis and Karoly are indicative that a human signal exists in observed heatwaves over smaller domains and shorter temporal scales. Indeed, the intensity and frequency of heatwaves during the 2012/2013 Australian summer increased in occurrence

**Fig. 2** Observed trends in Australian heatwave days over 1950–2013. A heatwave day must belong to a period of three or more consecutive days that have positive excess heat values (Nairn and Fawcett 2013). Hatching indicates statistical significance at the 5 % level. Updated from Perkins and Alexander (2013)



by respectively two- and three-fold respectively due to anthropogenic influence (Perkins et al. 2014a). While the aforementioned studies employed the same methodology (fraction of attributable risk, see Allen 2003), other methods also exist for determining anthropogenic influence (Allen and Tett 1999; Kovic et al. 2014). Such analyses have looked at long-term trends in daily extreme temperatures at global and continental scales (e.g. Kim et al. 2015), but have not yet been applied to Australian heatwaves.

### 3.2 Recent unprecedented heatwave events across Australia

Australia has experienced extreme heatwaves during the last decade. In late January and early February 2009, a severe heatwave occurred over Victoria that was followed by the most devastating bushfires in Australian history (the “Black Saturday” fires). Several records were set for high day- and night-time temperatures as well as for the duration of the event (National Climate Centre 2009). The heatwave occurred in association with a slow moving surface anticyclone and propagating Rossby waves. The land surface had also been particularly dry in the preceding weeks, which combined with the presence of a tropical low off northwest Western Australia and an active monsoon trough, provided ideal conditions for the advection of hot air towards southern Australia (Parker et al. 2013). Recent research also suggests that unprecedented Antarctic warming and a polar anticyclone over the Southern Ocean was at least partly responsible for the 2009 Victorian heatwave (Fiddes et al. 2015).

In January 2013, a record-breaking persistent heat event affected the majority of the continent, which was unprecedented spatially and temporally in observational records (Bureau of Meteorology 2013). Extremely hot air masses developed across north Australia that were driven southwards ahead of a series of cold fronts, creating a persistent hot air mass that sat over the continent (Bureau of Meteorology 2013). The event was associated with a delayed monsoon onset and slow moving weather systems over the continent, following a drier than average end to 2012. The heatwave affected set a new national daily maximum temperature record of 40.33 °C on 7th January 2013, consisting of seven consecutive days with maximum temperature above 39 °C (Bureau of Meteorology 2013).

## 4 Future changes

### 4.1 Projections of heatwave events

#### 4.1.1 Heatwave projections from global climate models

Heatwaves are expected to continue in a world under anthropogenic influence, with recent studies suggesting an increase in their frequency and duration (Orlowsky and Seneviratne 2012; Coumou and Robinson 2013; Fischer et al. 2013). Large efforts have been devoted to understanding the impact of anthropogenic climate change on heatwaves in North America and Europe (e.g. Lau and Nath 2012, 2014; Andrade et al. 2014), but this is less so for Australian heatwaves. Relevant work for Australia are summarised below.


Tryhorn and Risbey (2006) and Alexander and Arblaster (2009), employing a single climate model, and the Coupled Model Intercomparison Project (CMIP) phase 3 climate models respectively, found a projected increase in heatwave duration and warm nights in the 21st century under greenhouse forcing. Recently, revised regional climate change projections

for Australia provide a regional assessment of plausible future projections of extreme temperatures (CSIRO 2015). Projections based on 24 CMIP5 phase 5 (CMIP5) climate models for the Representative Concentration Pathways (RCP) 4.5 (medium-low) and 8.5 (high) emission scenarios (Taylor et al. 2012) show that changes in extremes are similar to changes in annual means, consistent with observations (Alexander et al. 2007). Projected changes in the frequency of warm spells (including heatwaves) by 2100 show a significant increase among the CMIP5 ensemble for both RCP4.5 and RCP8.5 (CSIRO 2015).

The CMIP5 models project heatwaves to become more frequent, hotter, and longer across Australia by the end of the 21st century (Cowan et al. 2014). Patterns of change are similar under RCP4.5 (Fig. 3a, b) and RCP8.5 (Fig. 3c, d), but scale with anthropogenic influence. Projections for northern Australia show the largest increase in heatwave days, due to the narrow temperature distribution in the tropics (e.g. Diffenbaugh and Scherer 2011). Increases in intensity and frequency across the southern regions are substantial (Fig. 3a, b, c, d). Under a moving-threshold heatwave definition, which removes the effect of an increase in mean temperature, future changes in frequency are minimal, indicating a similar rate of increase to mean temperature (Cowan et al. 2014). However, the intensity across central-southern Australia still increases, implying that heatwaves are getting hotter at a faster rate than mean temperature in this region.

#### 4.1.2 Regional and downscaled climate projections

Projected changes in temperature extremes have been quantified using dynamical downscaling techniques across Australia (Perkins et al. 2014b), resolution Tasmania (White et al. 2013b),



**Fig. 3** Austral summer heatwave increases compared to historical climatology. (*top*) Ensemble average heatwave frequency (HWF; days per summer), and (*bottom*) heatwave amplitude (HWA; °C). (**a,b**) CMIP5 RCP4.5, (**c,d**) CMIP5 RCP8.5, and (**e,f**) 50 km downscaled NARCIIM SRES A2. CMIP5 increases are calculated over 2081–2100 compared to 1950–2005 climatology. NARCIIM increases are calculated over 2060–2079 compared to 1990–2009 climatology. Heatwaves are based on the definition described in Pezza et al. (2012), where a heatwave occurs when at least three consecutive days are above the monthly 90th percentile for maximum temperature. Stippling indicates where future and historical climatologies are *not* significantly different at the 95 % confidence level. (**a-d**) adapted from Fig. 3 in Cowan et al. (2014) and based on 15 CMIP5 models



and for NSW and the Australian Capital Territory (Evans et al. 2014). White et al. (2013b) show a significant average increase in warm spell duration including by 2100, relative to the current baseline for a high-emissions scenario across Tasmania.

While regional climate ensemble projections agree with large-scale trends from their host models, such studies add spatial detail in extreme temperature frequency and intensity projections (Perkins et al. 2014b). For example, Fig. 3e and f illustrate this using 50 km regional simulations for heatwave intensity and frequency respectively (Evans et al. 2014).

## 4.2 Projected changes in atmospheric circulation

Currently, understanding the dynamic/thermodynamic components behind future projections of Australian heatwaves is limited. Purich et al. (2014) found that under climate change, a poleward shift and intensification of the most severe heatwave-inducing anticyclones can be expected, consistent with projected subtropical ridge and SAM changes (e.g. Timbal et al. 2010; Kent et al. 2013). However, the significant rise in the number of heatwave events across southern and central Australia is currently predominantly attributed to thermodynamic changes (Purich et al. 2014).

There are suggestions that SSTs influence synoptic conditions associated with heatwaves globally (Della-Marta et al. 2007; Trenberth and Fasullo 2012), although whether local SST anomalies are caused by, or are responsible for, Australian heatwaves remains uncertain (e.g. Pezza et al. 2012; Boschath et al. 2015). Moreover, evidence provided by observations is limited, and the CMIP5 models fail to capture observed SST patterns prior to southern Australian heatwaves (Purich et al. 2014). Further research is required on this topic, as well as how future changes in the large-scale modes will impact Australian heatwaves (see Parker et al. 2014a), given that models project significant increases in extreme El Niño and La Niña events (Cai et al. 2014, 2015) and a continuation of positive SAM trends in the RCP8.5 scenario (Zheng et al. 2013).

## 4.3 Projections of marine heatwaves

As marine heatwaves is an emerging field, there are few studies exploring future changes. Southeastern and southwestern Australia are identified as hotspots of ocean warming (Foster et al. 2014), with the Tasman Sea in particular experiencing surface warming three-to-four times the global rate (Holbrook and Bindoff 1997; Ridgway 2007). The CMIP5 models project a net warming (relative to 1986–2005) of SST in the Australia region of 0.6565 °C65 °C by 2050 under RCP2.6, rising to 0.99 °C9 °C and 1.22 °C2 °C under RCP4.5 and RCP8.5, respectively. The strongest signals are seen off the coasts of Tasmania and southwestern Australia, consistent with observed historical trends, and off the northwest shelf. This background warming is a significant driver of marine heatwaves, as the probability of large heat anomalies becomes much greater. In addition, changes in drivers such as ENSO can significantly impact marine heatwave occurrences off Western Australia (Feng et al. 2015). Changes in wind stress curl over the high-latitude South Pacific (e.g. through variations in the SAM) can impact eddy-driven marine heatwaves off southeastern Australia (Oliver et al. 2014a; Oliver and Holbrook 2014). However, there remains a large gap in understanding what future projections of marine heatwaves might entail.

## 5 Conclusions and remaining questions

As part of this Special Issue on Australian natural hazards, this paper summarizes scientific advances in the measurement and understanding of Australian atmospheric and marine heatwaves, and the state of our knowledge on future changes. While there is no single way to define heatwaves, it is clear that their intensity, frequency and duration has increased as anthropogenic influences on the climate has increased. Immediate research could focus on developing more impact-relevant projections on finer spatial scales. Moreover, investigating the human influence on observed trends in Australian heatwaves could be undertaken using internationally applied methods.

Considerable advance has been made in understanding the physical mechanisms driving Australian heatwaves, particularly relationships between ENSO and other modes of variability (Parker et al. 2014a; Perkins et al. 2015) and synoptic-scale dynamics (Pezza et al. 2012; Parker et al. 2014b; Boschhat et al. 2015). However, there is no comprehensive, Australia-wide study documenting the dynamics behind heatwaves. Consequently untangling causes and changes in Australian heatwaves should be prioritized, including land surface feedbacks and antecedent soil moisture, dynamic/thermodynamic components of heatwaves, and increases in the land-sea temperature gradient. The latter has not yet been studied in relation to Australian heatwaves, yet may be important, especially over coastal regions. Understanding the physical connections between heatwaves and drought would also be beneficial to stakeholders of both hazards. Such work is imperative towards a greater understanding of atmospheric heatwaves, as well as advancing Australia's international contribution towards this important field.

Much work remains to better understand marine heatwaves. Given local events in recent years (Pearce and Feng 2013) and the proposal of a measurement framework (Hobday et al. 2016), the Australian community is in a leading position. However, considerable work is required to generate future projections of marine heatwaves and understand interactions between driving mechanisms. Such work should be prioritized to bring our understanding of marine heatwaves in line with atmospheric events.

Lastly, there is a need for a more unified framework for identifying atmospheric events, following recent Australian efforts applied to marine events. The identification of events underpins subsequent research on dynamics, changes and impacts, thus a more unified framework allows for consistency across relevant studies and fields of research. This would require large collaboration across different research and industry sectors, and would need to be conducted at the global scale. This is an area that is likely to be active for many years to come, yet is imperative in addressing both regional and global grand challenges of heatwaves.

**Acknowledgments** S.E. Perkins-Kirkpatrick is supported by Australian Research Council grant number DE140100952. L.V. Alexander and E.C.J. Oliver are supported by Australian Research Council grant number CE110001028 and G. Boschhat by Australian Research Council grand number DP140102855. T. Cowan and A. Purich are supported by the Goyder Institute for Water Research, and the Australian Climate Change Science Program. J.P. Evans is supported by funding from the NSW Office of Environment and Heritage backed NSW/ACT Regional Climate Modelling (NARClM) Project and the Australian Research Council as part of the Future Fellowship FT110100576. This paper was a result of collaboration through the 'Trends and Extremes' working group as part of the Australian Water and Energy Exchanges Initiative (OzEWEX).

## References

- Alexander LV, Arblaster JM (2009) Assessing trends in observed and modelled climate extremes over Australia in relation to future projections. *Int J Climatol* 29:417–435. doi:10.1002/joc.1730
- Alexander LV, Tebaldi C (2012) Climate and weather extremes: observations, modelling and projections [in “the future of the World’s climate”]. Elsevier Science, ISBN 978–0–12–386917–3
- Alexander LV, Hope P, Collins D, Trewin B, Lynch A, Nicholls N (2007) Trends in Australia’s climate means and extremes: a global context. *Aust Meteorol Mag* 56:1–18
- Allen M (2003) Liability for climate change. *Nature* 421:891–892
- Allen MR, Tett SF (1999) Checking for model consistency in optimal fingerprinting. *Clim Dyn* 15:419–434
- Andrade C, Fraga H, Santos JA (2014) Climate change multi-model projections for temperature extremes in Portugal. *Atmos Sci Lett* 15:149–156
- Angéilil O, Stone DA, Tadross M, Tummon F, Wehner M, Knutti R (2014) Attribution of extreme weather to anthropogenic greenhouse gas emissions: sensitivity to spatial and temporal scales. *Geophys Res Lett* 41: 2150–2155
- Arblaster JM, Alexander LV (2012) The impact of the El Niño–Southern Oscillation on maximum temperature extremes. *Geophys Res Lett* 39(20)
- Argüeso D, Evans JP, Fita L, Bormann KJ (2014) Temperature response to future urbanization and climate change. *Clim Dyn* 42:2183–2199. doi:10.1007/s00382-013-1789-6
- Argüeso D, Evans JP, Pitman AJ, Di Luca A (2015) Effects of City expansion on heat stress under climate change conditions. *PLoS one* 10:e0117066. doi:10.1371/journal.pone.0117066
- Barlow KM, Christy BP, O’Leary GJ, Riffkin PA, Nuttall JG (2013) Simulating the impact of extreme heat and frost events on wheat production: the first steps. 20th International Congress on Modelling and Simulation, Adelaide, Australia, 1–6
- Benthuyens J, Feng M, Zhong L (2014) Spatial patterns of warming off Western Australia during the 2011 Ningaloo Niño: quantifying impacts of remote and local forcing. *Cont Shelf Res* 91:232–246
- Boschat G, Pezza AB, Simmonds I, Perkins SE, Cowan T, Purich A (2015) Large scale and sub-regional connections in the lead up to summer heat wave and extreme rainfall events in eastern Australia. *Clim Dyn* 44:1823–1840. doi:10.1007/s00382-014-2214-5
- Bureau of Meteorology (2012) Annual climate summary 2012. Bureau of Meteorology, Australia, Accessed at [http://www.bom.gov.au/climate/annual\\_sum/2012/index.shtml](http://www.bom.gov.au/climate/annual_sum/2012/index.shtml)
- Bureau of Meteorology (2013) Special climate statement 43 – extreme heat in January 2013. Bureau of Meteorology, Australia
- Cai W et al. (2014) Increasing frequency of extreme El Niño events due to greenhouse warming. *Nat Clim Chang* 4:111–116. doi:10.1038/nclimate2100
- Cai W et al. (2015) Increased frequency of extreme La Nina events under greenhouse warming. *Nat Clim Chang* 5:132–137. doi:10.1038/nclimate2492
- Coates L, Haynes K, O’Brien J, McAneney J, de Oliveira F (2014) Exploring 167 years of vulnerability: An examination of extreme heat events in Australia 1844–2010. *Environ Sci Pol* 42:33–44
- Coumou D, Robinson A (2013) Historic and future increase in the global land area affected by monthly heat extremes. *Environ Res Lett* 8:034018. doi:10.1088/1748-9326/8/3/034018
- Cowan T, Purich A, Perkins S, Pezza A, Boschat G, Sadler K (2014) More frequent, longer, and hotter heat waves for Australia in the twenty-first century. *J. Climate* 27:5851–3871
- CSIRO and Bureau of Meteorology 2015. Climate change in Australia information for Australia’s Natural Resource Management regions: technical report. Australia
- Davin EL, Seneviratne SI, Ciais P, et al. (2014) Preferential cooling of hot extremes from cropland albedo management. *PNAS* 111:9757–9761. doi:10.1073/pnas.1317323111
- Della-Marta P, Luterbacher J, von Weissenfluh H, Xoplaki E, Brunet M, Wanner H (2007) Summer heat waves over western Europe 1880–2003: their relationship to large-scale forcings and predictability. *Clim Dyn* 29: 251–275. doi:10.1007/s00382-007-0233-1
- Diffenbaugh NS, Scherer M (2011) Observational and model evidence of global emergence of permanent, unprecedented heat in the 20th and 21st centuries. *Clim Chang* 107:615–624. doi:10.1007/s10584-011-0112-y
- Donat MG, Alexander LV, Yang H, Durre I, Vose R, Caesar J (2013) Global land-based datasets for monitoring climatic extremes. *Bull Am Meteorol Soc* 96:997–1006. doi:10.1175/BAMS-D-12-00109.1
- Evans JP, Ji F, Lee C, Smith P, Argüeso D, Fita L (2014) Design of a regional climate modelling projection ensemble experiment – NARCLiM. *Geosci Model Dev* 7:621–629
- Feng M, Hendon HH, Xie SP, Marshall AG, Schiller A, Kosaka Y, Pearce A (2015) Decadal increase in Ningaloo Niño since the late 1990s. *Geophys Res Lett* 42(1):104–112

- Fiddes SL, Pezza AB, Renwick J (2015) Significant extra-tropical anomalies in the lead up to the Black Saturday fires. *Int J Climatol*. doi:[10.1002/joc.4387](https://doi.org/10.1002/joc.4387)
- Fischer EM, Schär C (2010) Consistent geographical patterns of changes in high-impact European heatwaves. *Nat Geosci* 3:398–403
- Fischer EM, Seneviratne SI, Lüthi D, Schär C (2007) Contribution of land-atmosphere coupling to recent European summer heat waves. *Geophys Res Lett* 34(6). doi:[10.1029/2006GL029068](https://doi.org/10.1029/2006GL029068)
- Fischer EM, Rajczak J, Schär C (2012) Changes in European summer temperature variability revisited. *Geophys Res Lett* 39:L19702. doi:[10.1029/2012GL052730](https://doi.org/10.1029/2012GL052730)
- Fischer EM, Beyerle U, Knutti R (2013) Robust spatially aggregated projections of climate extremes. *Nat Clim Chang* 3:1033–1038
- Foster SD, Griffin DA, Dunstan PK (2014) Twenty years of high-Resolution Sea surface temperature imagery around Australia: inter-annual and annual variability. *PLoS one* 9(7):e100762
- Furrer EM, Katz RW, Walter MD, Furrer R (2010) Statistical modelling of hot spells and heat waves. *Clim Res* 43:191–205. doi:[10.3354/cr00924](https://doi.org/10.3354/cr00924)
- Gleeson MW, Strong AE (1995) Applying MCSST to coral reef bleaching. *Adv Space Res* 16:151–154
- Grotjahn R et al. (2015) North American extreme temperature events and related large scale meteorological patterns: a review of statistical methods, dynamics, modeling, and trends. *Clim Dyn*. doi:[10.1007/s00382-015-2638-6](https://doi.org/10.1007/s00382-015-2638-6)
- Hobday A, Alexander LV, Perkins SE, Smale DA, Straub SC, Oliver E, Bentuysen J, Burrows MT, Donat MG, Feng M, Holbrook NJ, Moore PJ, Scannell H, Sen Gupta A, Wernberg T (2016) A hierarchical approach to defining marine heatwaves. *Prog Oceanogr* 141:227–238. doi:[10.1016/j.pocean.2015.12.014](https://doi.org/10.1016/j.pocean.2015.12.014)
- Holbrook NJ, Bindoff NL (1997) Interannual and decadal temperature variability in the southwest Pacific Ocean between 1955 and 1988. *J Clim* 10:1035–1049
- Horton DE, Johnson NC, Singh D, Swain DL, Rajaratnam B, Diffenbaugh NS (2015) Contribution of changes in atmospheric circulation patterns to extreme temperature trends. *Nature* 522:465–469
- Hudson D, Marshall A, Alves O (2011) Intraseasonal forecasting of the 2009 summer and winter Australian heat waves using POAMA. *Weather Forecast* 26:257–279
- Indian Ocean Climate Initiative (2012) Western Australia's Weather and Climate: a Synthesis of Indian Ocean climate initiative stage 3 research. In: Bryson Bates, Carsten Frederiksen and Janice Wormworth (eds) CSIRO and Bureau of Meteorology, Australia
- Jones DA, Trewin BC (2000) On the relationships between the El Niño–Southern Oscillation and Australian land surface temperature. *Int J Climatol* 20(7):697–719
- Kala J, Evans JP, Pitman AJ (2015) Influence of antecedent soil moisture conditions on the synoptic meteorology of the black Saturday bushfire event in southeast Australia. *Quarterly Journal of the Royal Meteorological Society*. Accepted 29 May 2015
- Kämpf J, Doubell M, Griffin D, Matthews RL, Ward TM (2004) Evidence of a large seasonal coastal upwelling system along the southern shelf of Australia. *Geophys Res Lett* 31(9)
- Kataoka T, Tozuka T, Behera S, Yamagata T (2014) On the Ningaloo Niño/Niña. *Clim Dyn* 43:1463–1482
- Kent D, Kirono D, Timbal B, Chiew F (2013) Representation of the Australian sub-tropical ridge in the CMIP3 models. *Int J Climatol* 33:48–57. doi:[10.1002/joc.3406](https://doi.org/10.1002/joc.3406)
- Kim Y-H, Min S-K, Zhang X, Ziwers F, Alexander LV, Donat MG, Tung Y (2015) Attribution of extreme temperature changes during 1951–2010. *Clim Dyn*. doi:[10.1007/s00382-015-2674-2](https://doi.org/10.1007/s00382-015-2674-2)
- Knutson TR, Zeng F, Wittenberg AT (2014) Multimodel assessment of extreme annual-mean warm anomalies during 2013 over regions of Australia and the western Tropical Pacific. [in “Explaining Extremes of 2013 from a Climate Perspective”]. *Bull Amer Meteor Soc* 95(9):S26–S30
- Kokic P, Crimp S, Howden M (2014) A probabilistic analysis of human influence on recent record global mean temperature changes. *Clim Risk Manag* 3:1–12
- Krueger O, Hegerl GC, Tett SFB (2015) Evaluation of mechanisms of hot and cold days in climate models over central Europe. *Environ Res Lett* 10. doi:[10.1088/1748-9326/10/1/014002](https://doi.org/10.1088/1748-9326/10/1/014002)
- Lau N-C, Nath M (2012) A model study of heat waves over North America: meteorological aspects and projections for the twenty-first century. *J Clim* 25:4761–4784. doi:[10.1175/JCLI-D-11-00575.1](https://doi.org/10.1175/JCLI-D-11-00575.1)
- Lau N-C, Nath M (2014) Model simulation and projection of European heat waves in present-day and future climates. *J Clim* 27(10). doi:[10.1175/JCLI-D-13-00284.1](https://doi.org/10.1175/JCLI-D-13-00284.1)
- Lewis SC, Karoly DJ (2013) Anthropogenic contributions to Australia's record summer temperatures of 2013. *Geophys Res Lett* 40:705–3709
- Lewis SC, Karoly DJ (2014) The role of anthropogenic forcing in the record 2013 Australia-wide annual and spring temperatures. [in “Explaining Extremes of 2013 from a Climate Perspective”]. *Bull Amer Meteor Soc* 95(9):S31–S34
- Lima FP, Wetthey DS (2012) Three decades of high-resolution coastal sea surface temperatures reveal more than warming. *Nat Commun* 3

- Loikith PC, Broccoli AJ (2012) Characteristics of observed atmospheric circulation patterns associated with temperature extremes over North America. *J Clim* 25:7266–7281
- Marshall AG, Hudson D, Wheeler MC, Alves O, Hendon HH, Pook MJ, Risbey JS (2013) Intra-seasonal drivers of extreme heat over Australia in observations and POAMA-2. *Clim Dyn* 43:1915–1937. doi:10.1007/s00382-013-2016-1
- McBride JL, Mills GA Wain AG (2009) The meteorology of Australian heatwaves. Understanding High Impact Weather, CAWCR Modelling Workshop, 30 Nov to 2 Dec 2009
- Mills KE, Pershing AJ, Brown CJ, Yong C, Fu-Sung C, Holland DS, Lehuta S, Nye JA, Sun JC, Thomas AC, Wahle RA (2013) Fisheries Management in a changing climate lessons from the 2012 ocean heat wave in the Northwest Atlantic. *Oceanography* 26:191–195
- Min SK, Cai W, Whetton P (2013) Influence of climate variability on seasonal extremes over Australia. *J Geophys Res Atmos* 118:643–654
- Miralles DG, Van Den Berg MJ, Teuling AJ, De Jeu RAM (2012) Soil moisture-temperature coupling: A multiscale observational analysis. *Geophys Res Lett*. doi:10.1029/2012GL053703
- Miralles DG, Teulingb, AJ, van Heerwaarden, CC, de Arellano, JVG (2014) Mega-heatwave temperatures due to combined soil desiccation and atmospheric heat accumulation. *Nat Geosci*, 7:345–349.
- Munich Re (2009) Heavy losses due to severe weather in the first six months of 2009, press release, 27 July 2009, [www.munichre.com/en/press/press\\_releases/2009/2009\\_07\\_27\\_press\\_release.aspx](http://www.munichre.com/en/press/press_releases/2009/2009_07_27_press_release.aspx)
- Naim, J, Fawcett, R (2013) Defining heatwaves: Heatwave defined as a heat-impact event servicing all community and business sectors in Australia. CAWCR Tech. Rep. 60, 96 pp. [Available online at [http://www.cawcr.gov.au/publications/technicalreports/CTR\\_060.pdf](http://www.cawcr.gov.au/publications/technicalreports/CTR_060.pdf)]
- National Climate Centre (2009). The exceptional January-February 2009 heatwave in southeastern Australia, Bureau of Meteorology, Special Climate Statement 17
- Nicholls N (2004) The changing nature of Australian droughts. *Clim Chang* 63:323–336
- Nicholls N, Lavery B, Frederiksen C, Drosowsky W, Torok S (1996) Recent apparent changes in relationships between the El Niño-Southern Oscillation and Australian rainfall and temperature. *Geophys Res Lett* 23: 3357–3360
- Olita A et al. (2007) Effects of the 2003 European heatwave on the Central Mediterranean Sea: surface fluxes and the dynamical response. *Ocean Sci* 3:273–289
- Oliver ECJ, Holbrook NJ (2014) Extending our understanding of South Pacific gyre ‘spin-up’: modeling the East Australian current in a future climate. *J Geophys Res* 119:2788–2805. doi:10.1002/2013JC009591
- Oliver ECJ, Wotherspoon SJ, Chamberlain MA, Holbrook NJ (2014a) Projected Tasman Sea extremes in sea surface temperature through the 21st century. *J Clim* 27:1980–1998. doi:10.1175/JCLI-D-13-00259.1
- Oliver ECJ, Wotherspoon SJ, Holbrook NJ (2014b) Estimating extremes from global ocean and climate models: A Bayesian hierarchical model approach. *Prog Oceanogr* 122:77–91. doi:10.1016/j.pocan.2013.12.004
- Orlowsky B, Seneviratne SI (2012) Global changes in extreme events: regional and seasonal dimension. *Clim Chang* 110:669–696. doi:10.1007/s10584-011-0122-9
- Parker TJ, Berry GJ, Reeder MJ (2013) The influence of tropical cyclones on heat waves in southeastern Australia. *Geophys Res Lett* 40(23):6264–6270
- Parker TJ, Berry GJ, Reeder MJ, Nicholls N (2014a) Modes of climate variability and heat waves in Victoria, southeastern Australia. *Geophys Res Lett* 41:6926–6934
- Parker TJ, Berry GJ, Reeder MJ (2014b) The structure and evolution of heat waves in Southeastern Australia. *J Clim* 27:5768–5785. doi:10.1175/JCLI-D-13-00740.1
- Pearce A, Feng M (2013) The rise and fall of the “marine heat wave” off Western Australia during the summer of 2010/2011. *J Mar Syst* 111-112:139–156
- Perkins SE, Alexander LV (2013) On the measurement of heatwaves. *J Clim* 26:4500–4517. doi:10.1175/JCLI-D-12-00383.1
- Perkins SE, Lewis SL, King AD, Alexander LV, (2014a). Increase simulated risk of the hot Australian summer of 2012/2013 due to anthropogenic activity as measured by heat wave frequency and intensity. [in “Explaining Extremes of 2013 from a Climate Perspective”]. *Bull Amer Meteor Soc* 95(9):S34–S36.
- Perkins SE, Moise A, Whetton P, Katzfey J (2014b) Regional changes of climate extremes over Australia – a comparison of regional dynamical downscaling and global climate model simulations. *Int J Climatol* 34: 3456–3478
- Perkins SE, Argüeso D, White CJ (2015) Relationships between climate variability, soil moisture, and Australian heatwaves. *J Geophys Res Atmos* 120(16):8144–8164
- Pezza AB, Van Rensch P, Cai W (2012) Severe heat waves in Southern Australia: synoptic climatology and large scale connections. *Clim Dyn* 38:209–224. doi:10.1007/s00382-011-1016-2
- Purich A, Cowan T, Cai W, van Rensch P, Uotila P, Pezza A, Bosch G, Perkins S (2014) Atmospheric and oceanic conditions associated with Southern Australian heat waves: A CMIP5 analysis. *J Clim* 27:7807–7829. doi:10.1175/JCLI-D-14-00098.1

- Quesada B, Vautard R, Yiou P, Hirschi M, Seneviratne SI (2012) Asymmetric European summer heat predictability from wet and dry southern winters and springs. *Nat Clim Chang* 2:736–741
- Ridgway KR (2007) Long-term trend and decadal variability of the southward penetration of the East Australian current. *Geophys Res Lett* 34(13)
- Risbey JS, Pook MJ, McIntosh PC, Wheeler MC, Hendon HH (2009) On the remote drivers of rainfall variability in Australia. *Mon Weather Rev* 137:3233–3253
- Roughan M, Middleton JH (2004) On the east Australian current: variability, encroachment, and upwelling. *J Geophys Res: Oceans*(1978–2012) 109(C7)
- Russo S, Dosio A, Graversen RG, Sillmann J, Carrao H, Dunbar MB, Vogt JV (2014) Magnitude of extreme heat waves in present climate and their projection in a warming world. *J Geophys Res Atmos* 119:12–500
- Selig ER, Casey KS, Bruno JF (2010) New insights into global patterns of ocean temperature anomalies: implications for coral reef health and management. *Glob Ecol Biogeogr* 19:397–411
- Seneviratne SI, Lüthi D, Litschi M, Schär C (2006) Land-atmosphere coupling and climate change in Europe. *Nature* 443:205–209
- Sorte CJB, Fuller A, Bracken MES (2010) Impacts of a simulated heat wave on composition of a marine community. *Oikos* 119:1909–1918
- Steffen W, Hughes, L, Perkins S, (2014) Heatwaves: Hotter, Longer, More Often. Special Report by the Climate Council of Australia, Sydney, Australia. 62 pp. [Available online at <http://www.climatecouncil.org.au/heatwaves-report/>]
- Stott PA, Stone DA, Allen MR (2004) Human contribution to the European heatwave of 2003. *Nature* 432(7017):610–614
- Taylor K, Stouffer R, Meehl G (2012) An overview of CMIP5 and the experiment design. *Bull Amer Meteor Soc* 93:485–498. doi:10.1175/BAMS-D-11-00094.1
- Timbal, B., et al., (2010) Understanding the anthropogenic nature of the observed rainfall decline across South Eastern Australia. CAWCR Tech. Rep. 026, Centre for Australian Weather and Climate Research, 202 pp. [Available online at [http://www.cawcr.gov.au/publications/technicalreports/CTR\\_026.pdf](http://www.cawcr.gov.au/publications/technicalreports/CTR_026.pdf).]
- Trenberth K, Fasullo J (2012) Climate extremes and climate change: the Russian heat wave and other climate extremes of 2010. *J Geophys Res* 117:D17103. doi:10.1029/2012JD018020
- Trewin BC (2009). A new index for monitoring changes in heatwaves and extended cold spells. In 9th International Conference on Southern Hemisphere Meteorology and Oceanography
- Trewin B, Vermont H (2010) Changes in the frequency of record temperatures in Australia, 1957–2009. *Aust Meteorol Oceanogr J* 60:113–119
- Tryhorn L, Risbey J (2006) On the distribution of heat waves over the Australian region. *Aus Meteorol Mag* 55: 169–182
- Wernberg T, Smale DA, Tuya F, Thomsen MS, Langlois TJ, De Bettignies T, Rousseaux CS (2013) An extreme climatic event alters marine ecosystem structure in a global biodiversity hotspot. *Nat Clim Chang* 3:78–82
- White CJ, Hudson D, Alves O (2013a) ENSO, the IOD and intraseasonal prediction of heat extremes across Australia using POAMA-2. *Clim Dyn* 43:1791–1810. doi:10.1007/s00382-013-2007-2
- White CJ, McInnes KL, Cechet RP, Corney SP, Grose MR, Holz G, Katzfey JJ, Bindoff NL (2013b) On regional dynamical downscaling for the assessment and projection of future temperature and precipitation extremes across Tasmania, Australia. *Clim Dyn* 41:3145–3165. doi:10.1007/s00382-013-1718-8
- Zander KK, Botzen WJW, Oppermann E, Kjellstrom T, Garnett ST (2015) Heat stress causes substantial labour productivity loss in Australia. *Nat Clim Chang* 5:647–651. doi:10.1038/nclimate2623
- Zheng F, Li J, Clark RT, Nnamchi HC (2013) Simulation and projection of the Southern Hemisphere annular mode in CMIP5 models. *J Clim* 26:9860–9879. doi:10.1175/JCLI-D-13-00204.1