

## Beyond Bonferroni: less conservative analyses for conservation genetics

Shawn R. Narum

Published online: 8 August 2006  
© Springer Science+Business Media B.V. 2006

**Erratum to: Conserv Genet**  
**DOI: 10.1007/s10592-005-9056-y**

In the example for the B-H FDR procedure, four incorrectly printed values create confusion for the reader. Corrections in the Methods section, second paragraph, third and fourth sentences: “The first  $p$ -value to satisfy  $p_i \leq i/k \times \alpha$  is  $p_{(10)}$  since  $p_{(10)} = 0.032 \leq 10/15 \times 0.05 = 0.0333$ . Thus pairwise tests in the experiment with  $p$ -values less than or equal to 0.0333 reject the null hypothesis. The gain in power

with the B-H method FDR over the Bonferroni procedure (critical values of 0.033 and 0.003 respectively in this example) are substantial.” Corrections in the Methods section, last paragraph, last sentence: “This critical value is intermediate relative to those calculated from Bonferroni (0.0033) and B-H method FDR (0.0333).”

---

The online version of the original article can be found at <http://www.dx.doi.org/10.1007/s10592-005-9056-y>

---

S. R. Narum (✉)  
Columbia River Inter-Tribal Fish Commission, 3059-F  
National Fish Hatchery Road, Hagerman, ID 83332, USA  
e-mail: nars@critfc.org