



Effects of atrazine and S-metolachlor on stream periphyton taxonomic and fatty acid compositions

Laura Malbezin¹ · Soizic Morin² · Isabelle Lavoie¹

Accepted: 5 February 2024 / Published online: 22 February 2024

© The Author(s), under exclusive licence to Springer Science+Business Media, LLC, part of Springer Nature 2024

Abstract

Extensive pesticide use for agriculture can diffusely pollute aquatic ecosystems through leaching and runoff events and has the potential to negatively affect non-target organisms. Atrazine and S-metolachlor are two widely used herbicides often detected in high concentrations in rivers that drain nearby agricultural lands. Previous studies focused on concentration-response exposure of algal monospecific cultures, over a short exposure period, with classical descriptors such as cell density, mortality or photosynthetic efficiency as response variables. In this study, we exposed algal biofilms (periphyton) to a concentration gradient of atrazine and S-metolachlor for 14 days. We focused on fatty acid composition as the main concentration-response descriptor, and we also measured chlorophyll a fluorescence. Results showed that atrazine increased cyanobacteria and diatom chlorophyll a fluorescence. Both herbicides caused dissimilarities in fatty acid profiles between control and high exposure concentrations, but S-metolachlor had a stronger effect than atrazine on the observed increase or reduction in saturated fatty acids (SFAs) and very long-chain fatty acids (VLCFAs), respectively. Our study demonstrates that two commonly used herbicides, atrazine and S-metolachlor, can negatively affect the taxonomic composition and fatty acid profiles of stream periphyton, thereby altering the nutritional quality of this resource for primary consumers.

Keywords Periphyton · Herbicides · Atrazine · S-metolachlor · Fatty acids · Fluorescence

Introduction

In 2020, worldwide pesticide use in agriculture was estimated at 2.7 million tons (FAO 2022). The application of these compounds on the landscape has resulted in the detection and persistence of pesticides in aquatic ecosystems. Even at low concentrations, pesticides can interact with other compounds and represent a serious risk to aquatic and terrestrial organisms (Groner and Relyea 2011; Relyea 2009). Pesticides that target autotrophs (i.e., herbicides) represent about 48% of the pesticides used globally, and they may comprise an even more substantial proportion, ranging from 63% to upwards of 80% in certain regions of the world such as in the United States of America (USA) (Brain and Anderson 2019; USEPA 2017). Atrazine and

S-metolachlor are two herbicides commonly applied for grain, legume and cereal crop production. Resultantly, these herbicides are frequently detected in nearby aquatic ecosystems with mean concentrations close to $1 \mu\text{g.L}^{-1}$ in surface waters in Argentina and in the USA (Bachetti et al. 2021; Hansen et al. 2019). Atrazine daily maximum concentrations reached hundreds $\mu\text{g.L}^{-1}$ in watersheds highly vulnerable to runoff in agricultural regions of the USA (see Perkins et al. 2021 for complete database) and can exceed water quality criteria in Europe (Parlakidis et al. 2022; Székács et al. 2015). S-metolachlor is also commonly applied for corn and soybean production, and can reach concentrations between $5 \mu\text{g.L}^{-1}$ and $50 \mu\text{g.L}^{-1}$ in agricultural regions of Europe (Griffini et al. 1997; Kapsi et al. 2019; Roubéix et al. 2012; Székács et al. 2015; Vryzas et al. 2011), and up to $100 \mu\text{g.L}^{-1}$ in agricultural regions of the USA (Battaglin et al. 2003, 2000).

Atrazine [2-chloro-4-(ethylamino)-6-(isopropylamino)-s-triazine] is a triazine compound marketed in the late 1950s but has been subsequently severely restricted in Europe since 2003–2004 (European Commission 2004) and banned in certain countries (e.g., France, Sénat de France 2003, and Germany, LAWA, 2019) due to its presence at

✉ Laura Malbezin
lmalbezin@outlook.fr

¹ Institut national de la recherche scientifique, centre Eau Terre Environnement, 490 rue de la Couronne, G1K 9A9 Quebec City, QC, Canada

² INRAE, EABX, 50 avenue de Verdun, 33612 Cestas Cedex, France

concentrations beyond water quality criteria. Although studies and government reports mentioned the potential risk of this molecule on non-target terrestrial and aquatic organisms (de Albuquerque et al. 2020; USEPA 2016), atrazine is still used in several countries worldwide, including in Canada and in the USA, albeit under increased regulation (e.g., Quebec, see Fortier 2018). Atrazine is a photosynthesis inhibitor herbicide that binds the D1 protein of photosystem II and blocks electron transport (Vallotton et al. 2008). By disrupting electron transport, atrazine leads to the production of reactive oxygen species (ROS), resulting to oxidative stress, peroxidation of membrane lipids and, ultimately, senescence of plant cells (de Albuquerque et al. 2020). When present in aquatic ecosystems, atrazine can be harmful for aquatic plants (Gao et al. 2019), micro-algae (Baxter et al. 2016), as well as non-phototrophic organisms such as bacteria (DeLorenzo et al. 1999). The effects of atrazine on amphibians, in particular, have long been disputed, but the USEPA mentioned a potential chronic risk to amphibians, fish, and aquatic invertebrates in locations where atrazine use is heaviest (USEPA 2016).

S-metolachlor (2-chloro-N-(2-ethyl-6-methylphenyl)-N-[(1S)-2-methoxy-1-methylethyl] acetamide) is an extensively used chloroacetamide herbicide available since the 1990s. S-metolachlor inhibits very long-chain fatty acids (VLCFAs) biosynthesis by binding with a synthase involved in fatty acid elongation (HRAC 2020; WSSA 2021). VLCFAs are an important component for the functioning of biological membranes. For example, Böger (2003) found that S-metolachlor inhibited 68% of VLCFAs biosynthesis in the green algae *Scenedesmus acutus* compared to control. Similarly, Debenest et al. (2009) found that this compound can directly affect cellular density of periphytic diatoms. In addition, S-metolachlor is highly soluble, mobile, can bioaccumulate in non-target organisms (Zemolin et al. 2014), and it is suspected to be an endocrine disruptor for certain fish species (Ou-Yang et al. 2022; Quintaneiro et al. 2017).

Freshwater biofilms or periphyton are a heterogeneous assemblage of algae, bacteria, fungi, archaea and viruses as well as micrometazoans trapped in a matrix of extracellular polymeric substances that develop on various submerged substrates (Wetzel 1983). Periphyton is an integral part to the function of aquatic ecosystems and provides services in nutrient cycling. In addition, it is the basal resource of aquatic food webs providing essential compounds such as proteins, lipids and fatty acids needed for the growth and metabolism of higher trophic levels (Thompson et al. 2002). Fatty acids (FAs), in particular, are an important compound transferred along the food chain from prey to consumers (Gladyshev et al. 2011). Polyunsaturated fatty acids (PUFAs) are involved in physiological processes and

maintain membrane structure (Huggins et al. 2004). While vegetal cells can synthesize PUFAs de novo, consumers must obtain them through dietary pathways (Brett and Müller-Navarra 1997). In particular, certain essential FAs such as linoleic acid (LIN; C18:2n6) and α -linoleic acid (ALA; C18:3n3) are almost exclusively produced by vegetal cells; therefore, algae represent an essential source of these molecules for animal consumers (Brett and Müller-Navarra 1997). In aquatic ecosystems, long-chain PUFAs (LCPUFAs) such as arachidonic acid (ARA; C20:4n6), eicosapentanoic acid (EPA; C20:5n3) and docosahexanoic acid (DHA; C22:6n3) are also mainly produced by microalgae (Li et al. 2014) and are transferred to consumers with high efficiency (Gladyshev et al. 2011). There is some evidence that herbicides may affect the FA composition of microalgae by interfering with vegetal lipid metabolism (Demailly et al. 2019; Gonçalves et al. 2021). Herbicides may also induce changes in microorganism community structure of periphyton by selecting for more tolerant species that differ in FA composition (Konschak et al. 2021). For example, diatoms are known to be rich in EPA, while green algae are characterized by high content of ALA and bacteria by C18:1n9, C16:0, and C18:0. Thus, there is considerable risk that herbicides reaching aquatic ecosystems may affect the structure of periphyton assemblages and consequently alter the nutritional quality of this basal resource to higher consumers. Indeed, it has been shown that food quality affects growth and development of consumers (Da Costa et al. 2023; Müller-Navarra et al. 2000; Rossoll et al. 2012).

This study investigated the effects of two herbicides frequently detected in aquatic ecosystems on complex biofilm communities. Fatty acid composition was used as the main response variable due to the key role FAs play in food webs. Most studies adopting a concentration-response exposure design have been carried out on monospecific cultures and over timescales of a few hours to a few days, with conventional descriptors such as cell numbers, mortality, or photosynthetic capacity as response variables. To our knowledge, this study is one of the first to adopt a concentration-response design with complex microorganism matrices (biofilms) in a chronic context (7 and 14 days of exposure) and focusing on fatty acids as a response variable to pesticide contamination. The primary aim was to provide information on the long-term effects of the tested herbicides. More specifically, we conducted a laboratory experiment to (1) determine the effects of atrazine and S-metolachlor on periphyton FA composition and to (2) relate possible modifications in FA profiles to changes in the community structure of autotrophic organisms monitored by chlorophyll a fluorescence measurements. For this purpose, we exposed cultured periphyton in microcosms to either atrazine or S-metolachlor along an environmentally relevant concentration gradient.

Table 1 Measured concentrations of atrazine and S-metolachlor (mean \pm standard deviation when treatment was replicated) at day 0, 7, and 14 before concentration adjustment. Italics: concentrations in abiotic microcosms

Day	Nominal concentration ($\mu\text{g.L}^{-1}$)			Measured concentration of atrazine ($\mu\text{g.L}^{-1}$)			Measured concentration of S-metolachlor ($\mu\text{g.L}^{-1}$)		
	0	7	14	0	7	14	0	7	14
Control (n = 4)	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD
5	1.0	4.9	5.3	3.4	1.9	2.2			
10 (n = 3)	8.7 \pm 0.4	8.8 \pm 1.1	8.9 \pm 0.9	3.0 \pm 0.8	3.7 \pm 2.4	4.5 \pm 1.3			
50	47.5	46.4	37.9	26.5	28.1	29.4			
<i>50 abiotic condition (n = 3)</i>				38.6 \pm 5.0	20.0 \pm 1.8				
100 (n = 3 for S-metolachlor)	19.1	80.6	73.8	109.5 \pm 13.3	73.7 \pm 15.8	74.6 \pm 9.1			
500	551.3	517.7	459.9	542.7	238.1	261.9			
<i>500 abiotic condition (n = 3)</i>	520.1 \pm 5.0	428.9 \pm 27.2							
1000	1259.3	1231.8	1266.1	998.2	272.5	567.8			

Materials and methods

Experimental setup and periphyton sampling

Periphyton inoculum was collected in a stream (watershed = 82 km²) with low to moderate anthropogenic activities (agricultural and urban) located a few kilometers west of Quebec City (Quebec, Canada; lat: 46°45'48.8"N, long: 71°21'24.0"W). The inoculum was acclimated in the laboratory in aquaria for 2 months under experimental conditions (temperature = 20–22 °C, natural photoperiod). Before the start of the experiment, acclimated periphyton was evenly transferred in suspension into 23 microcosms (dimensions: 30 \times 15 \times 20 cm) filled with 7.5 L of dechlorinated tap water enriched with nutrients (temperature = 20 °C, photoperiod = 16 h day/8 h night, average light flux = 54 $\mu\text{mol photons.m}^{-2}.\text{s}^{-1}$, nutrients summarized in Table S1) and equipped with an aeration pump. Each microcosm contained six glass slides (double-sided for a total of 141 cm²) to increase the surface area available for periphyton colonization. After a 1-month colonization period in the microcosms, periphyton were exposed to a gradient of atrazine and S-metolachlor concentrations (PESTANAL, analytical standard, Sigma Aldrich). The nominal concentrations of both herbicides tested were: 0, 5, 10, 50, 100, 500, and 1000 $\mu\text{g.L}^{-1}$. Treatments henceforth will be referred to by the first letter of the herbicide (A for Atrazine; S for S-metolachlor), followed by the nominal concentration (e.g., A5, A10, S5, S10, etc.) and the treatment that did not receive herbicide will be referred to as the control. We conducted a gradient study design where we chose to increase the number of treatments to the detriment of replication (Larras et al. 2018). However, experimental replicates were incorporated for the control (n = 4), A10 (n = 3), S10 (n = 3), and S100 (n = 3) treatments that we considered environmentally relevant concentrations. Increasing the number of treatments tested instead of testing replication is suggested to be

an advantageous strategy (Green et al. 2018). A limited number of replications may result in increased inter-treatment variability; however, this could be reduced by the number of measurements taken within each replication. Within each microcosm, samples were collected on three occasions, before exposure (day 0), and after 7 and 14 days of exposure. To ensure a homogeneous and representative sample, periphyton were scrapped from randomly collected glass slides as well as from the walls of the microcosms to make one composite sample per treatment which was preserved at -80 °C for FA analyses.

Chlorophyll a fluorescence of green algae, diatoms and cyanobacteria composing the periphyton was measured with the fluorometer probe Benthotorch (bbe BenthosTorch, Moldaenke, Germany) that uses the excitation-emission responses at several wavelengths (470 nm, 525 nm, and 610 nm) to determine chlorophyll a concentrations of attached autotrophic organisms. At each sampling time, six measurements were randomly taken per microcosm by placing the instrument directly onto the glass slides that were delicately and temporarily removed from the water. Biofilm samples were also collected and fixed with formaldehyde (3% from a stock formalin 37%) in order to qualitatively compare the relative composition of the main algal groups with fluorescence data provided by the BenthosTorch. This comparison was conducted only for six samples as the objective was simply to verify the fluorescence data. Despite the fact that green algae were observed under the microscope but were not very abundant as measured with the probe (leading to an underestimation by the BenthosTorch) their relative increase in the biofilm during the course of the experiment was measured by the probe and qualitatively verified by microscopy.

Throughout the experiment, pH = 8.2 \pm 0.1, conductivity = 318.3 \pm 25.6 $\mu\text{S.cm}^{-1}$ and water temperature = 18.9 \pm 0.3 °C (n = 66) were stable. Herbicide concentrations were determined by liquid chromatography (Finnigan Surveyor) with tandem mass spectrometry (TSQ

Quantum Access; Thermo Scientific) (LC-MS/MS) (Limit of detection = $0.1 \mu\text{g.L}^{-1}$, analytical standards: Atrazine-D5 and Metolachlor-D6). Herbicide concentrations were re-adjusted as needed over the course of the experiment. To determine any abiotic loss of atrazine and S-metolachlor in microcosms, three microcosms without periphyton were contaminated with atrazine at a nominal concentration of $500 \mu\text{g.L}^{-1}$ and three additional microcosms were contaminated with $50 \mu\text{g.L}^{-1}$ of S-metolachlor. Water was sampled after 7 days and analyzed by LC-MS/MS following the same method as described above. Measured atrazine concentrations were close to nominal concentration in biotic microcosms, while S-metolachlor concentrations were below targeted values (S2 for details). Despite the fact that measured concentrations deviated from the targeted nominal concentrations, a concentration gradient was observed for both herbicides as seen in Table 1.

Fatty acid analysis

Fatty acid extraction and analysis were performed according to Fadhlouli et al. (2020), where a 40 mg subsample of periphyton was homogenized in 8.4 mL of chloroform/methanol (2 v/v) solution for 1 min using a Homogenizer 850 (Fisherbrand™). A volume of 20 μL of trycosilic acid (C23:0) was added as an internal standard and the samples were then sonicated for 5 min using a Sonifier® (Branson). A 2 mL solution of NaCl (0.73%) was then added followed by centrifugation of the sample for 15 min at 3000 tr/min at 4 °C allowing for lipid separation in the lower phase. Lipids were recovered from this lower phase and evaporated using a TurboVap® (Caliper Life Sciences TurboVap II) for 15 min at 40 °C before being transferred to screw-capped tubes with 3 mL of BF_3 (boron trifluoride-methanol solution 14% in methanol). The BF_3 is used to esterified fatty acids and to facilitate analysis by gas chromatography. After 1 h of incubation at 75 °C, fatty acid methyl esters (FAMES) were extracted by adding 3 mL of ultra-pure water and 3 mL of petroleum ether. This step was repeated two more times to improve FAMES recovery. The top fraction of petroleum ether was recovered and dried using the TurboVap® for 15 min at 40 °C. Finally, FAMES were dissolved in 240 μL of hexane and then transferred into screw-capped vials to be analyzed by gas chromatography with a flame ionization detector (Agilent Technologies; 7890D GC system) equipped with a fused silica capillary column (DB-FATWAX from Agilent Technologies: 30 m [length], 0.250 mm [inner diameter], 0.25 μm [film thickness]). Injection was conducted at a constant pressure, and helium was used as the carrier gas. Temperature programming was as follows: initial temperature of 140 °C increased to 170 °C at a rate of $6.5 \text{ }^\circ\text{C.min}^{-1}$, then to 200 °C at a rate of $2.75 \text{ }^\circ\text{C.min}^{-1}$ for 14 min, and finally to 230 °C at a rate of $3 \text{ }^\circ\text{C.min}^{-1}$ for

12 min. Because the periphyton is highly heterogeneous, five subsamples from the one composite sample collected in each microcosm were analyzed (pseudo-replicates) to ensure a proper representation of fatty acid profiles within each microcosm.

Statistical analysis

Statistical analyses were performed in RStudio (R version 4.2.2). Water chemistry and chlorophyll a fluorescence data (μg of chlorophyll a.cm^{-2}) were expressed as mean \pm standard deviation. Due to inter-microcosm variability prior to exposure, photoautotroph fluorescence and FA composition changes (i.e., deltas Δ) between day 0 and the two sampling times (7 and 14) were used. Delta values were then used to perform linear regressions. For all statistical analyses, results were considered significant when the *p*-value was less than 0.05 and marginally significant where *p*-value was between 0.05 and 0.1. For photoautotroph chlorophyll a fluorescence, one-way ANOVAs were performed on raw data and only for replicated conditions. Pairwise t-tests with Bonferroni adjustment were used for post hoc comparisons.

Principal Component Analyses (PCA) were conducted on fatty acid data (including fatty acids with proportions >5% in at least one sample) from pseudo-replicates, allowing the representation of intra-condition variability. The “FactoMineR” and “factoextra” packages were used to explore patterns in FA profiles as a function of exposure concentrations. A PERMutational ANalysis Of VAriance (PERMANOVA) on dissimilarity matrix was performed on replicated conditions and was followed by a pairwise comparison to test for differences in FA profiles between conditions using the “adonis2” (method = “gower”) and “pairwise.adonis2” functions from the “vegan” package.


Results

Community structure of the autotrophic organisms

Effect of atrazine on chlorophyll a fluorescence

Diatoms and cyanobacteria were the two main groups of photoautotroph organisms in periphyton with green algae having a lower relative chlorophyll a fluorescence (see Fig. S1 for raw data). Chlorophyll a fluorescence data suggested inter-microcosms variability before contamination, in particular between controls and S100, A100, A500, and A1000. The fluorescence probe detected green algae in certain microcosms and diatoms and cyanobacteria exhibited lower levels under control condition. The data were standardized to reduce the effect of pre-exposure variability

Fig. 1 Chlorophyll a fluorescence variation (Δ) at day 7 and day 14 compared to day 0 (initial time of the experiment), based on chlorophyll a fluorescence of photoautotrophic groups as a function of measured atrazine concentrations ($\mu\text{g}\cdot\text{L}^{-1}$) (linear regression Df = 10)



where data for day 7 and day 14 were normalized using the data from day 0.

Under the control condition, the total chlorophyll a fluorescence significantly decreased between day 0 and day 14 (Df = 2, $F = 9.01$, $p = 0.01$). Especially, diatom specific fluorescence marginally decreased between 7 days and 14 days of exposure (Df = 2, $F = 3.67$, $p = 0.05$). Linear regressions showed some effect of atrazine on photoautotroph chlorophyll a fluorescence (Fig. 1). Specifically, cyanobacteria and diatoms specific fluorescence increased with atrazine concentration after 7 days (Df = 10, $F = 26.44$, $R^2 = 0.73$, $p < 0.001$ and Df = 10, $F = 28.98$, $R^2 = 0.75$, $p < 0.001$, respectively) and 14 days of exposure (Df = 10, $F = 20.36$, $R^2 = 0.67$, $p = 0.001$ and Df = 10, $F = 25.83$, $R^2 = 0.72$, $p < 0.001$, respectively). Green algae were a minor autotrophic group based on chlorophyll a fluorescence, and atrazine did not appear to affect its

chlorophyll a fluorescence as it remained stable between exposure concentrations and over time.

For all photoautotrophic groups, no significant differences were observed in ΔChla -fluorescence between control and $10 \mu\text{g}\cdot\text{L}^{-1}$ conditions after 7 days (Df = 5, with $F = 0.68$, $p = 0.45$ for cyanobacteria; $F = 3.25$, $p = 0.13$ for diatoms and $F = 5.19$, $p = 0.07$ for green algae) and 14 days of exposure (Df = 5, with $F = 0$, $p = 0.99$ for cyanobacteria; $F = 0.15$, $p = 0.72$ for diatoms and $F = 0.98$, $p = 0.37$ for green algae). Atrazine then appeared to have a significant effect on chlorophyll a fluorescence at concentrations higher than $10 \mu\text{g}\cdot\text{L}^{-1}$.

Effect of S-metolachlor on chlorophyll a fluorescence

As observed for atrazine, green algae remained the minor photosynthetic group. In contrast to atrazine, S-metolachlor

Fig. 2 Chlorophyll a fluorescence variation (Δ) at day 7 and day 14 compared to day 0 (initial time of the experiment), based on chlorophyll a fluorescence of photoautotrophic groups as a function of measured S-metolachlor concentrations ($\mu\text{g.L}^{-1}$) (linear regression Df = 12)





Fig. 3 Principal component analysis (PCA) of fatty acid profiles at day 14 for the different atrazine conditions. The left panel is the graph of individual FA and on the right panel corresponds to the circle of correlations. Ellipses have been plotted with a confidence level of 80%. All pseudo-replicates were considered to better represent the intra-condition variability






Fig. 4 Linear regressions based on differences in proportions ($\Delta\%$) for the main fatty acid groups as a function of measured concentrations of atrazine ($\mu\text{g.L}^{-1}$) after 7 days and 14 days of exposure

had no effect on photoautotroph chlorophyll a fluorescence (Fig. 2).


The one-way ANOVA showed no effect of S-metolachlor at $10 \mu\text{g.L}^{-1}$ and $100 \mu\text{g.L}^{-1}$ on photoautotrophic group fluorescence compared to the control condition after 7 days (Df = 2, with $F = 1.30$, $p = 0.33$ for cyanobacteria; $F = 3.56$, $p = 0.09$ for diatoms and $F = 1.23$, $p = 0.35$ for green algae) and 14 days of exposure (Df = 2, with $F = 1.06$, $p = 0.40$ for cyanobacteria; $F = 2.02$, $p = 0.20$ for diatoms and $F = 0.69$, $p = 0.54$ for green algae).

Effects of herbicides on periphyton fatty acid composition

Effect of atrazine on fatty acids

A total of 27 fatty acids were identified in the total lipid fraction of the periphyton. Average (\pm standard deviation) proportions of each FA are presented as supplementary information (Table S3). Unsaturated fatty acids (UFAs) were generally the predominant FA group in all treatments with a relative percentage of up to 62.9% comprised mostly

Fig. 5 Principal component analysis (PCA) of fatty acid profiles at day 14 for the different S-metolachlor conditions. The left panel is the graph of individual FA and on the right panel corresponds to the circle of correlations. Ellipses have been plotted with a confidence level of 80%. All pseudo-replicates were considered to better represent the intra-condition variability



of mono-unsaturated fatty acids (MUFAs; 23.0–41.1%) followed by poly-unsaturated fatty acids (PUFAs; 15.2–26.1%). Saturated fatty acids (SFAs) represented up to 54.3% of total lipid content in the periphyton samples.

PCA of periphyton FA relative percentages 14 days after atrazine exposure explained 66.9% of the variance in FA composition on two axes (dim1 = 42.9% and dim2 = 24%; Fig. 3). PCAs of FA composition on day 0 and day 7 are presented in the supplementary information (Fig. S2). The A5 and A100 treatments clustered together on the top left of the ordination and had higher proportions of MUFAs, in particular C16:1n7, compared to the A50, A500 and A1000 treatments that clustered in the lower portion of the PCA and were more characterized by SFAs and C18:0. A large dispersion of fatty acid data was observed, especially for the control and A10 conditions which overlapped all treatment groups.

The PERMANOVA and pairwise comparisons performed on replicated conditions (control and A10) revealed a significant difference in FA composition ($Df = 1$, $F = 3.92$; $p = 0.02$) after 14 days under atrazine exposure. PERMANOVA were also conducted at day 0 and day 7 and showed that differences were already present at day 7 ($Df = 1$, $F = 4.88$; $p = 0.008$) but not prior to exposure ($Df = 1$, $F = 2.33$, $p = 0.08$).

Linear regressions showed that atrazine concentration did not markedly affect the main FA groups (Fig. 4; linear regressions for individual FA are shown in Fig. S3). Only a regression marginally significant was observed for SFAs at 14 days ($Df = 10$, $R^2 = 0.27$; $p = 0.08$).

Effect of S-metolachlor on fatty acids

Mean (\pm standard deviation) proportions of each FA are presented in the supplementary information (Table S4).

Unsaturated fatty acids (UFAs) comprised up to 62.5% of the total FA content of periphyton among treatments, while mono-unsaturated fatty acids (MUFAs) varied from 27.3% to 38.8% and poly-unsaturated fatty acids (PUFAs) varied from 16.9% to 33.4%. Saturated fatty acids (SFAs) represented up to 54.3% of total FA content in the periphyton samples.

A PCA was performed to assess the effect of S-metolachlor after 14 days of exposure (Fig. 3) (See Fig. S4 for 0 and 7 days). The first two dimensions explained 62% of the variance (dim1 = 42.1% and dim2 = 19.9%). The two highest concentrations clustered on the left side of the ordination, while S10, S100 clustered on the right side. The S5 and S50 conditions clustered on the top portion of the ordination (dimension 2) and S100 clustered on the lower half of the ordination. The control condition clustered in the middle and showed high dispersion. High S-metolachlor concentrations (S500 and S1000) were more associated with SFAs such as C18:1n9 and C18:0, while lower S-metolachlor concentrations were rather characterized by PUFAs such as ALA, EPA, and C20:4n6 (Fig. 5).

At day 14, the PERMANOVA (performed only on replicated conditions; control, S10, and S100) showed a significant effect of S-metolachlor concentrations on the fatty acid composition of the periphyton. Indeed, there was a strong dissimilarity between the control and S100 ($Df = 2$, $F = 3.79$, $p = 0.001$). PERMANOVA conducted at day 0 and day 7 also revealed differences in FA profiles. Especially, the S10 condition ($Df = 1$, $F = 4.70$, $p = 0.008$) was already different from the control at day 0, while S100 had different FA composition from the control ($Df = 1$, $F = 7.35$, $p = 0.003$) after 7 days of S-metolachlor exposure. These results suggest that S-metolachlor affected the fatty acid profile of periphyton after only 7 days of exposure.

Linear regressions showed an effect of S-metolachlor contamination on the FA composition of the periphyton




Fig. 6 Linear regressions based on differences in proportions ($\Delta\%$) for the main fatty acid groups as a function of measured concentrations of S-metolachlor ($\mu\text{g.L}^{-1}$) after 7 days and 14 days of exposure

(Fig. 6; linear regressions for some specific FAs are shown in Fig. S5). Specifically, SFAs increased along the S-metolachlor gradient (Df = 12, $F = 18.70$, $R^2 = 0.60$; $p < 0.001$) after 14 days of exposure. MUFAs did not vary with exposure concentrations, while PUFAs marginally decreased with increasing S-metolachlor concentrations after 7 days of exposure (Df = 12, $F = 3.23$, $R^2 = 0.24$; $p = 0.08$) and then significantly decreased after 14 days (Df = 12, $F = 5.05$, $R^2 = 0.30$; $p = 0.04$). Finally, VLCFAs decreased with increasing herbicide concentration after

7 days (Df = 12, $F = 5.85$, $R^2 = 0.33$; $p = 0.03$). This relationship was stronger after 14 days (Df = 12, $F = 19.61$, $R^2 = 0.62$; $p < 0.001$), where a delta of 10% between the highest concentration and the control was observed.

Discussion

Despite high variability observed in photoautotroph community structure and FA composition, results showed some

effects of herbicide exposure. Photoautotroph related chlorophyll a fluorescence tended to increase with the increase of atrazine concentration after 7 days of exposure. In contrast, S-metolachlor did not clearly affect periphyton fluorescence. As periphytic biofilms are very heterogeneous, fluorescence and fatty acid data showed large intra-condition variability. Despite marked variability, results showed that the two herbicides, in particular S-metolachlor, affected fatty acid profiles. S-metolachlor had a stronger effect than atrazine, with a greater effect after 14 days of exposure compared to 7 days. In particular, the S500 condition showed a 30% increase in SFAs and a 34% decrease in VLCFAs proportions compared to the control condition.

Effects of herbicides on photoautotrophs chlorophyll a fluorescence

S-metolachlor exposure did not clearly affect periphyton biomass as measured by chlorophyll a fluorescence. Indeed, total chlorophyll a fluorescence and specific diatom fluorescence decreased over time under all concentrations including the control condition. Our finding of no significant herbicide effect is in contrast with several studies that showed chloroacetamide herbicides to decrease photoautotroph growth and chlorophyll a fluorescence. For example, Thakkar et al. (2013) showed that the exposure of the marine chlorophyte *Dunaliella tertiolecta* to a high metolachlor concentration (1 mg.L^{-1}) led to a decrease in chlorophyll a and b fluorescence and inhibited cell growth. Likewise, Coquillé et al. (2015) showed a decrease in chlorophyll a content in a freshwater diatom culture (*Gomphonema gracile*) after 7 days of exposure to $100 \text{ }\mu\text{g.L}^{-1}$ of S-metolachlor. The limited effect of S-metolachlor on photoautotrophic groups could be linked to the periphyton matrix that is composed by extracellular polymeric substances (EPS) which may represent up to 90% of the dry mass (Flemming and Wingender 2010). The EPS matrix has several functional groups allowing for the sorption of nutrients and xenobiotics, but can also form a protective layer for the biofilm cells against substances such as pesticides (Melo et al. 2022). The overall decrease in periphyton chlorophyll a fluorescence (i.e biomass) that we observed over time is likely due to the age of the periphyton in our experiment. The colonization time of periphyton varies between 2 and 4 weeks (Cattaneo and Amireault 1992), and is followed by a biomass loss phase after 4–5 weeks (Trbojević et al. 2017). In order to have sufficient biomass for fatty acid analyses, periphyton was contaminated after 4 weeks of colonization and growth. As the experiment lasted an additional 14 days, the periphyton may have started a senescence phase, with potential detachment of biomass under all treatment conditions (Boulêtreau et al. 2006).

In contrast, atrazine increased diatoms and cyanobacteria biomass measured by chlorophyll a fluorescence. The

increase in chlorophyll fluorescence can be linked to the mode of action of atrazine. When photosynthesis is proceeding normally, several steps contribute to the creation of an electron flow between different elements of the thylakoid membranes where the inhibition of photosystem II (PSII) by atrazine takes place. Atrazine competes with plastoquinone for the quinone binding site on the D1 protein (QB site) in PSII, interrupting the electron flow from plastoquinone QA to QB (Rea et al. 2009) leading to the re-emission of excitation energy as fluorescence (Muller et al. 2008) which is then captured by our measuring device. The increase in chlorophyll a fluorescence could also be linked to an increase of chlorophyll cell content. When exposed to atrazine, autotrophs within the periphyton may physiologically adapt to stress by increasing chlorophyll a content per cell (Pannard et al. 2009) to increase the number of photosystems. This “shade-adaptation” response may be a strategy to compensate for the inhibition of photosynthesis and has previously been documented to occur in response to other PSII inhibitor herbicides (e.g., diuron; Chesworth et al. 2004; Proia et al. 2011; Ricart et al. 2009). Given the mode of action of atrazine, the increase in chlorophyll a fluorescence could be taken as evidence of an atrazine effect on the periphyton.

In addition to affecting the fluorescence of photosynthetic organisms, the presence of herbicides may select for more resistant/tolerant taxa (Murdock et al. 2013), thus modifying the community structure of the periphyton (Schmitt-Jansen and Altenburger 2005). We found that atrazine exposure increased the chlorophyll a fluorescence of cyanobacteria in periphyton. Cyanobacteria may be more tolerant to atrazine and more competitive than diatoms and green algae as they have the potential to adapt to photosynthesis inhibition by the use of alternative carbon fixation pathways (Egorova and Bukhov 2006). This is consistent with Pannard et al. (2009) who showed that chronic exposure to atrazine (0.1, 1, and $10 \text{ }\mu\text{g.L}^{-1}$ for 7 weeks) led to a change in microalgal populations with the selection of opportunistic resistant species, some of which were cyanobacteria. Herbicides could also decrease competition for nutrients or increase labile carbon released after cell death further stimulating bacterial production (Downing et al. 2004).

The response of periphyton to herbicide contamination may be reversible as recovery from herbicide exposure as been observed (King et al. 2016; Morin et al. 2010; Prosser et al. 2013). For example, Laviale et al. (2011) showed that a 7 h exposure to atrazine affects the effective and optimal quantum yields of PSII photochemistry of periphyton, but that these effects were reversible within 12 h. However, Proia et al. (2011) mentioned that although the biofilm has a high recovery potential, pulses of contamination longer than a few hours could result in more persistent effects on the biofilm.

Effects of herbicides on periphyton fatty acid composition

Herbicide exposure caused different changes in the FA composition of periphyton with atrazine having little effect on FA composition compared to S-metolachlor. In particular, periphyton SFAs increased with S-metolachlor concentrations, while PUFAs and VLCFAs decreased. The decrease in VLCFAs proportion with the increase in S-metolachlor concentration exposure is consistent with previous results from Böger (2003), who showed a 68% inhibition in VLCFAs of *Scenedesmus acutus* (green algae) after exposure to 283 $\mu\text{g}\cdot\text{L}^{-1}$ of S-metolachlor. VLCFAs ($C \geq 20$) have a structural role in membranes (Bach et al. 2011; Vallotton et al. 2008). S-metolachlor binds to the fatty acid elongation synthase (FAE1-synthase) and inhibits the formation of VLCFAs, which can then affect the rigidity and permeability of cell membranes, resulting in increased cell size and impaired cell division (Matthes and Böger 2002; Thakkar et al. 2013). Even at lower concentrations of exposure (10 $\mu\text{g}\cdot\text{L}^{-1}$), Demailly et al. (2019) experimentally showed that S-metolachlor significantly increased the saturated fatty acid C16:0 and decreased PUFAs including C18:4n3 and C20:4n6 of the diatom *Gomphonema gracile* after 1 week of exposure. The loss of PUFAs observed here and in past studies may be due to the ability of S-metolachlor to increase ROS production (e.g., singlet oxygen $^1\text{O}_2$) resulting in the peroxidation of unsaturated fatty acids in lipid membranes. More specifically, these ROS remove hydrogen from the unsaturated chain of PUFAs constituting the lipids, leading to the loss of membrane integrity (Maronić et al. 2018), in turn jeopardizing the functioning of the cell (Garg and Manchanda 2009). In response to stress, algae often produce triacylglycerols (TAGs) (Nakamura and Li-Beisson, 2016; Shanta et al. 2021). TAGs are considered as carbon and energy storage products (Morales et al. 2021) and are used to maintain bioenergetic stability in the cell. SFAs and MUFAs such as C16:0, C18:0 and C18:1 are among the main components of triacylglycerols (TAGs). The increase in SFAs (e.g., C18:0) and the decrease in some long-chain UFAs that we observed in our experiment could therefore suggest a protective response of the cells against membrane S-metolachlor damages (Kabra et al. 2014).

Herbicides can also have an indirect effect on the FA composition of periphyton by altering the taxonomic composition of periphyton communities. Indeed, different taxonomic groups in the periphyton complex have different fatty acid profiles. For example, diatoms are particularly rich in EPA (C20:5n3) (Drerup and Vis, 2016) and green algae are rich in ALA (C18:3n3) (Genter and Lehman, 2000), while the SFA C16:0 (palmitic acid) is important for the structure of phospholipid membranes in prokaryotes

(Rock, 2008). Changes in the proportion of fatty acids may thus reflect herbicide-induced changes in the composition of the periphyton communities. More specifically, it is possible that the increase in SFAs with atrazine exposure may be due to an increase in bacteria resulting from reduced competition with photosynthetic organisms impacted by the contaminant as we observed increased cyanobacteria biomass (measured by chlorophyll a fluorescence and expressed in $\mu\text{gchl}\cdot\text{cm}^{-2}$) by this contaminant (Fig. S3). Nevertheless, this increase in cyanobacteria was hardly detectable in the FA profiles, where no significant changes in C18:2n6 and C18:3n3 were observed despite the fact that cyanobacteria are generally rich in these C18 PUFAs (Desvillettes et al. 1997). At present, it is still unclear what level of organization (i.e., from the cellular level to subtle changes at the community level) is responsible for the changes in the FA composition of periphyton highlighted by our experiment. It would then be useful to carry out further studies to use more endpoints such as specific composition and the number of cells per autotrophic group.

As previously mentioned, some studies suggest a recovery of periphyton structure and function after a post-exposure recovery phase to herbicides. Most studies used endpoints such as photosynthetic parameters, biomass, or taxonomic composition. However, to our knowledge, there are no studies using biofilm fatty acid composition to monitor recovery nor focusing on how recovery time may affect consumer organisms. In agricultural streams, pesticides contamination generally occurs by pulse, via surface runoff processes. Pulse exposure can occur in various scenarios, depending on its intensity (pulse height), duration (pulse width) and frequency/recovery time (Chèvre and Vallotton, 2013). As a result, reproducing realistic pulse exposure scenarios in a laboratory context may be complicated by logistical constraints. As only few studies if any have examined the influence of atrazine and S-metolachlor on fatty acid profiles in biofilms, using a continuous exposure approach provides a simpler way of controlling experimental conditions. However, to reproduce environmentally realistic conditions, it would be interesting to consider chronic multi-pulse exposure experiments including biofilm recovery phases (Giddings et al. 2018; King et al. 2016).

Conclusion

Periphyton plays a key role in the structure and function of aquatic ecosystems. The nutritional quality of periphyton is essential for the development of primary consumers and can be used as indicator of ecosystem health (Desvillettes et al. 1997). Fatty acids are key nutritional compounds transferred through trophic interactions that are sensitive to various environmental contaminants. We investigated the effects of two commonly used agricultural herbicides,

atrazine and S-metolachlor on periphyton and found that the two herbicides acted differently on the periphyton photoautotroph chlorophyll a fluorescence and fatty acid composition suggesting that there is no standard pattern of herbicide effects on stream periphytic communities. Fluorescence measurements provided information on changes in the relative proportion of the photoautotrophic groups (i.e., green algae, diatoms and cyanobacteria) within periphyton, however, we were limited in our quantification of heterotrophs. Considering that bacteria account for a large amount of biofilm mass (Ricart et al. 2009), are involved in nutrient cycles and can affect the fate of herbicides in water and within the biofilm, future studies should investigate the heterotrophic compartment of the biofilm, especially by DNA sequencing or the study of the metabolism of bacteria. The widespread presence of these two herbicides in rivers raises the question of their toxicity to non-target aquatic organisms and their interaction with the many other molecules present in water (i.e., antagonist, additive or synergistic effects) (Glinski et al. 2018). This study supports the interest to use fatty acids as biomarkers (Gugger 2002; Lang et al. 2011; Maltsev and Maltseva 2021; Shen et al. 2016) in the context of pesticide effect assessment (Filimonova et al. 2016; Gonçalves et al. 2021) but also as a tool for water quality biomonitoring (George et al. 2016).

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1007/s10646-024-02738-y>.

Acknowledgements The authors would like to thank Stéphane Moïse from the general laboratory at INRS-EET for his help on herbicide analysis. We would also like to thank Nolan Pearce for English revisions as well as the Groupe de recherche interuniversitaire en limnologie (GRIL).

Author contributions All authors made substantial contributions to this paper. L.M. was in charge of experimental conceptualization, laboratory experiments, sample collection, data analysis, and writing. S.M. was involved in experimental conceptualization, project management, reviewing, and editing. I.L. was responsible for funding acquisition project administration and was involved in the project conception, experimental conceptualization, reviewing, and editing.

Funding We would like to thank the Fonds de recherche du Québec (FRQNT) for a grant to IL (FRQNT Relève professorale; 2021-NC-285440) and the Centre de recherche en écotoxicologie du Québec (EcotoQ) for funding to LM.

Compliance with ethical standards

Conflict of interest The authors declare no competing interests.

References

- Bach L, Gissot L, Marion J, Tellier F, Moreau P, Satiat-Jeunemaitre B, Palauqui J-C, Napier JA, Faure J-D (2011) Very-long-chain fatty acids are required for cell plate formation during cytokinesis in *Arabidopsis thaliana*. *J Cell Sci* 124:3223–3234. <https://doi.org/10.1242/jcs.074575>
- Bachetti RA, Urseler N, Morgante V, Damilano G, Porporatto C, Agostini E, Morgante C (2021) Monitoring of Atrazine pollution and its spatial-seasonal variation on surface water sources of an agricultural river basin. *Bull Environ Contam Toxicol* 106:929–935. <https://doi.org/10.1007/s00128-021-03264-x>
- Battaglin WA, Furlong ET, Burkhardt MR, Peter CJ (2000) Occurrence of sulfonyleurea, sulfonamide, imidazolinone, and other herbicides in rivers, reservoirs and groundwater in the Midwestern United States, 1998. *Sci Total Environ* 248:123–133. [https://doi.org/10.1016/S0048-9697\(99\)00536-7](https://doi.org/10.1016/S0048-9697(99)00536-7)
- Battaglin WA, Thurman EM, Kalkhoff SJ, Porter SD (2003) Herbicides and transformation products in surface waters of the midwestern United States. *J Am Water Resour Assoc* 39:743–756. <https://doi.org/10.1111/j.1752-1688.2003.tb04402.x>
- Baxter L, Brain RA, Lissimore L, Solomon KR, Hanson ML, Prosser RS (2016) Influence of light, nutrients, and temperature on the toxicity of atrazine to the algal species *Raphidocelis subcapitata*: implications for the risk assessment of herbicides. *Ecotoxicol Environ Saf* 132:250–259. <https://doi.org/10.1016/j.ecoenv.2016.06.022>
- Böger P (2003) Mode of action for chloroacetamides and functionally related compounds. *J. Pestic. Sci.* 28:324–329. <https://doi.org/10.1584/jpestics.28.324>
- Boulétreau S, Garabetian F, Sauvage S, Sanchez-Perez J-M (2006) Assessing the importance of a self-generated detachment process in river biofilm models. *Freshw Biol* 51:901–912. <https://doi.org/10.1111/j.1365-2427.2006.01541.x>
- Brain RA, Anderson JC (2019) The agro-enabled urban revolution, pesticides, politics, and popular culture: a case study of land use, birds, and insecticides in the USA. *Environ Sci Pollut Res* 26:21717–21735. <https://doi.org/10.1007/s11356-019-05305-9>
- Brett M, Müller-Navarra D (1997) The role of highly unsaturated fatty acids in aquatic foodweb processes. *Freshw Biol* 38:483–499. <https://doi.org/10.1046/j.1365-2427.1997.00220.x>
- Cattaneo A, Amireault MC (1992) How artificial are artificial substrata for periphyton? *J N Am Benthol Soc* 11:244–256. <https://doi.org/10.2307/1467389>
- Chesworth JC, Donkin ME, Brown MT (2004) The interactive effects of the antifouling herbicides Irgarol 1051 and Diuron on the seagrass *Zostera marina* (L.). *Aquat Toxicol* 66:293–305. <https://doi.org/10.1016/j.aquatox.2003.10.002>
- Chèvre N, Vallotton N (2013) Pulse exposure in ecotoxicology. In: Féraud J-F, Blaise C (eds) *Encyclopedia of aquatic ecotoxicology*. Springer Netherlands, Dordrecht, pp. 917–926. https://doi.org/10.1007/978-94-007-5704-2_84
- Coquillé N, Jan G, Moreira A, Morin S (2015) Use of diatom motility features as endpoints of metolachlor toxicity. *Aquat Toxicol* 158:202–210. <https://doi.org/10.1016/j.aquatox.2014.11.021>
- Da Costa F, González-Araya R, Robert R (2023) Using combinations of microalgae to condition European flat oyster (*Ostrea edulis*) broodstock and feed the larvae: effects on reproduction, larval production and development. *Aquaculture* 568:739302. <https://doi.org/10.1016/j.aquaculture.2023.739302>
- de Albuquerque FP, de Oliveira JL, Moschini-Carlos V, Fraceto LF (2020) An overview of the potential impacts of atrazine in aquatic environments: Perspectives for tailored solutions based on nanotechnology. *Sci Total Environ* 700:134868. <https://doi.org/10.1016/j.scitotenv.2019.134868>
- Debenest T, Pinelli E, Coste M, Silvestre J, Mazzella N, Madigou C, Delmas F (2009) Sensitivity of freshwater periphytic diatoms to agricultural herbicides. *Aquat Toxicol* 93:11–17. <https://doi.org/10.1016/j.aquatox.2009.02.014>
- DeLorenzo ME, Lauth J, Pennington PL, Scott GI, Ross PE (1999) Atrazine effects on the microbial food web in tidal creek

- mesocosms. *Aquat Toxicol* 46:241–251. [https://doi.org/10.1016/S0166-445X\(98\)00132-5](https://doi.org/10.1016/S0166-445X(98)00132-5)
- Demailly F, Elfeky I, Malbezin L, Le Guédard M, Eon M, Bessoule J-J, Feurtet-Mazel A, Delmas F, Mazzella N, Gonzalez P, Morin S (2019) Impact of diuron and S-metolachlor on the freshwater diatom *Gomphonema gracile*: complementarity between fatty acid profiles and different kinds of ecotoxicological impact-endpoints. *Sci Total Environ* 688:960–969. <https://doi.org/10.1016/j.scitotenv.2019.06.347>
- Desvillettes CH, Bourdier G, Amblard CH, Barth B (1997) Use of fatty acids for the assessment of zooplankton grazing on bacteria, protozoans and microalgae. *Freshw Biol* 38:629–637. <https://doi.org/10.1046/j.1365-2427.1997.00241.x>
- Downing HF, Delorenzo ME, Fulton MH, Scott GI, Madden CJ, Kucklick JR (2004) Effects of the agricultural pesticides atrazine, chlorothalonil, and endosulfan on South Florida microbial assemblages. *Ecotoxicology* 13:245–260. <https://doi.org/10.1023/B:ECTX.0000023569.46544.9f>
- Drerup SA, Vis ML (2016) Responses of Stream biofilm phospholipid fatty acid profiles to acid mine drainage impairment and remediation. *Water Air Soil Pollut* 227:159. <https://doi.org/10.1007/s11270-016-2856-5>
- Egorova EA, Bukhov NG (2006) Mechanisms and functions of photosystem I-related alternative electron transport pathways in chloroplasts. *Russ J Plant Physiol* 53:571–582. <https://doi.org/10.1134/S1021443706050013>
- European Commission (2004) Commission Regulation (EC) No 775/2004 of 26 April 2004 amending Annex I to Regulation (EC) No 304/2003 of the European Parliament and of the Council concerning the export and import of dangerous chemicals (Text with EEA relevance), CE.
- Fadhlaoui M, Laderrière V, Lavoie I, Fortin C (2020) Influence of temperature and nickel on algal biofilm fatty acid composition. *Environ Toxicol Chem* 39:1566–1577. <https://doi.org/10.1002/etc.4741>
- FAO (2022) Pesticides use pesticides trade and pesticides indicators. FAO. <https://doi.org/10.4060/cc0918en>
- Filimonova V, Gonçalves F, Marques JC, De Troch M, Gonçalves AMM (2016) Fatty acid profiling as bioindicator of chemical stress in marine organisms: a review. *Ecol Indic* 67:657–672. <https://doi.org/10.1016/j.ecolind.2016.03.044>
- Flemming H-C, Wingender J (2010) The biofilm matrix. *Nat Rev Microbiol* 8:623–633. <https://doi.org/10.1038/nrmicro2415>
- Fortier A (2018) Règlement modifiant le Code de gestion des pesticides 5.
- Gao Y, Fang Jianguang, Li W, Wang X, Li F, Du M, Fang Jinghui, Lin F, Jiang W, Jiang Z (2019) Effects of atrazine on the physiology, sexual reproduction, and metabolism of eelgrass (*Zostera marina* L.). *Aquat Botany* 153:8–14. <https://doi.org/10.1016/j.aquabot.2018.10.002>
- Garg N, Manchanda G (2009) ROS generation in plants: boon or bane? *Plant Biosyst* 143:81–96. <https://doi.org/10.1080/11263500802633626>
- Genter RB, Lehman RM (2000) Metal toxicity inferred from algal population density, heterotrophic substrate use, and fatty acid profile in a small stream. *Environ Toxicol Chem* 19:869–878. <https://doi.org/10.1002/etc.5620190413>
- George SD, Ernst AG, Baldigo BP, Honeyfield DC (2016) Response of periphyton fatty acid composition to supplemental flows in the upper Esopus Creek, Catskill Mountains, New York: U.S. Geological Survey Scientific Investigations Report (Scientific Investigations Report). U.S. Geological Survey, Reston, Virginia. <https://doi.org/10.3133/sir20155161>
- Giddings JM, Campana D, Nair S, Brain R (2018) Data quality scoring system for microcosm and mesocosm studies used to derive a level of concern for atrazine: atrazine microcosm and mesocosm data quality scoring. *Integr Environ Assess Manag* 14:489–497. <https://doi.org/10.1002/ieam.4050>
- Gladyshev MI, Sushchik NN, Anishchenko OV, Makhutova ON, Kolmakov VI, Kalachova GS, Kolmakova AA, Dubovskaya OP (2011) Efficiency of transfer of essential polyunsaturated fatty acids versus organic carbon from producers to consumers in a eutrophic reservoir. *Oecologia* 165:521–531. <https://doi.org/10.1007/s00442-010-1843-6>
- Glinski DA, Purucker ST, Van Meter RJ, Black MC, Henderson WM (2018) Analysis of pesticides in surface water, stemflow, and throughfall in an agricultural area in South Georgia, USA. *Chemosphere* 209:496–507. <https://doi.org/10.1016/j.chemosphere.2018.06.116>
- Gonçalves AMM, Rocha CP, Marques JC, Gonçalves FJM (2021) Fatty acids as suitable biomarkers to assess pesticide impacts in freshwater biological scales—a review. *Ecol Indic* 122:107299. <https://doi.org/10.1016/j.ecolind.2020.107299>
- Green JW, Springer TA, Holbech H (2018) *Statistical Analysis of Ecotoxicity Studies*, 1st edn. John Wiley & Sons, Hoboken, NJ
- Griffini O, Bao ML, Barbieri D, Pantani F (1997) Occurrence of pesticides in the arno river and in potable water—a survey of the period 1992–1995. *Bull Environ Contam Toxicol* 59(2):202–209
- Groner ML, Relyea RA (2011) A tale of two pesticides: how common insecticides affect aquatic communities: a tale of two pesticides. *Freshw Biol* 56:2391–2404. <https://doi.org/10.1111/j.1365-2427.2011.02667.x>
- Gugger M (2002) Cellular fatty acids as chemotaxonomic markers of the genera *Anabaena*, *Aphanizomenon*, *Microcystis*, *Nostoc* and *Planktothrix* (cyanobacteria). *Int J Syst Evolut Microbiol* 52:1007–1015. <https://doi.org/10.1099/ijs.0.01917-0>
- Hansen SP, Messer TL, Mittelstet AR (2019) Mitigating the risk of atrazine exposure: identifying hot spots and hot times in surface waters across Nebraska, USA. *J Environ Manag* 250:109424. <https://doi.org/10.1016/j.jenvman.2019.109424>
- HRAC (2020) Global herbicide classification lookup. Herbicide Resistance Action Committee
- Huggins K, Frenette J-J, Arts MT (2004) Nutritional quality of biofilms with respect to light regime in Lake Saint-Pierre (Quebec, Canada). *Freshw Biol* 49:945–959. <https://doi.org/10.1111/j.1365-2427.2004.01236.x>
- Kabra AN, Ji M-K, Choi J, Kim JR, Govindwar SP, Jeon B-H (2014) Toxicity of atrazine and its bioaccumulation and biodegradation in a green microalga, *Chlamydomonas mexicana*. *Environ Sci Pollut Res* 21:12270–12278. <https://doi.org/10.1007/s11356-014-3157-4>
- Kapsi M, Tsoutsis C, Paschalidou A, Albanis T (2019) Environmental monitoring and risk assessment of pesticide residues in surface waters of the Louros River (N.W. Greece). *Sci Total Environ* 650:2188–2198. <https://doi.org/10.1016/j.scitotenv.2018.09.185>
- King RS, Brain RA, Back JA, Becker C, Wright MV, Toteu Djomte V, Scott WC, Virgil SR, Brooks BW, Hosmer AJ, Chambliss CK (2016) Effects of pulsed atrazine exposures on autotrophic community structure, biomass, and production in field-based stream mesocosms: pulsed atrazine exposures. *Environ Toxicol Chem* 35:660–675. <https://doi.org/10.1002/etc.3213>
- Konschak M, Zubrod JP, Duque Acosta TS, Bouchez A, Kroll A, Feckler A, Röder N, Baudy P, Schulz R, Bundschuh M (2021) Herbicide-induced shifts in the periphyton community composition indirectly affect feeding activity and physiology of the gastropod grazer *Physella acuta*. *Environ. Sci. Technol.* 55:14699–14709. <https://doi.org/10.1021/acs.est.1c01819>
- Lang I, Hodac L, Friedl T, Feussner I (2011) Fatty acid profiles and their distribution patterns in microalgae: a comprehensive analysis of more than 2000 strains from the SAG culture collection. *BMC Plant Biol* 11:124. <https://doi.org/10.1186/1471-2229-11-124>

- Larras F, Billoir E, Baillard V, Siberchicot A, Scholz S, Wubet T, Tarkka M, Schmitt-Jansen M, Delignette-Muller M-L (2018) DRomics: a turnkey tool to support the use of the dose–response framework for omics data in ecological risk assessment. *Environ. Sci. Technol.* 52:14461–14468. <https://doi.org/10.1021/acs.est.8b04752>
- Laviale M, Morin S, Créach A (2011) Short term recovery of periphyton photosynthesis after pulse exposition to the photosystem II inhibitors atrazine and isoproturon. *Chemosphere* 84:731–734. <https://doi.org/10.1016/j.chemosphere.2011.03.035>
- LAWA (2019) Rapport sur la qualité des eaux souterraines—produits phytosanitaires. Länderarbeitsgemeinschaft Wasser (LAWA)—Sous-comité Produits phytosanitaires dans les eaux souterraines
- Li H-Y, Lu Y, Zheng J-W, Yang W-D, Liu J-S (2014) Biochemical and genetic engineering of diatoms for polyunsaturated fatty acid biosynthesis. *Mar Drugs* 12:153–166. <https://doi.org/10.3390/md12010153>
- Maltsev Y, Maltseva K (2021) Fatty acids of microalgae: diversity and applications. *Rev Environ Sci Biotechnol* 20:515–547. <https://doi.org/10.1007/s11157-021-09571-3>
- Matthes B, Böger P (2002) Chloroacetamides affect the plasma membrane. *Zeitschrift für Naturforschung C* 57:843–852. <https://doi.org/10.1515/znc-2002-9-1015>
- Melo A, Quintelas C, Ferreira EC, Mesquita DP (2022) The role of extracellular polymeric substances in micropollutant removal. *Front. Chem. Eng.* 4:778469. <https://doi.org/10.3389/fceng.2022.778469>
- Morales M, Aflalo C, Bernard O (2021) Microalgal lipids: a review of lipids potential and quantification for 95 phytoplankton species. *Biomass Bioenergy* 150:106108. <https://doi.org/10.1016/j.biombioe.2021.106108>
- Morin S, Pesce S, Tlili A, Coste M, Montuelle B (2010) Recovery potential of periphytic communities in a river impacted by a vineyard watershed. *Ecol Indic* 10:419–426. <https://doi.org/10.1016/j.ecolind.2009.07.008>
- Muller R, Schreiber U, Escher BI, Quayle P, Bengtson Nash SM, Mueller JF (2008) Rapid exposure assessment of PSII herbicides in surface water using a novel chlorophyll a fluorescence imaging assay. *Sci Total Environ* 401:51–59. <https://doi.org/10.1016/j.scitotenv.2008.02.062>
- Müller-Navarra DC, Brett MT, Liston AM, Goldman CR (2000) A highly unsaturated fatty acid predicts carbon transfer between primary producers and consumers. *Nature* 403:74–77. <https://doi.org/10.1038/47469>
- Murdock JN, Shields FD, Lizotte RE (2013) Periphyton responses to nutrient and atrazine mixtures introduced through agricultural runoff. *Ecotoxicology* 22:215–230. <https://doi.org/10.1007/s10646-012-1018-9>
- Nakamura Y, Li-Beisson Y (eds) (2016) Lipids in plant and algae development, subcellular biochemistry. Springer International Publishing, Cham. <https://doi.org/10.1007/978-3-319-25979-6>
- Ou-Yang K, Feng T, Han Y, Li G, Li J, Ma H (2022) Bioaccumulation, metabolism and endocrine-reproductive effects of metolachlor and its S-enantiomer in adult zebrafish (*Danio rerio*). *Sci Total Environ* 802:149826. <https://doi.org/10.1016/j.scitotenv.2021.149826>
- Pannard A, Le Rouzic B, Binet F (2009) Response of phytoplankton community to low-dose atrazine exposure combined with phosphorus fluctuations. *Arch Environ Contam Toxicol* 57:50–59. <https://doi.org/10.1007/s00244-008-9245-z>
- Parlakidis P, Rodriguez MS, Gikas GD, Alexoudis C, Perez-Rojas G, Perez-Villanueva M, Carrera AP, Fernández-Cirelli A, Vryzas Z (2022) Occurrence of banned and currently used herbicides, in groundwater of Northern Greece: a human health risk assessment approach. *IJERPH* 19:8877. <https://doi.org/10.3390/ijerph19148877>
- Perkins DB, Chen W, Jacobson A, Stone Z, White M, Christensen B, Ghebremichael L, Brain R (2021) Development of a mixed-source, single pesticide database for use in ecological risk assessment: quality control and data standardization practices. *Environ Monit Assess* 193:827. <https://doi.org/10.1007/s10661-021-09596-9>
- Proia L, Morin S, Peipoch M, Romaní AM, Sabater S (2011) Resistance and recovery of river biofilms receiving short pulses of Triclosan and Diuron. *Sci Total Environ* 409:3129–3137. <https://doi.org/10.1016/j.scitotenv.2011.05.013>
- Prosser RS, Brain RA, Hosmer AJ, Solomon KR, Hanson ML (2013) Assessing sensitivity and recovery of field-collected periphyton acutely exposed to atrazine using PSII inhibition under laboratory conditions. *Ecotoxicology* 22:1367–1383. <https://doi.org/10.1007/s10646-013-1123-4>
- Quintaneiro C, Patrício D, Novais SC, Soares AMVM, Monteiro MS (2017) Endocrine and physiological effects of linuron and S-metolachlor in zebrafish developing embryos. *Sci Total Environ* 586:390–400. <https://doi.org/10.1016/j.scitotenv.2016.11.153>
- Rea G, Politicelli F, Antonacci A, Scognamiglio V, Katiyar P, Kulkarni SA, Johanningmeier U, Giardi MT (2009) Structure-based design of novel *Chlamydomonas reinhardtii* D1-D2 photosynthetic proteins for herbicide monitoring. *Protein Sci* 18:2139–2151. <https://doi.org/10.1002/pro.228>
- Relyea RA (2009) A cocktail of contaminants: how mixtures of pesticides at low concentrations affect aquatic communities. *Oecologia* 159:363–376. <https://doi.org/10.1007/s00442-008-1213-9>
- Ricart M, Barceló D, Geiszinger A, Guasch H, Alda MLde, Romaní AM, Vidal G, Villagrana M, Sabater S (2009) Effects of low concentrations of the phenylurea herbicide diuron on biofilm algae and bacteria. *Chemosphere* 76:1392–1401. <https://doi.org/10.1016/j.chemosphere.2009.06.017>
- Rock CO (2008) Fatty acid and phospholipid metabolism in prokaryotes. In: *Biochemistry of lipids, lipoproteins and membranes*. Elsevier, pp 59–96. <https://doi.org/10.1016/B978-044453219-0.50005-2>
- Rossoll D, Bermúdez R, Hauss H, Schulz KG, Riebesell U, Sommer U, Winder M (2012) Ocean acidification-induced food quality deterioration constrains trophic transfer. *PLoS ONE* 7:e34737. <https://doi.org/10.1371/journal.pone.0034737>
- Roubeix V, Fauvelle V, Tison-Rosebery J, Mazzella N, Coste M, Delmas F (2012) Assessing the impact of chloroacetanilide herbicides and their metabolites on periphyton in the Leyre River (SW France) via short term growth inhibition tests on autochthonous diatoms. *J. Environ. Monit.* 14:1655. <https://doi.org/10.1039/c2em10887a>
- Schmitt-Jansen M, Altenburger R (2005) Predicting and observing responses of algal communities to photosystem ii–herbicide exposure using pollution-induced community tolerance and species-sensitivity distributions. *Environ Toxicol Chem* 24:304. <https://doi.org/10.1897/03-647.1>
- Sénat de France, 2003. Annexe 47 (Atrazine) du rapport d’office parlementaire sur la qualité de l’eau et assainissement en France.
- Shanta PV, Li B, Stuart DD, Cheng Q (2021) Lipidomic Profiling of Algae with Microarray MALDI-MS toward Ecotoxicological Monitoring of Herbicide Exposure. *Environ. Sci. Technol.* 55:10558–10568. <https://doi.org/10.1021/acs.est.1c01138>
- Shen P-L, Wang H-T, Pan Y-F, Meng Y-Y, Wu P-C, Xue S (2016) Identification of characteristic fatty acids to quantify triacylglycerols in microalgae. *Front. Plant Sci* 7. <https://doi.org/10.3389/fpls.2016.00162>
- Špoljarić Maronić D, Štolfa Čamagajevac I, Horvatić J, Žuna Pfeiffer T, Stević F, Žarković N, Waeg G, Jaganjac M (2018) S-metolachlor promotes oxidative stress in green microalga *Parachlorella kessleri* - A potential environmental and health risk

- for higher organisms. *Sci Total Environ* 637–638:41–49. <https://doi.org/10.1016/j.scitotenv.2018.04.433>
- Székács A, Mörzl M, Darvas B (2015) Monitoring pesticide residues in surface and ground water in Hungary: surveys in 1990–2015. *J Chem* 2015:1–15. <https://doi.org/10.1155/2015/717948>
- Thakkar M, Randhawa V, Wei L (2013) Comparative responses of two species of marine phytoplankton to metolachlor exposure. *Aquat Toxicol* 126:198–206. <https://doi.org/10.1016/j.aquatox.2012.10.002>
- Thompson FL, Abreu PC, Wasielesky W (2002) Importance of biofilm for water quality and nourishment in intensive shrimp culture. *Aquaculture* 203:263–278. [https://doi.org/10.1016/S0044-8486\(01\)00642-1](https://doi.org/10.1016/S0044-8486(01)00642-1)
- Trbojević I, Jovanović J, Kostić D, Popović S, Krizmanić J, Karadžić V, Subakov Simić G (2017) Structure and succession of periphyton in an urban reservoir: artificial substrate specificity. *Oceano Hydrobiol Stud* 46:379–392. <https://doi.org/10.1515/ohs-2017-0038>
- USEPA (2017) Pesticides industry sales and usage, 2008–2012 Market estimates. United States Environmental Protection Agency.
- USEPA (2016) Refined ecological risk assessment for atrazine. United States Environmental Protection Agency.
- Vallotton N, Moser D, Eggen RIL, Junghans M, Chèvre N (2008) S-metolachlor pulse exposure on the alga *Scenedesmus vacuolatus*: effects during exposure and the subsequent recovery. *Chemosphere* 73:395–400. <https://doi.org/10.1016/j.chemosphere.2008.05.039>
- Vryzas Z, Alexoudis C, Vassiliou G, Galanis K, Papadopoulou-Mourkidou E (2011) Determination and aquatic risk assessment of pesticide residues in riparian drainage canals in northeastern Greece. *Ecotoxicol Environ Saf* 74:174–181. <https://doi.org/10.1016/j.ecoenv.2010.04.011>
- Wetzel RG (ed) (1983) *Periphyton of Freshwater Ecosystems: Proceedings of the First International Workshop on Periphyton of Freshwater Ecosystems held in Växjö, Sweden, 14–17 September 1982*. Springer Netherlands, Dordrecht. <https://doi.org/10.1007/978-94-009-7293-3>
- WSSA (2021) WSSA-Herbicide Site of Action (SOA) Classification List. Weed Science Society of America.
- Zemolin CR, Avila LA, Cassol GV, Massey JH, Camargo ER (2014) Environmental fate of S-Metolachlor: a review. *Planta Daninha* 32:655–664. <https://doi.org/10.1590/S0100-83582014000300022>
- Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.