

# Problem posing based on investigation activities by university students

João Pedro da Ponte · Ana Henriques

Published online: 8 November 2012

© Springer Science+Business Media Dordrecht 2012

**Abstract** This paper reports a classroom-based study involving investigation activities in a university numerical analysis course. The study aims to analyse students' mathematical processes and to understand how these activities provide opportunities for problem posing. The investigations were intended to stimulate students in asking questions, to trigger their thinking processes, to promote their ability to investigate and to support them in learning numerical analysis' concepts and procedures. The results show that the investigations provided opportunities for students to experience mathematical processes, including posing questions, formulating and testing conjectures and, to some extent, proving results. They also provide some understanding about the role of problem posing in these processes. Posing questions occurred mainly in an implicit way, in the interpretation of tasks and in identifying regularities, analysing graphs and testing cases. The conjectures were often based on pattern identification or data manipulation, and the students tended to accept them without testing or proving. The students also proposed alternative formulations for the initial questions and posed new problems from their explorations and attempts to refine previous conjectures.

**Keywords** Problem posing · Investigation activities · Mathematical processes · Reasoning

## 1 Introduction

Mathematics may be regarded as a body of knowledge or as a human activity (Freudenthal, 1973; Pólya, 1945). Whereas the notion of body of knowledge leads to the notion of mathematics teaching as a transmission process, the notion of mathematical activity suggests that students may be actively involved in doing mathematics (National Council of Teachers of Mathematics [NCTM], 1989). Pólya (2002) underlines the importance of such activity when he says: “To understand mathematics means to be able to do mathematics” (p. 7). At the heart of doing mathematics is posing problems and solving them (Brown & Walter, 1990).

---

J. P. da Ponte · A. Henriques (✉)  
Instituto de Educação, Universidade de Lisboa, Lisbon, Portugal  
e-mail: achenriques@ie.ul.pt

J. P. da Ponte  
e-mail: jpponte@ie.ul.pt

Problem posing may be regarded as emphasising the formulation of a key problem that hopefully will trigger an extended and productive mathematical activity, leading to its solution. But problem posing may also be regarded as the constant process of posing questions, which permeates any authentic mathematical activity and generates both key and subsidiary questions. One must note that sometimes subsidiary questions become central and sometimes key questions become dead ends. Adopting this second view, we see mathematical investigations as providing particularly rich opportunities for problem posing (Ponte & Matos, 1992). The close relation to problem posing and the increasing visibility of such activities in curriculum documents (such as NCTM, 2000) and in teachers' practices (Ruthven, Hofmann, & Mercer, 2011) make them an interesting and potentially fruitful field of study.

In this paper, we seek to identify the mathematical processes used by university students exploring investigations in the classroom, paying special attention to students' problem posing processes during this activity, from the interpretation of the situations to the justification of results.

## 2 Problem posing and investigation activities

In teaching and learning contexts, students may work in a way that is similar to professional mathematicians. Such is the view of Hadamard (1945):

Between the work of the student who tries to solve a problem in geometry or algebra and a work of invention, one can say that there is only a difference of degree, a difference of level, both works being of a similar nature. (p. 104)

Working on mathematical investigations provides students with the opportunity to experience important mathematical processes. Ernest (1991) points out problem posing as a first distinctive feature of mathematical investigations, as their statement often is not fully explicit and precise, requiring students to pose their own questions and to establish their own objectives. After exploring the situation, it is necessary to pose questions about the data, frequently based on the identification of patterns, and to formulate and test conjectures (Ponte, Ferreira, Brunheira, Oliveira, & Varandas, 1998). This may show the need to collect more data, to drop the initial conjectures and to formulate new ones. The analysis of the situation may lead to the modification of the initial questions (Silver, 1994) or to a change in some of the conditions to raise new problems. Then, one must establish plausible arguments and formal proofs to reject or validate this guesswork. The conjectures that resist the tests gain credibility, stimulating proofs that, once achieved, yield mathematical validity. Ernest (1991) emphasises that investigations provide stimulus to the students to justify and prove their conjectures and to explain mathematical arguments to their colleagues and to the teacher. Also, the processes of generalization and specialization may be sources not only of solutions but of new problems as well (Pólya, 1981). After obtaining a solution for a particular problem, the student can also "look back" to see how the solution may be affected by changes in the problem (Pólya, 1945). Thus, working on investigations, the students are led to formulate questions and conjectures, conduct tests and refutations and also present results, and discuss and argue with their colleagues and the teacher (Ponte, Brocardo & Oliveira, 2003). An additional feature of this process is that new questions arise along the way.

Carrying out mathematical investigations may significantly contribute to the understanding and consolidation of concepts and to the development of students' mathematical thinking (Henriques & Ponte, 2008; Ponte, 2007). Ponte et al. (1998) also argue that mathematical

investigations provide students with a more complete view of mathematics, thus opposing the narrow view of this subject as just carrying out procedures and algorithms. Moreover, investigations give students many opportunities to pose new problems (English, 2003; Kilpatrick, 1987). Those may arise *prior* to problem solving when students generate problems from a particular situation or *after* solving a problem when the goals or conditions of a solved problem are modified or applied to new situations. In addition, problem posing could occur *during* problem solving when the individuals intentionally change their goals while in the process of solving the problem (Silver, 1994).

Previous studies examined the conceptual benefits of problem posing. Silver (1994) argues that problem posing promotes students' mathematics thinking. English (1997) also points out that problem posing, reinforcing and enriching students' basic mathematics concepts, generates a more diverse and flexible thinking and improves their ability to solve problems. Moreover, Silver (1994) and English (1997) indicate that students who work on problem posing develop a more positive attitude towards mathematics, becoming more responsible and motivated for their learning. From a teaching perspective, problem posing can also become a useful assessment tool enabling teachers to develop a better understanding of students' cognitive processes, to find possible misconceptions early in time and to obtain information on levels of students' learning to adjust their teaching processes (English, 1997). In spite of this, few studies have examined the cognitive processes involved when students generate their own problems in the course of their problem solving activity or about instructional strategies that can effectively promote productive problem posing. Cai and Cifarelli (2005) explained several characteristics of the students' mathematical explorations in open-ended problem situations with particular emphasis on how students formulated and solved new problems that arose during their problem solving activity. The results obtained (also extended in Cifarelli & Cai, 2005) suggest a preliminary model of mathematical exploration processes that, as the authors consider, needs further refinement. In this paper, we seek to develop a more elaborated view concerning the mathematical processes used by students in working on investigations and address in a particular way how such tasks contribute to university students' problem posing processes.

### 3 Methodology

The methodological framework of this study is based on the interpretative research paradigm (Bogdan & Biklen, 1994). Examples in this paper come from a classroom-based study conducted in the first semester of 2008/2009 in a numerical analysis course taught by the second author, in order to promote students' experience of doing mathematics while learning numerical analysis concepts and procedures. The participants were all of the 36 second-year university students of Escola Naval, who had no prior experience in such activities.

The study offered a different (unusual) approach to a numerical analysis course. Classroom work included the realization of four investigation tasks along the course, which took over a significant part of the class time. The realization of each task was organized in three main moments: (1) presenting tasks, (2) exploring and (3) presenting and discussing students' conclusions (Ponte et al., 1998). Each task focused on a different numerical analysis topic (interval arithmetic, non-linear equations, curve fitting and numerical integration). Students were proposed situations for which they had neither theory nor routine

processes to handle, and therefore, they were challenged to pose their own questions and to develop and defend their own strategies.

Each task was presented to the students through a written statement. They worked in self-selected pairs or small groups, constituted at the beginning of the semester. Students could use graphic calculators, if they wanted. Upon finishing the exploration of the task, the students presented their work orally to the class. These discussions were important learning opportunities. The students presented their conclusions and explained their ideas and strategies, and, prompted by their colleagues and the teacher's strong questioning, they sought for justifications and discussed aspects to which they had only given little thought. These moments also provided the opportunity to introduce new topics progressively and to respond to students' questions, probing the understanding of the procedures of their colleagues and asking them to explain their reasoning. Between stages 2 and 3, outside the classroom, the students wrote a group report, explaining their strategies and presenting and justifying their conclusions. The work on this report encouraged students' reflection, since they had to articulate ideas, to explain procedures, and to review the processes used and the results obtained.

The investigation tasks were alternated with lectures to address theoretical aspects of numerical analysis arising during the tasks or to present other topics (numbers and errors, interpolation and differential equations). The classroom work also included moments for solving problems and doing practical exercises for consolidation of concepts and algorithms (see Fig. 1).

Due to the complexity of the complete study (Henriques, 2010), the results reported here focus on three students (Carlos, Gonçalo and Luís) working on two tasks, conveniently chosen in order to illustrate a variety of interesting aspects of problem posing processes.


Since the teacher was also a researcher, several difficulties and conflicts could arise from that dual role. Following the educational research ethical principles, the teacher–researcher informed all the students of the research goals and intended activities, asked their agreement to participate, keeping their identities confidential, and assured students that the research would not bring harm to any student in the classroom.

Data collection methods included observation of students exploring tasks, collection of their written reports (labelled WR#) and audio recording interviews after the exploration of each task (E#). The interviews were based on issues that emerged from the analysis of written reports in order to understand students' thinking processes and to provide data to clarify ambiguous issues.

For each task, we describe the students' work and document it via a set of their comments indicating both the students' goals and their subsequent actions oriented towards the solution. The basis for categorizing the mathematical processes used by the students during the task exploration were the mathematics processes discussed in Ponte et al. (2003), such as the formulation of questions, the formulation of particular/specific conjectures and their generalization to more general cases, the test of conjectures and their posterior justification. We also analysed students' actions to identify other instances that could be considered problem posing processes, such as posing questions about data, generating partial problems from a larger one or modifying the initial goals or conditions of an already solved problem.

#### 4 Students' mathematical processes

Task 1 was a starting point for the development and formalization of numerical methods for solving non-linear equations. In particular, we intended to create an opportunity to discuss



**Fig. 1** The classroom-based study

the bisection method, designing procedures (algorithms) for solving non-linear equations and understanding their rationale.

Consider the function  $f(x) = \ln(x) - e^{-x}$ . Look at the following sequence of real number intervals where the root of  $f$  is included,

[1.000, 2.000] [1.000, 1.500] [1.250, 1.500] [1.250, 1.375] [1.250, 1.313] [1.281, 1.313]

What is the next interval of the sequence? Explain clearly how you got that conclusion.

The three students started exploring the task by observing the sequence of consecutive intervals and looking for regularities. Gonalo explains it in his report: “We tried to find a pattern and understand what was really happening from one interval to another” (WR1). However, the students faced some difficulties since they did not use all of the available information (such as the root of  $f$ ): “Initially, we were a little unmotivated because we could not find a relationship that might look valid to us” (Gonalo, WR1). To solve the original problem, Gonalo formulated two more accessible subproblems: (1) Is there any decreasing pattern in the width of the intervals of the sequence? (2) What is the rule to find the interval bounds? The observation of the sequence of intervals led them to a first conjecture based on the

correct identification of a decreasing pattern of interval widths: “We realized immediately that the intervals were getting smaller and smaller, i.e., the width of the next interval is always half of the previous one and that happens for all the intervals” (Gonalo, E1).

Carlos and Lu s also began to pose a simpler question, trying to identify a pattern for interval widths, but they did not seek to identify a pattern for interval bounds. They calculated the range of each interval and correctly identified a decreasing pattern, formulating a first conjecture: “By looking at the sequence we found that the interval width decreases by half regarding the previous one” (Lu s, WR1).

The formulation of a criterion for deciding on which interval endpoints should be reduced raised some questions to the students because they did not recognize any regularity by observation. Carlos indicated that he and his colleagues made several attempts: “Several ideas came up till we got one that seemed correct” (E1). During this process, the students posed several questions that led them to formulate conjectures about the pattern of interval bounds, most of which were based on counting the number of times each endpoint changed or remained constant:

In these three that kept the upper endpoint of the interval, the lower bound was always decreasing. Three times. Then, it changed and it was the upper endpoint, twice. Thus, for the general case  $[a, b]$ , the minimum (a) varies in three intervals while the maximum (b) had the same value in two consecutive intervals, and then, in the next interval, had another value. (Lu s, E1)

However, these strategies were based on simple observations and counting, and did not allow the students to formulate a correct identification of a rule for the sequence of intervals since they did not use the necessary information about the root of  $f$ . Carlos and Lu s did not test their conjectures and did not realize that these were incorrect. Only when the students were induced to a closer reading of the task statement could they verify, through experimentation, that the intervals that they constructed did not verify all the given conditions: “Only from this reasoning we found that the root of  $f(x)$  was not included in all intervals” (Lu s, E1). This test of their conjecture prompted a reformulation:

It's the element of the interval (maximum or minimum) that was farthest from the root value that varies, while the other remains constant. Following this reasoning, the next interval (...) is:  $1.313 - 1.281 = 0.032$ , so in the following interval will be half of that width, i.e. 0.016. The farthest element from the root value, i.e. 1.281, will be added to 0.016 resulting in 1.297 and this being translated into  $[1.297, 1.313]$ . (Carlos, WR1)

The students now proceeded to verify this conjecture but only to the sequence of intervals presented in the task. However, they did not feel any need to justify it.

Gonalo also formulated his conjecture based on a counting scheme, considering the conditions given in the task:

In the interval 2, [a half range] had to be added, in the intervals 3 and 4 it had to be subtracted, in the intervals 5, 6 and 7 it had to be added again, so, I thought we have the following sequence (...)

1	2	3	4	5	6	7	8	9	10	11	
-	+	-	-	+	+	+	-	-	-	-	(Gonalo, E1)

Unlike his colleagues, Gonalo tested his conjecture and realized that it was incorrect. He used his knowledge on functions to identify the regularity: “As we know that the zero of the function is within the interval, from one interval to the next, we drop the half that does not contain our zero” (E1). Thus, this test allowed him to formulate a new conjecture.

Once the exploration of this task had been completed for the particular case of the given function, the next natural step would be to pose similar questions for any function. In the given function, from one interval to the next, a transformation occurs according to a rule consistent with some conditions. For Carlos and Luís, the generalization of this rule reflected, in part, the previous exploratory work. They considered this process a mere formalization and described their conjecture in symbolic notation:

From what I said earlier, we can define the following rule: If  $x - \max > (f(x) = 0)$ , then the next interval is  $[\min, x - \max]$  or  $[x + \min, \max]$  if and only if  $x - \max < (f(x) = 0)$ . (Luís, WR1)

The students did not realize that their conjecture depends on an unknown root (of this specific function). They felt no need to justify that rule, and therefore, they did not realize that their rule applies only to their particular case but not to the general case.

Gonçalo, in turn, realized that he must return to the previous questions for any function in general. Then he presented a rule for constructing intervals in an algorithmic form, with the elements of the sequence defined by recurrence, which would apply to any continuous function on a range with unknown roots:

As a rule and taking into account how we get the next intervals, given the interval  $[a, b]$  we perform the following steps:

- 1st Find the midpoint,  $vmed = (a+b)/2$
- 2nd Find  $f(vmed)$

If  $f(vmed) > 0$ , then we get the next interval  $[a, vmed]$

If  $f(vmed) < 0$ , then we get the next interval  $[vmed, b]$ . (Gonçalo, RT1)

Gonçalo seemed to understand that a generalization of this conjecture must be applicable to any function. He justified his algorithm when he described the rule for constructing intervals of the sequence based on known mathematical properties and theorems, including the Bolzano theorem and its corollaries:

As the function is continuous and  $f(a) < 0$  and  $f(b) > 0$ , we only had to maintain the [images of the] bounds of the intervals with opposite signs so that the next interval also contains the zero. As  $vmed$  is always an extreme it is enough to confirm the sign of  $f(vmed)$ . (E1)

Thus, intuitively, he built the bisection method for solving non-linear equations.

In summary, the mathematical processes used by the students in exploring this task included looking for regularities, formulating and testing conjectures, generalizing these conjectures and justifying them. All students formulated initial questions on given intervals, formulating conjectures involving patterns from data observation or manipulation. The students had difficulties in identifying patterns, as they did not take into account all information available. Only Gonçalo formulated a more general conjecture concerning all possible functions, and he was also the only one to produce justifications. Students' work on this task included the formulation of questions in an implicit way, identified by their conjectures. The questions arose when students formulated conjectures from examples and generated subproblems from a major problem. In the case of Gonçalo, questions also arose when he tried to generalize his conjecture to obtain a general solution.

In task 2, multiple sets of data, provided in tables, represent different behaviours to be modelled by different functions. Students were required to analyse and identify patterns and then to define criteria in order to complete some missing values in the tables, using known



mathematical models (e.g., linear, quadratic or exponential), together with what they just learned about interpolation.

From three distinct monitoring stations a researcher obtained some data on the evolution of a population of anaerobic bacterias in a lake, with which he intends to describe mathematically the growth of that population.

Station 1

t (hours)	2	3	4	5	6	8
p ( $\times 10^5$ )	90	140	---	240	---	390

Station 2

t (hours)	1	2	3	4	5	6	8
p ( $\times 10^5$ )	40	85	---	220	210	---	400

Station 3

t (hours)	2	3	4	5	6	7
p ( $\times 10^5$ )	85	140	---	250	380	600

- a) There are some missing values corresponding to the records of a few hours. How did you complete the above tables?
- b) Investigate what mathematical models could be adequate to describe the evolution of the bacteria population in the considered period of time.

Carlos and Luís began by asking themselves about the behaviour of the data. They calculated the differences between the values of the first table to identify a pattern: “Looking at the table station 1(...) we can realize that [the growth of bacteria population, p] is a linear function. The slope is constant  $\frac{140-90}{3-2} = \frac{240-140}{5-3} = \frac{390-140}{8-3}$ ” (Luís, E2). Based on these calculations, the students conjectured that these values were from a linear function. So, they posed another question: Assuming this linear behaviour for data, how do we find the missing values? For that, they have chosen to “use the rule of three” (Carlos, WR2). The students justified the conjecture by just referring to the slope.

Luís, working with his colleagues on station 1, believed that he could also use his recent knowledge about interpolation to find the missing values: “We think that one could also find the missing values by a polynomial interpolation method” (WR2). The student posed a question about whether Lagrange's method of (linear) polynomial interpolation would yield the same results. He compared the results of both methods: “We solved a problem with Lagrange's interpolation. We also used the rule of three to check whether there were differences but no, it gave the same” (E2). The analysis of his group work shows that the students also considered other possible explorations: “We can use any of the learned methods for polynomial interpolation. One can use either Newton's method with divided differences or the Lagrange's method” (WR2).

Gonçalo started exploring the task assuming that the data were in any position, and therefore, the missing values could be found by polynomial interpolation. He tried to find the missing values in station 1 by constructing a fourth-degree Newton's polynomial function based both on known properties of polynomials and on the idea (not always correct) that the greater the degree of the polynomial, the better the approximation to the data: “We used the four points because it's all we had. Moreover, the idea I had is that the more points we use more...” (E2). The choice of Newton's polynomial interpolation method with divided differences also drew on his recently acquired knowledge: “The nodes were not all at the same distance” (E2). The student explored the other tables in the same way and assumed that the constructed polynomials were suitable to represent data and to calculate the missing



values but did not make any test, possibly because he believed that he created a suitable option. Thus, he did not realize that some of the values obtained were hardly justifiable given the general behaviour of the data.

For the other two tables (stations 2 and 3), Carlos and Luís wondered again about the behaviour of data, but calculating the differences between each of two values, they verified that there is not a linear relationship:

For the tables station 2 and 3 we tried to apply the same as for table station 1. But:

$$\frac{85 - 40}{2 - 1} \neq \frac{220 - 40}{4 - 1} \neq \frac{210 - 85}{5 - 2} \text{ and } \frac{600 - 250}{7 - 5} \neq \frac{380 - 140}{6 - 3} \neq \frac{140 - 85}{3 - 2}.$$

So, they are not linear functions (...). (Luís, WR2)

Carlos and Luís chose to use the polynomial interpolation in order to find the missing values and, from this point on, reasoned very much like Gonçalo. However, before finishing the exploration of this task, Carlos proposed to his group a new problem assuming other conditions—to construct an algebraic expression for a second-degree polynomial. He took into account the data behaviour to pose this question and formulate a conjecture related to the degree of the polynomial. Then, he interpolated the function at 3 in station 2: “We obtained the algebraic expression of the polynomial:  $p_2(x)=22,5x+7,5x^2+10$ ” (WR2). He also considered “improving” their conjecture using different polynomials to describe the data trend, instead of the single expression previously found. His decisions on the degree of the polynomials to be used were based on identifying patterns in the data behaviour, the number and the monotony of the available values:

Looking at the data, I found 3 values growing up, a decreasing trend at 5 and then at 8 the values grow up again. Obviously the values could still be falling at 6, but we do not know... In this case, I thought I would get a function defined by three branches at the intervals, [1, 4[, [4, 5[, [5, 8]. (Carlos,E2)

When Carlos felt the need to justify their results, he tried to check whether such results agree with the expected ones and with the reasoning developed. He used a sophisticated graphical representation and properties of functions (monotony and derivative):

It was a kind of checking, if these polynomials were good for that range of values. (...) I even did a sketch here... It was through the observation of the table (...). It was growing from here to there (...). Therefore the polynomials had to be increasing or decreasing and their derivatives had to be positive or negative (...). (Carlos, E2)

An interesting extension of this question was raised by Luís when he questioned if, after obtaining an interpolated value, he could add these new data to the table and thus get a higher-degree polynomial to find the remaining missing values: “To discover the value of  $t=6$  [in station 2] we could now include the value of  $p(3)$  that we have just found and so the error was smaller” (E2). Here, we see an attempt to pose new questions.

Within this task, the students formulated conjectures about the most appropriate computation methods based on different assumptions about the data. They did not feel the need to test their results. However, raising questions on the global behaviour of data may lead to using different methods with quite different results. Students' problem posing is visible at several stages of their exploration. Initially, Carlos and Luís questioned themselves about the behaviour of data, and based on the identified pattern, they formulated conjectures. They also posed questions when selecting the appropriate methods to find the missing values,

taking into account the data, and when they sought to compare different methods. Before they finished, students were still posed new problems by considering the results or by modifying the initial conditions to refine the previous formulations. In contrast, Gonalo did not pose explicit questions about the data behaviour since he assumed that polynomial interpolation was the appropriate approach. However, even if only implicitly, he also posed questions about the appropriate polynomial to represent the data (degree and method).

## 5 Conclusion

During the study, the students were challenged to carry out investigation activities very different from the usual university mathematical tasks. This led them to experience mathematical processes such as looking for regularities, posing questions, formulating and testing conjectures, generalizing and providing justifications. These results extend to the classroom setting previous research by Cifarelli and Cai (2005). Furthermore, they show that, besides formulating conjectures and generalizing, students also engage in other mathematical processes.

Looking for regularities was based, mainly, on the observation of examples in visual and numerical diagrams or through counting or computation. As in former research by Silver (1994), the students' sketches were very important to help them in formulating conjectures. Difficulties in this process arose when they did not take into account all the available (and necessary) information.

All students in our study posed questions as they formulated initial questions about the data provided and sought to identify patterns that yielded conjectures. Sometimes, they also posed questions to simplify the original problem. However, in many cases, the posed questions were just implicit being identifiable from the conjectures.

The students mostly formulated conjectures based on identifying patterns from data observation or manipulation. However, they also drew on mathematical properties. Their conception of solving a task as a process of getting a result led them to formulate conjectures as statements, confirming the findings of Ponte and Matos (1992), and to accept them as conclusions, feeling no need for testing or proving. This result, also observed by Ponte et al. (1998), seemed to be more related to students' inexperience in conducting this type of task and the challenge of understanding the nature of the investigation process than to mathematical difficulties in testing their hypotheses/conjectures.

The work on the proposed tasks led the students to understand the importance of testing their conjectures, and from a certain point, this became a constant concern. Most of the time, testing conjectures was done through experimentation (often with the examples available in the statement). However, sometimes the testing was based on graphic representations and on mathematical concepts and properties. In these cases, testing conjectures coincided with the process of justifying them. When the students explained their reasoning, they unintentionally gave justifications based on mathematical properties or deductive reasoning. When, through testing, the students refuted a conjecture, they tried to reformulate it. In that way, the process of testing conjectures allowed not only its verification but also problem posing, reinforcing the perspective of Cifarelli and Cai (2005) of mathematical activity as a recursive process wherein students' reflection on the results can provide them with opportunities to formulate new problems to explore and solve.

An important step in the exploration of some tasks was the generalization of the initial conjectures, i.e., the formulation of a more general conjecture, which led to posing new questions. Carlos and Lu s seemed to be unaware that their conjectures might be valid for

particular cases but not for the general case and tended to consider the generalization a simple process of formalization. Rather, Gonçalo returned to questions previously raised and broadened the scope of his conjectures to obtain a general solution. The process of generalization carried out by this student resulted also in the construction of the bisection method. In a different way, Carlos and Luís also suggested possible extensions of the proposed questions. After obtaining a first solution, they posed new questions and formulated new conjectures or refined existing ones by changing the initial conditions (as suggested by Ponte et al., 1998, and Silver, 1994).

The results of this study highlight the potential of investigation tasks to promote problem posing and suggest that such tasks might be successful in university mathematics courses. Students do engage in problem posing when carrying out such tasks. This process occurs when they pose questions concerning the given data, look for patterns and formulate conjectures from a series of examples, generate more accessible subproblems from a larger problem, test their conjectures and reformulate them, try to generalize and/or refine their conjectures to obtain a general solution and propose alternative questions, expanding their explorations by changing the initial conditions. Thus, students' problem posing processes appear to permeate the problem solving activity, as suggested by Brown and Walter (1990) and Cifarelli and Cai (2005). As a key feature of authentic mathematical activity, our study shows that problem posing processes seem to be stimulated when students get involved in investigation activities.

## References

- Bogdan, R., & Biklen, S. (1994). *Investigação qualitativa em educação* [Qualitative investigation in education]. Porto: Porto Editora.
- Brown, S., & Walter, M. (1990). *The art of problem posing* (2nd ed.). Hillsdale: Lawrence Erlbaum.
- Cai, J., & Cifarelli, V. V. (2005). Exploring mathematical exploration: How two college students formulated and solved their own mathematical problems? *Focus on Learning Problems in Mathematics*, 27(3), 43–72.
- Cifarelli, V. V., & Cai, J. (2005). The evolution of mathematical explorations in open-ended problem-solving situations. *The Journal of Mathematical Behavior*, 24, 302–324.
- English, L. (1997). The development of fifth-grade children's problem-posing abilities. *Educational Studies in Mathematics*, 34(3), 183–217.
- English, L. D. (2003). Problem posing in elementary curriculum. In F. Lester & R. Charles (Eds.), *Teaching mathematics through problem solving*. Reston: National Council of Teachers of Mathematics.
- Ernest, P. (1991). *The philosophy of mathematics education*. New York: Falmer.
- Freudenthal, H. (1973). *Mathematics as an educational task*. Dordrecht: Reidel.
- Hadamard, J. (1945). *Psychology of invention in the mathematical field*. Princeton: Princeton University Press.
- Henriques, A. (2010). *O pensamento matemático avançado e a aprendizagem da análise numérica num contexto de actividades de investigação* [Advanced mathematical thinking and the learning of numerical analysis in the context of investigation activities] (Doctoral dissertation, University of Lisbon, available at <http://repositorio.ul.pt/>). Accessed 30 April 2011.
- Henriques, A., & Ponte, J. (2008). Actividades de investigação na aprendizagem de Análise Numérica [Investigation activities and the learning of numerical analysis]. *Revista de Educação*, 16(2), 5–32.
- Kilpatrick, J. (1987). Problem formulating: Were do good problems come from? In A. H. Schoenfeld (Ed.), *Cognitive science and mathematics education* (pp. 123–147). Hillsdale: Lawrence Erlbaum.
- National Council of Teachers of Mathematics. (1989). *Curriculum and evaluation standards for school mathematics*. Reston: NCTM.
- National Council of Teachers of Mathematics. (2000). *Principles and standards for school mathematics*. Reston: NCTM.
- Pólya, G. (1945). *How to solve it*. Princeton: Princeton University Press.
- Pólya, G. (1981). *Mathematical discovery: On understanding, learning and teaching problem solving*. New York: Wiley.

- Pólya, G. (2002). The goals of mathematical education. *Mathematics Teaching*, 181(6–7), 42–43.
- Ponte, J. (2007). Investigations and explorations in the mathematics classroom. *ZDM*, 39, 419–430.
- Ponte, J., Brocardo, J., & Oliveira, H. (2003). *Investigações matemáticas na sala de aula* [Mathematical investigations in the classroom]. Belo Horizonte: Autêntica.
- Ponte, J., Ferreira, C., Brunheira, L., Oliveira, H., & Varandas, J. (1998). Investigating mathematical investigations. *Proceedings of CIEAEM-49* (pp. 3–14), ESE Setúbal, Portugal.
- Ponte, J., & Matos, J. F. (1992). Cognitive processes and social interactions in investigation activities. In J. Ponte et al. (Eds.), *Mathematical problem solving and new information technologies: Research in contexts of practice* (pp. 239–254). Berlin: Springer.
- Ruthven, K., Hofmann, R., & Mercer, N. (2011). A dialogic approach to plenary problem synthesis. In B. Ubuz (Ed.), *Proceedings of the 35th Conference of the International Group for the Psychology of Mathematics Education* (Vol. 4, pp. 81–88). Ankara: PME.
- Silver, E. (1994). On mathematical problem posing. *For the Learning of Mathematics*, 14(1), 19–28.