



# Amazon deforestation: simulated impact of Brazil's proposed BR-319 highway project

Jerfferson L. Santos · Aurora M. Yanai ·  
Paulo M. L. A. Graça · Francis W. S. Correia ·  
Philip M. Fearnside

Received: 23 June 2023 / Accepted: 1 September 2023 / Published online: 16 September 2023  
© The Author(s), under exclusive licence to Springer Nature Switzerland AG 2023

**Abstract** The scenario of deforestation in the Amazon may change with the reconstruction of Highway BR-319, a long-distance road that will expand the region's agricultural frontier towards the north and west of the Western Amazon, stretches that until then have extensive areas of primary forest due to the hard access. We simulate the deforestation that would be caused by the reconstruction and paving of Highway BR-319 in Brazil's state of Amazonas for the period from 2021 to 2100. The scenarios were based on the historical dynamics of deforestation in the state of Amazonas (business as usual, or BAU). Two deforestation scenarios were developed: (a) BAU\_1, where

Highway BR-319 is not reconstructed, maintaining its current status, and (b) BAU\_2, where the reconstruction and paving of the highway will take place in 2025, favoring the advance of the deforestation frontier to the northern and western portion of the state of Amazonas. In the scenario where the highway reconstruction is foreseen (BAU\_2), the results show that deforestation increased by 60% by 2100 compared to the scenario without reconstruction (BAU\_1), demonstrating that paving would increase deforestation beyond the limits of the highway's official buffer area (40 km). The study showed that protected areas (conservation units and indigenous lands) help to maintain forest cover in the Amazon region. At the same time, it shows how studies like this one can help in decision-making.

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1007/s10661-023-11820-7>.

J. L. Santos  
Postgraduate program in Climate and the Environment,  
National Institute of Amazonian Research (INPA), Av.  
André Araújo, 2936, CEP, Manaus, Amazonas 69067-375,  
Brazil

J. L. Santos (✉) · F. W. S. Correia  
Postgraduate program in Climate and the Environment,  
State University of Amazonas (UEA), Av. Darcy Vargas,  
1200, CEP, Manaus, Amazonas 69050-020, Brazil  
e-mail: jlds.dcl19@uea.edu.br

A. M. Yanai · P. M. L. A. Graça · P. M. Fearnside  
Department of Environmental Dynamics, National  
Institute of Amazonian Research (INPA), Av. André  
Araújo, 2936, CEP, Manaus, Amazonas 69067-375, Brazil  
e-mail: yanai@inpa.gov.br

P. M. L. A. Graça  
e-mail: pmlag@inpa.gov.br

P. M. Fearnside  
e-mail: pmfearn@inpa.gov.br

F. W. S. Correia  
Laboratory of Terrestrial Climate System Modeling  
(LABCLIM), State University of Amazonas (UEA), Av.  
Darcy Vargas, 1200, CEP, Manaus, Amazonas 69050-020,  
Brazil

**Keywords** Environmental modeling · Land use change · Amazonia · Arc of deforestation · Human occupation

## Introduction

The Amazon basin covers an area of approximately 7 million km<sup>2</sup>, with 5.5 million km<sup>2</sup> covered by forests, which represents 40% of the global tropical forest area (Nobre, 2014; Weng et al., 2018). Amazon ecosystems host 15–20% of the planet’s species diversity (Lewinsohn & Prado, 2002) and store around 120 Gt of carbon (Saatchi et al., 2011). The Amazon rainforest plays an important role in the regional and global climate system through the storage and absorption of carbon (carbon cycle), transport of trace gases and aerosols, and water cycling, which provides moisture that is transported to other regions of the continent and contributes to maintaining the hydrological regime at regional and global scales (Rocha et al., 2015; Nobre et al., 2016; Marengo et al., 2018; Weng et al., 2018).


Deforestation, which is mostly for extensive cattle ranching, is a major contributor to greenhouse gas emissions and to climate change at both the regional and global scales (Fearnside et al., 2009; Moutinho, 2009; Marengo et al., 2018; Fearnside, 2022b). Deforestation in the Amazon has been monitored by satellite since 1988 and this monitoring is an important

tool for guiding public policies aimed at controlling the destruction of forests in the region (INPE, 2020).

Deforestation in the Amazon is one of the major problems that Brazil has been facing in recent decades, and the reconstruction of highway BR-319 (Fig. 1a) is a major issue that has drawn the attention of environmentalists and researchers. This highway would facilitate access to a large area of preserved forest, which could change the current scenario of deforestation in the Amazon (Fig. 1b) and cause substantial environmental and social impacts at the local, regional, and global levels.

Highway BR-319 was built in 1972 and 1973 but was only inaugurated in 1976 (DNIT, 2016), a period of military government. The highway was part of Brazil’s National Integration Program (PIN), under the motto “Security and Development,” uniting military concerns over perceived communist invasion with the developmental ideals promoted by President Juscelino Kubitschek in the 1950s (Lessa, 1991; Kohlhepp, 2002; Oliveira-Neto, 2014; Facundes, 2019). With the passage of time and lack of maintenance, the BR-319 became impassable in the late 1980s (DNIT, 2016) and its reconstruction became the focus of various local movements and governments (MPOG, 2004).

It was in the 1970s that the most critical period of changes in the Amazon landscape started in Brazil, when environmental impacts were intensified through colonization and development programs based on highways. These highways still have an



**Fig. 1** (a) Map of Highway BR-319, connecting the cities of Manaus, Amazonas, and Porto Velho, Rondônia, showing the main federal highways. (b) Official highways and the spatial


distribution of cumulative deforestation (1988 to 2021) with emphasis on the “arc of deforestation.” Map prepared by the authors. Data sources: IBGE, 2017; DNIT, 2021; INPE, 2020

important role in the occupation of space, attracting people in search of cheap land and natural resources and, consequently, increasing deforestation, fires, illegal logging, growth of cattle ranching, illegal mining, speculation and land grabbing, armed conflicts, and disease outbreaks, among other effects (Lessa, 1991; Loureiro, 2002; Fearnside, 2003; Graça et al., 2007; Laurance & Balmford, 2013; Brito & Castro, 2018).

Barber et al. (2014) showed that 94% of all deforestation in the Brazilian Amazon occurred around official and endogenous roads, demonstrating the role of highways as important drivers of deforestation. Reconstruction of Highway BR-319 is therefore the subject of growing concerns, as disorderly occupation and environmental degradation can extend the “arc of deforestation” (Fig. 1b) advancing to the northern part of the state of Amazonas and to the state of Roraima, reaching the border with Venezuela via Highway BR-174 (Manaus - Boa Vista) (Fearnside et al., 2009; Fearnside & Graça, 2009; Barni et al., 2015). Planned roads associated with BR-319 would extend the impact to the western portion of the state of Amazonas (Fearnside, 2018).

Even so, many politicians and enthusiasts for the reconstruction of Highway BR-319 have claimed that deforestation would not occur, contrary to the warnings of scientists. However, it is a fact that the simple announcement of the paving and improvement plans has already resulted in a disorderly pattern of occupation and an increase in deforestation and fires along the middle stretch of the highway, with rampant illegal logging and invasion of public lands for real estate speculation and extensive cattle ranching (Fearnside & Graça, 2009; Andrade et al., 2021; Ferrante et al., 2021a, b).

The situation is made more worrisome by the current Brazilian scenario in which there is a tendency for deforestation to increase, as can be seen in Fig. 2a. This trend is related to the economic pressures and political power of groups with interests in land-related businesses and infrastructure projects in the Amazon, which has led to the weakening of the Brazilian Forest Code (Supplementary Material, Appendix 1) and to other legislative changes that have been progressively eliminating restrictions on deforestation since 2012 (Fearnside, 2022a). The 2019–2022 Jair Bolsonaro presidential administration revoked many of



**Fig. 2** Deforestation in the Brazilian Legal Amazon (a) and in the State of Amazonas (b) from 1988 to 2021 in 103 km<sup>2</sup>. Source: INPE (Instituto Nacional de Pesquisas Espaciais) (2022)

the government's internal norms that had been established to combat deforestation (Barbosa et al., 2021). At least 401 of these changes can be reversed in 2023 by the incoming Luiz Inácio Lula da Silva administration (TALANOIA, 2022). Legislative changes, however, will face a National Congress with a new composition, indicating that it will be even more hostile to environmental protection than the Congress during the Bolsonaro administration (ClimaInfo, 2022).

According to data from the Project for Monitoring Deforestation in the Legal Amazon by Satellite (PRODES), of the National Institute for Space Research (INPE), the state of Amazonas resumed the increase of annual deforestation, from 523 km<sup>2</sup> in 2012 to 2306 km<sup>2</sup> in 2021, an increase of 440% (INPE, 2022), surpassing the historic record of 1995 (Fig. 2b). Furthermore, these data show that much of the deforestation in the state of Amazonas was concentrated in the southern part of the study area, which is under the direct influence of BR-230 (Transamazon Highway) and BR-364 (Porto Velho–Rio Branco).

Thus, given the possibility of the reconstruction and paving of BR-319 and the possible changes in the pattern of land use and cover, the question that the present study proposes to answer is: “What would be the impact of paving Highway BR-319 on deforestation in the state of Amazonas in 2050 and 2100?” The present study aims to evaluate the impact of BR-319 and other highways planned in the study area.

## Material and methods

### Study area

The study focuses on the federal highway BR-319, located in the interfluvium between the Madeira and Purus Rivers, connecting the cities of Manaus (Amazonas) and Porto Velho (Rondônia). The BR-319 is the main land access route to the municipalities of Careiro, Manaquiri, Careiro da Várzea, and Autazes, as well as facilitating access to Humaitá, Lábrea, and Manicoré. It provides the only land access to the communities of Vila Realidade (a district of the municipality of Humaitá) and Igapó-Açu (a district of the municipality of Borba). However, all of these locations are accessible from the two ends of the highway without reconstructing the critical “middle stretch” that would give access from the arc of deforestation

to all areas connected to Manaus by road, including the state of Roraima.


The official road network in the state of Amazonas that connects to the 885 km of BR-319 corresponds to 1934 km, comprising the federal highways BR-230 (827 km from Lábrea to the border between the states of Amazonas and Pará), BR-174 (85 km, stretch BR-319 - Manicoré), and state highways AM-254 (94 km, BR-319 - Autazes) and AM-354 (43 km, BR-319 - Manaquiri). In addition, there are other planned projects by the government of the state of Amazonas to build highways connecting BR-319 to other municipalities such as Borba (AM-356), Novo Aripuanã (AM-360), Tapauá, Tefé and Juruá (AM-366), and Coari (AM-343). The last two roads (AM-366 and AM-343) would advance into the vast area of forest to the west of the Purus River, facilitating deforestation in one of the most preserved forest areas in Amazonia, known as the “Trans-Purus” region (Fearnside et al., 2020) (Fig. 3). Very little of the area that would be accessed by these connecting roads is protected by designation as a “conservation unit” (Fig. 3).

The official area of influence used in Brazil's environmental licensing processes for highways in the Amazon region is 40 km of buffer area, defined by Interministerial Ordinance 60 of 24 March 2015 (Brazil, 2015). However, considering that the environmental impact of a paved highway in the Amazon can go beyond the minimum limit defined in the interministerial decree, the present study considered the state of Amazonas as the total area for modeling the impacts of deforestation, having as a “backbone” Highway BR-319, as well as its connecting highways and roads, including both existing and planned roads. The study area also includes a buffer zone of 20 km around the borders of the state of Amazonas to represent the influence of adjacent areas, especially the highways present in the states of Acre, Rondônia, Roraima, and Pará (Fig. 3).

### Land use modeling

Modeling deforestation was done using the environmental modeling platform DINAMICA-EGO (Environment for Geoprocessing Objects) (Soares-Filho et al., 2002; Leite-Filho et al., 2020). DINAMICA-EGO can be applied to a variety of types of studies, such as urban expansion modeling, economic ecological zoning proposals, and the simulation of





**Fig. 3** Study area, Highway BR-319, and road network planned around BR-319, federal- and state-protected areas, indigenous lands, federal settlement projects, and military areas. Map prepared by the authors. Data sources: IBGE (2017), ICMBio, INCRA, FUNAI

deforestation behavior (Soares-Filho et al., 2004; Rodrigues et al., 2007; Ramos et al., 2018; Santos et al., 2021). In addition, the software is open access and has a user-friendly interface, which can be used by people unfamiliar with programming languages such as R and Python. More details on the software can be found in the Supplementary Material (Appendix 2, Fig. S1).

**Deforestation modeling steps**

The modeling process was carried out through the following steps: input data, calibration, validation, and simulation (projection) of future deforestation. For the input and calibration data, the period from 2007 to 2013 was used. For validation, the period from 2014 to 2021 was used, while the simulation scenarios were for the period from 2021 to 2100.

*Input data*

All input cartographic data were in raster format with a spatial resolution of 100 m. The mapping used the *Brazil Policonic* cartesian coordinate system, Datum SIRGAS 2000.

In addition to land cover maps, maps of static and dynamic variables were used. Static variables are those for which the value of the class of each cell (pixel) does not change over the course of a simulation. For this category, maps of protected areas were used - indigenous lands (FUNAI, n.d.), federal-protected areas for integral protection and federal-protected areas for sustainable use (ICMBio, 2019), state-protected areas for integral protection and state-protected areas for sustainable use (SEMA, 2020), and military areas (ANM, 2021). A map of settlement projects (INCRA, n.d.) and official hydrography or watercourses (INPE, 2020) were also used.

Dynamic variables are those whose values change over the course of a simulation. These included distance from official and endogenous roads and distance from deforested areas. The Supplementary Material (Appendix 3) presents a summary of the variables used in the configurations (Table S1) and the map of static variables (Fig. S2).


### Regionalization of the study area

The model applied in this study used the regionalization approach, which consists of establishing different parameters for each region and modeling the regional context that influences a given phenomenon (Leite-Filho et al., 2020). The software uses a set of functors (tools or small subroutines) to divide a map into parts (i.e., regions) to process the dataset of each region separately and then combine them. For this, a regionalized map of the study area was added as input to the model.

Thus, considering that the regionalization of the area makes it possible to individually parameterize each region, in the present study the area was divided into nine regions (Fig. 4) that took into account the presence of highways (current and planned), human clusters, land use profile (contribution of social actors in deforestation), and hydrography. A summary of the parameters used to divide the study area into regions is provided in the Supplementary Material (Appendix 4, Table S2).

### Calibration

Calibration is the step of fitting the model parameters so that the simulation results are as similar as possible to the real study case (Campos et al., 2022). Therefore, in this phase, there is a continuous search to adjust these parameters until the simulation result is as close as possible to the real one. In this study, the reference period used to calibrate the model was from 2007 to 2013, with the goal of performing a



**Fig. 4** Regionalized map of the study area

validation simulation round for the period from 2014 to 2021, comparing the simulated map of 2021 with the satellite data for observed deforestation from the PRODES map for 2021.

Among the data needed to be applied in the simulation model are the weights of evidence of the variables, this being a measure of influence that each variable has to cause a change, in this case, the expansion of deforestation (Leite-Filho et al., 2020). The weights of evidence applied in DINAMICA-EGO are based on a Bayesian method where the effect of a spatial variable is calculated independently of any combination to produce maps that describe the most favorable areas for a change to occur (Soares-Filho et al., 2002, 2004; Leite-Filho et al., 2020).

To calculate the weights of evidence, a model was used in DINAMICA-EGO, which received the initial (2007) and final (2013) landscape maps, in addition to the maps of static and dynamic variables, followed by calculating the ranges and assigning transition-probability values for each variable used in the simulation model. An adjustment was necessary to achieve the desired result, by defining the interval and distance of the weights of evidence at 100 m and 1500 m, respectively, for the variables roads, deforestation, and hydrography. Such values were reached after several rounds of adjustments and the validation test indicated that the best result was in this influence range. The table of parameters used in the present study and a figure summarizing the calculation of the weights-of-evidence coefficients can be found in the Supplementary Material (Appendix 5, Table S3, Fig. S3).

Considering that the only assumption for the weights-of-evidence method is that the input maps be spatially independent, the next step is to analyze the correlation between the variable maps (Leite-Filho et al., 2020). After the analysis of correlated pairs between variables using Cramer’s test and joint-uncertainty information, values above 0.5 were considered dependent variables (Bonham-Carter, 1994). No dependent variables were observed in the present study.

Another parameter used in the model is the transition rate, which is necessary to determine the number of cells that transition between classes at each annual time step, in this case for forest to deforestation. The transition rate was calculated using a sub-model in DINAMICA-EGO called “Determine Transition

Matrix,” which uses maps of the initial state (cumulative deforestation by 2007) and final state (cumulative deforestation by 2013). This tool generates two matrices: the annual transition matrix (Multiple Step) and a global transition matrix (Single Step). “Multiple Step” portrays the process of change between the classes that occurs each year, while “Single Step” portrays the change over the whole analysis period (Leite-Filho et al., 2020). The simulation used the annual transition matrix (Multiple Step), which reflects the average annual transition in the calibration period (2007 to 2013).

However, simply applying the deforestation rate provided in the annual transition matrix would result in a constant rate across all model interactions. Thus, considering that deforestation rates actually fluctuate over time (increasing and decreasing), whether as a result of financial crises, conflicts, climatic events, political decisions, and other factors, this study included an increasing and reducing factor for deforestation rates, which was applied for interval periods of 6 years (period equal to the reference period used to calibrate the model).

To represent the increase in deforestation, an index was added to the transition rate (Multiple Step) that considered the deforested area in the previous year plus the average percentage increase in all years in which deforestation increased in the period from 2000 to 2014 in the state of Amazonas. This represented the increase in deforestation in the study area by means of the following equation:

$$\text{Ind.t} = ((\text{AD2} - \text{AD1}) / \text{AD1} \cdot 100) + \text{Md}_i \tag{1}$$

Ind.t = Transition Index

AD1 = area deforested in year 1 (km<sup>2</sup>)

AD2 = area deforested in year 2 (km<sup>2</sup>)

Md<sub>i</sub> = average annual deforestation during the years in which there was an increase (period from 2000 to 2014)

To represent the reduction in deforestation, Eq. 2 follows the same principle as Eq. 1, using the average percentage decrease in all years in which there was a reduction in deforestation during the period from 2000 to 2014.

$$\text{Ind.t} = ((\text{AD2} - \text{AD1}) / \text{AD1} \cdot 100) - \text{Md}_d \tag{2}$$

Ind.t = Transition Index

AD1 = area deforested in year 1 (km<sup>2</sup>)

AD2 = area deforested in year 2 (km<sup>2</sup>)

$Md_d$  = average annual deforestation during the years in which there was a reduction (period from 2000 to 2014)

The increase and decrease factors ( $Md_i$  and  $Md_d$ ) were calculated based on the average increase and decrease in deforestation during the period from 2001 to 2014, to better represent the trends of increase and decrease over time, which were defined as follows: 0.26 for increase and 0.20 for reduction. The years in which there were increases and decreases in deforestation in the state of Amazonas are shown in the Supplementary Material (Appendix 5, Fig. S4), as well as an example of the fluctuation of deforestation rates over time (Fig. S6). The present method allowed the transition rates to fluctuate with each iteration of the model, which means that as there is a change in the landscape at each time step, the (annual) transition rate is updated at each iteration in relation to the available forest area in each region. A summary and the input data are shown in Appendix 5 and Table S4 of the Supplementary Material.

The spatial allocation functions for the new deforestation patches used in the model were Patcher and Expander, where the Patcher function creates new areas (patches) of transition separate from the already-deforested areas, while the Expander function is responsible for enlarging already-deforested areas (Leite-Filho et al., 2020). In this study, several rounds of parameter adjustments were carried out and, in the validation test, the best result was found to be achieved using 30% as a value for the Expander function and 70% for the Patcher function. As for the size of the deforestation patches, the average range of the size of the deforestation polygons of each region defined in the study was calculated during the calibration period. The settings used to allocate deforestation patches through the Patcher and Expander functions, including the percentages adopted, are available in the Supplementary Material (Appendix 6, Table S6).

Considering that the model deals with the impact of roads on landscape change, the road builder module was coupled to the model, using the map of official and endogenous roads as input. This module calculates the relative cost that a road has in crossing a cell in the land use map, depending on the destination given to the cell (protected lands, non-destined forest areas, settlements, etc.). For this, we used an

attractiveness map (which indicates the most favorable areas for road construction) and a friction map (which indicates the areas with greater restrictions for road construction) (Leite-Filho et al., 2020). The settings used in the road builder module can be seen in the Supplementary Material (Appendix 7, Table S7).

### Model validation

After calibration (2007 to 2013), a simulation model was used for the period from 2014 to 2021 in order to calculate the change that occurred in this interval and validate the resulting map of the simulated model for 2021 by comparison with the real map from PRODES of 2021. For validation, this study simulated a period that was different from the calibration period in order to assess how good the model is at predicting changes in the landscape, based on the procedures used in past studies (Siqueira-Gay et al., 2022).

The validation method applied in this study was the fuzzy similarity method (Hagen, 2003), adapted by Leite-Filho et al. (2020). This method employs a constant decay function that measures the spatial adequacy between two maps through multiple-window similarity analysis, that is, if the same number of change cells is found in the window, the fit will be 1, regardless of their locations, and zero if the same number of change cells is not found (Leite-Filho et al., 2020). Simply put, the model makes the comparison through window sizes, that is, with the number of cells corresponding to the resolution used in the modeling. For example, in this study, the resolution adopted was 100 m, so window 1 (1 × 1) corresponds to 100 m × 100 m (0.01 km<sup>2</sup>), window 3 (3 × 3) = 300 m × 300 m (0.09 km<sup>2</sup>), and so on.

Because the comparison is made using both maps (simulated and observed), the results can generate rates with minimum and maximum similarity values, which can vary from 0 to 100% (0% indicates that the maps are completely different and 100% indicates they are identical). In this study, we adopted the minimum similarity value as a reference. We compared the simulation results with a null model, which uses the same maps and input rates but with weights-of-evidence values set to zero. The null map was also compared with the observed map (PRODES of 2021). To be considered efficient, the proposed model must win in all comparisons made with the null model.



Further details can be found in the Supplementary Material (Appendix 8).

### *Projection of future scenarios*

The current approach considers the trends in the expansion of territorial occupation by different local groups based on the dynamics of historical deforestation for the Amazon (Business as Usual, or BAU), which reflects occupation dynamics and conflicts that influence landscape change along highways (Castro et al., 2004; Brito & Castro, 2018; Fearnside, 2022a). Thus, deforestation rates were not projected based on the perspective of improving the environmental management in the area, such as strengthening and increasing the autonomy of public command-and-control institutions, public policies aimed at sustainability, or achieving the goal of reducing emissions stipulated in international agreements, as this depends on the long-term commitment of state and federal governments.

Two environmental prognosis scenarios were developed for the period from 2021 to 2100: (a) Scenario 1 (BAU\_1)—Highway BR-319 without paving (the current status with seasonal maintenance and with degradation in the rainy season, with the pending reconstruction and paving project not approved); (b) Scenario 2 (BAU\_2)—Highway BR-319 with paving (the reconstruction and paving project is assumed to be authorized and started in 2025).

For the BAU\_1 scenario, the averages of the historical transition rates from the calibration period (2007 to 2013) obtained from each region of the study area were applied according to the methodology presented in the item “model validation,” from 2021 to 2100. For the BAU\_2 scenario, the transition rates followed the same principles as the BAU\_1 scenario until the beginning of the paving of Highway BR-319 in 2025, when an increase in the deforestation rate begins as a result of the migratory flow resulting from the road improvement and the expansion of the planned road network until 2100. Post-paving rates were obtained from other regions within the study area itself, as defined below.

For Scenario BAU\_2, which considers Highway BR-319 to be paved from 2025 onwards, the rates found in Regions 3 and 4 (where the sections of the BR-319 are located) take on present the same rates found in Region 1 (area with a higher deforestation

rate). Regions 3 and 4 would be new frontiers for expansion of ranching if BR-319 is paved, and in Region 5 (Manaus), which will have the rate of Region 3, a region close to the capital of Rondônia (so that Region 5 has a rate similar to that near a state capital in the “arc of deforestation”).

After 2028, the transition rate found in Region 7 (providing Highway AM-366 is built as a result of the BR-319 highway) started to have the same rate as in Region 1 (same principle adopted to represent Regions 3 and 4, if AM-366 is built). The Region 1 rate was chosen because it represents a continuation of the expansion of deforestation towards the western part of the study area due to the influence of migration to Amazonas from the states of Pará, Rondônia, and Mato Grosso. We therefore chose Region 1 as a reference to represent the amount of deforestation.

Regardless of the applied rate, the model allows the use of weights-of-evidence coefficients from other regions that can better simulate what is intended to be represented. Thus, the weights-of-evidence coefficients were also replaced to better represent the influence of paved roads in the model, so Regions 3 and 4 (site of the BR-319 highway), and Region 5 (region with road connecting to BR-319, and therefore becoming a new agricultural frontier), started to have the same weights-of-evidence coefficient as in Region 6 (which is a region with the paved Highway BR-364 in the “arc of deforestation”).

Considering the construction plan for Highway AM-366 (without paving), Region 7 now has the same weight of evidence as Region 1 (which is a region with the unpaved Highway BR-230 in the “arc of deforestation” in the state of Amazonas). In addition, to complement the analysis of the impact of deforestation, a paving plan was made for Highway AM-366 for the year 2050, after which it started to change the weights-of-evidence coefficients to be more similar to those of Region 6 (i.e., to resemble region with a paved highway: part of the Porto Velho–Rio Branco stretch of BR-364).

The paving plan for Highway AM-366 is justified by the fact that the proposed road is located in a region planned for oil and gas extraction, which may favor financing or raising funds for construction, in addition to a greater possibility of political interference with the licensing body. However, it is worth noting that, considering the applied transition rates,



the result of the amount of deforestation does not change.

Patcher and Expander allocation followed the same principles as for the parameters used in road construction.

The plan for the construction and paving of the planned highways followed the principles of area availability and occupation opportunity because, regardless of government plans for building a highway, when there is an available area and opportunity, the illegal occupants of the area begin to follow the planned route of a highway, opening unofficial roads and branches on the proposed official highway. This fact can be observed in an area in Region 4, where an illegal road or “branch” is already being built on the route of the proposed Highway AM-366 (Fearnside, 2022b). Thus, for the present study, a 3-year construction schedule (whether official or not) was adopted to start after the paving of BR-319 (Table 1).

The application of transition rates in both scenarios followed the same methodology applied for the validation phase. However, the values for the “average of

years in which there was an increase and decrease in deforestation” ( $Md_i$  and  $Md_d$ ) were adjusted in both scenarios to better represent the trends, using the average increase and decrease over the period from 2000 to 2021. The value of 0.32 was adopted as an increase factor and 0.19 as a decrease factor, with intervals of 6 years starting in 2021 (Table S5, Supplementary Material).

## Results

### Validation

The validation compared the 2021 simulated deforestation map with the 2021 deforestation obtained by the PRODES mapping in 2021, which is considered as a reference for observed deforestation. This method considers the values of the similarity index of 50% sufficient for model validation (Soares-Filho et al., 2013). The value of the minimum similarity index obtained was 51% for the simulation model in a window of  $11 \times 11$  cells.

In addition to the validation for 2021, the results were compared to a null model. In the null model, the same input maps and transition rates were used, but with the weights-of-evidence coefficients set to zero, producing the result shown in Fig. 5.


Regarding the comparison of the simulated deforestation, the validation showed a difference of  $-0.54\%$  in relation to the reference deforestation for the year 2021, resulting in a difference of  $-313.92 \text{ km}^2$  (Supplementary Material, Appendix 8, Table S8). The results for each region are shown in the Supplementary Material (Fig. S8, Appendix 8).

**Table 1** The schedule for the construction and paving of the planned highways influenced by the implementation of BR-319

Road	Segment	Start
BR-319	Manaus - Porto Velho	2025*
AM-366 (Segment 1)	Tapauá - AM-343	2028
AM-343	Coari - AM-366	2028
AM-366 (Segment 2)	Entroncamento AM-366 - Tefé	2031
AM-366 (Segment 3)	Tefé - Jutaiá	2034
AM-356	BR-319 - Borba	2028
AM-360	BR-319 - Novo Aripuanã	2028
AM-366 (all segments) and AM-343	Tapauá - Coari - Jutaiá	2050*

\*Pavement estimate

**Fig. 5** Validation results for 2021 with minimum similarity and with the null model, using the constant decay method



## Deforestation prediction for the years 2050 and 2100

In this section, the results of the scenarios will be presented, highlighting the simulated changes by 2050 and by 2100. The results show that, for deforestation in BAU\_1, there is an increase of 200.24% up to 2050 and 607.42% up to 2100, in relation to that observed in the PRODES of 2021 map. For BAU\_2, there is an increase of 224.12% by 2050 and 711.33% up to 2100, for the entire modeled area, as shown in Fig. 6.

For the BAU\_1 scenario in the Madeira-Purus interfluvium (Regions 3 and 4) where the BR-319 Highway is located, there were increases of 197.37% up to 2050 and 600.95% up to 2100 in Region 3 and increases of 241.08% up to 2050 and 762.04% up to 2100 in Region 4. Especially for the northern stretch of Highway BR-319 (Region 4, which has more area available for deforestation) after paving (BAU\_2), there were increases of 260.08% up to 2050 and 843.65% up to 2100.

Another part of Amazonas that draws attention is the Trans-Purus region in the center of the state (Region 7). This is due to the possible construction of Highway AM-366, which would connect to BR-319 (BAU\_2). The BAU\_2 scenario shows an increase of 359.48% by 2050 and 1458.91% by 2100 (Fig. 7, panels D and E).

Region 5 (BR-174 from Manaus to the border with the state of Roraima) would have an increase of 225.36% by 2050 and 734.81% by 2100 due to the influence of the reconstruction of BR-319 (BAU\_2). Thus, for the regions influenced by Highway BR-319 (Regions 3, 4, 5, and 7), deforestation would have an increase of approximately 60% in BAU\_2 (159,961.31 km<sup>2</sup>) in relation to BAU\_1 (99,959.97 km<sup>2</sup>). The results for all regions are shown in Table 2.


Roads played an important role in the distribution and dispersion of deforestation over time in the proposed model. Figure 8 cuts out the study area to show how deforestation evolves around the simulated roads for the years 2050, 2060, 2070, 2080, 2090, and 2100. According to the model, a cluster of deforestation ends up attracting other deforestation, which can occur on the banks of rivers without the presence of roads. However, a large part of the deforestation is conducted along unofficial roads that branch off from the official roads (in Brazil, the pattern of these side roads is called the “fishbone”). This pattern develops along roads connecting to riverside towns and

cities, as can be seen in the evolution of deforestation shown in Fig. 8, corroborating the studies by Castro et al. (2004), Nepstad et al. (2006), Barber et al. (2014), dos Santos-Júnior et al. (2018), and Fearnside (2022a,b).

We can see that deforestation has increased in all protection categories (except for military areas, which have very low deforestation). When comparing the deforestation of protected areas in relation to the total forest loss (inside and outside protected areas) after 2021, an increase of deforestation in conservation units by 2153.60 km<sup>2</sup> up to 2050 can be observed in the BAU\_1 scenario, and 28,656.73 km<sup>2</sup> up to 2100, corresponding to 3.72% and 9.78%, respectively, in relation to total deforestation. In the BAU\_2 scenario, deforestation in the protected areas was 1960.65 km<sup>2</sup> in 2050 and 34,612.13 km<sup>2</sup> in 2100, corresponding to 2.73% and 9.80%, respectively, of the total deforested area.

In indigenous lands, projected deforestation after 2021 was 1042.81 km<sup>2</sup> in 2050 and 19,911.23 km<sup>2</sup> in 2100 for the BAU\_1 scenario, corresponding to 1.80% and 6.79%, respectively, in relation to total deforestation. For the BAU\_2 scenario, the total area of deforestation in indigenous lands was 964.44 km<sup>2</sup> in 2050 and 21,079.15 km<sup>2</sup> by 2100, respectively, from which 1.34% and 5.97% of the total deforested area were after 2021. Regarding the total area of protected areas, deforestation reaches 0.52% by 2050 and 7.91% by 2100, in the BAU\_1 scenario, and 0.48% by 2050 and 9.08% of the total area of conservation units and indigenous lands up to 2100 in the BAU\_2 scenario. Figure 9 presents the relationship between deforestation in protected and non-protected areas, showing the importance of protected areas for the conservation of forests in the Amazon.






For settlement projects, according to the results of the projection for the BAU\_1 scenario, the deforestation that occurred after 2021 was 16,897.26 km<sup>2</sup> by 2050 and 48,407.66 km<sup>2</sup> by 2100, corresponding to 41.22% and 19.79% in relation to the deforestation outside protected areas. For the BAU\_2 scenario, deforestation after 2021 was 21,660.76 km<sup>2</sup> by 2050 and 57,334.82 km<sup>2</sup>, which corresponds to 43.31% and 19.39% in relation to total deforestation (excluding protected areas), respectively (Fig. 10). Regarding the total area of settlements, deforestation reaches 22.76% up to 2050 of the total area of settlements and 65.19% up to 2100 in the BAU\_1 scenario, and it reaches 29.17% up to 2050, and 77.21% up to 2100 in the BAU\_2 scenario.



**Fig. 6** Evolution of cumulative deforestation for the period from 2021 (A) to 2050 and 2100 in the BAU\_1 (B and C) and BAU\_2 (D and E) scenarios. In this study, “non-forest” refers

to those areas not considered by PRODES/INPE in the calculation of deforestation in the Amazon (savannas, water, rocky outcrops, etc. - <http://terrabrasilis.dpi.inpe.br/>)





**Fig. 7** Evolution of cumulative deforestation for the period from 2021 (A) to 2050 and 2100, in scenarios BAU\_1 (B and C) and BAU\_2 (D and E) in Region 7 (Trans-Purus) as a result of the construction of Highways AM-366 and AM-248

**Table 2** Increase in cumulative deforestation by region and percentage of increase in cumulative deforestation over the simulated period in relation to 2021

Region	PRODES 2021	BAU_1 2050 (km <sup>2</sup> )	%	BAU_2 2050 (km <sup>2</sup> )	%	BAU_1 2100 (km <sup>2</sup> )	%	BAU_2 2100 (km <sup>2</sup> )	%
1	9042.42	27,569.06	304.89	27,569.06	304.89	92,897.55	1027.35	92,897.55	1027.35
2	5369.36	7272.21	135.44	7272.21	135.44	17,114.99	318.75	17,114.99	318.75
3	4469.53	9918.68	221.92	11,624.33	260.08	31,599.30	706.99	37,707.12	843.65
4	4713.67	8205.97	174.09	10,514.33	223.06	23,586.79	500.39	32,272.10	684.65
5	7634.83	12,083.39	158.27	17,205.73	225.36	33,927.84	444.38	56,101.63	734.81
6	19,040.05	38,864.17	204.12	38,864.17	204.12	117,380.29	616.49	117,380.29	616.49
7	2322.31	3694.81	159.10	8348.22	359.48	10,846.04	467.04	33,880.46	1458.91
8	3327.30	5046.21	151.66	5046.21	151.66	14,387.02	432.39	14,387.02	432.39
9	1825.68	2973.98	162.90	2973.98	162.90	9015.05	493.79	9015.05	493.79
Total	57,745.15	115,628.48	200.24	129,418.24	224.12	350,754.87	607.42	410,756.21	711.33

## Discussion

### Simulated deforestation

Although the method considers similarity index values above 50% to be enough to validate the model, which means that the amount of change correctly predicted is greater than the sum of the various types of error (Pontius-Jr et al., 2007; Soares-Filho et al., 2013), there is no general rule for calibration and validation in the land use modeling process (Rykiel, 1996; Mazzotti & Vinci, 2007). However, it is understood that the model must satisfactorily represent the spatial dynamics of deforestation in the study area.

In the current study, the model reached 51% in the 11 × 11 window, which corresponds to the similarity in an area of 1.21 km<sup>2</sup>. Some studies carried out in smaller areas in Amazonia also found similarity starting at 50% in the 11 × 11 window or smaller, such as Yanai et al. (2017) in the 5 × 5 window, Maeda et al. (2011) in the 11 × 11 window, Barni et al. (2015) in the 7 × 7 window, Roriz et al. (2017) in the 5 × 5 window, and Ramos et al. (2018) in the 11 × 11 window; dos Santos-Júnior et al. (2018) reached 49% in the 11 × 11 window, and Santos et al. (2021) reached 57% in the 7 × 7 window.

In addition, the accuracy was checked by comparison with a null model that, for the same window, reached 14% similarity. According to Pontius-Jr et al. (2004), a model becomes more accurate than the null model when the spatial resolution is increased, that is, the quality of the resolution scale influences the


result of a predictive model when compared to the null model. Considering the extent of the study area and the spatial resolution used, the validation results achieved in this study can be considered satisfactory.

In the model scenarios (BAU\_1 and BAU\_2), we sought to represent the current trend to increase deforestation rates in the Amazon. After the large reduction in annual deforestation from 2004 to 2012, a gradual and consistent increase in rates was observed beginning in 2012, when the Brazilian Forest Code was altered due to the strong political representation of agribusiness in the National Congress (Fearnside, 2022a). Many environmental regulations were also being revoked, especially during the 2019–2022 presidential administration of Jair Bolsonaro.

The results show that in both scenarios (BAU\_1 and BAU\_2), there is an evident increase in deforestation in the southern part of the Amazon, influenced by roads, settlements, and the “arc of deforestation.” Following this trend, the results show increases in deforestation in all of the modeled areas along Highway BR-319, as well as along connecting highways such as AM-366, especially for the BAU\_2 scenario due to the approval of the reconstruction and paving of Highway BR-319. This corroborates the predictions of Fearnside et al. (2009) and dos Santos-Júnior et al. (2018), in addition to models that considered projected road building in the Amazon region (Laurance et al., 2001; Soares Filho et al., 2004, 2006; Aguiar, 2006, 2016).

Deforestation of protected areas and Indigenous Lands can also increase considerably, according to






**Fig. 8** Evolution of deforestation around the simulated roads over time in the BAU\_2 scenario. The figure shows part of the region of influence of AM-366 (Trans-Purus)

various studies carried out in the region (Ferrante & Fearnside, 2019; Ferrante et al., 2021a,b). However, these areas continue to confer a certain resistance to environmental degradation by deforestation, as demonstrated by the current deforestation data available in the PRODES images from the National Institute for Space Research (INPE), as well as in the reports


of the programs of Ministry of Environment (MMA) to combat and control deforestation from the MMA (2016, 2018). Therefore, it is important to create, implement, maintain, monitor, and inspect protected areas in the Amazon.

Regarding settlement projects, the study shows that there is a significant increase in all categories,

**Fig. 9** Deforestation in protected areas (conservation units and indigenous lands) and non-protected areas (settlement projects are not considered to be protected areas)



**Fig. 10** Deforestation in settlement projects and non-designated public land



indicating that creating “sustainable-use settlements” in the region does not provide the desired protection (Yanai et al., 2017). Settlements currently represent 15.66% of the deforestation in the study area, but for deforestation up to 2100 this percentage rises to 65.19% in the BAU-1 scenario and 77.22% in the BAU\_2 scenario. This corroborates the studies by Yanai et al. (2017), who indicated that settlements play an important role in the dynamics of deforestation and future carbon emissions in the Brazilian Legal Amazon region.

Simply giving the news of a settlement approval starts a race in search of legalized lands made available by the government, according to the dynamics explained by Castro et al. (2004). This is exemplified by the Realidade Sustainable Development Project (PDS) that was created in 2009 around the BR-319 in the municipality of Humaitá (INCRA, 2015). The mere announcement of the approval of this PDS set off a race in search of land, promoting the invasion of the land and dividing it into small lots for sale to new arrivals, with no interference from the responsible government agency (The National Institute for Colonization and Agrarian Reform, or INCRA). Thus, making logging, agriculture, extensive livestock, speculation, and land grabbing grow in the settlement’s surroundings and along the highway,

as observed by Fearnside (2018), Andrade et al. (2021), and Ferrante et al. (2020, 2021a, b) in studies carried out in the region, demonstrating that the pattern of deforestation dynamics continues until the present day.

Another important issue is the proposed construction of State Highway AM-366, which would connect the BR-319 highway to the western part of the state of Amazonas (in this study represented by Region 7, see Fig. 5), one of the most preserved areas in Amazonia, and essential for the environmental services that the forest offers (Fearnside, 2020; Fearnside et al., 2020). An important source of impact would also be the advance of the “arc of deforestation” towards the north (Region 5) along the Federal Highway BR-174, which connects Manaus to Boa Vista and the border with Venezuela (Fearnside & Graça, 2009; Barni et al., 2015).

Although the roads are considered strategic and important because they reduce the isolation of the population and facilitate access, tourism, and the flow of products, the development model based on the expansion of road axes in the Amazon region is the main promoter of environmental degradation through its role in facilitating both the migration of population to the region and the expulsion of population to more distant frontiers as smallholdings are bought




up by large cattle ranchers. The forest is lost in this process, with major environmental impacts. We can say that Brazil has still not managed to find an action strategy that is efficient to reconcile the interests of the population that wants more highways, with the preservation of the environment. The BR-163 (Santarém-Cuiabá) Highway serves as an example: deforestation increased tremendously after the highway was reconstructed and paved, despite all attempts to develop policies, plans, and programs to reduce this environmental damage (Castro et al., 2004; Araújo et al., 2008; Brito & Castro, 2018).

As observed in the maps generated by the model, the impact of deforestation goes beyond the official 40-km influence area defined by Interministerial Ordinance 60, of 24 March 2015 for the environmental licensing processes of highways in the Amazon region. This demonstrates that the environmental licensing process would benefit from modeling the impact before defining the radius of influence in decision-making. Figure 11 shows the deforestation around Highway BR-319 and the buffer area of 40 km (for the stretch where the Installation License for the reconstruction of the highway is being requested), and we can observe the continuous deforestation beyond the limits of the 40-km buffer.

Thus, a more comprehensive modeling study similar to the current one could be used to define the probable area of a road project’s impact in the Amazon. This gives the environmental impact study more tools for decision-making, which makes it possible to

define the best mitigation measures to reduce negative impacts and to have a more realistic assessment of impacts for decisions on whether these highways should be built. While decisions on road building should consider all possible impacts, it is understood that environmental licensing is limited in its ability to require that the entrepreneur repair or mitigate the possible indirect impacts of an enterprise, such as the construction of connecting highways by the local authorities or negative influence on other states.

It is therefore urgent for Brazil to adopt tools such as the strategic environmental assessment (*Avaliação Ambiental Estratégica* = AAE), which is a planning and support instrument for strategic decision-making on the socio-environmental impacts of the Brazilian government’s Policies, Plans, and Programs (PPP) initiative (Partidário, 2011, 2003; Pellin et al., 2011), such as *Avança Brasil 2000* and the 2004–2007 Pluriannual Plan, which included the reconstruction of highways in the Amazon (Fearnside & Graça, 2009). Because, as we commonly see in the Amazon, a simple PPP announcement for the installation of any large enterprise is capable of promoting migration and irregular occupation of land by people in search of opportunities and cheap land, consequently leading to environmental degradation such as what is occurring around BR-319.



**Fig. 11** Official 40-km area of influence defined by Interministerial Ordinance 60 of 24 March 2015 for environmental licensing of highways in the Amazon region (a and b); the

expansion of deforestation in the BAU\_2 scenario is shown for 2100 (b) in relation to the reference year (a)

## Conclusion

The results presented in this study reflect the contribution of roads to advancing the agricultural frontier in Brazil's state of Amazonas, despite the limitations of environmental models in representing the complexity of the dynamics of deforestation in the Amazon. Given the assumptions of our model, we conclude that by 2100 reconstruction of Highway BR-319 (BAU\_2) would increase deforestation along the highway (Regions 3 and 4) and in the regions with roads directly connected to BR-319 (Regions 5 and 7) by 60% in relation to deforestation in the projected scenario without reconstruction (BAU\_1).

In relation to protected areas (indigenous lands and conservation units), despite deforestation increasing over time, these areas continue to play an important role in protecting the forest, and it is up to the government to increase protection, monitoring, and inspection, as well as to create new areas, in view of the advance of deforestation in non-designated public forests. Unlike protected areas, settlements do not provide environmental protection, regardless of their modality, and it is the government's responsibility to create environmental control mechanisms.

The results show that modeling the deforestation of a road enterprise can be part of the processes of environmental licensing and strategic environmental assessment for the formulation and implementation of policies, plans, and government investment programs in the Amazon region. Models of this type can better define the area of influence and expansion of socio-environmental impacts, as well as provide information for measures to mitigate and control negative impacts and to guide decision-making on whether or not to implement construction projects.

**Acknowledgments** We would like to thank the National Institute for Amazonian Research (INPA) and the State University of Amazonas (UEA) for supporting the Postgraduate Program in Climate and Environment (CLIAMB). The first author thanks the Brazilian Institute of the Environment and Renewable Natural Resources (IBAMA) for its support. We thank the Terrestrial Climate Systems Modeling Laboratory (LABCLIM/UEA) for the physical structure of numerical simulations; the Amazonas State Research Support Foundation (FAPEAM) (Resolution 003/2019) and the Coordination for the Improvement of Higher Education Personnel – CAPES (Finance Code 001) for institutional support. The PMF research is supported by FINEP/Rede CLIMA (01.13.0353-00), Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) (Process 2020/08916-8), FAPEAM

(Process 01.02.016301.000289/2021- 33), INPA (PRJ15.125) and National Council for Scientific and Technological Development (CNPq) (Process 312450/2021-4).

**Author contribution** Jerfferson Lobato dos Santos: conceptualization, data curation, calibration, validation, analysis, writing, original draft preparation.

Aurora Miho Yanai: conceptualization; methodology; writing—reviewing and editing.

Paulo Maurício Lima de Alencastro Graça: conceptualization; methodology; writing—reviewing and editing.

Francis Wagner Silva Correia: conceptualization; supervision; writing—reviewing and editing.

Philip Martin Fearnside: conceptualization; supervision; writing—reviewing and editing.

**Data availability** Datasets generated and/or analyzed during the current study are available from the corresponding author upon reasonable request.

## Declarations

**Ethics approval** All authors have read, understood, and complied as applicable with the statement on “Ethical responsibilities of Authors” as found in the Instructions for Authors.

**Competing interests** The authors declare no competing interests.

**Third-party material** All material is the property of the authors and no permissions are required.

**Double publication** The results/data/figures in this manuscript have not been published elsewhere, nor are they under consideration by another publisher.

## References

- Aguiar, A. P. D. (2006). *Modeling land use change in the Brazilian Amazon: Exploring intra-regional heterogeneity (Modelagem de Mudança do Uso da Terra na Amazonia: Explorando a Heterogeneidade Intrarregional)*. INPE Retrieved 30 November 2020, from <http://mtc-m16b.sid.inpe.br/col/sid.inpe.br/MTC-m13@80/2006/08.10.18.21/doc/publicacao.pdf>
- Aguiar, A. P. D., Vieira, I. C. G., Assis, T. O., Dalla-Nora, E. L., Toledo, P. M., Santos-Júnior, R. A. O., Batistela, M., Coelho, A. S., Savaget, E. K., Aragão, L. E. O. C., Nobre, C. A., & Ometto, J. P. H. (2016). Land use change emission scenarios: Anticipating a forest transition process in the Brazilian Amazon. *Global Change Biology*, 22, 1821–1840.
- Andrade, M. B. T., Ferrante, L., & Fearnside, P. M. (2021). Brazil's Highway BR-319 demonstrates a crucial lack of environmental governance in Amazonia. *Environmental*

- Conservation*, 48(3), 161–164. <https://doi.org/10.1017/S0376892921000084>
- ANM (Agência Nacional de Mineração). (2021). SHP. Retrieved 18 May 2021, from <https://dados.gov.br/datas-et/sistema-de-informacoes-geograficas-da-mineracao-sigmine>
- Araújo, R., Castro, E., Rocha, G., Sá, M. E., Matih, A., Monteiro, M., Puty, C., Monteiro, R., Canto, O., & Benatti, J. (2008). Estado e Sociedade na BR 163: desmatamento, conflitos e processos de ordenamento territorial. In Castro, E., *Sociedade, Território e Conflitos: Br 163 em Questão* (p. 297). NAEA.
- Barber, C. P., Cochrane, M. A., Souza, C. M., & Laurance, W. F. (2014). Roads, deforestation, and the mitigating effect of protected areas in the Amazon. *Biological Conservation*, 177, 203–209. <https://doi.org/10.1016/j.biocon.2014.07.004>
- Barbosa, L. G., Alves, M. A. S., & Grelle, C. E. V. (2021). Actions against sustainability: Dismantling of the environmental policies in Brazil. *Land Use Policy*, 104, 105384. <https://doi.org/10.1016/j.landusepol.2021.105384>
- Barni, P. E., Fearnside, P. M., & Graça, P. M. L. A. (2015). Simulating deforestation and carbon loss in Amazonia: Impacts in Brazil's Roraima State from reconstructing Highway BR-319 (Manaus-Porto Velho). *Environmental Management*, 55, 259–278. <https://doi.org/10.1007/s00267-014-0408-6>
- Bonham-Carter, G. F. (1994). *Geographic information systems for geoscientists: modelling with GIS*. Oxford: Pergamon. <https://doi.org/10.1016/C2013-0-03864-9>
- Brazil (2015). *Portaria Interministerial 60, de 24 de março de 2015*. Retrieved 9 September 2022, from [http://portal.iphan.gov.br/uploads/legislacao/Portaria\\_Interministerial\\_60\\_de\\_24\\_de\\_marco\\_de\\_2015.pdf](http://portal.iphan.gov.br/uploads/legislacao/Portaria_Interministerial_60_de_24_de_marco_de_2015.pdf)
- Brito, R., & Castro, E. R. (2018). Desenvolvimento e conflitos na Amazônia: um olhar sobre a colonialidade dos processos em curso na BR-163/Development and Conflict in the Amazon - a glimpse into the coloniality of on-going processes in BR-163. *REVISTA NERA*, 42, 51–73. <https://doi.org/10.47946/rnera.v0i42.5679>
- Campos, P. B. R., Almeida, C. M. D., & Queiroz, A. P. D. (2022). Spatial dynamic models for assessing the impact of public policies: The case of unified educational centers in the periphery of São Paulo city. *Land*, 11, 922. <https://doi.org/10.3390/land11060922>
- Castro, E. R., Monteiro, R., & Castro, C. P. (2004). Dinâmica de atores, uso da terra e desmatamento na rodovia Cuiabá-Santarém. *Paper do NAEA*, 179, 61. <https://doi.org/10.18542/papersnaea.v13i1.11558>
- ClimaInfo. (2022). *Desmonte ambiental: Próximo Congresso será "mais boiadeiro" que o atual*. ClimaInfo 6 October 2022. <https://climaInfo.org.br/2022/10/06/desmonte-ambiental-proximo-congresso-sera-mais-boiadeiro-que-o-atual/>
- DNIT (Departamento Nacional de Infraestrutura de Transporte) (2016). *BR-319/AM/RO Histórico do licenciamento ambiental da rodovia e situação dos instrumentos celebrados para o atendimento às condições do licenciamento*. Retrieved 25 May 2020, from <http://legis.senado.leg.br/sdleg-getter/documento/download/d3816d09-2e92-4f73-bf18-18e2c37b0589>
- DNIT (Departamento Nacional de Infraestrutura de Transporte) (2021). *DNIT Geo*. Retrieved 15 March 2022, from <http://servicos.dnit.gov.br/vgeo/>
- dos Santos-Junior, M. A., Yanai, A. M., Sousa-Junior, F. O., Freitas, I. S., Pinheiro, H. P., Oliveira, A. C. R., Silva, F. L., Graça, P. M. L. A., & Fearnside, P. M. (2018). *BR-319 como Propulsora de desmatamento: Simulando o Impacto da Rodovia Manaus-Porto Velho, Instituto de Desenvolvimento Sustentável da Amazônia (IDESAM)* (p. 56) <https://idesam.org/simula-desmatamento-br319>.
- Facundes, F. S., Lima, R. A. P. L., & Santos, V. F. (2019). Expansion des réseaux routiers en Amazonie orientale - Perimetral Norte, Amapá. *CONFINS - Revista Franco-Brasileira de Geografia*, 42. <https://doi.org/10.4000/confins.23789>
- Fearnside, P. M. (2003). *A Floresta Amazônica nas Mudanças Globais* (p. 134). Instituto Nacional de Pesquisas da Amazônia (INPA).
- Fearnside, P.M. (2018). *BR-319 e a destruição da floresta amazônica. Amazônia Real, 19 October 2018*. Retrieved 25 May 2020, from <http://amazoniareal.com.br/br-319-e-destruicao-da-floresta-amazonica>
- Fearnside, P. M. (2020). TransPurus: Amazonia's biogeochemical cycles depend on the fate of the region's largest block of intact forest. In *American Geophysical Union (AGU) Fall Meeting 2020*. Paper Number: GC009-0015.
- Fearnside, P. M. (2022a). Como sempre, os negócios: o ressurgimento do desmatamento na Amazônia brasileira. pp. 363-368. In P. M. Fearnside (Ed.), *Destruição e Conservação da Floresta Amazônica* (Vol. 1, p. 368). Editora do INPA <https://repositorio.inpa.gov.br/handle/1/38899>
- Fearnside, P. M. (2022b). Uso da terra na Amazônia e as mudanças climáticas globais. pp. 21-38. In P. M. Fearnside (Ed.), *Destruição e Conservação da Floresta Amazônica* (Vol. 1, p. 368). Editora do INPA. <https://repositorio.inpa.gov.br/handle/1/38899>
- Fearnside, P. M., & Graça, P. M. L. A. (2009). BR-319: A rodovia Manaus-Porto Velho e o impacto potencial de conectar o arco de desmatamento à Amazônia central. *Novos Cadernos NAEA*, 12(1), 19–50. <https://doi.org/10.5801/ncn.v12i1.241>
- Fearnside, P. M., Graça, P. M. L. A., Keizer, E. W. H., Maldonado, F. D., Barbosa, R. I., & Nogueira, E. M. (2009). Modelagem do desmatamento e emissões de gases do efeito estufa na região sob influência da Rodovia Manaus-Porto Velho (BR-319). *Revista Brasileira de Meteorologia*, 24(2), 208–233. <https://doi.org/10.1590/S0102-77862009000200009>
- Fearnside, P. M., Ferrante, L., Yanai, A. M., & Isaac-Júnior, M. A. (2020). Trans-Purus: Brazil's last intact Amazon forest at immediate risk (commentary). In *Mongabay*. Retrieved 22 May 2021, from <https://news.mongabay.com/2020/11/trans-purus-brazils-last-intact-amazon-forest-at-immediate-risk-commentary/>
- Ferrante, L., & Fearnside, P. M. (2019). Brazil's new president and "ruralists" threaten Amazonia's environment, traditional peoples and the global climate. *Environmental Conservation*, 46(4), 261–263. <https://doi.org/10.1017/S0376892919000213>



- Ferrante, L., Andrade, M. B. T., & Fearnside, P. M. (2021a). Land grabbing on Brazil's Highway BR-319 as a spearhead for Amazonian deforestation. *Land Use Policy*, *108*, 105559. <https://doi.org/10.1016/j.landusepol.2021.105559>
- Ferrante, L., Andrade, M. B. T., Leite, L., Silva-Júnior, C. A., Lima, M., Coelho-Junior, M. G., Silva-Neto, E. C., Campolina, D., Carolino, K., Diele-Viegas, L. M., Pereira, E. J. A. L., & Fearnside, P. M. (2021b). Brazil's Highway BR-319: The road to the collapse of the Amazon and the violation of indigenous rights. *Die Erde - Journal of the Geographical Society of Berlin*, *152*(1), 65–70. <https://doi.org/10.12854/erde-2021-552>
- FUNAI (Fundação Nacional do Índio). n.d. (no date information). *Download de dados geográficos: Terra Indígena (Regularizada, Homologada, Declarada, Delimitada e Área em Estudo)*. Retrieved 25 May 2020, from <http://www.funai.gov.br/index.php/shape>
- Graça, P. M. L. A., Maldonado, F. D., & Fearnside, P. M. (2007). Detecção de desmatamento em novas áreas de expansão agropecuária no sul do Amazonas utilizando imagens CBERS-2. In *Anais XIII Simpósio Brasileiro de Sensoriamento Remoto* (pp. 917–924). <https://repositorio.inpa.gov.br/handle/1/31089>
- Hagen, A. (2003). Fuzzy set approach to assessing similarity of categorical maps. *International Journal of Geographical Information Science*, *17*, 235–249. <https://doi.org/10.1080/13658810210157822>
- IBGE (Instituto Brasileiro de Geografia e Estatística) (2017). *Geociências*. Retrieved 25 May 2020, from <https://www.ibge.gov.br/geociencias/downloads-geociencias.html>
- ICMBIO (Instituto Chico Mendes de Conservação da Biodiversidade) (2019). *Limites das Unidades de Conservação Federais (atualizado em julho de 2019): Unidades de Conservação Federais – SHP (SIRGAS2000)*. Retrieved 25 May 2020, from <https://www.icmbio.gov.br/portal/geoprocessamento/151-menu-servicos/4004-downloads-mapa-tematico-e-dados-geoestatisticos-das-uc-s>
- INCRA (Instituto Nacional de Colonização e Reforma Agrária). (2015). *Relatório de Assentamentos*. Retrieved 25 September 2020, from <https://painel.incra.gov.br/sistemas/Painel/>
- INCRA (Instituto Nacional de Colonização e Reforma Agrária). n.d. (no update date information). *Exportar shapefile*. Retrieved 25 May 2020, from [http://certificacao.incra.gov.br/csv\\_shp/export\\_shp.py](http://certificacao.incra.gov.br/csv_shp/export_shp.py)
- INPE (Instituto Nacional de Pesquisas Espaciais) (2020). *PRODES – Amazônia*. Retrieved 6 July 2022, from <http://www.obt.inpe.br/OBT/assuntos/programas/amazonia/prodes>
- INPE (Instituto Nacional de Pesquisas Espaciais). (2022). *PRODES*. Retrieved 25 May 2022, from [http://terrabrasilis.dpi.inpe.br/app/dashboard/deforestation/biomes/legal\\_amazon/rates](http://terrabrasilis.dpi.inpe.br/app/dashboard/deforestation/biomes/legal_amazon/rates)
- Kohlhepp, G. (2002). Conflitos de interesses no ordenamento territorial da Amazônia brasileira. *Estudos Avançados*, *16*(45). <https://doi.org/10.1590/S0103-4014200200020004>
- Laurance, W. F., Cochrane, M. A., Bergen, S., Fearnside, P. M., Delamônica, P., Barber, C., D'Angelo, S., & Fernandes, T. (2001). The future of the Brazilian Amazon. *Science*, *291*, 438–439. <https://doi.org/10.1126/science.291.5503.438>
- Laurance, W. F., & Balmford, A. (2013). A global map for road building. *Nature*, *495*, 308–309. <https://doi.org/10.1038/495308a>
- Leite-Filho, A.T., Soares-Filho, B.S., Davis, J.L. & Rodrigues, H.O. (2020). *Modeling Environmental Dynamics with Dinamica EGO*. Retrieved 20 October 2020, from [https://www.csr.ufmg.br/dinamica/dokuwiki/doku.php?id=guide\\_book\\_start](https://www.csr.ufmg.br/dinamica/dokuwiki/doku.php?id=guide_book_start)
- Lessa, R. (1991). *Amazônia: as raízes da destruição*. Ed. Atual. Edição.
- Lewinsohn, T. M., & Prado, P. I. (2002). *Biodiversidade brasileira: síntese do estado atual do conhecimento*. Contexto.
- Loureiro, V. R. (2002). Amazônia: uma história de perdas e danos, um futuro a (re)construir. *Estud. Avançados*, *16*(4). <https://doi.org/10.1590/S0103-40142002000200008>
- Maeda, E. E., Almeida, C. M., Ximenes, A. C., Formaggio, A. R., Shimabukuro, Y. E., & Pellikka, P. (2011). Dynamic modeling of forest conversion: Simulation of past and future scenarios of rural activities expansion in the fringes of the Xingu National Park, Brazilian Amazon. *International Journal of Applied Earth Observation and Geoinformation*, *13*(3), 435–446. <https://doi.org/10.1016/j.jag.2010.09.008>
- Marengo, J. A., Souza, C. M., Thonicke, K., Burton, C., Halladay, K., Betts, R. A., Alves, L. M., & Soares, W. R. (2018). Changes in climate and land use over the Amazon region: Current and future variability and trends. *Frontiers in Earth Science*, *6*, 228. <https://doi.org/10.3389/feart.2018.00228>
- Mazzotti, F. J., & Vinci, J. J. (2007). *Validation, verification, and calibration: Using standardized terminology when describing ecological models*. IFAS Extension, University of Florida <https://edis.ifas.ufl.edu/pdf/UW/UW25600.pdf>
- MMA (Ministério do Meio Ambiente). (2016). *Plano de ação para prevenção e controle do desmatamento da Amazônia Legal: pelo uso sustentável e conservação da floresta. 3ª fase (2012-2015)*. Casa Civil, Brasília Retrieved 6 July 2020, from [https://www.mma.gov.br/images/arquivo/80120/PPCDAmPPCerrado-EncartePrincipal-GPTI\\_psite.pdf](https://www.mma.gov.br/images/arquivo/80120/PPCDAmPPCerrado-EncartePrincipal-GPTI_psite.pdf)
- MMA (Ministério do Meio Ambiente) (2018). *Balço de execução 2018: PPCDAm e PPCerrado 2016-2020*. Retrieved 6 July 2020, from <https://www2.camara.leg.br/atividade-legislativa/comissoes/comissoes-temporarias/externas/56a-legislatura/politicas-para-integracao-meio-ambiente-e-economia/expedientes-recebidos/ric-1577-2019-ministerio-do-meio-ambiente>
- Moutinho, P. (2009). *Desmatamento na Amazônia: desafios para reduzir as emissões de gases de efeito estufa do Brasil*. Retrieved 10 June 2020, from <http://www.fbd.org.br/IMG/pdf/doc-411.pdf>. (accessed: 10 June 2020).
- MPOG (Ministério do Planejamento Orçamento e Gestão). (2004). *PPA 2004-2007 - Lista Geral de Projetos de Infra-estrutura. Setembro 2004*. MPOG.
- Nepstad, D., Stickler, C. M., & Almeida, O. T. (2006). Globalization of the Amazon soy and beef industries: Opportunities for conservation. *Conservation Biology*. <https://doi.org/10.1111/j.1523-1739.2006.00510.x>

- Nobre, A. D. (2014). *The Future Climate of Amazonia: Scientific Assessment Report*. CCST-INPE.
- Nobre, C. A., Sampaio, G., Borma, L. S., Castilla-Rubio, J. C., Silva, J. S., Cardoso, M., et al. (2016). The fate of the Amazon forests: Land-use and climate change risks and the need of a novel sustainable development paradigm. *Proceedings of the National Academy of Sciences of the United States of America*, 113, 10759–10768. <https://doi.org/10.1073/pnas.1605516113>
- Oliveira-Neto, T. (2014). A Geopolítica rodoviária na Amazônia: BR-319. *Revista de Geopolítica*, 5(2), 109–128.
- Partidário, M. R. (2011). From EIA to SEA. chapter 14. In F. Indovina & L. Fregolent (Eds.), *Environmental Sustainability, Monographic issue*, n. 71/72-2001. Archivio di Studi Urbani e Regionali.
- Partidário, M. R. (2003). Avaliação de Impactes Ambientais de Políticas, Planos e Programas. *Ambiente*, 21(8).
- Pellin, A., Lemos, C. C., Tachard, A., Oliveira, I. S. D., & Souza, M. P. (2011). Avaliação ambiental estratégica no Brasil: considerações a respeito do papel das agências multilaterais de desenvolvimento. *Artigos Técnicos Eng Sanit Ambient*, 16(1).
- Pontius-Jr, R. G., Boersma, W., Castella, J.-C., Clarke, K., de Nijs, T., Dietzel, C., et al. (2007). Comparing the input, output, and validation maps for several models of land change. *The Annals of Regional Science*, 42(1), 11–37. <https://doi.org/10.1007/s00168-007-0138-2>
- Pontius-Jr, R. G., Huffaker, D., & Denman, K. (2004). Useful techniques of validation for spatially explicit land change models. *Ecological Modelling*, 179(4), 445–461. <https://doi.org/10.1016/j.ecolmodel.2004.05.010>
- Ramos, C. J. P., Graça, P. M. L. A., & Fearnside, P. M. (2018). Deforestation dynamics on an Amazonian peri-urban frontier: Simulating the influence of the Rio Negro Bridge in Manaus, Brazil. *Environmental Management*, 62(6), 1134–1149. <https://doi.org/10.1007/s00267-018-1097-3>
- Rocha, V. M., Correia, F. W. S., Satyamurty, P., de Freitas, S. R., Moreira, D. S., da Silva, P. R. T., & Fialho, E. S. (2015). Impacts of land cover and greenhouse gas (GHG) concentration changes on the hydrological cycle in amazon basin: A regional climate model study. *Revista Brasileira de Climatologia*, 15, 7–27. <https://doi.org/10.5380/abclima.v15i0.36386>
- Rodrigues, H. O., Soares-Filho, B. S., & Costa, W. L. S. (2007). Dinamica EGO, uma plataforma para modelagem de sistemas ambientais. In *Anais XIII Simpósio Brasileiro de Sensoriamento Remoto, Florianópolis, Brasil, 21-26 abril 2007* (pp. 3089–3096). INPE.
- Roriz, P. A. C., Yanai, A. M., & Fearnside, P. M. (2017). Deforestation and carbon loss in southwest Amazonia: Impact of Brazil’s revised forest code. *Environmental Management*, 60, 367–382. <https://doi.org/10.1007/s00267-017-0879-3>
- Rykiel-Jr, E. J. (1996). Testing ecological models: The meaning of validation. *Ecological Modelling*, 90, 229–244.
- Santos, Y. L. F., Yanai, A. M., Ramos, C. J. P., Graça, P. M. L. A., Veiga, J. A. P., Correia, F. W. S., & Fearnside, P. M. (2021). Amazon deforestation and urban expansion: Simulating future growth in the Manaus Metropolitan Region. *Brazil. Journal of Environmental Management*, 304(1), 114279. <https://doi.org/10.1016/j.jenvman.2021.114279>
- Saatchi, S. S., Harris, N. L., Brown, S., Lefsky, M., Mitchard, E., Salas, W., Zutta, B., Buermann, W., Lewis, S., Hagen, S., Petrova, S., White, L., Silman, M. A., & Morel, A. (2011). Benchmark map of forest carbon stocks in tropical regions across three continents. *Proceedings of the National Academy of Sciences of the United States of America*, 108(24), 9899–9904. <https://doi.org/10.1073/pnas.1019576108>
- SEMA (Secretaria do Meio Ambiente). (2020). UNIDADE DE CONSERVAÇÃO (atualizado em 2020). Retrieved 25 May 2020, from <https://meioambiente.am.gov.br/unidade-de-conservacao/>
- Siqueira-Gay, J., Metzger, J. P., Sánchez, L. E., et al. (2022). Strategic planning to mitigate mining impacts on protected areas in the Brazilian Amazon. *Nature Sustainability*, 5, 853–860. <https://doi.org/10.1038/s41893-022-00921-9>
- Soares Filho, B. S., Pennachin, C. L., & Cerqueira, G. (2002). DINAMICA – a stochastic cellular automata model designed to simulate the landscape dynamics in an Amazonian colonization frontier. *Ecological Modelling*, 154(3), 217–235. [https://doi.org/10.1016/S0304-3800\(02\)00059-5](https://doi.org/10.1016/S0304-3800(02)00059-5)
- Soares Filho, B. S., Alencar, A., Nepstad, D., Cerqueira, G., Dias, M., Rivero, S., Solórzanos, L., & Voll, E. (2004). Simulating the response of land-cover change to road paving and governance along a major Amazon highway: The Santarém-Cuiabá corridor. *Global Change Biology*, 10, 745–764. <https://doi.org/10.1111/j.1529-8817.2003.00769.x>
- Soares-Filho, B. S., Nepstad, D., Curran, L., Voll, E., Cerqueira, G., Garcia, R. A., Ramos, C. A., McDonald, A., Lefebvre, P., & Schlesinger, P. (2006). Modelling conservation in the Amazon basin. *Nature*, 440, 520–523. <https://doi.org/10.1038/nature04389>
- Soares-Filho, B., Rodrigues, H., & Follador, M. (2013). Um método híbrido analítico-heurístico para calibrar modelos de mudança de uso da terra. *Ambiente. Modelo. Softw.*, 43, 80–87.
- TALANOIA. (2022). *Reconstrução: 401 atos do Poder Executivo Federal (2019 - 2022) a serem revogados ou revisados para a reconstrução da agenda climática e ambiental brasileira* (p. 169). Instituto Talanoia <https://www.politicaprinteiro.org/2022/11/03/reconstrucao/>
- Weng, W., Luedeke, M. K., Zemp, D. C., & Lakes, T. (2018). Aerial and surface rivers: Downwind impacts on water availability from land use changes in Amazonia. *Hydrology and Earth System Sciences*, 22(1), 911–927. <https://doi.org/10.5194/hess-22-911-2018>
- Yanai, A. M., Nogueira, E. M., Graça, P. M. L. A., & Fearnside, P. M. (2017). Deforestation and carbon stock loss in Brazil’s Amazonian settlements. *Environmental Management*, 59, 393–409. <https://doi.org/10.1007/s00267-016-0783-2>

**Publisher’s Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner)

holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article

is solely governed by the terms of such publishing agreement and applicable law.