

Modulatory role of L-carnitine against microcystin-LR-induced immunotoxicity and oxidative stress in common carp

Xiu-Mei Chen · Gui-Liang Guo · Li Sun ·
Qiu-Shi Yang · Gui-Qin Wang · Dong-Ming Zhang

Received: 12 December 2016 / Accepted: 5 February 2017 / Published online: 17 March 2017
© Springer Science+Business Media Dordrecht 2017

Abstract Microcystin-LR (MCLR), one of the most popular microcystins (MCs) found in many field water bodies around the world, poses great health risks to animals and humans. In the present study, healthy common carp (initial weight 24.8 ± 2.3 g) were randomly assigned to five groups. Group I was fed on normal diet as control. Group II was maintained on normal diet and received MCLR intraperitoneal injection ($150 \mu\text{g kg}^{-1}$ BW). Common carp in groups III, IV, and V were daily pretreated with L-carnitine (LC) at doses of 0.5, 1.0, and 2.0 g kg^{-1} of the diet for 4 weeks prior to MCLR intraperitoneal injection. The results showed that MCLR alone led to a significant downregulation in immune response, including serum complement C3, lysozyme, and bactericidal activity. However, oxidative stress response: catalase (CAT), superoxide dismutase (SOD), glutathione (GSH), glutathione peroxidase (GPx), and lipid peroxidation (LPO) levels were significantly increased. Similarly, gene expressions of inflammatory IL-1 β , TNF- α , IFN I, and heat shock proteins (HSP70 and HSP90) were also upregulated after challenged with MCLR. However, LC pretreated group caused a significant elevation in immune

response (C3, lysozyme, and bactericidal activity) and gene expressions of inflammatory IL-1 β , TNF- α , IFN I, and heat shock proteins (HSP70 and HSP90) after MCLR stress. Antioxidant activities (CAT, SOD, GSH, GPx, and LPO) were returned to background levels at 96 h after MCLR challenge. Strikingly, LC supplementation at 2.0 g kg^{-1} has been considered the optimum for common carp since it exhibited enhancement of immune response and antioxidant activity over the level 0.5 and 1.0 g kg^{-1} , and even better than that of control level. It was concluded that LC as a functional feed additive significantly inhibited the progression of MCLR-induced immunotoxicity and oxidative stress in common carp.

Keywords L-Carnitine · MCLR-induced oxidative stress · Antioxidant enzymes · Common carp

Introduction

Cyanobacterial blooms and the associated cyanotoxins are being increasingly reported worldwide (Palus et al. 2007; Graham et al. 2010; Davis et al. 2012). These toxins can be accumulated in aquatic organisms and transferred to higher trophic levels, representing a health hazard to animals and humans (Chen et al. 2009; Campos and Vasconcelos 2010). Among all the cyanotoxins, microcystins (MCs) are the most frequently studied due to their wide distribution and high toxicity. Microcystin-LR (MCLR) is generally recognized as one of the most toxic microcystin variants, and the concentration in surface waters often exceeds the World Health Organization

X.-M. Chen · Q.-S. Yang · G.-Q. Wang (✉) ·
D.-M. Zhang (✉)
College of Animal Science and Technology, Jilin Agricultural
University, Changchun 130118, China
e-mail: wangguiqing131@163.com
e-mail: zhangdongming1961@163.com

G.-L. Guo · L. Sun
Testing Center of Quality and Safety in Aquatic Product, 777
CaiYu Road, Changchun, Jilin, China

advisory level of $1 \mu\text{g l}^{-1}$ (Organization 2004). MCs are released from the cyanobacterial cells into the water bodies where aquatic organisms especially fish spend their whole life stage, including growth, reproduction, and embryonic development (Zhang et al. 2009).

Although MCs have been demonstrated to cause damages to fish intestine, kidney, gills, heart, and brain (Qiu et al. 2009; Li et al. 2012; Chen et al. 2012; Trinchet et al. 2011; Li et al. 2013), the liver is the most affected organ in fish, with symptoms of hepatocyte dissociation, degeneration, and necrosis (Malbrouck et al. 2003; Li and Xie 2009). A classic toxic mechanism of MCs is their inhibition of protein phosphatase 1 and 2A, leading to increased protein phosphorylation, which is directly related to their cytotoxicity and tumor-promoting activity (Honkanen et al. 1990). Several evidences showed that oxidative stress played an important role in the pathogenesis of MC toxicity in aquatic organisms such as crab (Pinho et al. 2003) and tilapia *Oreochromis niloticus* (Prieto et al. 2006, 2007).

Because of the rapid, irreversible, and severe damage to the liver caused by MCs, therapy is likely to have little or no value; effective prophylaxis is critical. In spite of the potential human hazards associated with MCs, very little work has been done on the development of effective chemoprotectants or antidotes against these toxins. L-Carnitine (LC) is synthesized from the essential amino acids (lysine and methionine) with the assistance of vitamin C and other secondary compounds produced in the body (Harpaz et al. 2005). The supplementation of LC in fish diets has been advocated in aquaculture for multiple reasons: LC can be used to improved growth performance indices including specific growth rate (SGR), feed conversion ratio (FCR), and survival rate (Saliny et al. 1994), to protect against toxic levels of xenobiotics (Schreiber et al. 1997), to ameliorate stress that is related to water temperature extremes, to facilitate better acclimation to water temperature changes (Harpaz et al. 1999), and to improve immune responses (Safari et al. 2015). Of late, the role of LC as an antioxidant agent has been confirmed on ischemia–reperfusion injury, adriamycin-induced membrane damage, and diphtheria toxins (Ma et al. 2008). However, there is little information about the protective effects of LC on MCLR-induced immunotoxicity and oxidative stress in aquaculture. The aim of the present study is to evaluate the protective effect of LC on the prevention of MCLR stress by evaluating the activities of antioxidant-

related enzyme and gene expressions of inflammatory and immune response in common carp.

Materials and methods

Feed and experimental design

The experimental design was completely randomized with four treatment diets, each of which was replicated three times. For each treatment replicate, 30 common carp ($24.8 \pm 2.3 \text{ g}$) were randomly chosen and placed in 300-l cycling-filtered plastic tanks containing continuously circulating aerated water. The four treatment diets were as follows: group I (control group I) and group II (control group II) were always fed with control (basic) diet throughout the feeding trial; groups III, IV, and V were fed 0.5, 1.0, and 2.0 g kg^{-1} LC (Sigma, $\geq 98\%$), respectively. Fish were fed twice a day (9:00 and 15:00) at a rate of 3% of the bodyweight and kept in glass aquaria at $23 \pm 1 \text{ }^\circ\text{C}$ with laboratory conditions as mentioned above for 4 weeks. Tank bottom debris was removed by siphon daily, and about one third of the water was replaced daily.

MCLR challenge experiments

After 4 weeks of feeding the fish, one set of fish ($n = 25$) from groups II, III, IV, and V was injected i.p. with MCLR at dose of $150 \mu\text{g kg}^{-1}$ BW. Doses of MCLR used in the experiment were based on the result from 48-h LD 50 study of MCLR in our previous experiment, which calculated the LD 50 value ($310.5 \mu\text{g kg}^{-1}$ BW) with a 95% confidence interval ($256.8\text{--}364.2 \mu\text{g kg}^{-1}$ BW). The control (group I) fish were injected i.p. with equal volume of physiological saline solution (0.85% NaCl). In the experiment, sampling points were set at 0, 12, 24, 48, and 96 h (two and one fish died at 12 and 24 h, and no more dead fish was found at 48 and 96 h. However, no mortality was found in the control.). At each sampling point, five fish from each dose group were anesthetized with 0.02% tricaine methane sulfonate (MS-222) solution. Live of five individuals were excised and immediately frozen in liquid nitrogen and then stored at $-80 \text{ }^\circ\text{C}$ for analysis of gene expression and enzyme activities. Blood of all fish from each group was also analyzed for serum

complement C3, lysozyme, and bactericidal activity.

Antioxidant enzyme activity assays

Total superoxide dismutase (SOD) activity was determined following the methods of Beauchamp and Fridovich (Lawrence et al. 1976). Catalase (CAT) activity was determined by measuring the decrease in H₂O₂ concentration. Glutathione peroxidase (GPx) activity was assayed by following the rate of NADPH oxidation at 340 nm by the coupled reaction with glutathione reductase according to the method by Lawrence and Burk (1976). The contents of GSH were determined using commercial kits (Nanjing, Jiangsu, China) following the manufacturer's instructions.

Lipid peroxidation level assays

The thiobarbituric acid (TBA) method of Esterbauer and Cheeseman (1990) was used to determine the lipid peroxidation by determining the amount of TBA reactive substances present in the liver homogenates obtained from common carp.

Serum immune parameter assays

Blood was sampled from the caudal vasculature using a 2.5-ml syringe after the fish were euthanized by overdose of MS-222. Individual fish was sampled only once to avoid the influence on the assays due to multiple bleeding and handling stress on the fish. Lysozyme activity measurement was based on the turbidimetric method described by Ellis (1990). A unit of lysozyme activity was defined as the amount of serum lysozyme that caused a decrease in absorbance of 0.001 per minute at 530 nm.

The serum bactericidal activity was determined according to the method which previously described (Barnes et al. 2003) with some modification. *Staphylococcus aureus* isolates were adjusted to 1.0 (OD₅₄₀; 3.0×10^7 CFU ml⁻¹). Bacterial suspension and serum samples were mixed with 1:1 ratio and incubated for 90 min at 25 °C. Thereafter, 10 ml of serum and bacteria mixtures were transferred to the Shieh broth medium and cultured for 24 h at 25 °C. Phosphate-buffered saline (PBS, Sigma) was used instead of serum as negative control. Viable colonies were counted, and

results expressed as percentage of survival in the PBS controls.

The serum complement C3 level was assayed using Complement C3 assay kit (Jiancheng, Nanjing, Jiangsu, China) (Wang et al. 2011). Results are presented as complement C3 milligram per milliliter.

Immune related gene expression

The gene expressions of inflammatory IL-1 β , TNF- α , IFN I, and heat shock proteins (HSP70 and HSP90) were tested according to our previous study (Chen et al. 2015). Total RNA was extracted from blood samples by TRIzol Reagent (SimGEN). Complementary DNA (cDNA) was then synthesized using the Reverse Transcriptase M-MLV Kit (TaKaRa) following the instructions. The real-time quantitative PCR was performed using THUNDERBIRD SYBR qPCR Mix Kit (TOYOBO) and carried out in a Stratagene MxProSystem (stratagene mx3005p, USA) in 96-well reaction plates. The β -actin gene was used as a housekeeping gene. The reaction mixture included 10 μ l of THUNDERBIRD SYBR qPCR Mix, 1 μ l of forward and reverse primer (10 mM), and 1 μ l of cDNA and was then filled up with ultra-pure water to a final total volume of 20 μ l. The cycling conditions were as follows: 95 °C for 30 s, followed by 40 cycles of 95 °C for 5 s and 60 °C for 34 s. All PCRs were performed at least three times. Additional dissociation curve analysis was performed and showed a single melting curve in all cases. The PCR efficiency of each primer was between 98.5 and 99.6%. Data were analyzed by the stratagene MxPro software (stratagene mx3005p, USA).

Data analysis

The data in this study were analyzed by Statistical Product and Service Solutions (SPSS 16.0) and expressed as the arithmetic mean \pm standard deviation. Data were analyzed by repeated measures analysis of variance (ANOVA) and the LSD post hoc test. The homogeneity of the replicates of the samples was checked by the Mann–Whitney *U* test. Differences between the three groups were measured and considered statistically different at $P < 0.05$ or $P < 0.01$.

Table 1 CAT (nmol substrate min⁻¹ ml⁻¹ blood) activity of common carp fed with LC-containing diet after MCLR stress

Groups	0 h	12 h	24 h	48 h	96 h
Group I	97.3 ± 5.5a	98.4 ± 6.3a	96.8 ± 2.3a	98.7 ± 5.4a	97. ± 4.3a
Group II	98.4 ± 3.9a	147.6 ± 7.5b	155.3 ± 6.7b	167.8 ± 4.9b	179.2 ± 5.7b
Group III	97.9 ± 7.1a	138.3 ± 4.3c	128.8 ± 5.1c	112.6 ± 5.8c	103.4 ± 3.8c
Group IV	99.5 ± 8.6a	139.0 ± 6.5c	122.2 ± 6.9c	107.3 ± 6.2c	104.8 ± 6.3c
Group V	97.8 ± 6.8a	123.5 ± 1.8d	109.8 ± 5.5d	103.8 ± 4.9d	104.6 ± 5.2c

Data are expressed as mean ± SD at the same sampling time with different letters as significant difference ($P < 0.05$). Group I: fish fed with basic diets and challenged with no MCLR. Group II: fish fed with basic diets and challenged with 150 µg kg⁻¹ BW MCLR. Group III: fish fed with 0.5 g kg⁻¹ LC and challenged with 150 µg kg⁻¹ BW MCLR. Group IV: fish fed with 1.0 g kg⁻¹ and challenged with 150 µg kg⁻¹ BW MCLR. Group V: fish fed with 2.0 g kg⁻¹ LC and challenged with 150 µg kg⁻¹ BW MCLR

Results

Activity of antioxidant enzyme after MCLR stress

The results of antioxidant enzyme activity assays were presented in Tables 1, 2, 3, 4 and 5. As shown in tables, antioxidant activity (CAT, SOD, GSH, GPx, LPO) in group II (MCLR treated but with no LC pretreatment) was significantly increased as compared with the control group, indicating severe oxidative stress in common carp.

Common carp fed the diets containing 0.5, 1.0, and 2.0 g kg⁻¹ LC had significantly higher CAT, SOD, and GPx activities compared to fish fed the control diets ($P < 0.05$) at 0 h (before MCLR stress) (Tables 1, 2 and 3); however, only SOD activities showed the statistical difference ($P < 0.05$) (Table 3). After MCLR stress, activities of CAT, SOD, and GPx were increased firstly and then decreased, the highest values were observed at 12 h, and group V had returned to the original

values at 48–96 h post-stress. As for GSH, slight increases in GSH content were detected after MCLR stress at 0 h (Table 4), but there was no significant difference among them. High dose of LC pretreated group (group V) showed a significant increase in the GSH levels when compared with MCLR ($P < 0.01$). Better preventive effects were observed at dose of 2.0 g kg⁻¹ than other LC pretreated groups.

LPO activity

The results of LPO activity assays are presented in Table 5. LPO levels in group II (MCLR treated with no LC pretreatment) were significantly elevated in response to MCLR treatment as compared with fish fed the control (basal) diets (group I, $P < 0.05$) at 0 h (before MCLR stress); it indicated that MCLR caused obvious oxidative damages on common carp. We found that the increase was

Table 2 GPx (U mg⁻¹) activity of common carp fed with LC-containing diet after MCLR stress

Groups	0 h	12 h	24 h	48 h	96 h
Group I	7.8 ± 0.8a	7.9 ± 1.1a	7.8 ± 1.4a	8.0 ± 0.9a	7.9 ± 0.7a
Group II	7.7 ± 0.6a	14.6 ± 1.3b	18.8 ± 0.5b	20.3 ± 0.3b	18.2 ± 0.4b
Group III	7.9 ± 0.7a	11.1 ± 0.9c	8.4 ± 0.8c	7.6 ± 0.4a	7.4 ± 0.6a
Group IV	8.1 ± 1.5a	10.5 ± 1.3c	8.5 ± 0.9c	7.7 ± 0.8a	7.8 ± 0.8a
Group V	7.7 ± 0.4a	9.2 ± 1.1c	8.8 ± 0.5c	8.0 ± 0.6a	7.6 ± 0.9a

Data are expressed as mean ± SD at the same sampling time with different letters as significant difference ($P < 0.05$). Group I: fish fed with basic diets and challenged with no MCLR. Group II: fish fed with basic diets and challenged with 150 µg kg⁻¹ BW MCLR. Group III: fish fed with 0.5 g kg⁻¹ LC and challenged with 150 µg kg⁻¹ BW MCLR. Group IV: fish fed with 1.0 g kg⁻¹ and challenged with 150 µg kg⁻¹ BW MCLR. Group V: fish fed with 2.0 g kg⁻¹ LC and challenged with 150 µg kg⁻¹ BW MCLR

Table 3 SOD (U ml^{-1}) activity of common carp fed with LC-containing diet after MCLR stress

Groups	0 h	12 h	24 h	48 h	96 h
Group I	92.3 ± 1.56a	92.9 ± 1.56a	93.2 ± 0.66a	93.8 ± 1.27a	93.4 ± 1.06a
Group II	93.2 ± 1.43a	112.7 ± 1.88b	118.1 ± 1.71b	114.2 ± 1.43b	111.2 ± 1.13b
Group III	98.4 ± 0.98b	106.5 ± 1.54a	86.2 ± 0.33c	88.4 ± 1.74c	90.4 ± 1.66a
Group IV	100.2 ± 1.87b	101.2 ± 1.29c	90.1 ± 1.25c	92.1 ± 1.89c	94.5 ± 1.05a
Group V	112.5 ± 0.67c	105.2 ± 0.35b	89.9 ± 0.77c	92.5 ± 1.05c	94.0 ± 1.24a

Data are expressed as mean ± SD at the same sampling time with different letters as significant difference ($P < 0.05$). Group I: fish fed with basic diets and challenged with no MCLR. Group II: fish fed with basic diets and challenged with $150 \mu\text{g kg}^{-1}$ BW MCLR. Group III: fish fed with 0.5 g kg^{-1} LC and challenged with $150 \mu\text{g kg}^{-1}$ BW MCLR. Group IV: fish fed with 1.0 g kg^{-1} and challenged with $150 \mu\text{g kg}^{-1}$ BW MCLR. Group V: fish fed with 2.0 g kg^{-1} LC and challenged with $150 \mu\text{g kg}^{-1}$ BW MCLR.

dramatically diminished by LC pretreatment at 2.0 g kg^{-1} dosage.

Immune parameter assays

Serum complement C3, bactericidal, and lysozyme activities of common carp among any treatment groups at day 0 had no significant changes. However, all data in group II were significantly decreased as compared to control group ($P < 0.05$) after MCLR stress. After treated with MCLR, serum complement C3, bactericidal, and lysozyme activities in groups III, IV, and V were increased, and group V had returned to the original values at 96 h post-stress (Figs. 1, 2 and 3).

Immune relates gene expression

The transcriptional changes of IL-1 β , TNF- α , and IFN I genes in the blood of common carp are shown in (Figs. 4, 5 and 6). All the test gene transcriptions were

markedly increased by MCLR stress as compared with group II ($P < 0.05$). The administration of LC alone (at 0 h) did not affect gene expression of messenger RNA (mRNA) of all the tested genes.

Compared with group II, transcriptional level of IL-1 β and IFN I in groups III, IV, and V was significantly decreased in most cases after MCLR treatment ($P < 0.05$). The TNF- α mRNA levels were downregulated in the fish fed the diets containing 0.5, 1.0, and 2.0 g kg^{-1} LC as compared to the control after MCLR stress; however, only group V showed the statistical difference ($P < 0.05$); it had recovered to the normal level at 96 h (Table 5).

The effects of LC in the expression of HSP70 and HSP90 genes after MCLR stress were determined. The relative expression of HSP70 gene (Fig. 7) transcript level was increased before 24 h but fell back to the blank levels at 96 h. The expression of HSP90 in groups III, IV, and V was increased firstly and then decreased; the highest values were observed at 12 h and had returned to the original values at 96 h (Fig. 8).

Table 4 GSH (U mg^{-1}) activity of common carp fed with LC-containing diet after MCLR stress

Groups	0 h	12 h	24 h	48 h	96 h
Group I	15.7 ± 2.2a	16.4 ± 1.7a	15.8 ± 1.2a	16.7 ± 1.8a	16.2 ± 1.3a
Group II	15.4 ± 1.5a	24.6 ± 0.6b	28.3 ± 0.3b	30.8 ± 1.1b	33.2 ± 0.9b
Group III	15.9 ± 2.0b	23.1 ± 0.8b	27.4 ± 1.3b	27.6 ± 0.8c	26.8 ± 1.8c
Group IV	16.1 ± 1.6b	24.2 ± 1.5b	24.5 ± 0.9c	22.7 ± 1.2d	23.8 ± 0.5c
Group V	15.8 ± 1.8c	22.5 ± 1.8b	19.8 ± 1.25d	16.8 ± 1.5a	17.6 ± 1.4a

Data are expressed as mean ± SD at the same sampling time with different letters as significant difference ($P < 0.05$). Group I: fish fed with basic diets and challenged with no MCLR. Group II: fish fed with basic diets and challenged with $150 \mu\text{g kg}^{-1}$ BW MCLR. Group III: fish fed with 0.5 g kg^{-1} LC and challenged with $150 \mu\text{g kg}^{-1}$ BW MCLR. Group IV: fish fed with 1.0 g kg^{-1} and challenged with $150 \mu\text{g kg}^{-1}$ BW MCLR. Group V: fish fed with 2.0 g kg^{-1} LC and challenged with $150 \mu\text{g kg}^{-1}$ BW MCLR.

Table 5 LPO (nmol TBARs g⁻¹) activity of common carp fed with LC-containing diet after MCLR stress

Groups	0 h	12 h	24 h	48 h	96 h
Group I	2.4 ± 0.3a	2.2 ± 0.4a	2.5 ± 0.4a	2.2 ± 0.2a	2.3 ± 0.2a
Group II	2.4 ± 0.6a	3.2 ± 0.1b	3.5 ± 0.3b	4.1 ± 0.2b	4.6 ± 0.3b
Group III	2.5 ± 0.4a	3.1 ± 0.2b	2.9 ± 0.2c	2.9 ± 0.3c	2.8 ± 0.2c
Group IV	2.4 ± 1.0a	2.9 ± 0.3b	2.7 ± 0.3c	2.7 ± 0.5c	2.8 ± 0.3c
Group V	2.4 ± 0.6a	2.9 ± 0.2b	2.6 ± 0.2d	2.5 ± 0.4d	2.6 ± 0.3c

Data are expressed as mean ± SD at the same sampling time with different letters as significant difference ($P < 0.05$). Group I: fish fed with basic diets and challenged with no MCLR. Group II: fish fed with basic diets and challenged with 150 µg kg⁻¹ BW MCLR. Group III: fish fed with 0.5 g kg⁻¹ LC and challenged with 150 µg kg⁻¹ BW MCLR. Group IV: fish fed with 1.0 g kg⁻¹ and challenged with 150 µg kg⁻¹ BW MCLR. Group V: fish fed with 2.0 g kg⁻¹ LC and challenged with 150 µg kg⁻¹ BW MCLR

Discussion

The monocyclic heptapeptide MCs are the most common and abundant cyanotoxins and pose a world health threat to humans and animals (de Figueiredo et al. 2004). Oxidative stress is considered the major cytotoxic mechanism of MCLR. Various antioxidants and detoxicants have been studied to identify new compounds that can regulate the oxidative stress and toxic effects caused by MCLR (Prieto et al. 2008). Recently, there is an increasing interest in natural feed additives as potent antioxidants (Ma et al. 2008). In the present study, we explored the usefulness of dietary pretreatment with LC

(0.5, 1.0, and 2.0 g kg⁻¹ of fish over a 4-week period) as prophylaxis for MCLR-induced oxidative stress in common carp for the first time.

Increasing evidences suggest that oxidative stress as a result of excessive reactive oxygen species (ROS) production may play an important role in the toxic mechanism of MCs (Li et al. 2003; Amado and Monserrat 2010; Chen et al. 2012). Under normal physiological situations, there is a balance between ROS production and the antioxidant defense system. Cellular oxidative stress occurs when the physiological antioxidant protection does not counteract the elevated ROS levels (Prieto et al. 2008; Jiang et al. 2012), and the




Fig. 1 Bactericidal activity (U ml⁻¹) of common carp fed with LC-containing diet after MCLR stress. Values followed by the different letters are significantly different ($P < 0.05$). Fish fed with basic diets and challenged with no MCLR (Group I). Fish fed with basic diets and challenged with 150 µg kg⁻¹ BW MCLR (Group

II). Fish fed with 0.5 g kg⁻¹ LC and challenged with 150 µg kg⁻¹ BW MCLR (Group III). Fish fed with 1.0 g kg⁻¹ and challenged with 150 µg kg⁻¹ BW MCLR (Group IV). Fish fed with 2.0 g kg⁻¹ LC and challenged with 150 µg kg⁻¹ BW MCLR (Group V)




Fig. 2 Lysozyme activity ($U\ ml^{-1}$) of common carp fed with LC-containing diet after MCLR stress. Values followed by the different letters are significantly different ($P < 0.05$). Fish fed with basic diets and challenged with no MCLR (*Group I*). Fish fed with basic diets and challenged with $150\ \mu g\ kg^{-1}$ BW MCLR (*Group II*).

Fish fed with $0.5\ g\ kg^{-1}$ LC and challenged with $150\ \mu g\ kg^{-1}$ BW MCLR (*Group III*). Fish fed with $1.0\ g\ kg^{-1}$ and challenged with $150\ \mu g\ kg^{-1}$ BW MCLR (*Group IV*). Fish fed with $2.0\ g\ kg^{-1}$ LC and challenged with $150\ \mu g\ kg^{-1}$ BW MCLR (*Group V*)

antioxidant system can counteract the ROS and reduce the oxidative stress with the antioxidant enzymes (SOD, CAT, and GPx). SOD catalyzes the conversion of superoxide to hydrogen peroxide, while CAT or GPx reduces hydrogen peroxide to H_2O (Cadenas 1989).

The results of our study indicate that MCLR induces oxidative stress in live common carp and altered activity of antioxidant enzymes. Concretely, the activity of SOD, GPx, and GST was increased after MCLR exposure, possibly indicating their scavenging activity




Fig. 3 Complement C3 of common carp fed with LC-containing diet after MCLR stress. Values followed by the different letters are significantly different ($P < 0.05$). Fish fed with basic diets and challenged with no MCLR (*Group I*). Fish fed with basic diets and challenged with $150\ \mu g\ kg^{-1}$ BW MCLR (*Group II*). Fish fed with

$0.5\ g\ kg^{-1}$ LC and challenged with $150\ \mu g\ kg^{-1}$ BW MCLR (*Group III*). Fish fed with $1.0\ g\ kg^{-1}$ and challenged with $150\ \mu g\ kg^{-1}$ BW MCLR (*Group IV*). Fish fed with $2.0\ g\ kg^{-1}$ LC and challenged with $150\ \mu g\ kg^{-1}$ BW MCLR (*Group V*)




Fig. 4 The relative expression of IFN I in the livers of common carp fed with LC-containing diet after MCLR stress. Values followed by the different letters are significantly different ($P < 0.05$). Fish fed with basic diets and challenged with no MCLR (Group I). Fish fed with basic diets and challenged with

150 $\mu\text{g kg}^{-1}$ BW MCLR (Group II). Fish fed with 0.5 g kg^{-1} LC and challenged with 150 $\mu\text{g kg}^{-1}$ BW MCLR (Group III). Fish fed with 1.0 g kg^{-1} and challenged with 150 $\mu\text{g kg}^{-1}$ BW MCLR (Group IV). Fish fed with 2.0 g kg^{-1} LC and challenged with 150 $\mu\text{g kg}^{-1}$ BW MCLR (Group V)

against ROS. In line with our study, Li et al. (2003) also reported an enhancement of SOD activity in hepatocytes of common carp (*Cyprinus carpio* L.) exposed to MCLR. Prieto et al. (2006) showed that i.p.-administered pure MCLR (500 $\mu\text{g kg}^{-1}$) induced a significant increase in the activity of CAT, SOD, and glutathione reductase (GR) in the liver of tilapia fish. In the case of

oral uptake route, a time-dependent increase in the activity of CAT, SOD, and GPx was also observed in the liver of tilapia fed with crush lyophilized cyanobacterial cells (approximately 60.0 $\mu\text{g MCLR fish}^{-1} \text{ day}^{-1}$) for 21 days (Jos et al. 2005). On the contrary, Liu et al. (2014) found that activities of SOD, CAT, and GPx drastically decreased in parental MCLR-treated groups




Fig. 5 The relative expression of IL-1 β in the livers of common carp fed with LC-containing diet after MCLR stress. Values followed by the different letters are significantly different ($P < 0.05$). Fish fed with basic diets and challenged with no MCLR (Group I). Fish fed with basic diets and challenged with

150 $\mu\text{g kg}^{-1}$ BW MCLR (Group II). Fish fed with 0.5 g kg^{-1} LC and challenged with 150 $\mu\text{g kg}^{-1}$ BW MCLR (Group III). Fish fed with 1.0 g kg^{-1} and challenged with 150 $\mu\text{g kg}^{-1}$ BW MCLR (Group IV). Fish fed with 2.0 g kg^{-1} LC and challenged with 150 $\mu\text{g kg}^{-1}$ BW MCLR (Group V)




Fig. 6 The relative expression of TNF- α in the liver of common carp fed with LC-containing diet after MCLR stress. Values followed by the different letters are significantly different ($P < 0.05$). Fish fed with basic diets and challenged with no MCLR (Group I). Fish fed with basic diets and challenged with

150 $\mu\text{g kg}^{-1}$ BW MCLR (Group II). Fish fed with 0.5 g kg^{-1} LC and challenged with 150 $\mu\text{g kg}^{-1}$ BW MCLR (Group III). Fish fed with 1.0 g kg^{-1} and challenged with 150 $\mu\text{g kg}^{-1}$ BW MCLR (Group IV). Fish fed with 2.0 g kg^{-1} LC and challenged with 150 $\mu\text{g kg}^{-1}$ BW MCLR (Group V)

compared with the control. The activities of GPx, GR, SOD, and CAT enzymes decreased after fish were orally exposed to a single dose of cyanobacterial cells containing 120 g per fish MCLR (Prieto et al. 2007). Under balneation conditions, Pavagadhi et al. (2012) found

that enzyme activities including GST, GPx, and SOD increased at lower concentrations ($\leq 5.0 \mu\text{g l}^{-1}$) and decreased at higher concentrations ($\geq 5.0 \mu\text{g l}^{-1}$) in the liver of adult zebrafish after MCLR exposure. The higher MC dose may damage the enzyme proteins,




Fig. 7 The relative expression of HSP70 in the liver of common carp fed with LC-containing diet after MCLR stress. Values followed by the different letters are significantly different ($P < 0.05$). Fish fed with basic diets and challenged with no MCLR (Group I). Fish fed with basic diets and challenged with

150 $\mu\text{g kg}^{-1}$ BW MCLR (Group II). Fish fed with 0.5 g kg^{-1} LC and challenged with 150 $\mu\text{g kg}^{-1}$ BW MCLR (Group III). Fish fed with 1.0 g kg^{-1} and challenged with 150 $\mu\text{g kg}^{-1}$ BW MCLR (Group IV). Fish fed with 2.0 g kg^{-1} LC and challenged with 150 $\mu\text{g kg}^{-1}$ BW MCLR (Group V)




Fig. 8 The relative expression of HSP90 in the liver of common carp fed with LC-containing diet after MCLR stress. Values followed by the different letters are significantly different ($P < 0.05$). Fish fed with basic diets and challenged with no MCLR (Group I). Fish fed with basic diets and challenged with

150 $\mu\text{g kg}^{-1}$ BW MCLR (Group II). Fish fed with 0.5 g kg^{-1} LC and challenged with 150 $\mu\text{g kg}^{-1}$ BW MCLR (Group III). Fish fed with 1.0 g kg^{-1} and challenged with 150 $\mu\text{g kg}^{-1}$ BW MCLR (Group IV). Fish fed with 2.0 g kg^{-1} LC and challenged with 150 $\mu\text{g kg}^{-1}$ BW MCLR (Group V)

while the lower dose, given for a longer period of time, may induce a defensive response. The antioxidant response to MC exposure varied largely and mainly depended upon the dose of the toxins and the exposure route (Malbrouck and Kestemont 2006).

GSH is one of the major antioxidant proteins, protecting the cell against the effects of reactive oxygen species (Nanda et al. 1996; Shila et al. 2005). It is also an essential protein in maintaining other antioxidant proteins, since it is responsible for preserving the reductive nature of the cell and regulating the binding of xenobiotics with cellular thiol. It is present in higher concentrations in metabolizing organs, such as the liver (Shila et al. 2005). In the present study, GSH level was drastically increased in MCLR-treated groups compared with the control. Lipid peroxidation is particularly important for aquatic animals since they normally contain greater amounts of highly unsaturated fatty acids (HUFAs) than other species (Huang et al. 2003). In the present study, the increase of LPO, as assessed by the formation of MDA, found in the liver of common carp treated with MCLR, suggests oxidative stress during MCLR intoxication. LC pretreated fish showed significantly decreased levels as compared to MCLR. This observation directly demonstrates the antiperoxidative and antioxidant effects of LC. Similarly, Ma et al. (2008) reported that dietary L-carnitine decreased the iron-

induced lipid peroxidation in liposomes through formation of free iron complexes. This advised that dietary L-carnitine lowered lipid peroxidation and improved resistance to oxidative stress in crayfish.

Some studies reported that the toxicity of MCLR was related to the stimulation of immune system (Li et al. 2012; Rymuszka 2013; Qiao et al. 2013). Recently, some cytokines have been studied to explore the immunomodulatory effects of MCs. In the present study, after the fish were fed with LC for 4 weeks, serum complement C3, lysozyme, and bactericidal activity were significantly ($P < 0.05$) increased after treated with MCLR as compared to the control group (group II). It indicated that LC enhances the protective effect against MCLR-induced immune response. TNF- α is considered to be an important component in innate immunity and inflammatory responses in fish (Rosa et al. 2008) and essential for inflammatory response to pathogenic germs or toxicants (Liew 2003). IFN 1 provides an important first line of defense against toxicants (McBeath et al. 2007). Some evidences demonstrated that MCLR affected the transcription of TNF- α and IFN I in fish (Wei et al. 2009; Rymuszka and Adaszek 2012). In the present study, after the fish were fed with LC for 4 weeks,

a strong downregulation in IL-1 β , TNF- α , and IFN I expression was observed as compared to the control group after treated with MCLR (group II). The downregulated genes might have controlled the inflammatory response of the stimulated pro-inflammatory cytokines, thereby minimizing damage to the host due to an excessive response (Raida et al. 2008)

HSPs are another protection system to protect the organisms from oxidative stress by preventing the irreversible loss of vital proteins and facilitating their subsequent regeneration (Jiang et al. 2012). When organisms are exposed to a variety of stress factors such as cold, heat, CO₂, heavy metal, and various chemicals (Hoffmann and Parsons 1991; Ferrando et al. 1995), they synthesize a set of HSPs, which usually act as molecular chaperones, and play diverse roles in transporting, folding, and assembling of degraded or misfolded proteins (Johnston et al. 1998; Sørensen et al. 2003). Heat shock proteins, particularly HSP70, have been proposed as biochemical markers of environmental stress. The HSP induction may rely on perturbation of the cellular redox status (Rai et al. 2005). In the present study, HSP70 and HSP90 gene expressions in the serum of common carp were significantly higher as compared to control group after treated with MCLR. The dramatically increased transcription of HSP90 and HSP70 may indicate their important roles as molecular chaperones under oxidative stress caused by MCLR, and the expression level recovered to original level at 96 h. In agreement with the results of the present study, HSP70 gene expression in serum was significantly ($P < 0.05$) downregulated in the treatment group fed with Immunogen® in rainbow trout as compared with control group. Downregulation of HSP70 expression is possibly due to elevated tolerance toward usual stresses during culture-like stresses caused by sampling for monitoring water quality, fish biometry, or other unwanted stresses (Wang et al. 2011).

In conclusion, dietary supplementation with LC induced expression of several antioxidant enzyme-related genes, stimulated PO, CAT, SOD, AST, and ALT activity, and improved the survival rate against ammonia stress. The results obtained indicated that LC has a protective effect against ammonia-induced oxidative stress in common carp.

Acknowledgments The research was supported by the National Natural Science Foundation of China (31372540) and Foundation projects in Jilin Province (20140303049ZY).

References

- Amado LL, Monserrat JM (2010) Oxidative stress generation by microcystins in aquatic animals: why and how. *Environ. Int.* 36:226–235
- Barnes AC, Young FM, Horne MT, Ellis AE (2003) Streptococcus iniae: serological differences, presence of capsule and resistance to immune serum killing. *Dis Aquat Org* 53:241–247
- Cadenas E (1989) Biochemistry of oxygen toxicity. *Annu Rev Biochem* 58:79–110
- Chen XM, Ru HM, Niu XT, Wang GQ, Zhang DM (2015) Enhancement of secondary metabolites from *Bacillus Licheniformis* XY-52 on immune response and expression of some immune-related genes in common carp, *Cyprinus carpio*. *Fish Shellfish Immunol* 45(1):124–131
- Campos A, Vasconcelos V (2010) Molecular mechanisms of microcystin toxicity in animal cells. *Int J Mol Sci* 11:268–287
- Chen J, Zhang D, Xie P, Wang Q, Ma Z (2009) Simultaneous determination of microcystin contaminations in various vertebrates (fish, turtle, duck and water bird) from a large eutrophic Chinese lake, Lake Taihu, with toxic Microcystis blooms. *Sci Total Environ* 407:3317–3322
- Chen Y, Zeng SF, Cao YF (2012) Oxidative stress response in zebrafish (*Danio rerio*) gill experimentally exposed to sub-chronic microcystin-LR. *Environ Monit Assess* 184:6775–6787
- Davis TM, Koch F, Marcoval MA, Wilhelm SW, Gobler CJ (2012) Mesozooplankton and microzooplankton grazing during cyanobacterial blooms in the western basin of Lake Erie. *Harmful Algae* 15:26–35
- de Figueiredo DR, Azeiteiro UM, Esteves SM, Goncalves FJM, Pereira JM (2004) Microcystin-producing blooms—a serious global public health issue. *Ecotox Environ Saf* 59(2):151–163
- Esterbauer H, Cheeseman KH (1990) Determination of aldehydic lipid peroxidation products: malonaldehyde and 4-hydroxynonenal. *Methods Enzymol* 186:407–421
- Ellis AE, Stolen JS, Fletcher TC, Anderson DP, Roberson BS, van Muiswinkel WB (1990) Techniques in fish immunology. SOS Publications, NJ, pp 101–103
- Ferrando RE, Schuschereba ST, Quong JA, Bowman PD (1995) Carbon dioxide laser induction of heat shock protein 70 synthesis: comparison with high temperature treatment. *Lasers Med Sci* 10:207–212
- Graham JL, Loftin KA, Meyer MT, Ziegler AC (2010) Cyanotoxin mixtures and taste-and-odor compounds in cyanobacterial blooms from the Midwestern United States. *Comp Biochem Physiol B: Environ Sci Technol* 44:7361–7368
- Harpaz S (2005) L-Carnitine and its attributed functions in fish culture and nutrition—a review. *Aquacult* 249:3–21
- Harpaz S, Becker K, Blum R (1999) The effect of dietary l-carnitine supplementation on cold tolerance and growth of

- ornamental cichlid fish (*Pelvicachromis pulcher*)—preliminary results. *J Thermal Biology* 24:57–62
- Honkanen RE, Zwiller JEMR, Moore RE, Daily SL, Khatra BS, Dukelow M, Boynton AL (1990) Characterization of microcystin-LR, a potent inhibitor of type 1 and type 2A protein phosphatases. *J Biol Chem* 265:19401–19404
- Hoffmann AA, Parsons PA (1991) Evolutionary genetics and environmental stress. Oxford University Press, New York
- Huang CH, Chang RJ, Huang SL, Chen W (2003) Dietary vitamin E supplementation affects tissue lipid peroxidation of hybrid tilapia. *Oreochromis niloticus* × *O. aureus*. *Comp Biochem Physiol B* 134:265–270
- Jiang JL, Shi Y, Shan ZJ, Yang LY, Wang XR, Shi LL (2012) Bioaccumulation, oxidative stress and HSP70 expression in *Cyprinus carpio* L. exposed to microcystin-LR under laboratory conditions. *Comp Biochem Phys C* 155:483–490
- Johnston JA, Ward CL, Kopito RR (1998) Aggresomes: a cellular response to misfolded proteins. *J Cell Biol* 143:1883–1898
- Jos Á, Pichardo S, Prieto AI, Repetto G, Vázquez CM, Moreno I (2005) Toxic cyanobacterial cells containing microcystins induce oxidative stress in exposed tilapia fish (*Oreochromis* sp.) under laboratory conditions. *Aquat Toxicol* 72:261–271
- Lawrence A, Burk RF (1976) Glutathione peroxidase activity in selenium deficient rat liver. *Biochem Biophys Res Commun* 71:952–958
- Li G, Yan W, Qiao Q, Chen J, Cai F, He Y, Zhang X (2012) Global effects of subchronic treatment of microcystin-LR on rat splenic protein levels. *J Proteome* 77:383–393
- Li L, Xie P (2009) Hepatic histopathological characteristics and antioxidant response of phytoplanktivorous silver carp intraperitoneally injected with extracted microcystins. *Biomed Environ Sci* 22:297–302
- Li L, Xie P, Lei H, Zhang X (2013) Renal accumulation and effects of intraperitoneal injection of extracted microcystins in omnivorous crucian carp (*Carassius auratus*). *Toxicol* 49:1150–1157
- Li X, Liu Y, Song L, Liu J (2003) Responses of antioxidant systems in the hepatocytes of common carp (*Cyprinus carpio* L.) to the toxicity of microcystin-LR. *Toxicol* 42:85–89
- Liew FY (2003) The role of innate cytokines in inflammatory response. *Immunol Lett* 85:131–134
- Liu WJ, Qiao Q, Chen YY, Wu K, Zhang XZ (2014) Microcystin-LR exposure to adult zebrafish (*Danio rerio*) leads to growth inhibition and immune dysfunction in F1 offspring, a parental transmission effect of toxicity. *Aquatic Toxicol* 155:360–367
- Ma JJ, Xu ZRQ, Shao JJ, Xu Z, Hung SSO, Hu WL, Zhou LY (2008) Effect of dietary supplemental L-carnitine on growth performance, body composition and antioxidant status in juvenile black sea bream, *Sparus macrocephalus*. *Aquac Nutr* 14:464–471
- Malbrouck C, Trausch G, Devos P, Kestemont P (2003) Hepatic accumulation and effects of microcystin-LR on juvenile goldfish *Carassius auratus* L. *Comp. Biochem. Physiol. C: Toxicol. Pharmacol.* 135:39–48
- Malbrouck C, Kestemont P (2006) Effects of microcystins on fish. *Environ Toxicol Chem* 25:72–86
- McBeath A, Snow M, Secombes C, Ellis A, Collet B (2007) Expression kinetics of interferon and interferon-induced genes in Atlantic salmon (*Salmo salar*) following infection with infectious pancreatic necrosis virus and infectious salmon anaemia virus. *Fish. Shellfish. Immun.* 22:230–241
- Nanda D, Tolputt J, Collard KJ (1996) Changes in brain glutathione levels during postnatal development in the rat. *Dev Brain Res* 94:238–241
- Organization, W.H 2004. Guidelines for Drinking Water Quality, vol. 1 Recommendations. World Health Organization, Geneva
- Palus J, Dziubałtowska E, Stanczyk M, Lewinska D, Mankiewicz-Boczek J, Izydorczyk K, Bonisławska A, Jurczak T, Zalewski M, Wasowicz W (2007) Biomonitoring of cyanobacterial blooms in polish water reservoir and the cytotoxicity and genotoxicity of selected cyanobacterial extracts. *Int J Occup Med Environ* 20:48–65
- Pavagadhi S, Gong Z, Hande MP, Dionysiou DD, de la Cruz AA, Balasubramanian R (2012) Biochemical response of diverse organs in adult *Danio rerio* (zebrafish) exposed to sub-lethal concentrations of microcystin-LR and microcystin-RR: a balneation study. *Aquat Toxicol* 109:1–10
- Pinho GLL, Moura da Rosa C, Yunes JS, Luquet CM, Bianchini A, Monserrat JM (2003) Toxic effects of microcystins in the hepatopancreas of the estuarine crab *Chasmagnathus granulatus* (*Decapoda, Grapsidae*). *Comp Biochem Physiol C: Toxicol Pharmacol* 135:459–468
- Prieto AI, Jos Á, Pichardo S, Moreno I, Cameán AM (2006) Differential oxidative stress responses to microcystins LR and RR in intraperitoneally exposed tilapia fish (*Oreochromis* sp.). *Aquat Toxicol* 77:314–321
- Prieto AI, Pichardo S, Jos Á, Moreno I, Cameán AM (2007) Time-dependent oxidative stress responses after acute exposure to toxic cyanobacterial cells containing microcystins in tilapia fish (*Oreochromis niloticus*) under laboratory conditions. *Aquat Toxicol* 84:337–345
- Prieto AI, Jos Á, Pichardo S, Moreno I, Cameán AM (2008) Protective role of vitamin E on the microcystin-induced oxidative stress in tilapia fish (*Oreochromis niloticus*). *Environ Toxicol Chem* 27:1152–1159
- Qiu T, Xie P, Liu Y, Li G, Xiong Q, Hao L, Li H (2009) The profound effects of microcystin on cardiac antioxidant enzymes, mitochondrial function and cardiac toxicity in rat. *Toxicology* 257:86–94
- Qiao Q, Liang H, Zhang X (2013) Effect of cyanobacteria on immune function of crucian carp (*Carassius auratus*) via chronic exposure in diet. *Chemosphere* 90:1167–1176
- Raida MK, Buchmann K (2008) Development of adaptive immunity in rainbow trout, *Oncorhynchus mykiss* (Walbaum) surviving an infection with *Yersinia ruckeri*. *Fish Shellfish Immunol* 25:533–541
- Rai R, Richardson C, Flecknell P, Robertson H, Burt A, Manas DM (2005) Study of apoptosis and heat shock protein (HSP) expression in hepatocytes following radiofrequency ablation (RFA). *J Surg Res* 129:147–151
- Rosa CE, Figueiredo MA, Lanes CFC, Almeida DV, Monserrat JM, Marins L (2008) Metabolic rate and reactive oxygen species production in different genotypes of GH-transgenic zebrafish. *Comp Biochem Phys B* 149:209–214
- Rymuszka A, Adaszek Ł (2012) Pro- and anti-inflammatory cytokine expression in carp blood and head kidney leukocytes exposed to cyanotoxin stress—an in vitro study. *Fish Shellfish Immun* 33:382–388
- Rymuszka A (2013) Microcystin-LR induces cytotoxicity and affects carp immune cells by impairment of their

- phagocytosis and the organization of the cytoskeleton. *J Appl Toxicol* 33:1294–1302
- Safari O, Atash MMS, Paolucci M (2015) Effects of dietary L-carnitine level on growth performance, immune responses and stress resistance of juvenile narrow clawed crayfish, *Astacus leptodactylus leptodactylus* Eschscholtz, 1823. *Aquaculture* 439:20–28
- Saliny, G. 1994. Effect of L-carnitine on growth responses of the fresh water prawn, *Macrobrachium idella idella* (Hilgendorf). University of Kerala, India. PhD thesis (213 pp)
- Schreiber S, Becker K, Bresler V, Fishelson L (1997) Dietary l-carnitine protects the gills and skin of guppies (*Poecilia reticulata*) against anionic xenobiotics. *Comp Biochem Physiol* 117C:99–102
- Sørensen JG, Kristensen TN, Loeschcke V (2003) The evolutionary and ecological role of heat shock proteins. *Ecol Lett* 6: 1025–1037
- Shila S, Subathra M, Devi MA, Panneerselvam C (2005) Arsenic intoxication induced reduction of glutathione level and of the activity of related enzymes in rat brain regions: reversal by DL-alpha-lipoic acid. *Arch Toxicol* 79:140–146
- Trinchet I, Djediat C, Huet H, Dao SP, Edery M (2011) Pathological modifications following sub-chronic exposure of medaka fish, *Oryzias latipes* to microcystin-LR. *Reprod Toxicol* 32:329–340
- Wang GX, Wang Y, Wu ZF, Jiang HF, Dong RQ, Li FY et al (2011) Immunomodulatory effects of secondary metabolites from the rmophilic *Anoxybacillus kamchatkensis* XA-1 on carp, *Cyprinus carpio*. *Fish Shellfish Immunol* 30:1331–1338
- Wei L, Sun B, Chang M, Liu Y, Nie P (2009) Effects of cyanobacterial toxin microcystin-LR on the transcription levels of immune-related genes in grass carp *Ctenopharyngodon idella*. *Environ Biol Fish* 85:231–238
- Zhang DW, Xie P, Liu YQ, Qiu T (2009) Transfer, distribution and bioaccumulation of microcystins in the aquatic food web in Lake Taihu, China, with potential risks to human health. *Sci Total Environ* 407:2191–2199