

Super Quantum Discord for X-type States

Tao Li · Teng Ma · Yaokun Wang · Shaoming Fei ·
Zhixi Wang

Received: 16 February 2014 / Accepted: 7 July 2014 / Published online: 24 July 2014
© Springer Science+Business Media New York 2014

Abstract Weak measurement is a new way to manipulate and control quantum systems. Different from projection measurement, weak measurement only makes a small change in status. Applying weak measurement to quantum discord, Singh and Pati proposed a new kind of quantum correlations called “super quantum discord (SQD)” [Ann. Phys. **343**,141(2014)]. Unfortunately, the super quantum discord is also difficult to calculate. There are only few explicit formulae about SQD. We derive an analytical formula of SQD for general X-type two-qubit states, which surpass the conclusion for Werner states and Bell diagonal states. Furthermore, our results reveal more knowledge about the new insight of quantum correlation and give a new way to compare SQD with normal quantum discord. Finally, we analyze its dynamics under nondissipative channels.

Keywords Weak measurement · Super quantum discord · Quantum correlations · X-type state

1 Introduction

The quantum entanglement plays important roles in quantum information processing [1]. However, besides quantum entanglement there are other quantum correlations also useful for quantum information processing. It is found that many tasks can be carried out with quantum correlations other than entanglement [2–4]. In particular, the quantum discord [5–21] plays an important role in some quantum information processing like to assist optimal

T. Li · T. Ma · Y. Wang · S. Fei · Z. Wang (✉)
School of Mathematical Sciences, Capital Normal University, Beijing 100048, China
e-mail: zhixiwang@gmail.com

Y. Wang
Department of Mathematics, Tonghua Normal University, Tonghua 134001, China

S. Fei
Max-Planck-Institute for Mathematics in the Sciences, 04103 Leipzig, Germany

state discrimination, in which only one side discord is required in the optimization process of assisted state discrimination, while another side discord and entanglement is not necessary [22, 23].

Unfortunately, underlying quantum measurements process quantum states are fragile. When we measure a quantum state in some orthogonal basis, since quantum discord is defined by the POVM quantum measurement, the coherence of the state has been loosened. Taking account of quantum states’ potential privacy, it is reasonable to find a solution to deal with this problem. Such a solution was making use of weak measurement which induced by Aharonov-Albert-Vaidman[24]. Applying such a scheme, we can replace the POVM measurement by weak measurement in the definition of quantum discord, which gives rise to so called super quantum discord (SQD) [25].

Super quantum discord sheds a new insight on the nature of quantum correlation. It also has vivid properties, such as the monotony. Super quantum discord not only a new insight in fundamental physics but also useful in applications. But super quantum discord is difficult to calculate. There are only few explicit formulae about SQD. The analytical formulae of Werner states [25] and Bell diagonal states [26] are only two results. In order to obtain more useful results, we take a new method to compute more general states—X-type states, which including Werner states and Bell diagonal states. Obviously, our result include the results in [25] and [26]. Furthermore, in order to show the potential property of technological implications, we analyze the dynamics of two-qubit X-states under non-dissipative channels and compare super quantum discord with discord by using the explicit formulae. This is exemplified by the fact that the super quantum discord often larger than the quantum discord defined by projective measurement. Also, the super quantum discord sometime captures more quantum correlations.

This paper is organized as follows. In Section 2, we derive analytical formulae of super quantum discord for X-states. In Section 3, we compute the super quantum discord of some concrete examples, and analyze their dynamics under nondissipative channels, we further compare it with discord and present some new property of super quantum discord.

2 Super Quantum Discord for Two-qubit X-states

Super quantum discord of some special states has been computed recently, the Werner states and the Bell diagonal states are included. Now, we extend the results about super quantum discord in [25] and [26] to the whole two-qubit X-states. Let us consider a two-qubit X-state:

$$\rho_X = \begin{pmatrix} a_{11} & 0 & 0 & a_{14} \\ 0 & a_{22} & a_{23} & 0 \\ 0 & a_{23}^* & a_{33} & 0 \\ a_{14}^* & 0 & 0 & a_{44} \end{pmatrix}, \tag{1}$$

where $\sum_{i=1}^4 a_{ii} = 1, |a_{23}^2| \leq a_{22}a_{33}, |a_{14}^2| \leq a_{11}a_{44}$. The density matrix ρ_X can be written as [27]:

$$\rho_X = \frac{1}{4} \begin{pmatrix} 1 + d_1 & 0 & 0 & c_1 - c_2 \\ 0 & 1 + d_2 & c_1 + c_2 & 0 \\ 0 & c_1^* + c_2^* & 1 + d_3 & 0 \\ c_1^* - c_2^* & 0 & 0 & 1 + d_4 \end{pmatrix}, \tag{2}$$

where c_1 and c_2 are complex, d_1, d_2, d_3 and d_4 are real, $d_1 = c_3 + a_3 + b_3, d_2 = -c_3 + a_3 - b_3, d_3 = -c_3 - a_3 + b_3, d_4 = c_3 - a_3 - b_3$. These parameters are determined by the entries of the density matrix, $a_3 = a_{11} - a_{44} + a_{22} - a_{33}, b_3 = a_{11} - a_{44} - a_{22} + a_{33}, c_3 = a_{11} + a_{44} - a_{22} - a_{33}, c_1 = 2(a_{23} + a_{14}), c_2 = 2(a_{23} - a_{14})$.

Let $\{\Pi_i^B\}, i = 0, 1$, be the projective measurements. The discord of a bipartite quantum state ρ_{AB} with the measurement $\{\Pi_i^B\}$ on the subsystem B is the dissimilarity between the mutual information $I(\rho_{AB})$ [28] and the classical correlation $J_B(\rho_{AB})$ [29]:

$$D(\rho_{AB}) = \min_{\{\Pi_i^B\}} \sum_i p_i S(\rho_{A|i}) + S(\rho_B) - S(\rho_{AB}), \tag{3}$$

where the minimization goes over all projective measurements $\{\Pi_i^B\}, S(\rho) = -\text{tr}(\rho \log_2 \rho)$ is the von Neumann entropy of a quantum state ρ, ρ_B is the reduced density matrices of ρ_{AB} and

$$p_i = \text{tr}_{AB} \left[(I_A \otimes \Pi_i^B) \rho_{AB} (I_A \otimes \Pi_i^B) \right], \tag{4}$$

$$\rho_{A|i} = \frac{1}{p_i} \text{tr}_B \left[(I_A \otimes \Pi_i^B) \rho_{AB} (I_A \otimes \Pi_i^B) \right]. \tag{5}$$

The weak measurement operators are given by [30],

$$P(+x) = \sqrt{\frac{1 - \tanh x}{2}} \Pi_0 + \sqrt{\frac{1 + \tanh x}{2}} \Pi_1, \tag{6}$$

$$P(-x) = \sqrt{\frac{1 + \tanh x}{2}} \Pi_0 + \sqrt{\frac{1 - \tanh x}{2}} \Pi_1, \tag{7}$$

where Π_0 and Π_1 are two orthogonal projectors satisfying $\Pi_0 + \Pi_1 = I, x$ is the strength parameter of measurement. The weak measurement operators satisfy: (i) $P^\dagger(+x)P(+x) + P^\dagger(-x)P(-x) = I$, (ii) $\lim_{x \rightarrow \infty} P(+x) = \Pi_0$ and $\lim_{x \rightarrow \infty} P(-x) = \Pi_1$.

The super quantum discord is defined by [25]:

$$D_w(\rho_{AB}) = \min_{\{\Pi_i^B\}} S_w \left(A|\{P^B(x)\} \right) + S(\rho_B) - S(\rho_{AB}),$$

where

$$S_w \left(A|\{P^B(x)\} \right) = p(+x)S(\rho_{A|P^B(+x)}) + p(-x)S(\rho_{A|P^B(-x)}), \tag{8}$$

$$p(\pm x) = \text{tr}_{AB} \left[\left(I_A \otimes P^B(\pm x) \right) \rho_{AB} \left(I_A \otimes P^B(\pm x) \right) \right],$$

$$\rho_{A|P^B(\pm x)} = \frac{\text{tr}_B \left[\left(I_A \otimes P^B(\pm x) \right) \rho_{AB} \left(I_A \otimes P^B(\pm x) \right) \right]}{\text{tr}_{AB} \left[\left(I_A \otimes P^B(\pm x) \right) \rho_{AB} \left(I_A \otimes P^B(\pm x) \right) \right]},$$

where $\{P^B(x)\}$ is the weak measurement operators on subsystem B .

The weak measurement operators on subsystem B can be expressed as

$$I_A \otimes P^B(\pm x) = \sqrt{\frac{1 \mp \tanh x}{2}} I \otimes V \Pi_0 V^\dagger + \sqrt{\frac{1 \pm \tanh x}{2}} I \otimes V \Pi_1 V^\dagger, \tag{9}$$

where $\Pi_k = |k\rangle\langle k|, k = 0, 1, |k\rangle$ is the computational base, and V is a 2×2 unitary transformation. V can be generally expressed as [15]:

$$V = tI + iy \cdot \sigma, \tag{10}$$

where $\mathbf{y} = (y_1, y_2, y_3)$ and $t, y_1, y_2, y_3 \in \mathbb{R}^1, t^2 + y_1^2 + y_2^2 + y_3^2 = 1$.

Table 1 The minimum value of $S(\rho_{A|P^B(+x)})$ and $p(+x)$

(z_3, z_2, z_1)	$p(+x)$	λ_{\pm}
(1,0,0)	$\frac{1-b_3 \tanh x}{2}$	$\frac{1}{2} \left[1 \pm \frac{a_3 - c_3 \tanh x}{1 - b_3 \tanh x} \right]$
(0,0,1)	$\frac{1}{2}$	$\frac{1}{2} [1 \pm a_3 \cdot \tanh x]$
(0,1,0)	$\frac{1}{2}$	$\frac{1}{2} \left[1 \pm \sqrt{a_3^2 + (c_2 ^2 - c_1 - c_2 ^2 - c_1 + c_2 ^2 + (\Re(c_1 - c_2) + \Im(c_1 + c_2))^2 + (\Re(c_1 + c_2) + \Im(c_1 - c_2))^2) \tanh^2 x} \right]$
(0,-1,0)	$\frac{1}{2}$	$\frac{1}{2} \left[1 \pm \sqrt{a_3^2 + (c_2 ^2 - c_1 - c_2 ^2 - c_1 + c_2 ^2 + (\Re(c_1 - c_2) - \Im(c_1 + c_2))^2 + (\Re(c_1 + c_2) - \Im(c_1 - c_2))^2) \tanh^2 x} \right]$

To evaluate the super quantum discord of ρ_X , let us first express ρ_X in terms of the bases $I \otimes I, \sigma_i \otimes \sigma_j, i, j = 0, 1, 2$ (Tables 1 and 2).

$$\rho_X = \frac{1}{4} \left(I + \sum_i \Re(c_i) \sigma_i \otimes \sigma_i \right) + \frac{1}{4} [(b_3 - a_3) I \otimes \sigma_3 + (\Im(c_1) + \Im(c_2) \sigma_1 \otimes \sigma_2)],$$

where $\Re(c_i), \Im(c_i)$ are the real and complex parts of c_i . By using the relations

$$V^\dagger \sigma_1 V = (t^2 + y_1^2 - y_2^2 - y_3^2) \sigma_1 + 2(ty_3 + y_1y_2) \sigma_2 + 2(-ty_2 + y_1y_3) \sigma_3,$$

$$V^\dagger \sigma_2 V = (t^2 + y_2^2 - y_1^2 - y_3^2) \sigma_2 + 2(ty_1 + y_2y_3) \sigma_3 + 2(-ty_3 + y_1y_2) \sigma_1,$$

$$V^\dagger \sigma_3 V = (t^2 + y_3^2 - y_1^2 - y_2^2) \sigma_3 + 2(ty_2 + y_1y_3) \sigma_1 + 2(-ty_1 + y_2y_3) \sigma_2,$$

$\Pi_0 \sigma_3 \Pi_0 = \Pi_0, \Pi_1 \sigma_3 \Pi_1 = -\Pi_1, \Pi_j \sigma_k \Pi_j = 0$ for $j = 0, 1, k = 1, 2$ in [15]. Setting $a_1 = z_1 \Re(c_1) + z_2 \Im(c_2), a_2 = z_2 \Re(c_2) - z_1 \Im(c_1)$, with $z_1 = 2(-ty_2 + y_1y_3), z_2 = 2(ty_1 + y_2y_3), z_3 = t^2 + y_3^2 - y_1^2 - y_2^2$, we have the ensemble $\{\rho_{A|P^B(\pm x)}, p(\pm x)\}$ after weak measurement, from (4) and (5)

$$p(+x) = \frac{1}{2} (1 - b_3 z_3 \tanh x), \tag{11a}$$

$$\rho_{A|P^B(+x)} = \frac{1}{2} \left[I + \frac{\tanh x (-a_1 \sigma_1 - a_2 \sigma_2) + (a_3 - c_3 z_3 \tanh x) \sigma_3}{1 - b_3 z_3 \tanh x} \right], \tag{11b}$$

$$p(-x) = \frac{1}{2} (1 + b_3 z_3 \tanh x), \tag{12a}$$

$$\rho_{A|P^B(-x)} = \frac{1}{2} \left[I + \frac{\tanh x (a_1 \sigma_1 + a_2 \sigma_2) + (a_3 + c_3 z_3 \tanh x) \sigma_3}{1 + b_3 z_3 \tanh x} \right]. \tag{12b}$$

Table 2 The minimum value of $S(\rho_{A|P^B(-x)})$ and $p(-x)$

(z_3, z_2, z_1)	$p(-x)$	λ'_{\pm}
(1,0,0)	$\frac{1+b_3 \tanh x}{2}$	$\frac{1}{2} \left[1 \pm \frac{a_3 + c_3 \tanh x}{1 + b_3 \tanh x} \right]$
(0,0,1)	$\frac{1}{2}$	$\frac{1}{2} [1 \pm a_3 \cdot \tanh x]$
(0,1,0)	$\frac{1}{2}$	$\frac{1}{2} \left[1 \pm \sqrt{a_3^2 + (c_2 ^2 - c_1 - c_2 ^2 - c_1 + c_2 ^2 + (\Re(c_1 - c_2) + \Im(c_1 + c_2))^2 + (\Re(c_1 + c_2) + \Im(c_1 - c_2))^2) \tanh^2 x} \right]$
(0,-1,0)	$\frac{1}{2}$	$\frac{1}{2} \left[1 \pm \sqrt{a_3^2 + (c_2 ^2 - c_1 - c_2 ^2 - c_1 + c_2 ^2 + (\Re(c_1 - c_2) - \Im(c_1 + c_2))^2 + (\Re(c_1 + c_2) - \Im(c_1 - c_2))^2) \tanh^2 x} \right]$

The eigenvalues of $\rho_{A|PB(+x)}$ and $\rho_{A|PB(-x)}$ are given by

$$\frac{1}{2} \left[1 \pm \frac{\sqrt{(a_3 - c_3 z_3 \tanh x)^2 + (a_2^2 + a_1^2) \tanh^2 x}}{1 - b_3 z_3 \tanh x} \right], \tag{13}$$

$$\frac{1}{2} \left[1 \pm \frac{\sqrt{(a_3 + c_3 z_3 \tanh x)^2 + (a_2^2 + a_1^2) \tanh^2 x}}{1 + b_3 z_3 \tanh x} \right]. \tag{14}$$

We now compute the minimum value of $S(\rho_{A|PB(+x)})$ and the corresponding $p(+x)$ by using the method of Hessian matrix and the symmetries in (9–13). In order to avoid redundant narrating, we only give the result in the following tables, and the minimum value lies at $z_3 = 0$ or $z_3 = 1$. The extremum lies at the following points:

Similarly for the minimum value of $S(\rho_{A|PB(-x)})$ and $p(-x)$, we have:

From Tables 1 and 2, for a given state ρ_X , one can get the minimum values of λ_{\pm} and λ'_{\pm} , which give rise to

$$S_w(\rho_{A|PB(+x)}) = -\lambda_+ \log_2 \lambda_+ - \lambda_- \log_2 \lambda_-, \tag{15}$$

$$S_w(\rho_{A|PB(-x)}) = -\lambda'_+ \log_2 \lambda'_+ - \lambda'_- \log_2 \lambda'_-, \tag{16}$$

and the super quantum discord

$$D_w(\rho_X) = p(+x)S_w(\rho_{A|PB(+x)}) + p(-x)S_w(\rho_{A|PB(-x)}) + S(\rho_X^B) - S(\rho_X). \tag{17}$$

3 Dynamics of Super Quantum Discord Under Nondissipative Channels

In this section, firstly we will verify our formulae with examples and illustrate that it is an extension of results in [22-23]. The first one is Werner state [31] which is known to be a special X-state,

$$\rho_W = \begin{pmatrix} \frac{1+z}{4} & 0 & 0 & \frac{z}{2} \\ 0 & \frac{1-z}{4} & 0 & 0 \\ 0 & 0 & \frac{1-z}{4} & 0 \\ \frac{z}{2} & 0 & 0 & \frac{1+z}{4} \end{pmatrix}.$$

Based on formulae of the previous section, we are able to calculate eigenvalues $\lambda_{\pm i} = \lambda'_{\pm i} = \frac{1 \pm z \tanh x}{2}$, $i = 1, 2, 3, 4$. As everyone knows, for the Werner state, all eigenvalues get the same results for any measurement basis. The eigenvalues of ρ_W^B are $\frac{1}{2}, \frac{1}{2}$, and the eigenvalues of ρ_W are $\frac{1+3z}{4}, \frac{1-z}{4}, \frac{1-z}{4}, \frac{1-z}{4}$. From (16) the super quantum discord of ρ_W is

$$D_w = -\frac{1-z \tanh x}{2} \log_2 \frac{1-z \tanh x}{2} - \frac{1+z \tanh x}{2} \log_2 \frac{1+z \tanh x}{2} + 1 + \frac{3(1-z)}{4} \log_2 \frac{1-z}{4} + \frac{1+3z}{4} \log_2 \frac{1+3z}{4},$$

which is in coincident with the result in [25].

As another example, we consider the Bell diagonal states [32]

$$\rho = \begin{pmatrix} \frac{1+c_3}{4} & 0 & 0 & \frac{c_1-c_2}{4} \\ 0 & \frac{1-c_3}{4} & \frac{c_1+c_2}{4} & 0 \\ 0 & \frac{c_1+c_2}{4} & \frac{1-c_3}{4} & 0 \\ \frac{c_1-c_2}{4} & 0 & 0 & \frac{1+c_3}{4} \end{pmatrix}.$$

From (10–13) we get $\lambda_{\pm 1} = \lambda'_{\pm 1} = \frac{1 \pm c_1 \tanh x}{2}$, $\lambda_{\pm 2} = \lambda'_{\pm 2} = \frac{1 \pm c_2 \tanh x}{2}$, $\lambda_{\pm 3} = \lambda'_{\pm 3} = \frac{1 \pm c_3 \tanh x}{2}$, $\lambda_{\pm 4} = \lambda'_{\pm 4} = \frac{1 \pm c_3 \tanh x}{2}$. It is also easy to calculate the eigenvalues of ρ^B are $\frac{1}{2}, \frac{1}{2}$ and the eigenvalues of ρ are $\frac{1-c_1-c_2-c_3}{4}, \frac{1-c_1+c_2+c_3}{4}, \frac{1+c_1-c_2+c_3}{4}, \frac{1+c_1+c_2-c_3}{4}$. Let $c = \max\{c_1, c_2, c_3\}$, by (16), we have the super quantum discord

$$\begin{aligned} D_w = & -\frac{1-c \tanh x}{2} \log_2 \frac{1-c \tanh x}{2} - \frac{1+c \tanh x}{2} \log_2 \frac{1+c \tanh x}{2} \\ & + 1 + \frac{1-c_1-c_2-c_3}{4} \log_2 \frac{1-c_1-c_2-c_3}{4} \frac{1-c_1+c_2+c_3}{4} \\ & + \frac{1+c_1-c_2+c_3}{4} \log_2 \frac{1+c_1-c_2+c_3}{4} \frac{1+c_1+c_2-c_3}{4} \log_2 \\ & + \frac{1+c_1+c_2-c_3}{4}, \end{aligned}$$

which coincides with the result in [26].

By above examples we illustrated how to apply the main result and recover the results in Refs.[22-23] as special cases.

Due to the fundamental significance and potential applications of super quantum discord, the evolution of super quantum discord under bit-flip noise which characterized by Kraus operators

$$E_0 = \sqrt{p} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad E_1 = \sqrt{1-p} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \tag{18}$$

has been considered. We have the channel “local bit-flip(Λ_{bfi})”:

$$\Lambda_{\text{bfi}}(\rho_X) = (I \otimes E_0)\rho_X(I \otimes E_0)^\dagger + (I \otimes E_1)\rho_X(I \otimes E_1)^\dagger. \tag{19}$$

Under this channel, the entries of the density matrix have the following transformations:

channel	a_{11}	a_{14}	a_{22}
local bit-flip	$a_{22} + pa_{11} - pa_{22}$	$a_{23} + pa_{14} - pa_{23}$	$a_{11} - pa_{11} + pa_{22}$
channel	a_{23}	a_{33}	a_{44}
local bit-flip	$a_{14} - pa_{14} + pa_{23}$	$a_{44} + pa_{33} - pa_{44}$	$a_{33} - pa_{33} + pa_{44}$
channel	a_{23}^*	a_{14}^*	
local bit-flip	$a_{14}^* - pa_{14}^* + pa_{23}^*$	$a_{23}^* + pa_{14}^* - pa_{23}^*$	

As an illustrative example, we choose a subfamily of X-types. Let us consider

$$\rho_X = \begin{pmatrix} 0.25 & 0 & 0 & 0.0625 \\ 0 & 0.25 & 0.125 & 0 \\ 0 & 0.125 & 0.25 & 0 \\ 0.0625 & 0 & 0 & 0.25 \end{pmatrix}. \tag{20}$$




Fig. 1 Super quantum discord (dashed line) and quantum discord (solid line) as function of x

Since quantum discord has also been employed in the study of quantum computation as an important resource, we prefer to compare super quantum discord and discord under noisy channels. It can be seen from Fig. 1 that the super quantum discord attains the maximum value at $x = 0$, where the weak measurement is the weakest. When $x \rightarrow \infty$, the super quantum discord approaches quantum discord.

Interestingly, the above relation motives us to introduce the super quantum discord and discord when the signal through noisy channels. From (14–16), we get $p(+x) = p(-x) = 0.5$, $\lambda_+ = \lambda'_+ = \max\{0.5, 0.5 + 0.1875 \cdot \tanh x, 0.5 + 0.0625 \cdot \tanh x\}$, $\lambda_- = \lambda'_- = \min\{0.5, 0.5 - 0.1875 \cdot \tanh x, 0.5 - 0.0625 \cdot \tanh x\}$. Due to the symmetry of $\tanh x$, we take $x > 0$. Namely, $p(+x) = p(-x) = 0.5$, $\lambda_+ = \lambda'_+ = 0.5 + 0.1875 \cdot \tanh x$, $\lambda_- = \lambda'_- = 0.5 - 0.1875 \cdot \tanh x$. Under the local bit-flip channel, we have $p(+x) = p(-x) = 0.5$, $\lambda_+ = \lambda'_+ = 0.5 + 0.1875 \cdot \tanh x$, $\lambda_- = \lambda'_- = 0.5 - 0.1875 \cdot \tanh x$.

On the left side of Fig. 2 we find that the super quantum discord which not through the bit-flip noise channel is larger than the super quantum discord which through the bit-flip noise channel. We can get more detail properties from the right side of Fig. 2. The five curves from top to bottom are obtained by choosing the controlling parameters as the weak measurement parameter $x = 0, 0.5, 1, 2$, and 1000. It can be seen that at $p = 0$ or $p = 1$, the super quantum discord is invariant under local bit-flip channel.




Fig. 2 Super quantum discord (dotted line) and super quantum discord (solid line) under local bit-flip channel of (20)




Fig. 3 Quantum discord (red plane) and super quantum discord (blue surface)

From a practical point of view, we sent signals through the bit-flip channel which leaves the qubit untouched with probability p , and flips the qubits with probability $1 - p$. So through bit-flip channel the state $|0\rangle$ was taken to $|1\rangle$ for $p = 0$. On the contrary, for $p = 1$ the state keeps invariant. In this view, the state ρ_x was not taken $|0\rangle$ to $|1\rangle$, the super quantum discord also did not change when $p = 0$ or $p = 1$.

In view of the above argument, we then conclude that the super quantum discord will decay after through noisy channels. It means that we will lose information after the signal through noisy, hence have to control the noise probability. Furthermore, the affection of local bit-flip channel for the super quantum discord is symmetric and attains the minimum at $p = 0.5$, so the noise probability plays a symmetric role in this noisy channel. It will disappear when the noise probability attains half of one.

When considering quantum correlations captured by the super quantum discord, it is usually known that the weak measurement captures more information than POVM measurement. However, there are some counterintuitive phenomena in our example when we compare the super quantum discord after noisy channel with discord after noisy channel.

On the Fig. 3 we can see that the red plane through the blue curve, some blue surface above red plane and others under red plane. That means after local bit-flip channel, super quantum discord is smaller than the normal quantum discord, and this difference is most obvious when the noise probability p near 0.5. Although the super quantum discord is smaller than the quantum discord only in the region of 0.01 to 0.02, it shows in this example, the weak measurement does not capture more information than POVM measurement. Thus, super quantum discord is a different resource than quantum discord.

4 Conclusions and discussions

Super quantum discord is a fundamental resource in quantum information. We have studied it for X-type states which including the Werner states and Bell diagonal states. Explicit formulae of super quantum discord for X-type states have been derived. The evolution of these states under local bit-flip channel has been investigated and reveal more different properties between super quantum discord and quantum discord. The relations between the super quantum discord and discord, evolution of super quantum discord and the weak measurement strength have been analyzed.

Acknowledgments The work is supported by NSFC11275131, NSFC11305105 and KZ201410028033.

References

1. Nielsen, M.A., Chuang, I.L.: Quantum computation and quantum information. Cambridge University Press, Cambridge, UK (2000)
2. Bennett, C.H., DiVincenzo, D.P., Fuchs, C.A., Mor, T., Rains, E., Shor, P.W., Smolin, J.A., Wootters, W.K.: *Phys Rev A* **59**, 1070 (1999)
3. Horodecki, M., Horodecki, P., Horodecki, R., Oppenheim, J., Sen, A., Sen, U., Synak-Radke, B.: *Phys. Rev. A* **71**, 062307 (2005)
4. Niset, J., Cerf, N.J.: *Phys. Rev. A* **74**, 052103 (2006)
5. Ollivier, H., Zurek, W.H.: *Phys. Rev. Lett* **88**, 017901 (2001)
6. Bylicka, B., Chruściński, D.: vol. 81. 062102 (2010)
7. Werlang, T., Souza, S., Fanchini, F.F., Villas Boas, C.J.: *Phys. Rev. A* **80**, 024103 (2009)
8. Sarandy, M.S.: *Rev. Phys. A* **80**, 022108 (2009)
9. Ferraro, A., Aolita, L., Cavalcanti, D., Cucchietti, F.M., Acín, A.: *Phys. Rev. A* **81**, 052318 (2010)
10. Fanchini, F.F., Werlang, T., Brasil, C.A., Arruda, L.G.E., Caldeira, A.O.: *Phys. Rev. A* **81**, 052107 (2010)
11. Dakić, B., Vedral, V., Brukner, Č.: *Lett. Phys. Rev* **105**, 190502 (2010)
12. Modi, K., Paterek, T., Son, W., Vedral, V., Williamson, M.: *Phys. Rev. Lett* **104**, 080501 (2010)
13. Li, N., Luo, S.: *Phys. Rev. A* **76**, 032327 (2007)
14. S. Luo: *Ibid* **77**, 022301 (2008)
15. Luo, S.: vol. 77, p. 042303 (2008)
16. Li, B., Wang, Z.X., Fei, S.M.: *Phys. Rev. A* **83**, 022321 (2011)
17. Lang, M.D., Caves, C.M.: *Phys. Rev. Lett* **105**, 150501 (2010)
18. Ali, M., Rau, A.R.P., Alber, G.: *Phys. Rev. A* **81**, 042105 (2010)
19. Ali, M., Rau, A.R.P., Alber, G.: *Ibid* **82**, 069902 (2010)
20. Mazzola, L., Piilo, J., Maniscalco, S.: *Phys. Rev. Lett* **104**, 200401 (2010)
21. Maziero, J., Céleri, L.C., Serra, R.M., Vedral, V.: *Phys. Rev. A* **80**, 044102 (2009)
22. Roa, L., Retamal, J.C., Alid-Vaccarezza, M.: *Phys. Rev. Lett* **107**, 080401 (2011)
23. Li, B., Fei, S.M., Wang, Z.X., Fan, H.: *Phys. Rev. A* **85**, 022328 (2012)
24. Aharonov, Y., Albert, D.Z., Vaidman, L.: *Phys. Rev. Lett* **60**, 1351 (1998)
25. Singh, U., Pati, A.K.: *Ann. Phys.* **343**, 141 (2014)
26. Wang, Y.K., Ma, T., Fan, H., Fei, S.M., Wang, Z.X.: *Quantum Inf. Process* **13**, 283 (2014)
27. Shi, M., Sun, C., Jiang, F., Yan, X., Du, J.: *Phys. Rev. A* **85**, 064104 (2012)
28. Partovi, M.H.: *Lett. Phys. A* **137**, 455 (1989)
29. Henderson, L., Vedral, V.: *J. Phys. A* **34**, 6899 (2001)
30. Oreshkov, O., Brun, T.A.: *Phys. Rev. Lett* **95**, 110409 (2005)
31. Werner, R.F.: *Rev. Phys. A* **40**, 4277 (1989)
32. Bennett, C.H., DiVincenzo, D.P., Smolin, J.A., Wootters, W.K.: *Phys. Rev. A* **54**, 3824 (1996)