


Quality of Life, Coping Styles, Stress Levels, and Time Use in Mothers of Children with Autism Spectrum Disorders: Comparing Single Versus Coupled Households

Tomomi McAuliffe¹  · Reinie Cordier¹ · Sharmila Vaz¹ · Yvonne Thomas² · Torbjorn Falkmer^{1,3,4,5}

Published online: 20 July 2017
© Springer Science+Business Media, LLC 2017

Abstract This study aimed to examine the influence of differences in household status on the parental stress, coping, time use and quality of life (QoL) among mothers of children with autism spectrum disorders. Forty-three single and 164 coupled mothers completed the survey. Data were analysed using multivariate logistic regression. We found that single mothers were 1.05 times more likely to report lower levels of environmental QoL. Whilst they were 1.73 times more likely to use acceptance coping style, this association did not persist after adjusting for total number of children, household income and employment status. There was no difference in time use and stress between these mothers. Possible environmental issues for single mothers and implications for future research are discussed.

Keywords ASD · Single and coupled mothers · Time use · Quality of life · Stress · Coping

Introduction

Parenting a child with autism spectrum disorders (ASD) can be a challenging task (Nicholas et al. 2016). It is well documented that parents of children with ASD experience higher levels of stress than parents of children without ASD (Griffith et al. 2010; Montes and Halterman 2007; Zablotsky et al. 2013). Limited social skills, challenging behaviour of the child with ASD (Bromley et al. 2004; Griffith et al. 2010) and low levels of family support (Bromley et al. 2004) are some of the factors associated with the increased parental stress levels.

Previous studies with families of children with ASD have developed knowledge on the parental experiences, such as parental stress levels (Griffith et al. 2010), coping styles (Marshall and Long 2010), time use (Sawyer et al. 2010), and quality of life (QoL) (Dardas and Ahmad 2014a). In the study of the parental experiences, the experiences of mothers of children with ASD are commonly examined (Ekas et al. 2010; Ekas and Whitman 2010; Kuhaneck et al. 2010;

✉ Tomomi McAuliffe
tomomi.mcauliffe@curtin.edu.au

Reinie Cordier
reinie.cordier@curtin.edu.au

Sharmila Vaz
s.vaz@curtin.edu.au

Yvonne Thomas
y.thomas@worc.ac.uk

Torbjorn Falkmer
t.falkmer@curtin.edu.au

¹ School of Occupational Therapy and Social Work, Faculty of Health Sciences, Curtin University, GPO Box U1987, Perth, WA 6845, Australia

² Allied Health and Social Sciences, Institute of Health & Society, University of Worcester, Henwick Grove, Worcester WR2 6AJ, UK

³ School of Occupational Therapy, La Trobe University, Melbourne, VIC, Australia

⁴ Rehabilitation Medicine, Department of Medicine and Health Sciences (IMH), Faculty of Health Sciences, Linköping University & Pain and Rehabilitation Centre, UHL, County Council, Linköping, Sweden

⁵ Cooperative Research Centre for Living with Autism Spectrum Disorders (Autism CRC), Long Pocket, Brisbane, QLD, Australia

Marshall and Long 2010; Seymour et al. 2013). Mothers face an increased risk of ill health (Allik et al. 2006), and are required to assume multiple roles to cater for the unique demands of the child with ASD (Safe et al. 2012). Studies that specifically examined the experiences of fathers are somewhat limited (Burrell et al. 2017; Cheuk and Lashewicz 2016); however, some studies have compared the experiences between mothers and fathers of children with ASD (Allik et al. 2006; McStay et al. 2014; Mugno et al. 2007). For example, there is emerging evidence to suggest mothers of children with ASD experience lower levels of QoL than fathers (McStay et al. 2014).

Previous studies have used various parental characteristics in investigating the experiences of parents of children with ASD. These characteristics include age (Gray 2006; Pruitt et al. 2016), gender (Allik et al. 2006; Lee 2009; McStay et al. 2014; Mugno et al. 2007), employment status (Dardas and Ahmad 2014a), and household income (Dardas and Ahmad 2014a). In fact, a recent review on coping styles among parents of children with ASD reported that age and gender of parents are the most commonly discussed factors (Lai and Oei 2014). Spousal relationship factors, such as spousal support (Dunn et al. 2001), marital satisfaction (Higgins et al. 2005), and marital quality (Harper et al. 2013), are other parental characteristics used to investigate the experiences of parents of children with ASD. Parents of children with ASD were found to experience spousal relationship issues when they used maladaptive coping strategies, such as escape-avoidance (Dunn et al. 2001).

Few studies have compared the experiences of parents of children with ASD using a parental characteristic of single versus coupled household status. Household status is common demographic information to be collected in the study of parents of children with ASD; however, it is not always used as the primary variable of interest (Mugno et al. 2007). One study reported insufficient statistical power to compare parental coping strategies based on the single versus coupled household status (Tway et al. 2007). Some studies have used the household status in their statistical analyses (Benjak 2011; Bromley et al. 2004; Reed et al. 2016). For example, one study that compared subjective QoL between parents of children with ASD and parents of children without a disability in Croatia conducted a logistic regression analysis with an independent variable of single versus coupled household status (Benjak 2011). However, the focus of the previous studies that used the variable of the household status was not to investigate the parental experiences due to single versus coupled household status. This current study, therefore, intends to fill the gap in the study of parents of children with ASD through analysing various parental experiences, such as stress levels, coping styles, time use, and QoL that relate to single versus coupled household status.

Mothers' Experiences When There is a Child with ASD in Their Families

To date, parental experiences, such as stress levels, coping styles, time use and QoL have been examined among mothers of children with ASD (Benson 2010, 2014; Dardas and Ahmad 2014a, b, 2015; Mugno et al. 2007; Sawyer et al. 2010). Families of children with ASD tend to assume traditional family roles where mothers commonly hold the primary caregiver's role (Nealy et al. 2012; Pepperell et al. 2016). As primary caregivers, mothers are more likely to experience parenting related difficulties. One study that examined child related stressors, availability of resources, and coping strategies among parents of children with ASD found that mothers experienced higher levels of stress than fathers (McStay et al. 2014). Although parents of children with ASD use various coping styles, such as acceptance, humour, and gaining hope from child's improvement (Marshall and Long 2010), some coping styles used may be counterproductive or maladaptive (Hall and Graff 2011; Sivberg 2002). A recent longitudinal study, which was conducted over a 7-year period, on maternal coping styles and adjustment among mothers of children with ASD found that increased use of maladaptive coping styles was related to increased stress proliferation and distress, and decreased parental efficacy (Benson 2014). It is therefore important to examine the parental coping styles, in particular those of mothers, given that mothers are commonly the primary organisers of the family schedules (Larson 2000).

The role of being a mother is time intensive, especially when children have a disability (Olson and Esdaile 2000). Compared with parents of typically developing children, parents of children with special needs spend longer time in childcare activities (Crowe and Florez 2006; McCann et al. 2012). One study found that mothers of children with a disability engaged in childcare activities on average 13.1 h more per week than mothers of children without a disability (Crowe and Florez 2006). Parents of children with ASD were found to spend on average 43 h per week in caring for their children (Järbrink et al. 2003). A consequence of increased childcare hours is a reduction of time available for other daily activities among parents, including personal leisure, work, and personal care (Brandon 2007). Mothers feel physically and mentally exhausted by meeting this increased care needs, while not spending much time in personal care, such as sleeping (McGuire et al. 2004).

A recent systematic review found that the QoL of parents of children with ASD is lower than that of parents of typically developing children or the general population (Vasilopoulou and Nisbet 2016). Various factors are associated with parents of children with ASD reporting lower QoL. These factors include increased challenging behaviours of the child with ASD (Dardas and Ahmad 2014a;

McStay et al. 2014; Tung et al. 2014), severity of ASD (Pozo et al. 2014), lower household income (Dardas and Ahmad 2014a), decreased levels of social support (Pozo et al. 2014), parental distress (Dardas and Ahmad 2014a), and parental coping style (Dardas and Ahmad 2014a; Pozo et al. 2014). The complex interplay between these factors should, therefore, be considered when describing the QoL among parents of children with ASD. Parents of children with ASD are often required to adjust many aspects in their life, including their family interactions, family lifestyles, marital relationships, work arrangement, coping styles, and their perspective of life (Ooi et al. 2016). QoL is useful in capturing the complexity of the lives of these parents and should be considered in understanding the impact of having a child with ASD on parents (Dardas and Ahmad 2014a).

Parental Experiences Among Single Mothers

The family structure has changed over the years in Western society due to the decreased marriage rates, the increased divorce rates, and the increased number of cohabiting couples (The Organisation for Economic Co-operation and Development [OECD] 2012). One of the notable family structural changes is an increase of single parent families. In Australia, by 2031 the rate of the single-parent families is projected to increase by between 40 and 77% (Australian Bureau of Statistics 2010). Other OECD countries, such as UK and Norway, have projected similar trends of an increase of single parent families (OECD, 2012). In 2015, there were 617,800 single parent families with dependants living with them in Australia and of those, 84% were led by single mothers (Australian Bureau of Statistics 2016). This over-representation of single parent families that are led by mothers is predicted to remain unchanged over the next 15 years (Australian Bureau of Statistics 2010).

Single motherhood, in general, presents some unique challenges, for example, additional time pressure in managing daily family responsibilities (Compas and Williams 1990; Sachs et al. 1995) and financial pressures (Cooper et al. 2008; Sachs et al. 1995). One study that compared the experiences of single and coupled first-time mothers of typically developing infants found that single mothers experienced higher levels of stress than their counterpart (Copeland and Harbaugh 2010). Both global (Witvliet et al. 2014) and Australian (Crosier et al. 2007) studies indicate that single mothers experience poorer physical and mental health outcomes compared to coupled mothers.

Single motherhood has been previously explored particularly within the area of social welfare and employment research (Cook 2012b, c; Herbst and Tekin 2014; Wu et al. 2014); however, there is a paucity of research on single mothers of children with chronic health conditions,

including children with a disability (Brown et al. 2008). Even fewer studies have been conducted with single mothers of children with ASD (Dyches et al. 2016). Given the challenges that single mothers face, combined with the unique factors identified among mothers of children with ASD, such as increased stress levels and lower QoL, there is a need to investigate the experiences of single mothers of children with ASD.

The Aim of the Study

The aim of the current study was to examine the influence of differences in household status (single versus coupled) on the stress levels, coping styles, time use and QoL among mothers of children with ASD. For the purpose of this study, we employed the definition of lone parent provided by the Australian Bureau of Statistics (2015) to define single mothers. Single mothers in this study are those who have “no spouse or partner usually resident in the household but who forms a parent–child relationship with at least one child usually resident in the household” (Australian Bureau of Statistics 2015, para. 21).

Methods

Study Design

The present study was part of a larger cross-sectional study that investigated the experiences of families of children with ASD in Western Australia. Researchers at Curtin University conducted this study in collaboration with the Disability Services Commission of Western Australia (DSC) and South-West Autism Network (SWAN). The larger study examined a number of topics related to the experiences of families of children with ASD, with a cross-sectional survey that consisted of the following four sections: (1) demographic information of the family, (2) the process of ASD diagnosis and access to health services, (3) the impact of having a child with ASD on family life, and (4) possible behavioural difficulties related to the characteristics of ASD. The survey was pilot tested with a convenient sample for clarify of questions, the flow of content, and estimating completion timeframe. The survey was refined following the pilot testing.

Participants

A parent or a primary caregiver of a 2–18 year old child or young person diagnosed with ASD living in Western Australia was eligible to participate in this study. The participants were recruited through the existing internal lists of

families of children with ASD who agreed to participate in future research held at Curtin University and the Telethon Kids Institute; service providers, such as DSC and SWAN; ASD related events held at Curtin University; health professionals working with families of children with ASD; and media release on radio.

Instruments

Four measures were used in this current study: (1) The World Health Organisation Quality of Life BREF (The WHOQOL Group 1998), (2) Autism Parental Stress Index (Silva and Schalock 2012), (3) The Brief COPE (Carver 1997), and (4) time use estimate table. All four measures were included in section three of the survey (i.e., the impact of having a child with ASD on family life).

The World Health Organisation Quality of Life Brief (WHOQOL BREF)

The WHOQOL BREF that is a short form of the WHOQOL-100 is a convenient and concise tool to use for studies in which QoL is of interest (The WHOQOL Group 1998). It is possible to calculate four domain scores: physical, psychological, social relationships, and environment. Social relationships and environmental domains are often not included in other short version of QoL assessments (The WHOQOL Group 1998). There are 26 items in total. Some examples of the items in the physical domain include pain and discomfort, energy and fatigue, mobility, and work capacity. In the psychological domain, items, such as positive and negative feelings, self-esteem, bodily image, and spirituality, are included. The social relationships domain includes personal relationships, social support, and sexual activity. In the environment domain, items, such as physical safety and security, home environment, financial resources, accessibility to services and its quality, and opportunities for leisure activity, are included (The WHOQOL Group 1998). Two items are independent from these four domains: respondents' overall perception of QoL and their overall perception of health (The WHOQOL Group 1996). A five-point Likert scale is used in WHOQOL BREF, and the respondents are asked to answer each item reflecting their experiences in the last 2 weeks (Skevington et al. 2004). Four domain scores are calculated using the mean score of each item within each domain, and in order to make the domain scores comparable with the WHOQOL-100, the mean scores are then multiplied by four (The WHOQOL Group 1996). Each domain is reported to have acceptable internal consistency. The alpha levels reported for each domain are as follows: for the physical domain 0.82, for the psychological domain 0.81, for the social relationships domain 0.68, and for the environmental

domain 0.80 (Skevington et al. 2004). Likewise, acceptable test–retest reliability of 0.66 for the physical domain, 0.72 for the psychological domain, 0.76 for the social relationships domain, and 0.87 for the environment domain over a 2–4 week period was reported (The WHOQOL Group 1998).

Autism Parental Stress Index (APSI)

The APSI has an overall parental stress scale and three subscales that measure stress levels of parents of a child with ASD, specifically identifying the areas with which parents require further support (Silva and Schalock 2012). These three subscales include (1) the core autism symptoms, (2) comorbid behaviours, and (3) comorbid physical issues. The results of the APSI validation suggests good internal consistency for the overall scale (Cronbach's $\alpha=0.83$). The acceptable internal consistency of 0.79 for the core autism symptoms; 0.76 for the co-morbid behaviours; and 0.67 for the co-morbid physical issue was reported. A good test–retest reliability of 0.88 was reported for the overall scale over a 4-month period (Silva and Schalock 2012).

Brief COPE

The 28-item Brief COPE was developed as a shorter version of the full COPE, and it measures 14 different coping styles (Carver 1997). The 14 theoretically derived subscales that consist of two items per subscale represent different coping styles: (1) active coping, (2) planning, (3) positive reframing, (4) acceptance, (5) humour, (6) religion, (7) emotional support, (8) instrumental support, (9) self-distraction, (10) denial, (11) venting, (12) substance use, (13) behavioural disengagement, and (14) self-blame. The Brief COPE is a useful tool for researchers to assess a range of coping styles, whilst reducing participant response burden. Respondents are asked to rate each statement that describes a coping style on a 4-point scale that ranges from 'I haven't been doing this at all' to 'I've been doing this a lot'. On average, adequate internal reliabilities (α ranged from 0.50 to 0.90) have been reported (Carver 1997).

Time Use Estimate Table

This constituted a 24-h format table that participants completed to describe their average weekday and weekend day by providing estimated hours on each of the 14 activities presented. We chose these 14 activities from previous publications on time use of mothers of children with a disability (Gevir et al. 2006; McCann et al. 2012). The fourteen activities listed in this study were (1) sleep, (2) grooming and personal hygiene, (3) meal preparations and clean up, (4) having a meal, (5) taking care of child, (6) travel

time to and from school or work, (7) time spent at work, (8) personal leisure time, (9) house duties, (10) shopping, (11) having quiet time or down time, (12) visiting family/friends, (13) studying, and (14) voluntary work.

Procedures

The survey was made available in three different formats for completion: (1) an online version, (2) a telephone interview to complete the survey, or (3) a paper and pencil version to be returned via post. The online survey was available from January 2015 to December 2015. The paper and pencil version of the survey was distributed to prospective participants at several events, which were organised by the Curtin Autism Research Group (CARG) in 2015. Participants who completed the online survey ($n=150$) provided consent to participate in this study by completing the survey. For those participants who completed the survey ($n=2$) by telephone, a researcher read the scripted consent prior to starting the survey, and the participants' response was recorded in the electronic data entry form. The participants who completed the paper and pencil survey ($n=55$) were asked to return the signed consent form to the research team. We have conducted a sensitivity analysis with all variables that went into the univariate analysis between those participants who completed the online survey and those who completed the paper and pencil version. There were no meaningful differences between the two forms of data collection on any of the variables. Due to the procedure used to collect data, it was impossible to calculate how many potential participants who received the survey, opted not to participate.

All data were entered directly by participants online or the researchers entered data from the telephone and the paper and pencil versions online. These data were stored on Curtin University's Qualtrics Web Server, which was password protected. A total of 248 surveys were completed. We removed those double or triple entries made by the same participants ($n=22$) and only kept their latest entry in this current study. We also removed the entries made by fathers ($n=16$) and by grandparents ($n=3$) as the targeted population of the current study was mothers. In total, 207 were completed by mothers, including biological, step and foster, and thus were deemed relevant to the current study. For the purpose of this study, those 207 participants were categorised into coupled mothers and single mothers.

Data Analyses

The Statistical Package for the Social Sciences (SPSS) version 22 (IBM Corp. 2013) was used to analyse data. The question on participant's household status was used to create a dichotomous variable, a coupled mother or a single mother. The participants chose their household status from

the following seven options: (1) two-parent, (2) single parent, (3) extended family (grandparents), (4) two-parent plus extended family, (5) single parent plus extended family, (6) foster situation, and (7) other. The household status category of the coupled mother was formed by collapsing the two parent ($n=156$) or the two-parent plus extended family options ($n=3$). The household status category of the single mother was formed by combining the single parent ($n=39$) and the single parent plus extended family options ($n=4$). Those participants who chose the other option were included if further information to identify their household status (i.e. two-parent or single parent) was provided ($n=5$). Those participants who chose the extended family option were not included in this current study. The newly created household dichotomous variable, a coupled mother or a single mother, was used as a dependent variable.

Although the structure of the coping styles with Brief COPE among parents of children with ASD has been previously explored (Benson 2010; Hastings et al. 2005), both studies yielded different factor structures and had smaller sample size ($N=113$; $N=135$ respectively) than the current study ($N=207$). Therefore, factor analysis was conducted to explore the structure of the coping styles that best fit to the current study participants. A principal component analysis of the 28 items using oblimin rotation was undertaken. The scree-test was used to determine the number of factors with the eigenvalue greater than 1.0 (Field 2005; Osborne and Costello 2009) and factor loading greater than 0.40. Eigenvalues greater than one indicated that the first five factors, which are listed in Table 1, explained 18, 15, 9, 8, and 6% of the variance respectively. The sixth and the seventh factors had eigenvalues of just over 1%, and they explained 5 and 4% of the variance respectively. Following the initial examination of the correlation matrix, the two items that form a subscale of substance use and another two items that form a subscale of use of religion were excluded, as their correlations were high ($r>0.8$). The Kaiser-Meyer-Olkin measure indicated good sampling adequacy ($KMO=0.75$) (Field 2005). Bartlett's test of sphericity [$\chi^2(276)=1724.67$, $p=0.001$] indicated that the correlation between each item was sufficient (Field 2005). As a result, the five-factor solution explaining 56% of the total variance was used in the current study. Cronbach's alpha for each of the five-factor is reported in Table 1.

Unlike the previous two studies (Benson 2010; Hastings et al. 2005) that reported four factors, our analysis yielded five factors with one item dropped. The dropped items was 'I have been turning to work or other activities to take mind off things', which is an item for the self-distraction subscale. The result of the factor analysis is presented in Table 1. Our factors somewhat resembled the result of Hastings et al. (2005) and hence their labels suited the current extracted factors. Factor one (problem focused)

Table 1 Factor loadings for analysis of brief COPE items

Factor	Problem focused	Active avoidance	Positive coping	Limited use of relationship support	Acceptance
% Variance	18.44	15.34	9.02	7.74	5.74
Cronbach's alpha	0.77	0.78	0.74	0.75	0.70
Coping taking action to try to make the situation better	0.722	-0.052	0.028	-0.001	0.172
Coping concentrating my efforts on doing something about the situation I am in	0.700	0.081	0.033	-0.068	-0.087
Coping trying to come up with a strategy about what to do	0.652	0.208	0.041	-0.046	0.244
Coping thinking hard about what steps to take	0.578	0.117	-0.036	-0.028	0.450
Coping blaming myself for things that happened	0.147	0.738	-0.009	0.028	-0.035
Coping giving up the attempt to cope	-0.205	0.735	-0.184	0.048	0.186
Coping criticizing myself	0.151	0.721	-0.002	0.119	-0.102
Coping saying to myself this isnt real	0.096	0.646	0.131	0.120	-0.340
Coping refusing to believe that it has happened	0.231	0.615	-0.011	0.040	-0.414
Coping giving up trying to deal with it	-0.470	0.543	-0.017	0.109	0.335
Coping saying things to let my unpleasant feelings escape	-0.076	0.523	-0.122	-0.370	-0.123
Coping doing something to think about it less, such as going to movies, watching TV, reading, daydreaming, sleeping, or shopping	-0.260	0.409	0.233	-0.245	0.188
Coping turning to work or other activities to take my mind off things	0.010	0.349	0.090	-0.023	0.172
Coping making jokes about it	-0.080	0.009	0.894	0.042	-0.039
Coping making fun of the situation	-0.161	0.009	0.861	0.001	-0.134
Coping trying to see it in a different light, to make it seem more positive	0.325	0.069	0.594	-0.050	0.127
Coping looking for something good in what is happening	0.211	-0.150	0.536	-0.083	0.195
Coping getting emotional support from others	-0.075	-0.058	-0.047	-0.836	0.015
Coping getting comfort and understanding from someone	-0.043	-0.231	0.139	-0.726	-0.029
Coping getting help and advice from other people	0.366	-0.199	0.100	-0.611	-0.064
Coping expressing my negative feelings	-0.055	0.409	0.144	-0.570	0.002
Coping trying to get advice or help from other people about what to do	0.359	0.084	-0.081	-0.566	0.040
Coping learning to live with it	0.137	-0.062	0.097	0.104	0.715
Coping accepting the reality of the fact that it has happened	0.234	-0.099	0.007	-0.081	0.655

Significant loadings are highlighted in bold

included all items of the planning and active coping subscales. This factor describes mothers' proactive coping styles to deal with stressors. Factor two (active avoidance) included all items of the self-blame, behavioural disengagement, denial and venting and one item from self-distraction subscales. This factor describes mothers' coping style to avoid stressors physically or emotionally. Factor three (positive coping) included all items of the humour and positive reframing subscales. This factor describes mothers' coping style to perceive stressors through their constructive lens. Factor four (lack of relational support) included all items

of the emotional support and instrumental support and one item from venting subscales. This factor describes mothers' limited use of external support in dealing with stressors. Factor five (acceptance) included both items of the subscale of acceptance, and one item of the subscale of planning. This factor describes mothers' coping style to deal with stressors while adopting the situation.

A *chi-square* test was conducted to compare the relationship between single versus coupled household status and demographic variables. A univariate analysis was conducted to identify significant associations between single

versus coupled household status and the scores for the WHOQOL BREF and the APSI, the extracted factors of the Brief COPE, and the number of hours reported in the time use estimate table. Binary logistic regression analyses were then performed with the significant variables to identify key factors associated with single mothers' stress levels, coping styles, time use, QoL, and demographic variables. A backward elimination method was used to develop the final model. In this method, all variables that were found significant in the univariate analysis were entered in the regression model initially and the least significant variable was removed one at a time until it reached the final model.

Results

Demographic Information of Participants

Table 2 provides an overall descriptive profile of the participants by single versus coupled household status. There were 43 single mothers (20.8%) and 164 coupled mothers (79.2%). Significant differences were found between single and coupled mothers on the following five demographic variables: (1) respondents' self-reported presence of cognitive impairment or intellectual disability with the child with ASD was higher among single mothers ($p=0.009$), (2) total number of children was higher among coupled mothers ($p=0.011$), (3) more single mothers were unemployed ($p<0.001$), (4) household income was lower among single mothers ($p<0.001$), and (5) single mothers had lower education levels ($p=0.027$). No significant differences were found on any other demographic variables.

Univariate Analysis

Univariate analysis was conducted to identify independent variables significantly associated with single versus coupled household status. The following independent variables were found to be significant: (1) WHOQOL BREF physical domain ($p=0.003$), (2) WHOQOL BREF psychological domain ($p=0.028$), (3) WHOQOL BREF social relationships domain ($p=0.012$), (4) WHOQOL BREF environment domain ($p<0.001$), (5) WHOQOL BREF overall QoL ($p=0.001$), (6) WHOQOL BREF health satisfaction ($p=0.39$), (7) factor two (active avoidance) coping style ($p=0.001$), (8) factor five (acceptance) coping style ($p=0.008$), (9) estimated hours completing house duties during the week ($p=0.04$), (10) estimated hours visiting family or friends during the weekend ($p=0.049$), (11) APSI core autism symptoms ($p=0.013$), (12) APSI comorbid behaviour ($p=0.027$), (13) APSI comorbid physical ($p=0.014$), and (14) APSI overall scores for parental stress levels ($p=0.002$). These significant independent variables

were entered into the multivariate logistic regression analysis. Apart from the two variables, estimated hours completing house duties during the week and visiting family or friends during the weekend, there were no significant difference between single and coupled mothers in their estimated time use. No significant difference was found with the variables of coping style factor three (positive coping) and factor four (lack of relational support). These non-significant independent variables were excluded in the multivariate logistic regression analysis.

Multivariate Logistic Regression Analysis

The model was tested for goodness of fit against a constant only model and was found to be statistically significant ($\chi^2=29.29$, $DF=4$, $p<0.001$). The included factors explained 21% of the variance (Nagelkerke $R^2=0.21$). The overall prediction success rate of the model was 80.3%. The Wald criterion was used to identify factors that were significantly associated with single mothers, and Exp (B) was used to predict the direction of the outcomes. Table 3 shows the results of the multivariate logistic regression analysis.

The multivariate logistic regression analysis revealed two factors that significantly contributed towards being single mothers: (1) WHOQOL BREF environment domain ($p<0.001$); and (2) factor five, acceptance as a coping style ($p=0.006$). The current results showed that single mothers were 1.05 times more likely to report lower levels of QoL in the environmental domain (OR 1.05; 95% CI 1.04, 1.09). Although factor five (acceptance) was found significant ($p<0.05$) in the multivariate logistic regression analysis (OR 1.73; 95% CI 1.17, 2.25), which indicates single mothers were 1.73 times more likely to use acceptance as a coping style, this result did not persist after adjusted for total number of children, household income and employment status in the analysis. The WHOQOL BREF environmental domain remained significant ($p=0.004$) following this adjustment.

The following factors did not significantly contribute to the multivariate model: (1) WHOQOL BREF Physical health domain, (2) WHOQOL BREF Psychological health domain, (3) WHOQOL BREF Social relationships domain, (4) estimated hours completing house duties during the week, (5) estimated hours visiting family or friends during the weekend, (6) overall scores for parental stress levels, and (7) factor two (active avoidance) coping style.

Discussion

The aim of this current study was to examine the influence of differences in single versus coupled household status on

Table 2 Characteristics of the mothers and their children for the total sample, the sample of single mothers and the sample of coupled mothers

Characteristics	Total sample		Single mothers		Coupled mothers	
	<i>N</i> =207	%	<i>N</i> =43	%	<i>N</i> =164	%
Children's age (month) mean (standard deviation)			10.23 (4.7)		9.95 (4.9) <i>t</i> =0.73, <i>p</i> =0.78	
Children's gender						
Male	173	83.6	37	86	136	82.9
Female	34	16.4	6	14	28	17.1
				$\chi^2=0.24, DF=1, p=0.62$		
Official ASD diagnosis						
Autistic disorder (autism)	111	55.8	23	57.5	88	55.3
HFA, AS, PDD-NOS	88	44.2	17	42.5	71	44.7
				$\chi^2=0.06, DF=1, p=0.81$		
Presence of cognitive impairments/intellectual disability						
Yes	55	27	18	42.9	37	22.8
No	149	73	24	57.1	125	77.2
				$\chi^2=6.79, DF=1, p=0.009^*$		
Children's age when noticed 1st sign of something not right						
<3 years old	166	80.2	35	81.4	131	79.9
3 years old and older	41	19.8	8	18.6	33	20.1
				$\chi^2=0.49, DF=1, p=0.82$		
Children's age when formally diagnosed						
<4 years old	84	41	16	37.2	68	42
4 years old or older	121	59	27	62.8	94	58
				$\chi^2=0.32, DF=1, p=0.57$		
Number of children with ASD						
One child	189	91.3	40	93	149	90.9
More than one	18	8.7	3	7	15	9.1
				$\chi^2=0.20, DF=1, p=0.65$		
Total number of children						
One child	32	15.5	12	27.9	20	12.2
Two or more children	175	84.5	31	72.1	144	87.8
				$\chi^2=6.44, DF=1, p=0.011^*$		
Employment status						
Employed	170	88.5	19	52.8	151	96.8
Unemployed	22	11.5	17	47.2	5	3.2
				$\chi^2=55.86, DF=1, p<0.001^{**}$		
Household income ^a						
Up to \$51,999	46	22.9	31	73.8	15	9.4
\$52,000 and over	155	77.1	11	26.2	144	90.6
				$\chi^2=78.01, DF=1, p<0.001^{**}$		
Mothers' education levels						
Did not complete school	25	12.1	5	11.6	20	12.2
Year 12	25	12.1	5	11.6	20	12.2
Post TAFE or certificate	73	35.3	23	53.5	50	30.5
University	84	40.6	10	23.3	74	45.1
				$\chi^2=9.14, DF=3, p=0.027^*$		

p* < 0.05; *p* < 0.001^aThe cut-off point was set based on the median household income in Western Australia at the time of the study

Table 3 (a) Variables associated with single mothers of children with ASD, and (b) variables associated with single mothers of children with ASD with demographic variables

Single mothers							
Independent variables	B	SE	Wald	p	Exp (B)	95% CI for Exp (B)	
						Lower	Upper
(a)							
WHOQOL environmental domain	-0.57	0.013	18.88	<0.001	0.945	0.92	0.97
Brief COPE factor five (acceptance)	0.55	0.199	7.60	0.006	1.73	1.17	2.55
(b)							
WHOQOL environmental domain	-0.57	0.013	18.88	<0.001	0.945	0.92	0.97
Brief COPE factor five (acceptance)	0.236	0.286	0.845	0.358	1.30	0.743	2.28
Total number of children	-1.30	0.642	4.11	0.043	0.272	0.77	0.958
Household income	-2.22	0.618	11.82	0.001	0.119	0.35	0.399
Employment status	2.21	0.787	7.85	0.005	9.08	1.94	42.50

the stress levels, coping styles, time use and QoL among mothers of children with ASD. The finding revealed that single mothers experience lower QoL in the environmental domain, whilst they use an adaptive coping style (acceptance) more to deal with everyday stressors.

Maternal Quality of Life

In this current study, single mothers reported lower levels of QoL in the environmental domain than coupled mothers. Financial hardship is a common issue among single mothers (Crosier et al. 2007; Dyches et al. 2016; Sachs et al. 1995; OECD 2012), in particular if they are from working class background (Rowlingson and McKay 2005). One of the reasons for single mothers to experience financial hardship may be due to a lack of suitable jobs, including those with flexible employment opportunity, which prevents those single mothers from obtaining full time employment. A study that examined the work force participation of single parents across 30 European countries found that single mothers were less likely to be employed full time than their counterpart, particularly while their children were young (Riggeri and Bird 2014). Other factors, such as a lack of suitable childcare services, may also prohibit single mothers’ abilities to obtain full time employment (Broussard et al. 2012). In a meta-synthesis of single mothers’ experiences of transitioning from welfare to work, the issues of the affordability, accessibility, and practicability of childcare services were found to exacerbate the already difficult transition process among these mothers in USA and Canada (Cook 2012a). Having access to suitable childcare support can be a challenging issue among mothers of children with ASD (Bromley et al. 2004). It is, therefore, speculated that this issue maybe even more pronounced among single mothers of children with ASD.

Mothers are generally able to increase working hours as children mature and start attending school, and the

difference in full time employment rates between single and coupled mothers narrows (Riggeri and Bird 2014). However, mothers of children with complex needs, such as ASD, may need to seek part-time employment as the demand to cater for their children’s unique needs (McCann et al. 2012; Sawyer et al. 2010) may continue through to young adulthood (Smith et al. 2010). In fact, parents of children with ASD report that reduced working hours and the associated loss in income are the single largest contributing factor to the cost of having a child with ASD (Horlin et al. 2014).

In Australia, social security payments were the largest source of income for 61% of single parent families in 2003–2004 (Australian Bureau of Statistics 2007). A legislative change in Australia, which shifted some single parents to lower social security payments, resulted in these single parent families facing increased risk of poverty (The Australian Council of Social Service 2012). In the current study, the single mothers reported lower household income than coupled mothers ($p < 0.001$), and more single mothers were unemployed ($p < 0.001$). Financial insecurity derived from a lack of flexible employment opportunity and insufficient social security payment among single mothers may have contributed to their lowered perceptions of QoL in the environmental domain.

This current study did not find differences between single and coupled mothers in the physical, psychological, and social relationships domains of QoL. This result may indicate that the status of being a single versus coupled household with a child with ASD does not have much impact on their QoL. It is also possible that both groups of mothers experience lower QoL compared to the general population, but there was no marked difference between the two groups. Indeed, one study has found that parents of children with ASD had lower QoL in every domain compared with the general population (Tung et al. 2014). Given that the current study did not have a control group, future research

should consider comparing the QoL of these mothers with that of a control group to thoroughly understand the relative influence of being a single versus coupled household with a child with ASD.

Acceptance Coping Style

Single mothers were found to use the acceptance coping style more than coupled mothers. This finding is similar to a study by Compas and Williams (1990) where single mothers of typically developing young adolescents were found to use the acceptance and positive reappraisal coping style. Single mothers often need to cope with daily stressors without the support of an intimate partner, unlike coupled parents who have an option to share parental burden while supporting each other (Gray 2006). In fact, coupled mothers consider spousal support critical in coping with stressors (Kuhaneck et al. 2010). Coupled parents can access a dyadic coping strategy, where they combine individual skills and capacities to deal with daily stressors or to support their partner in managing stressors (Bodenmann 1995). In positive dyadic coping, couples can express and share stressful situations, receive both practical and emotional support from their partner, request for help in managing the stressors, and use a collaborative approach to cope with the stressors (Falconier et al. 2015). However, single mothers simply do not have access to spousal support. Instead, family and other social supports play a crucial role in managing stressors and sharing parental responsibilities for single mothers of children with a disability (Levine 2009; Muir et al. 2008). Some of these mothers consciously build and maintain their social support networks, in preparation for the future needs of their children (Levine 2009). However, finding reliable social support can be challenging due to multiple factors, for example, having to travel large geographical distances to access the support networks and having to train and empower people who lack knowledge and skills in caring for the child with complex needs (Muir et al. 2008). It may be plausible that given the complexity to establish and maintain reliable social networks, for single mothers of children with ASD, dealing situations by using the acceptance coping style maybe an easier option. While a lack of spousal and social support may be forcing these mothers to adapt the acceptance coping style, other factors may also be able to explain the current result.

The use of acceptance coping style did not remain significant after adjusting for total number of children, household income, and employment status. Our results suggest that those three demographic variables are critical in influencing the coping styles that single mothers adopt. Complex interactions between parental characteristics should be considered as coping is a process and is influenced by personal and situational factors (Folkman and Moskowitz

2004). Another possible explanation of this result is that both single and coupled mothers of children with ASD are resilient. There is emerging evidence to show that families of children with ASD are resilient (Bayat 2007; Bekhet et al. 2012; Tunali and Power 2002). Despite the unique daily challenges, such as lack of public or family understanding towards ASD (Higgins et al. 2005; Nealy et al. 2012) and challenging behaviour of the child (Bromley et al. 2004), mothers of children with ASD show extreme commitments in caring for their children (van Tongerloo et al. 2015). One study that explored the experiences of mothers of preschool children with ASD found that mothers developed hope and positive optimism over the years and their positive perspectives contributed towards their resilience (Bultas and Pohlman 2014). In a study that compared the psychological functioning and coping styles of the mothers of children with ASD ($n=364$) and the mothers of children without ASD ($n=61,408$), it was found that mothers of children with ASD were 1.8 times more likely to report that they were coping better with daily parenting issues (Montes and Halterman 2007). Parents of children with ASD develop more adaptable coping styles, such as acceptance over the years (Ooi et al. 2016), and the coping style of acceptance has been found to be a mediating factor between stress and QoL among parents of children with ASD (Dardas and Ahmad 2015). Parents are able to deal with stressors in a practical manner by accepting situations (Marshall and Long 2010). Acceptance is an indicator of resilience (Bekhet et al. 2012) that helps to unite families, which further strengthen their family relationships (Bayat 2007).

Time Use

This study did not find much difference in time use between single and coupled mothers. The current result may indicate that single versus coupled household status does not contribute to the difference in time use among mothers of children with ASD. A study conducted in Australia, found no statistically significant difference between single and coupled mothers in everyday time use (Craig and Mullan 2012; Le and Miller 2013). However, an earlier study that compared family life experiences between single and coupled mothers of children with a disability found that single mothers of children with a disability experienced more time pressures than coupled mothers (Schilling et al. 1986). In the current study, participants estimated hours for listed activities. Although time estimation method has previously been used with mothers of children with a disability (Padiadiu 1998), it may not have fully captured the experiences of mothers or account for day-to-day variations. Future study should use more comprehensive data collection methods, such as an experience sampling method, to obtain

more accurate information of mothers' time use (Chen et al. 2015; Cordier et al. 2016).

Maternal Stress Levels

This study did not find difference in parental stress levels between single and coupled mothers, which was similar to a previous research finding for mothers of children with a disability (Schilling et al. 1986). The instrument used in this study, the APSI (Silva and Schalock 2012), measures parental stress levels that are specifically related to the characteristics of ASD and do not measure other parental stress factors, such as lowered income. Therefore, the results of the current study indicate that parents of children with ASD, regardless of single versus coupled household status, experience similar levels of parenting stress related to managing the characteristics of ASD. The demographic characteristics of the single mothers in this current study were reflective of common parental stress factors among single mothers in general (Cooper et al. 2009). Our single mother participants had on average lower household incomes, higher unemployment rates and lower educational levels compared with coupled mothers. Future research comparing single and coupled mothers of a child with ASD should consider using other instruments that measure broader stressors, such as financial pressures and lack of employment opportunities, as single mothers may be more prone to these stressors than their counterpart (Cook 2012a; Sachs et al. 1995).

Limitation

The current study has several limitations. The data collected were only from mothers who resided in Western Australia. Several recruitment strategies were employed in this study and the researchers did not have access to the actual number of surveys that were distributed or the pool of participants that were approached. Although we incorporated drop out analyses in the online version of the survey, none of the participants provided data. Hence, we were unable to determine whether the profile of the non-respondents differed from the participants and this limits our ability to calculate the true response rate. Therefore, caution should be taken when generalising the findings. The current study did not have a control group. The use of a control group would be useful to understand the impact of the single versus and coupled household status on parental experiences. The focus of the current study was to compare the broader parental experiences between single versus coupled mothers, rather than the impact of the unique characteristics inherent to ASD. However, factors, such as communication difficulties and limited social skills, have been found to be associated with parental QoL (Baghdadli

et al. 2014). Future studies examining parental experiences should consider these aspects. Lastly, the method used to collect time use data may have allowed participants to over or underestimate their time, which may have resulted in an inaccurate reflection of their time allocation. For example, in a previous study, time spent on childcare was overestimated when time estimate format was used to collect data (Järbrink et al. 2003). However, due to the design of this current study, we were unable to use other methods, such as daily diary entry, to collect data.

Conclusion

This current study discussed possible environmental issues surrounding single mothers of children with ASD, including a lack of flexible employment opportunity, insufficient social security payment and the resultant financial disadvantage. The environmental factors that are associated with single mother's QoL should be further explored with a focus on the economic aspect. Prior to adjusting for demographic variables, our results demonstrated that single mothers use acceptance coping style more in managing stressors. These differences should be further investigated, in particular the reason for the use of this adaptive coping style among single mothers in order to understand the underlying factors that may assist coupled mothers in turn. We did not find much difference in time use and maternal stress levels between these two mothers. However, time pressure is a common issue in single motherhood and previous studies have identified unique parental stressors among single mothers. The use of different methods, such as experience sampling method and the use of other stress measures, maybe beneficial in comparing single and coupled mothers whilst highlighting the vital differences.

Acknowledgments We would like to acknowledge the founder of this study, the Lishman Health Foundation to conduct the study. We would like to acknowledge the contributions of the Disability Services Commission and South West Autism Network and supports provided by other ASD related service providers and health professionals in Western Australia. We would also like to thank the families who participated in this study.

Author Contributions TM assisted the data collection, performed the statistical analysis and interpreted the data, and drafted the final manuscript. RC obtained grant to conduct the larger study, participated in the design and coordination of the study and assisted the statistical analysis, interpretation of the data and drafting the manuscript. SV obtained grant to conduct the larger study, participated in the design and coordination of the study and assisted interpretation of the data and drafting the manuscript. TF obtained grant to conduct the larger study, assisted interpretation of the data and drafting the manuscript. YT assisted interpretation of the data and drafting the manuscript. All authors read and approved the final manuscript.

Compliance with Ethical Standards

Conflict of interest The authors declare that there is no conflict of interest. The authors conducted this research project, independent of the Lishman Health Foundation and those view expressed are not necessarily those of the founder's. All researchers take full responsibility for the integrity of the work.

Ethical Approval The study was approved by the Curtin University Human Research Ethics Committee (HR123/2014) and all procedures performed in this study were in accordance with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed Consent Informed consent was obtained from all individual participants included in this study.

References

- Allik, H., Larsson, J. O., & Smedje, H. (2006). Health-related quality of life in parents of school-age children with Asperger syndrome or high-functioning autism. *Health and Quality of Life Outcomes*, 4(1), 1477–7525. doi:10.1186/1477-7525-4-1.
- Australian Bureau of Statistics. (2007). *Australian social trends 2007: One-parent families*. Canberra: Australian Bureau of Statistics. Retrieved from <http://www.abs.gov.au>.
- Australian Bureau of Statistics. (2010). *Household and family projections Australia 2006 to 2031*. (cat. no. 3236.0). Canberra: Australian Bureau of Statistics. Retrieved from <http://www.abs.gov.au>.
- Australian Bureau of Statistics. (2015). *Family characteristics and transitions, Australia, 2012–2013*. (cat. no. 4442.0). Canberra: Australian Bureau of Statistics. Retrieved from <http://www.abs.gov.au>.
- Australian Bureau of Statistics. (2016). *Labour force, Australia: Labour force status and other characteristics of families*. (cat. no. 6224.0.55.001). Canberra: Australian Bureau of Statistics. Retrieved from <http://www.abs.gov.au>.
- Baghdadli, A., Pry, R., Michelon, C., & Rattaz, C. (2014). Impact of autism in adolescents on parental quality of life. *Quality of Life Research*, 23, 1859–1868. doi:10.1007/s11136-014-0635-6.
- Bayat, M. (2007). Evidence of resilience in families of children with autism. *Journal of Intellectual Disability Research*, 51(9), 702–714. doi:10.1111/j.1365-2788.2007.00960.x.
- Bekhet, A. K., Johnson, N. L., & Zauszniewski, J. A. (2012). Resilience in family members of persons with autism spectrum disorder: A review of the literature. *Issues in Mental Health Nursing*, 33, 650–656. doi:10.3109/01612840.2012.671441.
- Benjak, T. (2011). Subjective quality of life for parents of children with autism spectrum disorders in Croatia. *Applied Research in Quality of Life*, 6(1), 91–102. doi:10.1007/s11482-010-9114-6.
- Benson, P. (2010). Coping, distress, and well-being in mothers of children with autism. *Research in Autism Spectrum Disorders*, 4(2), 217–228. doi:10.1016/j.rasd.2009.09.008.
- Benson, P. (2014). Coping and psychological adjustment among mothers of children with ASD: An accelerated longitudinal study. *Journal of Autism and Developmental Disorders*, 44, 1793–1807. doi:10.1007/s10803-014-2079-9.
- Bodenmann, G. (1995). A systemic-transactional conceptualization of stress and coping in couples. *Swiss Journal of Psychology*, 54(1), 34–49.
- Brandon, P. (2007). Time away from “smelling the roses”: Where do mothers raising children with disabilities find the time to work? *Social Science & Medicine*, 65, 667–679. doi:10.1016/j.socscimed.2007.04.007.
- Bromley, J., Hare, D. J., Davison, K., & Emerson, E. (2004). Mothers supporting children with autistic spectrum disorders: Social support, mental health status and satisfaction with services. *Autism: The International Journal of Research and Practice*, 8(4), 409–423. doi:10.1177/1362361304047224.
- Broussard, C. A., Joseph, A. L., & Thompson, M. (2012). Stressors and coping strategies used by single mothers living in poverty. *Affilia: Journal of Women & Social Work*, 27(2), 190–204. doi:10.1177/0886109912443884.
- Brown, R. T., Wiener, L., Kupst, M. J., Brennan, T., Behrman, R., Compas, B. E., ... Zeltzer, L. (2008). Single parents of children with chronic illness: An understudied phenomenon. *Journal of Pediatric Psychology*, 33(4), 408–421. doi:10.1093/jpepsy/jsm079.
- Bultas, M. W., & Pohlman, S. (2014). Silver linings. *Journal of Pediatric Nursing*, 29(6), 596–605. doi:10.1016/j.pedn.2014.03.023.
- Burrell, A., Ives, J., & Unwin, G. (2017). The experiences of fathers who have offspring with autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 47(4), 1135–1147. doi:10.1007/s10803-017-3035-2.
- Carver, C. (1997). You want to measure coping but your protocol's too long: Consider the brief COPE. *International Journal of Behavioral Medicine*, 4(1), 92–100.
- Chen, Y.-W., Cordier, R., & Brown, N. (2015). A preliminary study on the reliability and validity of using experience sampling method in children with autism spectrum disorders. *Developmental Neurorehabilitation*, 18(6), 383–389. doi:10.3109/17518423.2013.855274.
- Cheuk, S., & Lashewicz, B. (2016). How are they doing? Listening as fathers of children with autism spectrum disorder compare themselves to fathers of children who are typically developing. *Autism: The International Journal of Research and Practice*, 20(3), 343–352. doi:10.1177/1362361315584464.
- Compas, B. E., & Williams, R. A. (1990). Stress, coping, and adjustment in mothers and young adolescents in single- and two-parent families. *American Journal of Community Psychology*, 18(4), 525–545.
- Cook, K. E. (2012a). Neoliberalism, welfare policy and health: A qualitative meta-synthesis of single parents' experience of the transition from welfare to work. *Health: An Interdisciplinary Journal for the Social Study of Health, Illness & Medicine*, 16(5), 507–530. doi:10.1177/1363459311434650.
- Cook, K. E. (2012b). Single parents' subjective wellbeing over the welfare to work transition. *Social Policy and Society*, 11(2), 143–155. doi:10.1017/S1474746411000546.
- Cook, K. E. (2012c). Social support in single parents' transition from welfare to work: Analysis of qualitative findings I. *International Journal of Social Welfare*, 21(4), 338–350. doi:10.1111/j.1468-2397.2011.00844.x.
- Cooper, C., Bebbington, P. E., Meltzer, H., Bhugra, D., Brugha, T., Jenkins, R., ... King, M. (2008). Depression and common mental disorders in lone parents: Results of the 2000 National Psychiatric Morbidity Survey. *Psychological Medicine*, 38(3), 335–342. doi:10.1017/S0033291707001493.
- Cooper, C., McLanahan, S., Meadows, S., & Brooks-Gunn, J. (2009). Family structure transitions and maternal parenting stress. *Journal of Marriage and Family*, 71(3), 558–574. doi:10.1111/j.1741-3737.2009.00619.x.
- Copeland, D. B., & Harbaugh, B. L. (2010). Psychosocial differences related to parenting infants among single and married mothers. *Issues in Comprehensive Pediatric Nursing*, 33(3), 129–148. doi:10.3109/01460862.2010.498330.
- Cordier, R., Brown, N., Chen, Y.-W., Wilkes-Gillan, S., & Falkmer, T. (2016). Piloting the use of experience sampling method to

- investigate the everyday social experiences of children with Asperger syndrome/high functioning autism. *Developmental Neurorehabilitation*, 19(2), 103–110. doi:10.3109/17518423.2014.915244.
- Craig, L., & Mullan, K. (2012). Lone and partnered mothers' child-care time within context in four countries. *European Sociological Review*, 28(4), 512–526. doi:10.1093/esr/jcr01.
- Crosier, T., Butterworth, P., & Rodgers, B. (2007). Mental health problems among single and partnered mothers. *Social Psychiatry and Psychiatric Epidemiology*, 42(1), 6–13. doi:10.1007/s00127-006-0125-4.
- Crowe, T. K., & Florez, S. I. (2006). Time use of mothers with school-age children: A continuing impact of a child's disability. *American Journal of Occupational Therapy*, 60, 194–203.
- Dardas, L. A., & Ahmad, M. M. (2014a). Predictors of quality of life for fathers and mothers of children with autistic disorder. *Research in Developmental Disabilities*, 35, 1326–1333. doi:10.1016/j.ridd.2014/03/009.
- Dardas, L. A., & Ahmad, M. M. (2014b). Validation of the World Health Organization's quality of life questionnaire with parents of children with autistic disorder. *Journal of Autism and Developmental Disorders*, 44, 2257–2263. doi:10.1007/s10803-014-2110-1.
- Dardas, L. A., & Ahmad, M. M. (2015). Coping strategies as mediators and moderators between stress and quality of life among parents of children with autistic disorder. *Stress and Health*, 31(1), 5–12. doi:10.1002/smi.2513.
- Dunn, M. E., Burbine, T., Bowers, C. A., & Tantleff-Dunn, S. (2001). Moderators of stress in parents of children with autism. *Community Mental Health Journal*, 37(1), 39–52. doi:10.1023/a:1026592305436.
- Dyches, T., Christensen, R., Harper, J., Mandleco, B., & Roper, S. (2016). Respite care for single mothers of children with autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 46(3), 812–824. doi:10.1007/s10803-015-2618-z.
- Ekas, N. V., Lickenbrock, D. M., & Whitman, T. L. (2010). Optimism, social support, and well-being in mothers of children with autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 40, 1274–1284. doi:10.1007/s10803-010-0986-y.
- Ekas, N. V., & Whitman, T. L. (2010). Adaptation to daily stress among mothers of children with an autism spectrum disorder: The role of daily positive affect. *Journal of Autism and Developmental Disorders*, 41, 1202–1213. doi:10.1007/s10803-010-1142-4.
- Falconier, M. K., Jackson, J. B., Hilpert, P., & Bodenmann, G. (2015). Dyadic coping and relationship satisfaction: A meta-analysis. *Clinical Psychology Review*, 42, 28–46. doi:10.1016/j.cpr.2015.07.002.
- Field, A. (2005). *Discovering statistics using SPSS* (2nd ed.). London: SAGE Publications Ltd.
- Folkman, S., & Moskowitz, J. T. (2004). COPING: Pitfalls and promise. *Annual Review of Psychology*, 55(1), 745–774. doi:10.1146/annurev.psych.55.090902.141456.
- Gevir, D., Goldstand, S., Weintraub, N., & Parush, S. (2006). A comparison of time use between mothers of children with and without disabilities. *OTJR: Occupation, Participation & Health*, 26(3), 117–127.
- Gray, D. E. (2006). Coping over time: The parents of children with autism. *Journal of Intellectual Disability Research*, 50(12), 970–976.
- Griffith, G. M., Hastings, R. P., Nash, S., & Hill, C. (2010). Using matched groups to explore child behavior problems and maternal well-being in children with Down syndrome and autism. *Journal of Autism and Developmental Disorders*, 40(5), 610–619. doi:10.1007/s10803-009-0906-1.
- Hall, H. R., & Graff, J. C. (2011). The relationships among adaptive behaviors of children with autism, family support, parenting stress, and coping. *Issues in Comprehensive Pediatric Nursing*, 34(1), 4–25. doi:10.3109/01460862.2011.555270.
- Harper, A., Dyches, T., Harper, J., Roper, S., & South, M. (2013). Respite care, marital quality, and stress in parents of children with autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 43(11), 2604–2616. doi:10.1007/s10803-013-1812-0.
- Hastings, R. P., Kovshoff, H., Brown, T., Ward, N. J., Espinosa, F. D., & Remington, B. (2005). Coping strategies in mothers and fathers of preschool and school-age children with autism. *Autism: The International Journal of Research and Practice*, 9(4), 377–391. doi:10.1177/1362361305056078.
- Herbst, C. M., & Tekin, E. (2014). Child care subsidies, maternal health and child-parent interactions: Evidence from three nationally representative datasets. *Health Economics*, 23(8), 894–916. doi:10.1002/hec.2964.
- Higgins, D. J., Bailey, S. R., & Pearce, J. C. (2005). Factors associated with functioning style and coping strategies of families with a child with an autism spectrum disorder. *Autism: The International Journal of Research and Practice*, 9(2), 125–137. doi:10.1177/1362361305051403.
- Horlin, C., Falkmer, M., Parsons, R., Albrecht, M. A., & Falkmer, T. (2014). The cost of autism spectrum disorders. *PLoS ONE*, 9(9), 1–11. doi:10.1371/journal.pone.0106552.
- IBM Corp. (2013). *IBM SPSS Statistics for Windows (Version 22.0)*. Armonk: IBM Corp.
- Järbrink, K., Fombonne, E., & Knapp, M. (2003). Measuring the parental, service and cost impacts of children with autistic spectrum disorder: A pilot study. *Journal of Autism and Developmental Disorders*, 33(4), 395–402. doi:10.1023/a:1025058711465.
- Kuhaneck, H. M., Burroughs, T., Wright, J., Lemanczyk, T., & Darragh, A. R. (2010). A qualitative study of coping in mothers of children with an autism spectrum disorder. *Physical & Occupational Therapy in Pediatrics*, 30(4), 340–350. doi:10.3109/0194638.2010.481662.
- Lai, W. W., & Oei, T. P. S. (2014). Coping in parents and caregivers of children with autism spectrum disorders (ASD): A review. *Review Journal of Autism and Developmental Disorders*, 1(3), 207–224. doi:10.1007/s40489-014-0021-x.
- Larson, E. A. (2000). The orchestration of occupation: The dance of mothers. *The American Journal of Occupational Therapy*, 54(3), 269–280.
- Le, A. T., & Miller, P. W. (2013). Lone mothers' time allocations: Choices and satisfactions. *Australian Journal of Social Issues*, 48(1), 57–82.
- Lee, G. K. (2009). Parents of children with high functioning autism: How well do they cope and adjust? *Journal of Developmental and Physical Disabilities*, 21(2), 93–114. doi:10.1007/s10882-008-9128-2.
- Levine, K. A. (2009). Against all odds: Resilience in single mothers of children with disabilities. *Social Work in Health Care*, 48(4), 402–419. doi:10.1080/00981380802605781.
- Marshall, V., & Long, B. C. (2010). Coping processes as revealed in the stories of mothers of children with autism. *Qualitative Health Research*, 20(1), 105–116. doi:10.1177/1049732309348367.
- McCann, D., Bull, R., & Winzenberg, T. (2012). The daily patterns of time use for parents of children with complex needs: A systematic review. *Journal of Child Health Care*, 16(1), 26–52. doi:10.1177/1367493511420186.
- McStay, R., Trembath, D., & Dissanayake, C. (2014). Stress and family quality of life in parents of children with autism spectrum disorder: Parent gender and the double ABCX model. *Journal of Autism and Developmental Disorders*, 44, 3101–3118. doi:10.1007/s10803-014-2178-7.

- Montes, G., & Halterman, J. S. (2007). Psychological functioning and coping among mothers of children with autism: A population-based study. *Pediatrics*, *119*(5), e1040–e1046. doi:10.1542/peds.2006-2819.
- Mugno, D., Ruta, L., D'Arrigo, V. G., & Mazzone, L. (2007). Impairment of quality of life in parents of children and adolescents with pervasive developmental disorder. *Health and Quality of Life Outcomes*, *5*(22), 1–9. doi:10.1186/1477-7525-5-22.
- Muir, K., Tudball, J., & Robinson, S. (2008). *Family resilience where families have a child (0–8 years) with disability: Final report. (SPRC Report 10/08)*. Sydney: Commonwealth State/Territory Disability Agreement Australian Government, Social Policy Research Centre UNSW.
- Nealy, C. E., O'Hare, L., Powers, J. D., & Swick, D. C. (2012). The impact of autism spectrum disorders on the family: A qualitative study of mothers' perspectives. *Journal of Family Social Work*, *15*(3), 187–201. doi:10.1080/10522158.2012.675624.
- Nicholas, D. B., Zwaigenbaum, L., Ing, S., MacCulloch, R., Roberts, W., McKeever, P., & McMorris, C. A. (2016). "Live it to understand it". *Qualitative Health Research*, *26*, 921–934. doi:10.1177/1049732315616622.
- Olson, J., & Esdaile, S. (2000). Mothering young children with disabilities in a challenging urban environment. *American Journal of Occupational Therapy*, *54*, 307–314.
- Ooi, K. L., Ong, Y. S., Jacob, S. A., & Khan, T. M. (2016). A meta-synthesis on parenting a child with autism. *Neuropsychiatric Disease and Treatment*, *12*, 745–762. doi:10.2147/NDT.S100634.
- Osborne, J. W., & Costello, A. B. (2009). Best practises in exploratory factor analysis: Four recommendations for getting the most from your analysis. *Pan-Pacific Management Review*, *12*, 131–146.
- Padeliadu, S. (1998). Time demands and experienced stress in Greek mothers of children with Down's syndrome. *Journal of Intellectual Disability Research*, *42*(2), 144–153.
- Pepperell, T. A., Paynter, J., & Gilmore, L. (2016). Social support and coping strategies of parents raising a child with autism spectrum disorder. *Early Child Development and Care*. doi:10.1080/03004430.2016.1261338.
- Pozo, P., Sarria, E., & Brioso, A. (2014). Family quality of life and psychological well-being in parents of children with autism spectrum disorders: A double ABCX model. *Journal of Intellectual Disability Research*, *58*, 442–458. doi:10.1111/jir.12042.
- Pruitt, M. M., Willis, K., Timmons, L., & Ekas, N. V. (2016). The impact of maternal, child, and family characteristics on the daily well-being and parenting experiences of mothers of children with autism spectrum disorder. *Autism: The International Journal of Research and Practice*, *20*(8), 973–985. doi:10.1177/1362361315620409.
- Reed, P., Sejunaite, K., & Osborne, L. A. (2016). Relationship between self-reported health and stress in mothers of children with autism spectrum disorders. *Journal of Autism and Developmental Disorders*, *46*, 934–941. doi:10.1007/s10803-015-2638-8.
- Riggeri, K., & Bird, C. E. (2014). *Single parents and employment in Europe: Short statistical report no. 3*. Belgium: RAND EUROPE.
- Rowlingson, K., & McKay, S. (2005). Lone motherhood and socio-economic disadvantage: Insights from quantitative and qualitative evidence. *The Sociologica Review*, *53*, 30–49. doi:10.1111/j.1467-954X.2005.00502.x.
- Sachs, B., Hall, L. A., & Pietrukowicz, M. A. (1995). Moving beyond survival: Coping behaviours of low-income single mothers. *Journal of Psychiatric and Mental Health Nursing*, *2*(4), 207–215. doi:10.1111/j.1365-2850.1995.tb00059.x.
- Safe, A., Joosten, A., & Molineux, M. (2012). The experiences of mothers of children with autism: Managing multiple roles. *Journal of Intellectual & Developmental Disability*, *37*(4), 294–302. doi:10.3109/13668250.2012.736614.
- Sawyer, M. G., Bittman, M., La Greca, A. M., Crettenden, A. D., Harchak, T. F., & Martin, J. (2010). Time demands of caring for children with autism: What are the implications for maternal mental health? *Journal of Autism and Developmental Disorders*, *40*, 620–628. doi:10.1007/s10803-009-0912-3.
- Schilling, R. F., Kirkham, M. A., Snow, W. H., & Schinke, S. P. (1986). Single mothers with handicapped children: Differer from their married counterparts? *Family Relations*, *35*(1), 66–77.
- Seymour, M., Wood, C., Giallo, R., & Jellett, R. (2013). Fatigue, stress and coping in mothers of children with an autism spectrum disorder. *Journal of Autism and Developmental Disorders*, *43*(7), 1547–1554. doi:10.1007/s10803-012-1701-y.
- Silva, L. M., & Schalock, M. (2012). Autism parenting stress index: Initial psychometric evidence. *Journal of Autism and Developmental Disorders*, *42*(4), 566–574. doi:10.1007/s10803-011-1274-1.
- Sivberg, B. (2002). Family system and coping behaviors. *Autism: The International Journal of Research and Practice*, *6*(4), 397–409.
- Skevington, S. M., Lofty, M., & O'Connell, K. A. (2004). The World Helath Organization's WHOQOL-BREF quality of life assessment: Psychometric properties and results of the international field trial a report from the WHOQOL Group. *Quality of Life Research*, *13*, 299–310.
- Smith, L. E., Hong, J., Seltzer, M. M., Greenberg, J. S., Almeida, D. M., & Bishop, S. L. (2010). Daily experiences among mothers of adolescents and adults with autism spectrum disorder. *Journal of Autism and Developmental Disorders*, *40*(2), 167–178. doi:10.1007/s10803-009-0844-y.
- The Australian Council of Social Service. (2012). *Poverty in Australia 2014*. Strawberry Hills, NSW: The Australian Council of Social Service. Retrieved from http://www.acoss.org.au/images/uploads/ACOSS_Poverty_in_Australia_2014.pdf.
- The Organisation for Economic Co-operation and Development [OECD]. (2012). *The future of families to 2030*. Paris: OECD Publishing.
- The WHOQOL Group. (1996). *WHOQOL-BREF: Introduction, administration, scoring and generic version of the assessment*. Geneva: World Health Organization.
- The WHOQOL Group. (1998). Development of the World Health Organization WHOQOL-BREF quality of life assessment. *Psychological Medicine*, *28*, 551–558.
- Tunali, B., & Power, T. G. (2002). Coping by redefinition: Cognitive appraisals in mothers of children with autism and children without autism. *Journal of Autism and Developmental Disorders*, *32*(1), 25–34.
- Tung, L., Huang, C., Tseng, M., Yen, M., Tsai, Y., Lin, Y., & Chen, K. (2014). Correlates of health-related quality of life and the perception of its importance in caregivers of children with autism. *Research in Autism Spectrum Disorders*, *8*(9), 1235–1242. doi:10.1016/j.rasd.2014/06/010.
- Twoy, R., Connolly, P. M., & Novak, J. M. (2007). Coping strategies used by parents of children with autism. *Journal of the American Academy of Nurse Practitioners*, *19*(5), 251–260. doi:10.1111/j.1745-7599.2007.00222.x.
- van Tongerloo, M. A. M. M., van Wijngaarden, P. J. M., van der Gaag, R. J., & Lagro-Janssen, A. L. M. (2015). Raising a child with an autism spectrum disorder: 'If this were a partner relationship, I would have quit ages ago'. *Family Practice*, *32*(1), 88–93. doi:10.1093/fampra/cmu076.
- Vasilopoulou, E., & Nisbet, J. (2016). The quality of life of parents of children with autism spectrum disorder: A systematic review. *Research in Autism Spectrum Disorders*, *23*, 36–49. doi:10.1016/j.rasd.2015.11.008.

- Witvliet, M. I., Arah, O. A., Stronks, K., & Kunst, A. E. (2014). A global study on lone mothers: Exploring the Associations of self-assessed general health with motherhood types and gender inequality in 32 countries. *Women's Health Issues, 24*(2), e177–e185. doi:[10.1016/j.whi.2013.12.001](https://doi.org/10.1016/j.whi.2013.12.001).
- Wu, C.-F., Wang, M.-S., & Eamon, M. K. (2014). Employment hardships and single mothers' self-rated health: Evidence from the panel study of income dynamics. *Social Work in Health Care, 53*(5), 478–502. doi:[10.1080/00981389.2014.896846](https://doi.org/10.1080/00981389.2014.896846).
- Zablotsky, B., Bradshaw, C. P., & Stuart, E. A. (2013). The association between mental health, stress, and coping supports in mothers of children with autism spectrum disorders. *Journal of Autism and Developmental Disorders, 43*, 1380–1393. doi:[10.1007/s10803-012-1693-7](https://doi.org/10.1007/s10803-012-1693-7).