



Complexity of the Universal Theory of Residuated Ordered Groupoids

Dmitry Shkatov¹ · C. J. Van Alten¹

Accepted: 19 December 2022 / Published online: 13 January 2023
© The Author(s), under exclusive licence to Springer Nature B.V. 2023

Abstract

We study the computational complexity of the universal theory of residuated ordered groupoids, which are algebraic structures corresponding to Nonassociative Lambek Calculus. We prove that the universal theory is coNP-complete which, as we observe, is the lowest possible complexity for a universal theory of a non-trivial class of structures. The universal theories of the classes of unital and integral residuated ordered groupoids are also shown to be coNP-complete. We also prove the coNP-completeness of the universal theory of classes of residuated algebras, algebraic structures corresponding to Generalized Lambek Calculus.

Keywords Residuated ordered groupoid · Universal theory · Partial algebra · Residuated algebra

1 Introduction

Residuated ordered groupoids¹ (Fuchs, 1963; Blyth & Janowitz, 1972), for short *rogs*, are algebraic models of the Nonassociative Lambek Calculus NL (Lambek, 1961), a version of the Lambek Calculus (Lambek, 1958) obtained by dropping the implicit rule of associativity. The atomic theory of *rogs*, that is, the set of sentences of the form $\forall x_1 \dots \forall x_n (s \leq t)$ valid on every *rog*, is, in essence, a notational variant of NL, with atomic sentences corresponding to sequents $s \vdash t$. Similarly, the Horn theory of *rogs*, which is the set of Horn sentences $\forall x_1 \dots \forall x_n (u_1 \leq v_1 \& \dots \& u_m \leq v_m \Rightarrow s \leq t)$ valid on every *rog*, is a notational variant of NL with nonlogical axioms (Buszkowski,

¹ In alternative terminology, residuated groupoids (Buszkowski, 2005).

✉ C. J. Van Alten
clint.vanalten@wits.ac.za
Dmitry Shkatov
shkatov@gmail.com

¹ School of Computer Science and Applied Mathematics, University of the Witwatersrand, Johannesburg, South Africa

2005). In this modification of NL, proofs are allowed that commence not only with identity sequents, but also with nonlogical assumptions $u \vdash v$, where u and v are arbitrary formulas. These assumptions correspond to the algebraic inequalities $u \leq v$ that are premisses of the corresponding Horn sentence. It is worth noting that, in contrast with NL, the cut-rule is, in general, not eliminable in NL with nonlogical axioms.

The universal theory of *rogs* is the set of sentences of the form $\forall x_1 \dots \forall x_n \varphi$, where φ is a Boolean combination of algebraic inequalities, valid on every *rog*. The universal theory is a natural extension of the Horn theory; a proof-theoretic analog thereof would be a deductive system allowing deductions of a disjunctively interpreted set of sequents from a conjunctively interpreted set of sequents. Such deductive formalisms, in the guise of multi-conclusion systems (Sambin et al., 2000; Jeřábek, 2009; Iemhoff, 2015, 2016; Bezhanishvili et al., 2016; Jalali, 2021) and hypersequents (Avron, 1996; Lellmann, 2016), have recently received considerable attention, even though no such system capturing the universal theory of *rogs* has, as far as we know, yet been constructed.

Both the atomic and Horn theory of *rogs* are decidable in polynomial time, as shown by Aarts and Trautwein (1995) and Buszkowski (2005), respectively. The universal theory of *rogs* is shown to be decidable by Farulewski (2008), however, the complexity has, to date, not been established. In the present paper we fill this gap by showing that the universal theory of *rogs* is coNP-complete which, as we observe, is the lowest possible complexity for a universal theory of a non-trivial class of structures.

In addition to *rogs* we consider two variations of *rogs* that have attracted attention, namely *unital* and *integral rogs*. By a *unital residuated ordered groupoid*, or *urog*, we mean a *rog* that has an identity for the groupoid operation. The class of *urogs* is an algebraic semantics for NL with unit. It is shown by Bulińska (2009) that the Horn and, hence, atomic theory of *urogs* is decidable in polynomial time. By an *integral residuated ordered groupoid*, or *irog*, we mean a *urog* in which the identity is the greatest element with respect to the order. The class of *irogs* is considered by Blok and Van Alten (2005), where it is shown to have the ‘finite embeddability property’, or *fep*, meaning that every finite partial substructure of an *irog* is embeddable into a finite *irog*. The *fep*, together with finite axiomatizability, implies decidability of the universal theory of *irogs*. In the present paper, we show that the universal theory of both *urogs* and *irogs* is coNP-complete. We note that decidability of the universal theory of *rogs* was established by Farulewski (2008) by proving the *fep* for the class of *rogs*. It is shown there that NL has the strong finite model property, from which the *fep* for the class of *rogs* then follows by Blok and Van Alten (2002).

We shall also consider classes of *residuated algebras*. These algebras were introduced by Buszkowski (1989) as algebraic semantics for the Generalized Lambek Calculus [see also (Buszkowski, 1998; Kołowska-Gawiejnowicz, 1997; Kandulski, 1997; Jäger, 2004)]. It is shown by Buszkowski (2005) that the Horn and, hence, atomic theory of each class of residuated algebras is decidable in polynomial time. The *fep*, and hence decidability of the universal theory, for each class of residuated algebras is proved by Buszkowski (2011). In this paper, we extend the methods used for residuated ordered groupoids to show that the universal theory of each class of residuated algebras is coNP-complete.

The upper complexity bounds obtained herein are established using a technique of partial structures described by Van Alten (2013). For *rogs*, for example, the technique requires an intrinsic characterization of partial substructures of *rogs*, or ‘partial *rogs*’. A universal sentence fails to hold in some *rog* if, and only if, it fails to hold in some partial *rog* of size not greater than the size of the given universal sentence. The characterization of partial *rogs* presented in Theorem 1 can be checked in polynomial time for a given partial structure; hence, if an arbitrary partial structure is guessed, it can be decided in polynomial time if it is a partial *rog*. In this way we obtain a non-deterministic polynomial-time algorithm for deciding if a partial structure is a partial *rog*, from which it follows that the universal theory of *rogs* is in coNP—the full details are given in Sect. 4. For the lower complexity bounds we use the observation, stated in Theorem 4, that the universal theory of any non-trivial class of structures is coNP-hard.

2 Residuated Ordered Groupoids

We consider structures corresponding to the signature σ containing binary operation symbols \circ , \backslash and $/$, as well as a binary relation symbol \leq ; we call such structures σ -structures. We assume throughout the paper that terms are built out of a fixed countable set *Var* of variables, whose members are typically denoted by x, y, z, x_1, y_1, \dots .

A *residuated ordered groupoid*, or *rog*, is a σ -structure $\mathbf{A} = \langle A, \circ, \backslash, /, \leq \rangle$, where $\langle A, \leq \rangle$ is a partially ordered set and \circ, \backslash and $/$ are binary operations on A such that, for all $a, b, c \in A$,

$$a \circ b \leq c \iff b \leq a \backslash c \iff a \leq c / b. \quad (1)$$

The class of all *rogs* is denoted by \mathcal{ROG} . It follows from (1) that \circ is order-preserving in both coordinates, \backslash is order-reversing in the first coordinate and order-preserving in the second, while $/$ is order-preserving in the first coordinate and order-reversing in the second.

If $\mathbf{A} = \langle A, \circ^A, \backslash^A, /^A, \leq^A \rangle$ and $\mathbf{B} = \langle B, \circ^B, \backslash^B, /^B, \leq^B \rangle$ are *rogs*, then by an *embedding* of \mathbf{B} into \mathbf{A} we mean a map $\alpha : B \rightarrow A$ such that $a \leq^B b$ if, and only if, $\alpha(a) \leq^A \alpha(b)$, and $\alpha(a \star^B b) = \alpha(a) \star^A \alpha(b)$ for every $\star \in \{\circ, \backslash, /\}$ and all $a, b \in B$. Observe that every embedding is injective since, for all $a, b \in B$, if $a \neq b$ then either $a \not\leq^B b$ or $b \not\leq^B a$, so $\alpha(a) \not\leq^A \alpha(b)$ or $\alpha(b) \not\leq^A \alpha(a)$; in particular, $\alpha(a) \neq \alpha(b)$.

We shall use the following terminology and notation. Let $\langle Q, \leq \rangle$ be a partially ordered set. A subset X of Q is called an *upset* of $\langle Q, \leq \rangle$ if $a \in X$ and $a \leq b$ imply $b \in X$, for all $a, b \in Q$. If the order relation on Q is clear from the context, we shall use $U(Q)$ to denote the set of all upsets of $\langle Q, \leq \rangle$ and, for $a \in Q$, by the *principal upset* of a , we shall mean the set $[a] := \{b \in Q \mid a \leq b\}$.

We shall rely on the frame representation theory for *rogs* due to Dunn (1993). A *rog-frame* is a structure $\mathfrak{F} = \langle P, \leq, R \rangle$, where $\langle P, \leq \rangle$ is a partially ordered set and R is a ternary relation on P that is monotone in the last coordinate and antitone in the first two coordinates, i.e., such that, for all $f, f', g, g', h, h' \in P$,

$$R(f, g, h) \ \& \ f' \leq f \ \& \ g' \leq g \ \& \ h \leq h' \implies R(f', g', h'). \quad (2)$$

Starting from a *rog*-frame $\mathfrak{F} = \langle P, \leq, R \rangle$, we obtain a *rog* that has as its universe the set $U(P)$, as follows. Define, for all $X, Y \in U(P)$,

$$X \circ Y := \{h \in P \mid (\exists f, g \in P) [f \in X \ \& \ g \in Y \ \& \ R(f, g, h)]\}; \tag{3}$$

$$X \setminus Y := \{g \in P \mid (\forall f, h \in P) [f \in X \ \& \ R(f, g, h) \implies h \in Y]\}; \tag{4}$$

$$Y/X := \{f \in P \mid (\forall g, h \in P) [g \in X \ \& \ R(f, g, h) \implies h \in Y]\}. \tag{5}$$

Since \mathfrak{F} satisfies (2), so defined \circ, \setminus and $/$ are operations on $U(P)$. The definitions (3–5) ensure that (1) is satisfied with respect to the partial order \subseteq on $U(P)$. Thus, $\mathbf{A}_{\mathfrak{F}} = \langle U(P), \circ, \setminus, /, \subseteq \rangle$ is a *rog*, which we call the *rog associated with \mathfrak{F}* .

Conversely, starting from a *rog* $\mathbf{A} = \langle A, \circ, \setminus, /, \leq \rangle$, we can obtain a *rog*-frame, as follows. Define a ternary relation R on $U(A)$ by

$$R(f, g, h) \iff (\forall a, b \in A) [a \in f \ \& \ b \in g \implies a \circ b \in h]. \tag{6}$$

Then R and \subseteq satisfy condition (2), hence $\mathfrak{F}_{\mathbf{A}} = \langle U(A), \subseteq, R \rangle$ is a *rog*-frame. Thus, we may obtain $\mathbf{A}_{\mathfrak{F}_{\mathbf{A}}}$, the *rog associated with $\mathfrak{F}_{\mathbf{A}}$* . It is straightforward to check that the map $\mu : A \rightarrow U(U(A))$ defined by $\mu(a) = \{f \in U(A) \mid a \in f\}$ is an embedding of \mathbf{A} into $\mathbf{A}_{\mathfrak{F}_{\mathbf{A}}}$.

3 Partial σ -Structures and Partial *rogs*

In this section, we introduce partial *rogs* and obtain an intrinsic characterization thereof; this characterization will play a key role in establishing our complexity results.

A *partial σ -structure* is a structure $\mathbf{B} = \langle B, \circ^{\mathbf{B}}, \setminus^{\mathbf{B}}, /^{\mathbf{B}}, \leq^{\mathbf{B}} \rangle$, where $\leq^{\mathbf{B}}$ is a binary relation on B and $\circ^{\mathbf{B}}, \setminus^{\mathbf{B}}$ and $/^{\mathbf{B}}$ are partial binary operations on B , i.e., partial functions from $B \times B$ into B . The domains of $\circ^{\mathbf{B}}, \setminus^{\mathbf{B}}$ and $/^{\mathbf{B}}$ are denoted by, respectively, $\text{dom}(\circ^{\mathbf{B}})$, $\text{dom}(\setminus^{\mathbf{B}})$ and $\text{dom}(/^{\mathbf{B}})$. For clarity, we note that for each $\star \in \{\circ, \setminus, /\}$, $\text{dom}(\star^{\mathbf{B}}) \subseteq B \times B$.

A *partial *rog** is a partial σ -structure $\mathbf{B} = \langle B, \circ^{\mathbf{B}}, \setminus^{\mathbf{B}}, /^{\mathbf{B}}, \leq^{\mathbf{B}} \rangle$ which is a partial substructure of a *rog*, that is, for which there exists a *rog* $\mathbf{A} = \langle A, \circ^{\mathbf{A}}, \setminus^{\mathbf{A}}, /^{\mathbf{A}}, \leq^{\mathbf{A}} \rangle$ such that $B \subseteq A$, $\leq^{\mathbf{B}} = \leq^{\mathbf{A}} \upharpoonright_B$ and $a \star^{\mathbf{B}} b = a \star^{\mathbf{A}} b$ for each $\star \in \{\circ, \setminus, /\}$ and $(a, b) \in \text{dom}(\star^{\mathbf{B}})$.

We note that if \mathbf{B} is a partial *rog* that is a partial substructure of a *rog* \mathbf{A} , then $\star^{\mathbf{B}}$ is not necessarily the restriction of $\star^{\mathbf{A}}$ to B ; that is, there may exist $a, b \in B$ and $\star \in \{\circ, \setminus, /\}$ for which $a \star^{\mathbf{A}} b \in B$ but $(a, b) \notin \text{dom}(\star^{\mathbf{B}})$. This definition is consistent with that of partial substructure used by Van Alten (2013), but contrasts with the closely related notion of partial subalgebra used, for example, by Blok and Van Alten (2002).

By an *embedding* of a partial σ -structure $\mathbf{B} = \langle B, \circ^{\mathbf{B}}, \setminus^{\mathbf{B}}, /^{\mathbf{B}}, \leq^{\mathbf{B}} \rangle$ into a σ -structure $\mathbf{A} = \langle A, \circ^{\mathbf{A}}, \setminus^{\mathbf{A}}, /^{\mathbf{A}}, \leq^{\mathbf{A}} \rangle$ we mean a map $\alpha : B \rightarrow A$ such that $a \leq^{\mathbf{B}} b$ if, and only if, $\alpha(a) \leq^{\mathbf{A}} \alpha(b)$, and $\alpha(a \star^{\mathbf{B}} b) = \alpha(a) \star^{\mathbf{A}} \alpha(b)$ for each $\star \in \{\circ, \setminus, /\}$ and $(a, b) \in \text{dom}(\star^{\mathbf{B}})$.

Observe that if \mathbf{B} is a partial σ -structure and there exists an embedding of \mathbf{B} into a *rog* \mathbf{A} , then \mathbf{B} is isomorphic to a partial substructure of \mathbf{A} , that is, \mathbf{B} is (isomorphic

to) a partial *rog*. Using this observation, in the following theorem we give an intrinsic characterization of partial *rogs*. The construction used in the proof of the theorem resembles the construction of the *rog* associated with \mathfrak{F}_A , i.e., $\mathbf{A}_{\mathfrak{F}_A}$, from a *rog* \mathbf{A} .

Theorem 1 *A partial σ -structure $\mathbf{B} = \langle B, \circ^{\mathbf{B}}, \backslash^{\mathbf{B}}, /^{\mathbf{B}}, \leq^{\mathbf{B}} \rangle$ is a partial *rog* if, and only if, it satisfies the following conditions:*

- (i) $\langle B, \leq^{\mathbf{B}} \rangle$ is a partially ordered set;
- (ii) $(\forall \langle a, b \rangle, \langle c, d \rangle \in \text{dom}(\circ^{\mathbf{B}})) [a \leq^{\mathbf{B}} c \ \& \ b \leq^{\mathbf{B}} d \implies a \circ^{\mathbf{B}} b \leq^{\mathbf{B}} c \circ^{\mathbf{B}} d]$;
- (iii) $(\forall \langle a, b \rangle \in \text{dom}(\circ^{\mathbf{B}})) (\forall \langle c, d \rangle \in \text{dom}(\backslash^{\mathbf{B}}))$
 $[a \leq^{\mathbf{B}} c \ \& \ b \leq^{\mathbf{B}} c \backslash^{\mathbf{B}} d \implies a \circ^{\mathbf{B}} b \leq^{\mathbf{B}} d]$;
- (iv) $(\forall \langle a, b \rangle \in \text{dom}(\circ^{\mathbf{B}})) (\forall \langle c, d \rangle \in \text{dom}(/^{\mathbf{B}}))$
 $[a \leq^{\mathbf{B}} c /^{\mathbf{B}} d \ \& \ b \leq^{\mathbf{B}} d \implies a \circ^{\mathbf{B}} b \leq^{\mathbf{B}} c]$;
- (v) $(\forall \langle a, b \rangle \in \text{dom}(\backslash^{\mathbf{B}})) (\forall \langle c, d \rangle \in \text{dom}(\circ^{\mathbf{B}}))$
 $[a \leq^{\mathbf{B}} c \ \& \ c \circ^{\mathbf{B}} d \leq^{\mathbf{B}} b \implies d \leq^{\mathbf{B}} a \backslash^{\mathbf{B}} b]$;
- (vi) $(\forall \langle a, b \rangle, \langle c, d \rangle \in \text{dom}(\backslash^{\mathbf{B}})) [a \leq^{\mathbf{B}} c \ \& \ d \leq^{\mathbf{B}} b \implies c \backslash^{\mathbf{B}} d \leq^{\mathbf{B}} a \backslash^{\mathbf{B}} b]$;
- (vii) $(\forall \langle a, b \rangle \in \text{dom}(\backslash^{\mathbf{B}})) (\forall \langle c, d \rangle \in \text{dom}(/^{\mathbf{B}}))$
 $[a \leq^{\mathbf{B}} c /^{\mathbf{B}} d \ \& \ c \leq^{\mathbf{B}} b \implies d \leq^{\mathbf{B}} a \backslash^{\mathbf{B}} b]$;
- (viii) $(\forall \langle a, b \rangle \in \text{dom}(/^{\mathbf{B}})) (\forall \langle c, d \rangle \in \text{dom}(\circ^{\mathbf{B}}))$
 $[b \leq^{\mathbf{B}} d \ \& \ c \circ^{\mathbf{B}} d \leq^{\mathbf{B}} a \implies c \leq^{\mathbf{B}} a /^{\mathbf{B}} b]$;
- (ix) $(\forall \langle a, b \rangle \in \text{dom}(/^{\mathbf{B}})) (\forall \langle c, d \rangle \in \text{dom}(\backslash^{\mathbf{B}}))$
 $[d \leq^{\mathbf{B}} a \ \& \ b \leq^{\mathbf{B}} c \backslash^{\mathbf{B}} d \implies c \leq^{\mathbf{B}} a /^{\mathbf{B}} b]$;
- (x) $(\forall \langle a, b \rangle, \langle c, d \rangle \in \text{dom}(/^{\mathbf{B}})) [c \leq^{\mathbf{B}} a \ \& \ b \leq^{\mathbf{B}} d \implies c /^{\mathbf{B}} d \leq^{\mathbf{B}} a /^{\mathbf{B}} b]$.

Proof It is straightforward to check that conditions (i–x) hold in every *rog*. Since these conditions are universal sentences with quantifiers relativized by the domains of the partial operations, it follows that (i–x) hold in every partial *rog*.

Conversely, suppose $\mathbf{B} = \langle B, \circ^{\mathbf{B}}, \backslash^{\mathbf{B}}, /^{\mathbf{B}}, \leq^{\mathbf{B}} \rangle$ is a partial σ -structure satisfying (i–x). We construct a *rog* into which \mathbf{B} is embeddable. Define a ternary relation $R^{\mathbf{B}}$ on the set $U(B)$ of all upsets of $\langle B, \leq^{\mathbf{B}} \rangle$ by:

$$\begin{aligned} R^{\mathbf{B}}(f, g, h) \iff & (\forall \langle a, b \rangle \in \text{dom}(\circ^{\mathbf{B}})) [a \in f \ \& \ b \in g \implies a \circ^{\mathbf{B}} b \in h] \\ & \& (\forall \langle a, b \rangle \in \text{dom}(\backslash^{\mathbf{B}})) [a \in f \ \& \ a \backslash^{\mathbf{B}} b \in g \implies b \in h] \\ & \& (\forall \langle a, b \rangle \in \text{dom}(/^{\mathbf{B}})) [a /^{\mathbf{B}} b \in f \ \& \ b \in g \implies a \in h]. \end{aligned}$$

It is straightforward to check that the structure $\mathfrak{F} = \langle U(B), \subseteq, R^{\mathbf{B}} \rangle$ is a *rog*-frame. Let $\mathbf{A}_{\mathfrak{F}} = \langle U(U(B)), \circ, \backslash, /, \subseteq \rangle$ be the *rog* associated with \mathfrak{F} and let $\mu: B \rightarrow U(U(B))$ be the map defined by $\mu(a) = \{f \in U(B) \mid a \in f\}$. We show that μ is an embedding of \mathbf{B} into $\mathbf{A}_{\mathfrak{F}}$.

If $a, b \in B$ with $a \leq^{\mathbf{B}} b$, then every element of $\mu(a)$ contains b ; hence, $\mu(a) \subseteq \mu(b)$. If $a \not\leq^{\mathbf{B}} b$, then the principal upset $[a]$ belongs to $\mu(a)$ but not to $\mu(b)$; hence $\mu(a) \not\subseteq \mu(b)$. Thus, $a \leq^{\mathbf{B}} b$ if, and only if, $\mu(a) \subseteq \mu(b)$.

Let $\langle a, b \rangle \in \text{dom}(\circ^{\mathbf{B}})$. We show that $\mu(a \circ^{\mathbf{B}} b) = \mu(a) \circ \mu(b)$, where

$$\mu(a \circ^{\mathbf{B}} b) = \{h \in U(B) \mid a \circ^{\mathbf{B}} b \in h\}$$

and

$$\mu(a) \circ \mu(b) = \{h \in U(B) \mid (\exists f, g \in U(B)) [a \in f \ \& \ b \in g \ \& \ R^{\mathbf{B}}(f, g, h)]\}.$$

Assume $h \in \mu(a) \circ \mu(b)$. Then there exist $f, g \in U(B)$ such that $a \in f, b \in g$ and $R^{\mathbf{B}}(f, g, h)$. It follows from the definition of $R^{\mathbf{B}}$ that $a \circ^{\mathbf{B}} b \in h$, hence $h \in \mu(a \circ^{\mathbf{B}} b)$. Thus, $\mu(a) \circ \mu(b) \subseteq \mu(a \circ^{\mathbf{B}} b)$.

Next, assume $h \in \mu(a \circ^{\mathbf{B}} b)$, i.e., $a \circ^{\mathbf{B}} b \in h$. Let $f := [a]$ and $g := [b]$. Clearly, $f, g \in U(B)$, $a \in f$ and $b \in g$. We show that $R^{\mathbf{B}}(f, g, h)$. Suppose that c and d are arbitrary elements of B for which $\langle c, d \rangle \in \text{dom}(\circ^{\mathbf{B}})$, $c \in f$ and $d \in g$. Then, $a \leq^{\mathbf{B}} c$ and $b \leq^{\mathbf{B}} d$, hence, by (ii), $a \circ^{\mathbf{B}} b \leq^{\mathbf{B}} c \circ^{\mathbf{B}} d$. Since $a \circ^{\mathbf{B}} b \in h$ and h is an upset, $c \circ^{\mathbf{B}} d \in h$. Suppose that c and d are arbitrary elements of B for which $\langle c, d \rangle \in \text{dom}(\backslash^{\mathbf{B}})$, $c \in f$ and $c \backslash^{\mathbf{B}} d \in g$. Then, $a \leq^{\mathbf{B}} c$ and $b \leq^{\mathbf{B}} c \backslash^{\mathbf{B}} d$, hence, by (iii), $a \circ^{\mathbf{B}} b \leq^{\mathbf{B}} d$, whence $d \in h$. Lastly, suppose that c and d are arbitrary elements of B for which $\langle c, d \rangle \in \text{dom}(/^{\mathbf{B}})$, $c /^{\mathbf{B}} d \in f$ and $d \in g$. Then, $a \leq^{\mathbf{B}} c /^{\mathbf{B}} d$ and $b \leq^{\mathbf{B}} d$, hence, by (iv), $a \circ^{\mathbf{B}} b \leq^{\mathbf{B}} c$, whence $c \in h$. Thus, $R^{\mathbf{B}}(f, g, h)$. Since $a \in f$ and $b \in g$, we obtain $h \in \mu(a) \circ \mu(b)$, hence $\mu(a \circ^{\mathbf{B}} b) \subseteq \mu(a) \circ \mu(b)$.

Let $\langle a, b \rangle \in \text{dom}(\backslash^{\mathbf{B}})$. We show that $\mu(a \backslash^{\mathbf{B}} b) = \mu(a) \backslash \mu(b)$, where

$$\mu(a \backslash^{\mathbf{B}} b) = \{g \in U(B) \mid a \backslash^{\mathbf{B}} b \in g\}$$

and

$$\mu(a) \backslash \mu(b) = \{g \in U(B) \mid (\forall f, h \in U(B)) [a \in f \ \& \ R^{\mathbf{B}}(f, g, h) \implies b \in h]\}.$$

Assume $g \in \mu(a \backslash^{\mathbf{B}} b)$, i.e., $a \backslash^{\mathbf{B}} b \in g$. Suppose that, for some $f, h \in U(B)$, both $a \in f$ and $R^{\mathbf{B}}(f, g, h)$. Since $a \backslash^{\mathbf{B}} b \in g$, by definition of $R^{\mathbf{B}}$, we obtain $b \in h$ and so $g \in \mu(a) \backslash \mu(b)$. Thus, $\mu(a \backslash^{\mathbf{B}} b) \subseteq \mu(a) \backslash \mu(b)$.

Next, assume $g \notin \mu(a \backslash^{\mathbf{B}} b)$, i.e., $a \backslash^{\mathbf{B}} b \notin g$. To prove that $g \notin \mu(a) \backslash \mu(b)$, we show that there exist $f, h \in U(B)$ such that $a \in f$ and $R^{\mathbf{B}}(f, g, h)$, but $b \notin h$. Let $f := [a]$ and $h := \{e \in B \mid e \not\leq^{\mathbf{B}} b\}$. Clearly, $f, h \in U(B)$, $a \in f$ and $b \notin h$. We show that $R^{\mathbf{B}}(f, g, h)$. Suppose that c and d are arbitrary elements of B for which $\langle c, d \rangle \in \text{dom}(\circ^{\mathbf{B}})$, $c \in f$ and $d \in g$. Then, $a \leq^{\mathbf{B}} c$. Since g is an upset and $a \backslash^{\mathbf{B}} b \notin g$, surely $d \not\leq^{\mathbf{B}} a \backslash^{\mathbf{B}} b$, hence, by (v), $c \circ^{\mathbf{B}} d \not\leq^{\mathbf{B}} b$, and so $c \circ^{\mathbf{B}} d \in h$. Suppose that c and d are arbitrary elements of B for which $\langle c, d \rangle \in \text{dom}(\backslash^{\mathbf{B}})$, $c \in f$ and $c \backslash^{\mathbf{B}} d \in g$. Then $a \leq^{\mathbf{B}} c$ and $c \backslash^{\mathbf{B}} d \not\leq^{\mathbf{B}} a \backslash^{\mathbf{B}} b$, hence, by (vi), $d \not\leq^{\mathbf{B}} b$, and so $d \in h$. Lastly, suppose that c and d are arbitrary elements of B for which $\langle c, d \rangle \in \text{dom}(/^{\mathbf{B}})$, $c /^{\mathbf{B}} d \in f$ and $d \in g$. Then, $a \leq^{\mathbf{B}} c /^{\mathbf{B}} d$ and $d \not\leq^{\mathbf{B}} a \backslash^{\mathbf{B}} b$, hence, by (vii), $c \not\leq^{\mathbf{B}} b$, and so $c \in h$. Thus, $R^{\mathbf{B}}(f, g, h)$. Since $a \in f$ and $b \notin h$, we obtain $g \notin \mu(a) \backslash \mu(b)$ hence $\mu(a) \backslash \mu(b) \subseteq \mu(a \backslash^{\mathbf{B}} b)$.

For $\langle a, b \rangle \in \text{dom}(/^{\mathbf{B}})$, the proof that $\mu(a /^{\mathbf{B}} b) = \mu(a) / \mu(b)$, which relies on conditions (viii-x), is analogous.

Thus, μ is an embedding of \mathbf{B} into $\mathbf{A}_{\mathfrak{F}}$. It follows that \mathbf{B} is a partial *rog*, which completes the proof. □

4 Complexity of the Universal Theory of \mathcal{ROG}

In this section, we consider the universal theory of \mathcal{ROG} and prove its coNP-completeness.

We recall that universal σ -sentences are formulas of the form $\forall x_1 \dots \forall x_n \varphi$, where φ is a quantifier-free (first-order) σ -formula, i.e., a formula defined by the BNF expression

$$\varphi := (t \leq t) \mid \text{not } \varphi \mid (\varphi \ \& \ \varphi) \mid (\varphi \ \text{or} \ \varphi),$$

with t ranging over σ -terms, and containing no variables other than x_1, \dots, x_n .

We omit parentheses around formulas where this does not lead to ambiguity. Henceforth, given a quantifier-free σ -formula φ , we write $\varphi(x_1, \dots, x_n)$ to indicate that φ contains no variables except x_1, \dots, x_n . We denote by $\text{Var}(\varphi)$ and $\text{Ter}(\varphi)$ the sets of, respectively, variables and terms occurring in φ .

An *assignment* on a *rog* \mathbf{A} is a map $v : \text{Var} \rightarrow A$. The definition of the satisfaction relation $\mathbf{A} \models^v \varphi$ between *rogs* \mathbf{A} , assignments v and quantifier-free σ -formulas φ is standard—see, e.g., Chang and Keisler (1990). A quantifier-free σ -formula φ is *satisfiable in \mathcal{ROG}* if $\mathbf{A} \models^v \varphi$ for some *rog* \mathbf{A} and some assignment v on \mathbf{A} ; φ is *valid on \mathcal{ROG}* if $\mathbf{A} \models^v \varphi$ for every *rog* \mathbf{A} and every assignment v on \mathbf{A} . A universal σ -sentence $\forall x_1 \dots \forall x_n \varphi$ is *valid on \mathcal{ROG}* if φ is valid on \mathcal{ROG} . The *universal theory of \mathcal{ROG}* is the set of all universal σ -sentences valid on \mathcal{ROG} . Clearly, a universal sentence $\forall x_1 \dots \forall x_n \varphi$ is valid on \mathcal{ROG} if, and only if, $\text{not } \varphi$ is not satisfiable in \mathcal{ROG} . Thus, satisfiability of quantifier-free σ -formulas in \mathcal{ROG} and validity of universal σ -sentences on \mathcal{ROG} (i.e., membership in the universal theory of \mathcal{ROG}) are complementary computational problems.

We shall also need the notion of satisfaction of a quantifier-free σ -formula in a partial *rog* under a partial assignment, i.e., a partial function from Var into the universe of a partial *rog*.

For a partial assignment v on a partial *rog* \mathbf{B} , we recursively define, for every σ -term t , the relation $\mathbf{B} \downarrow v(t)$ (“the value of t in \mathbf{B} is defined under v ”) as follows:

$$\begin{aligned} \mathbf{B} \downarrow v(x_i) & \iff x_i \in \text{dom}(v); \\ \mathbf{B} \downarrow v(t_1 \star t_2) & \iff \mathbf{B} \downarrow v(t_1), \mathbf{B} \downarrow v(t_2) \text{ and } \langle v(t_1), v(t_2) \rangle \in \text{dom}(\star^{\mathbf{B}}), \\ & \text{where } \star \in \{\circ, \backslash, /\}. \end{aligned}$$

Next, for a partial assignment v on a partial *rog* \mathbf{B} , we recursively define, for every quantifier-free σ -formula φ , the relations $\mathbf{B} \models^v \varphi$ (“ φ is satisfied in \mathbf{B} under v ”), $\mathbf{B} \not\models^v \varphi$ (“ φ is not satisfied in \mathbf{B} under v ”) and $\mathbf{B} \approx^v \varphi$ (“the truth value of φ in \mathbf{B} ”).

under v is undefined”) as follows:

$$\begin{aligned}
 \mathbf{B} \models^v (t_1 \leq t_2) & \iff \mathbf{B} \downarrow v(t_1), \mathbf{B} \downarrow v(t_2) \text{ and } v(t_1) \leq^{\mathbf{B}} v(t_2); \\
 \mathbf{B} \not\models^v (t_1 \leq t_2) & \iff \mathbf{B} \downarrow v(t_1), \mathbf{B} \downarrow v(t_2) \text{ and } v(t_1) \not\leq^{\mathbf{B}} v(t_2); \\
 \mathbf{B} \approx^v (t_1 \leq t_2) & \text{ otherwise;} \\
 \mathbf{B} \models^v \text{not } \varphi & \iff \mathbf{B} \not\models^v \varphi; \\
 \mathbf{B} \not\models^v \text{not } \varphi & \iff \mathbf{B} \models^v \varphi; \\
 \mathbf{B} \approx^v \text{not } \varphi & \text{ otherwise;} \\
 \mathbf{B} \models^v (\varphi_1 \& \varphi_2) & \iff \mathbf{B} \models^v \varphi_1 \text{ and } \mathbf{B} \models^v \varphi_2; \\
 \mathbf{B} \not\models^v (\varphi_1 \& \varphi_2) & \iff \mathbf{B} \not\models^v \varphi_1 \text{ or } \mathbf{B} \not\models^v \varphi_2; \\
 \mathbf{B} \approx^v (\varphi_1 \& \varphi_2) & \text{ otherwise;} \\
 \mathbf{B} \models^v (\varphi_1 \text{ or } \varphi_2) & \iff \mathbf{B} \models^v \varphi_1 \text{ or } \mathbf{B} \models^v \varphi_2; \\
 \mathbf{B} \not\models^v (\varphi_1 \text{ or } \varphi_2) & \iff \mathbf{B} \not\models^v \varphi_1 \text{ and } \mathbf{B} \not\models^v \varphi_2; \\
 \mathbf{B} \approx^v (\varphi_1 \text{ or } \varphi_2) & \text{ otherwise.}
 \end{aligned}$$

We say that a quantifier-free σ -formula φ is *satisfiable* in a partial *rog* \mathbf{B} if there exists a partial assignment v on \mathbf{B} such that $\mathbf{B} \models^v \varphi$.

Intuitively, $\mathbf{B} \models^v \varphi$ (respectively, $\mathbf{B} \not\models^v \varphi$) means that the relation $\mathbf{B} \downarrow v(t)$ holds for enough terms $t \in \text{Ter}(\varphi)$ for us to be able to compute the ‘truth value’ of φ in \mathbf{B} under v as true (respectively, false).

More precisely, we use, in metalanguage, a 3-valued logic with the values ‘true’, ‘false’ and ‘undefined’ (for short, u). The 3-valued logic we use agrees with the 3-valued Łukasiewicz logic in the definitions of **not**, **&** and **or**, but differs in the definition of implication: $u \rightarrow u$ is u in the logic we use, but true in the Łukasiewicz logic. In our earlier papers (Shkatov & Van Alten, 2019, 2020, 2021), satisfaction in partial structures was defined slightly differently, in effect using a 3-valued logic where the result of a logical operation is undefined whenever at least one of the operands is undefined. The difference between the two approaches has no impact on any of our results.

We shall use the following observation in the proof of Theorem 2 below. Let \mathbf{B} be a partial *rog*, φ a quantifier-free σ -formula and v be a partial assignment on \mathbf{B} such that $\mathbf{B} \models^v \varphi$. Let v' be a (full) assignment on \mathbf{B} extending v . Observe that, if $x \notin \text{Var}(\varphi)$, then the value of x under v' is irrelevant to the satisfaction of φ in \mathbf{B} under v' . Also observe that, if $x \in \text{Var}(\varphi)$, but $x \notin \text{dom}(v)$, then the value of x under v' is, again, irrelevant to the satisfaction of φ in \mathbf{B} under v' . Formally, it can be shown by induction that if $\mathbf{B} \models^v \varphi$ (respectively, $\mathbf{B} \not\models^v \varphi$), then $\mathbf{B} \models^{v'} \varphi$ (respectively, $\mathbf{B} \not\models^{v'} \varphi$).

We now relate satisfiability in *rogs* and partial *rogs*. We define the *size* of a quantifier-free σ -formula φ , denoted $\text{size}(\varphi)$, to be the number of variables in φ plus the total number of occurrences of operation symbols in φ . To be consistent with the literature, we use the standard measure of the complexity of an input formula φ , its length, defined in the standard way as the number of occurrences of symbols in φ , and denoted by $\text{len}(\varphi)$. The running time of our algorithms is, however, more appropriately measured

by the *size* of a formula. Since, evidently, $size(\varphi) \leq len(\varphi)$, whenever we prove the existence of an algorithm running in $\mathcal{O}(f(size(\varphi)))$ we, thereby, establish the existence an algorithm running in $\mathcal{O}(f(len(\varphi)))$, whatever function f is.

The following theorem is a special case of Van Alten (2013, Theorem 2.1); we include a proof of this special case for the reader's convenience.

Theorem 2 *A quantifier-free σ -formula φ is satisfiable in \mathcal{ROG} if, and only if, it is satisfiable in a partial *rog* whose cardinality does not exceed $size(\varphi)$.*

Proof Assume $\mathbf{A} \models^v \varphi$ for some *rog* $\mathbf{A} = \langle A, \circ^{\mathbf{A}}, \backslash^{\mathbf{A}}, /^{\mathbf{A}}, \leq^{\mathbf{A}} \rangle$ and some assignment v on \mathbf{A} . Let $B = \{v(t) \mid t \in Ter(\varphi)\}$. Observe that $|B| \leq size(\varphi)$. For all $a, b \in B$ and $\star \in \{\circ, \backslash, /\}$, let $(a, b) \in \text{dom}(\star^{\mathbf{B}})$ if there exists $t_1 \star t_2 \in Ter(\varphi)$ with $a = v(t_1)$ and $b = v(t_2)$. Then, for every $\star \in \{\circ, \backslash, /\}$ and all $(a, b) \in \text{dom}(\star^{\mathbf{B}})$, set $a \star^{\mathbf{B}} b := a \star^{\mathbf{A}} b$. Lastly, set $\leq^{\mathbf{B}} = \leq^{\mathbf{A}} \upharpoonright_B$. Then $\mathbf{B} := \langle B, \circ^{\mathbf{B}}, \backslash^{\mathbf{B}}, /^{\mathbf{B}}, \leq^{\mathbf{B}} \rangle$ is a partial *rog*. Let $w := v \upharpoonright_{Var(\varphi)}$. It should be clear that $\mathbf{B} \models^w \varphi$. Thus, φ is satisfiable in a partial *rog* of the required cardinality.

Conversely, assume $\mathbf{B} \models^{v'} \varphi$ for some partial *rog* \mathbf{B} and some partial assignment v' on \mathbf{B} . Then \mathbf{B} is a partial substructure of some *rog*, say \mathbf{A} . Let v' be a (full) assignment on \mathbf{B} extending v' . Then, as observed above, $\mathbf{B} \models^{v'} \varphi$. Since \mathbf{B} is a partial substructure of \mathbf{A} , it follows that $\mathbf{A} \models^{v'} \varphi$. \square

Using the above theorem, together with the characterization of partial *rogs* given in Theorem 1, we obtain the upper bound for complexity of satisfiability in \mathcal{ROG} and hence also of the universal theory of \mathcal{ROG} .

Theorem 3 *Satisfiability of quantifier-free σ -formulas in \mathcal{ROG} is in NP. The universal theory of \mathcal{ROG} is in coNP.*

Proof Let φ be a quantifier-free σ -formula. Due to Theorem 2, to determine whether φ is satisfiable in \mathcal{ROG} it suffices to check if it is satisfiable in a partial *rog* of cardinality not exceeding $size(\varphi)$. To that end, we use the following nondeterministic algorithm.

Guess a partial σ -structure $\mathbf{B} = \langle B, \circ^{\mathbf{B}}, \backslash^{\mathbf{B}}, /^{\mathbf{B}}, \leq^{\mathbf{B}} \rangle$ with $|B| \leq size(\varphi)$ and a partial assignment v on \mathbf{B} . Check whether \mathbf{B} is a partial *rog* and whether $\mathbf{B} \models^v \varphi$. If both checks succeed, return “yes”; otherwise, return “no.”

In view of Theorem 1, to check whether \mathbf{B} is a partial *rog* it suffices to check properties (i–x), which can be done in time polynomial in $|B|$ and, hence, $size(\varphi)$. Checking whether $\mathbf{B} \models^v \varphi$ can be done in time polynomial in $size(\varphi)$. Thus, the outlined algorithm runs in time polynomial in $size(\varphi)$.

Satisfiability of quantifier-free σ -formulas in \mathcal{ROG} is, therefore, in NP. Since, as we have observed, membership of universal σ -sentences in the universal theory of \mathcal{ROG} is a complementary problem, the universal theory of \mathcal{ROG} is in coNP. This completes the proof. \square

For the lower bound, we use the following observation stating a simple condition under which the universal theory of a class of structures is coNP-hard. We say that a k -ary predicate P on a structure with domain A is *non-trivial* if $P \neq \emptyset$ and $P \neq A^k$; we say that a structure is *non-trivial* if it has a non-trivial predicate definable in its signature.

Theorem 4 *Let \mathcal{K} be a class of structures that contains a non-trivial structure. Then, satisfiability of quantifier-free first-order formulas in \mathcal{K} is NP-hard and the universal theory of \mathcal{K} is coNP-hard.*

Proof The proof is by reduction from the NP-hard satisfiability problem for Boolean formulas (Cook, 1971).

Let \mathbf{A} be a non-trivial structure in \mathcal{K} with non-trivial k -ary predicate P . Then there exist $a_1, \dots, a_k, b_1, \dots, b_k$ in the domain of \mathbf{A} with $\langle a_1, \dots, a_k \rangle \in P$ and $\langle b_1, \dots, b_k \rangle \notin P$, that is, $P(a_1, \dots, a_k)$ is true and $P(b_1, \dots, b_k)$ is false.

Let $\beta(p_1, \dots, p_n)$ be a Boolean formula. For every $i \in \{1, \dots, n\}$ and every $j \in \{1, \dots, k\}$, let q_j^i be a new variable. Recursively define the translation \cdot^* :

$$\begin{aligned} p_i^* &:= P(q_1^i, \dots, q_k^i), \quad \text{for every } i \in \{1, \dots, n\}; \\ (\neg \gamma_1)^* &:= \text{not } \gamma_1^*; \\ (\gamma_1 \wedge \gamma_2)^* &:= \gamma_1^* \ \& \ \gamma_2^*; \\ (\gamma_1 \vee \gamma_2)^* &:= \gamma_1^* \ \text{or} \ \gamma_2^*. \end{aligned}$$

Then β^* is a quantifier-free formula in the signature of \mathcal{K} . We show that β is satisfiable in the two-element Boolean algebra if, and only if, β^* is satisfiable in \mathcal{K} .

Assume β is satisfiable in the two-element Boolean algebra $\mathbf{2}$ with universe $\{0, 1\}$ under assignment v . Let w be an assignment on \mathbf{A} such that, for all $i \in \{1, \dots, n\}$ and $j \in \{1, \dots, k\}$,

$$w(q_j^i) := \begin{cases} a_j & \text{if } v(p_i) = 1; \\ b_j & \text{if } v(p_i) = 0. \end{cases}$$

Then, $\mathbf{A} \models^w p_i^*$ if $v(p_i) = 1$, and $\mathbf{A} \not\models^w p_i^*$ if $v(p_i) = 0$. It follows, recursively, that $\mathbf{A} \models^w \beta^*$.

Conversely, suppose β^* is satisfiable in \mathcal{K} . Then, $\mathbf{B} \models^w \beta^*$ for some $\mathbf{B} \in \mathcal{K}$ and assignment w on \mathbf{B} . Let v be an assignment on the Boolean algebra $\mathbf{2}$ such that

$$v(p_i) := \begin{cases} 1 & \text{if } \mathbf{B} \models^w P(q_1^i, \dots, q_k^i); \\ 0 & \text{if } \mathbf{B} \not\models^w P(q_1^i, \dots, q_k^i). \end{cases}$$

It follows that $\mathbf{2} \models^v \beta$, as required. □

From Theorems 3 and 4, which is applicable since the signature σ contains the relation symbol \leq , we obtain the following:

Theorem 5 *Satisfiability of quantifier-free σ -formulas in \mathcal{ROG} is NP-complete. The universal theory of \mathcal{ROG} is coNP-complete.*

5 Unital and Integral Residuated Ordered Groupoids

In this section we extend the methods of the previous sections to show that the universal theories of the classes of unital and integral *rogs* are coNP-complete.

We denote by σ' the signature that extends the signature σ by a constant 1. A *unital residuated ordered groupoid*, or *urog*, is a σ' -structure $\mathbf{A} = \langle A, \circ, \backslash, /, 1, \leq \rangle$, where $\langle A, \circ, \backslash, /, \leq \rangle$ is a *rog* and 1 is an identity for \circ , i.e., \mathbf{A} satisfies $x \circ 1 = x = 1 \circ x$. The class of all *urogs* is denoted by \mathcal{UROG} . An *integral residuated ordered groupoid*, or *irog*, is a *urog* satisfying $x \leq 1$. The class of all *irogs* is denoted by \mathcal{IROG} .

An embedding of a *urog* (respectively, *irog*) into another *urog* (respectively, *irog*) is an embedding of an underlying *rog* that, additionally, preserves the identity.

We extend the frame representation theory for *rogs* given in Sect. 2 to *urogs* and *irogs*.

First, we consider *urogs*. A *urog-frame* is a structure $\mathfrak{F} = \langle P, e, \leq, R \rangle$, where $\langle P, \leq, R \rangle$ is a *rog-frame* and $e \in P$ such that the following hold:

$$(\forall f \in P)[R(f, e, f) \ \& \ R(e, f, f)]; \quad (7)$$

$$(\forall f, h \in P)[R(f, e, h) \implies f \leq h]; \quad (8)$$

$$(\forall g, h \in P)[R(e, g, h) \implies g \leq h]. \quad (9)$$

Let $\langle P, e, \leq, R \rangle$ be a *urog-frame*. Since $\langle P, \leq, R \rangle$ is a *rog-frame* we may construct, as in Sect. 3, the *rog* associated with it, that is, $\langle U(P), \circ, \backslash, /, \subseteq \rangle$, where the operations \circ, \backslash and $/$ are as defined in (3–5). We observe the following result.

Lemma 1 *If $\langle P, e, \leq, R \rangle$ be a *urog-frame*, then $[e]$ is the identity for the operation \circ on $U(P)$ defined in (3).*

Proof Let $X \in U(P)$; then by (3),

$$X \circ [e] = \{h \in P \mid (\exists f \in X)(\exists g \in [e]) R(f, g, h)\}.$$

If $h \in X \circ [e]$, then there exists $f \in X$ and $g \in [e]$ such that $R(f, g, h)$. Since $e \leq g$ we have $R(f, e, h)$, by (2), hence $f \leq h$, by (8). Thus, $h \in X$, so $X \circ [e] \subseteq X$. Conversely, if $h \in X$, then, since $e \in [e]$ and $R(h, e, h)$, by (7), we get that $h \in X \circ [e]$. Thus, $X \subseteq X \circ [e]$. Therefore, $X \circ [e] = X$. We can show, similarly, that $[e] \circ X = X$, so $[e]$ is an identity for \circ . By standard algebraic methods it follows that $[e]$ is the unique identity for \circ . \square

Thus, if $\mathfrak{F} = \langle P, e, \leq, R \rangle$ is a *urog-frame*, then $\mathbf{A}_{\mathfrak{F}} := \langle U(P), \circ, \backslash, /, [e], \subseteq \rangle$ is a *urog* (with $[e] = 1$); we call it the *urog associated with \mathfrak{F}* .

If $\mathbf{A} = \langle A, \circ, \backslash, /, 1, \leq \rangle$ is a *urog*, then we can obtain a *urog-frame*, as follows. Let R be the ternary relation on $U(A)$ defined as in (6). Then the structure $\langle U(A), \subseteq, R \rangle$ is a *rog-frame*. It is straightforward to check that $\mathfrak{F}_{\mathbf{A}} := \langle U(A), [1], \subseteq, R \rangle$ satisfies (7–9); hence, it is a *urog-frame*.

We next consider the class of *irogs*. An *irog-frame* is a *urog-frame* $\mathfrak{F} = \langle P, e, \leq, R \rangle$ satisfying

$$(\forall f \in P)[e \leq f]. \quad (10)$$

For any *irog*-frame $\mathfrak{F} = \langle P, e, \leq, R \rangle$, the *urog* associated with \mathfrak{F} , namely, $\mathbf{A}_{\mathfrak{F}} = \langle U(P), \circ, \backslash, /, [e], \subseteq \rangle$, has $[e] = P$, by (10), which is clearly the greatest element of $U(P)$. Thus, $\mathbf{A}_{\mathfrak{F}}$ is an *irog*, which we call the *irog associated with \mathfrak{F}* .

Let $\mathbf{A} = \langle A, \circ, \backslash, /, 1, \leq \rangle$ be an *irog*. Let $U'(A)$ be the set of all nonempty elements of $U(A)$ and R the ternary relation on $U'(A)$ defined as in (6). Then $\mathfrak{F}'_{\mathbf{A}} = \langle U'(A), \{1\}, \subseteq, R \rangle$ is a *urog*-frame. As for *urogs*, the constant element e of $\mathfrak{F}'_{\mathbf{A}}$ is $\{1\}$, which is just $\{1\}$ in this case. In addition, every non-empty upset of A contains 1 , so $\{1\}$ is the smallest element of $U'(A)$. Thus, $\mathfrak{F}'_{\mathbf{A}}$ satisfies (10), hence it is an *irog*-frame.

A *partial σ' -structure* is a structure $\mathbf{B} = \langle B, \circ^{\mathbf{B}}, \backslash^{\mathbf{B}}, /^{\mathbf{B}}, 1^{\mathbf{B}}, \leq^{\mathbf{B}} \rangle$, where $\leq^{\mathbf{B}}$ is a binary relation on B , while $\circ^{\mathbf{B}}, \backslash^{\mathbf{B}}$ and $/^{\mathbf{B}}$ are partial binary operations on B and $1^{\mathbf{B}} \in B$. (Note that the partiality of the binary operations on \mathbf{B} does not apply to the constant $1^{\mathbf{B}}$ which is always assumed to be an element of B .) A *partial urog* is a partial σ' -structure $\mathbf{B} = \langle B, \circ^{\mathbf{B}}, \backslash^{\mathbf{B}}, /^{\mathbf{B}}, 1^{\mathbf{B}}, \leq^{\mathbf{B}} \rangle$ that is a partial substructure of a *urog*, i.e., for which there exists a *urog* $\mathbf{A} = \langle A, \circ^{\mathbf{A}}, \backslash^{\mathbf{A}}, /^{\mathbf{A}}, 1^{\mathbf{A}}, \leq^{\mathbf{A}} \rangle$ such that $B \subseteq A$, $\leq^{\mathbf{B}} = \leq^{\mathbf{A}} \upharpoonright_B$, $1^{\mathbf{B}} = 1^{\mathbf{A}}$ and $a \star^{\mathbf{B}} b = a \star^{\mathbf{A}} b$ for each $\star \in \{\circ, \backslash, /\}$ and $\langle a, b \rangle \in \text{dom}(\star^{\mathbf{B}})$. Partial *irogs* are defined analogously.

An *embedding* of a partial σ' -structure $\mathbf{B} = \langle B, \circ^{\mathbf{B}}, \backslash^{\mathbf{B}}, /^{\mathbf{B}}, \leq^{\mathbf{B}} \rangle$ into a σ' -structure $\mathbf{A} = \langle A, \circ^{\mathbf{A}}, \backslash^{\mathbf{A}}, /^{\mathbf{A}}, \leq^{\mathbf{A}} \rangle$ is a map $\alpha : B \rightarrow A$ such that: $a \leq^{\mathbf{B}} b$ if, and only if, $\alpha(a) \leq^{\mathbf{A}} \alpha(b)$, $\alpha(1^{\mathbf{B}}) = 1^{\mathbf{A}}$ and $\alpha(a \star^{\mathbf{B}} b) = \alpha(a) \star^{\mathbf{A}} \alpha(b)$ for each $\star \in \{\circ, \backslash, /\}$ and $\langle a, b \rangle \in \text{dom}(\star^{\mathbf{B}})$. As in the case of *rogs*, embeddings can easily be shown to be injective. In addition, if \mathbf{B} is a partial σ' -structure and there exists an embedding of \mathbf{B} into a *urog* (respectively, *irog*) \mathbf{A} , then \mathbf{B} is isomorphic to a partial substructure of \mathbf{A} , that is, \mathbf{B} is a partial *urog* (respectively, *irog*).

In the following two theorems we give intrinsic characterizations of, respectively, partial *urogs* and partial *irogs*.

Theorem 6 A partial σ' -structure $\mathbf{B} = \langle B, \circ^{\mathbf{B}}, \backslash^{\mathbf{B}}, /^{\mathbf{B}}, 1^{\mathbf{B}}, \leq^{\mathbf{B}} \rangle$ is a partial *urog* if, and only if, it satisfies conditions (i–x) of Theorem 1 and

- (xi) $(\forall a \in B)[\langle a, 1^{\mathbf{B}} \rangle \in \text{dom}(\circ^{\mathbf{B}}) \implies a \circ^{\mathbf{B}} 1^{\mathbf{B}} = a]$;
- (xii) $(\forall a \in B)[\langle 1^{\mathbf{B}}, a \rangle \in \text{dom}(\circ^{\mathbf{B}}) \implies 1^{\mathbf{B}} \circ^{\mathbf{B}} a = a]$.

Proof As observed in the proof of Theorem 1, conditions (i–x) hold in every partial *rog*, hence also in every partial *urog*. Conditions (xi) and (xii) hold in every *urog*; since they are universal sentences with quantifiers relativized by the domains of the partial operations, it follows that they hold in every partial *urog*.

Conversely, suppose $\mathbf{B} = \langle B, \circ^{\mathbf{B}}, \backslash^{\mathbf{B}}, /^{\mathbf{B}}, 1^{\mathbf{B}}, \leq^{\mathbf{B}} \rangle$ is a partial σ' -structure satisfying the requirements of the theorem. For every $a \in B$, if $\langle 1^{\mathbf{B}}, a \rangle \notin \text{dom}(\circ^{\mathbf{B}})$, then we extend the domain of $\circ^{\mathbf{B}}$ to include $\langle 1^{\mathbf{B}}, a \rangle$ and set $1^{\mathbf{B}} \circ^{\mathbf{B}} a = a$. We deal similarly with every $a \in B$ for which $\langle a, 1^{\mathbf{B}} \rangle \notin \text{dom}(\circ^{\mathbf{B}})$. Then, by (xi) and (xii), $a \circ^{\mathbf{B}} 1^{\mathbf{B}} = a = 1^{\mathbf{B}} \circ^{\mathbf{B}} a$ for every $a \in B$.

As in Theorem 1, we define a ternary relation $R^{\mathbf{B}}$ on $U(B)$ as follows:

$$\begin{aligned}
 R^{\mathbf{B}}(f, g, h) \iff & (\forall \langle a, b \rangle \in \text{dom}(\circ^{\mathbf{B}})) [a \in f \ \& \ b \in g \implies a \circ^{\mathbf{B}} b \in h] \\
 & \& (\forall \langle a, b \rangle \in \text{dom}(\backslash^{\mathbf{B}})) [a \in f \ \& \ a \backslash^{\mathbf{B}} b \in g \implies b \in h] \\
 & \& (\forall \langle a, b \rangle \in \text{dom}(/^{\mathbf{B}})) [a /^{\mathbf{B}} b \in f \ \& \ b \in g \implies a \in h].
 \end{aligned}$$

Let $\mathfrak{F} := \langle U(B), [1^{\mathbf{B}}], \subseteq, R^{\mathbf{B}} \rangle$; we show that \mathfrak{F} is a *urog*-frame. As noted in the proof of Theorem 1, $\langle U(B), \subseteq, R^{\mathbf{B}} \rangle$ is a *rog*-frame. We next show that \mathfrak{F} is a *urog*-frame, that is, it satisfies (7–9).

To see that \mathfrak{F} satisfies (7), let $f \in U(B)$; we must show that $R^{\mathbf{B}}(f, [1^{\mathbf{B}}], f)$ and $R^{\mathbf{B}}([1^{\mathbf{B}}], f, f)$. Suppose that $\langle a, b \rangle \in \text{dom}(\circ^{\mathbf{B}})$ for some $a \in f$ and $b \in [1^{\mathbf{B}}]$. Then, $1^{\mathbf{B}} \leq^{\mathbf{B}} b$, so $a \circ^{\mathbf{B}} 1^{\mathbf{B}} \leq^{\mathbf{B}} a \circ^{\mathbf{B}} b$, by (ii), hence $a \leq^{\mathbf{B}} a \circ^{\mathbf{B}} b$, so $a \circ^{\mathbf{B}} b \in f$. Next, suppose that $\langle a, b \rangle \in \text{dom}(\backslash^{\mathbf{B}})$, where $a \in f$ and $a \backslash^{\mathbf{B}} b \in [1^{\mathbf{B}}]$. By an application of (iii), since $a \leq^{\mathbf{B}} a$ and $1^{\mathbf{B}} \leq^{\mathbf{B}} a \backslash^{\mathbf{B}} b$, we obtain that $a \circ^{\mathbf{B}} 1^{\mathbf{B}} \leq^{\mathbf{B}} b$, i.e., $a \leq^{\mathbf{B}} b$. Thus, $b \in f$, as required. Lastly, suppose that $\langle a, b \rangle \in \text{dom}(/^{\mathbf{B}})$, where $a /^{\mathbf{B}} b \in f$ and $b \in [1^{\mathbf{B}}]$. By an application of (iv), since $a /^{\mathbf{B}} b \leq^{\mathbf{B}} a /^{\mathbf{B}} b$ and $1^{\mathbf{B}} \leq^{\mathbf{B}} b$, we obtain that $(a /^{\mathbf{B}} b) \circ^{\mathbf{B}} 1^{\mathbf{B}} \leq^{\mathbf{B}} a$, i.e., $a /^{\mathbf{B}} b \leq^{\mathbf{B}} a$. Thus, $a \in f$, as required. This shows that $R^{\mathbf{B}}(f, [1^{\mathbf{B}}], f)$; a similar proof shows that $R^{\mathbf{B}}([1^{\mathbf{B}}], f, f)$.

To see that \mathfrak{F} satisfies (8), let $f, h \in U(B)$ such that $R^{\mathbf{B}}(f, [1^{\mathbf{B}}], h)$; we must show that $f \subseteq h$. Let $a \in f$. Since $1^{\mathbf{B}} \in [1^{\mathbf{B}}]$ and $\langle a, 1^{\mathbf{B}} \rangle \in \text{dom}(\circ^{\mathbf{B}})$, we get that $a \circ^{\mathbf{B}} 1^{\mathbf{B}} \in h$. Since $a \circ^{\mathbf{B}} 1^{\mathbf{B}} = a$, it follows that $a \in h$, as required. A similar argument shows that \mathfrak{F} satisfies (9).

Let $\mathbf{A}_{\mathfrak{F}} = \langle U(U(B)), \circ, \backslash, /, [[1^{\mathbf{B}}]], \subseteq \rangle$ be the *urog* associated with \mathfrak{F} . It can be shown, as in the proof of Theorem 1, that the map $\mu: B \rightarrow U(U(B))$ defined by $\mu(a) = \{f \in U(B) \mid a \in f\}$ is an embedding of \mathbf{B} into $\mathbf{A}_{\mathfrak{F}}$; we need only check additionally that $\mu(1^{\mathbf{B}}) = [[1^{\mathbf{B}}]]$, which is immediate. This completes the proof. \square

Theorem 7 A partial σ' -structure $\mathbf{B} = \langle B, \circ^{\mathbf{B}}, \backslash^{\mathbf{B}}, /^{\mathbf{B}}, 1^{\mathbf{B}}, \leq^{\mathbf{B}} \rangle$ is a partial *irog* if, and only if, it satisfies conditions (i–x) of Theorem 1, conditions (xi) and (xii) of Theorem 6 and

$$(xiii) (\forall a \in B)[a \leq^{\mathbf{B}} 1^{\mathbf{B}}].$$

Proof Conditions (i–xiii) hold in any *irog*, hence also in every partial *irog*.

Conversely, suppose $\mathbf{B} = \langle B, \circ^{\mathbf{B}}, \backslash^{\mathbf{B}}, /^{\mathbf{B}}, 1^{\mathbf{B}}, \leq^{\mathbf{B}} \rangle$ is a partial σ' -structure satisfying the requirements of the theorem. For every $a \in B$, if $\langle 1^{\mathbf{B}}, a \rangle \notin \text{dom}(\circ^{\mathbf{B}})$, then we extend the domain of $\circ^{\mathbf{B}}$ to include $\langle 1^{\mathbf{B}}, a \rangle$ and set $1^{\mathbf{B}} \circ^{\mathbf{B}} a = a$. We deal similarly with every $a \in B$ for which $\langle a, 1^{\mathbf{B}} \rangle \notin \text{dom}(\circ^{\mathbf{B}})$.

Let $U'(B)$ be the set of all nonempty upsets of $\langle B, \leq^{\mathbf{B}} \rangle$. Define a ternary relation $R^{\mathbf{B}}$ on $U'(B)$ as in Theorems 1 and 6. It can be shown, as in the proof of Theorem 6, that $\mathfrak{F}' = \langle U'(B), \{1^{\mathbf{B}}\}, \subseteq, R^{\mathbf{B}} \rangle$ is a *urog*-frame. By (xiii), every element of $U'(B)$ contains $1^{\mathbf{B}}$, hence $\{1^{\mathbf{B}}\}$ is the smallest set in $U'(B)$, that is, \mathfrak{F}' satisfies (10); hence, \mathfrak{F}' is an *irog*-frame.

Let $\mathbf{A}_{\mathfrak{F}'} = \langle U(U'(B)), \circ, \backslash, /, [[1^{\mathbf{B}}]], \subseteq \rangle$ be the *irog* associated with \mathfrak{F}' . Then the map $\mu: B \rightarrow U(U'(B))$ defined by $\mu(a) = \{f \in U'(B) \mid a \in f\}$ is an embedding of \mathbf{B} into $\mathbf{A}'_{\mathfrak{F}'}$. This follows as in the proofs of Theorems 1 and 6; we need only check additionally that the upsets utilised in the proofs are nonempty. In the proof of Theorem 1, where it is shown that $g \notin \mu(a \backslash^{\mathbf{B}} b)$ implies $g \notin \mu(a) \backslash \mu(b)$, the set $h = \{e \in B \mid e \not\leq^{\mathbf{B}} b\}$ is used. This set is empty in the case that $b = 1^{\mathbf{B}}$, however, we show that it will never be the case that $g \notin \mu(a \backslash^{\mathbf{B}} 1^{\mathbf{B}})$. Using property (v), since $a \leq^{\mathbf{B}} a$ and $a \circ^{\mathbf{B}} 1^{\mathbf{B}} \leq^{\mathbf{B}} 1^{\mathbf{B}}$, we get $1^{\mathbf{B}} \leq^{\mathbf{B}} a \backslash^{\mathbf{B}} 1^{\mathbf{B}}$. Thus, $a \backslash^{\mathbf{B}} 1^{\mathbf{B}} = 1^{\mathbf{B}}$, so $a \backslash^{\mathbf{B}} 1^{\mathbf{B}} \in g$ for every nonempty upset g , and so $g \in \mu(a \backslash^{\mathbf{B}} 1^{\mathbf{B}})$. A similar situation exists for $/^{\mathbf{B}}$. \square

Quantifier-free σ' -formulas and universal σ' -sentences are defined analogously to quantifier-free σ -formulas and universal σ -sentences; these may contain the constant 1. The notations and conventions used in this section agree with those introduced in Sect. 4.

Assignments on *urogs* and *irogs*, as well as partial assignments on partial *urogs* and *irogs*, are required to assign the identity of the structure to the constant 1. The definitions of satisfaction of quantifier-free σ' -formulas in *urogs* and *irogs*, as well as in partial *urogs* and *irogs*, are analogous to those given in Sect. 4 for *rogs*.

The *size* of a quantifier-free σ' -formula φ , denoted $size(\varphi)$, is the number of variables in φ plus the total number of occurrences of operation symbols in φ , plus one (for the constant). The following result is an analogue of Theorem 2.

Theorem 8 *A quantifier-free σ' -formula φ is satisfiable in $UROG$ (respectively, $IROG$) if, and only if, it is satisfiable in a partial *urog* (respectively, *irog*) whose cardinality does not exceed $size(\varphi)$.*

Relying on the characterizations of partial *urogs* and partial *irogs* from Theorems 6 and 7, we obtain the NP-upper bound for satisfiability in $UROG$ and $IROG$ using an argument analogous to the proof of Theorem 5. The NP-lower bound for satisfiability in $UROG$ and $IROG$ follows from Theorem 4.

Theorem 9 *Satisfiability of quantifier-free σ' -formulas in both $UROG$ and $IROG$ is NP-complete. The universal theory of both $UROG$ and $IROG$ is coNP-complete.*

6 Residuated k -Algebras

In this section and the next, we consider generalizations of *rogs* in which the \circ operation is replaced by an operation \mathbf{t} of arity k , where $k \geq 1$, and $\backslash, /$ are replaced by k operations $\mathbf{r}_1, \dots, \mathbf{r}_k$ of arity k , such that k -ary versions of the residuation property (1) hold. We shall refer to such a \mathbf{t} as a (k -ary) ‘fusion’ operation and $\mathbf{r}_1, \dots, \mathbf{r}_k$ as its ‘residuals’. In Sect. 8 we shall consider ‘residuated algebras’ that may contain finitely many fusion operations of varying arities together with their residuals. Such algebras were originally defined by Buszkowski (1989) [see also (Buszkowski, 1998, 2011; Kołowska-Gawiejnowicz, 1997; Kandulski, 1997; Jäger, 2004)].

Let k be an integer with $k \geq 1$. We use σ_k to denote the signature consisting of $k + 1$ operation symbols $\mathbf{t}, \mathbf{r}_1, \dots, \mathbf{r}_k$, each of arity k , together with a binary relation symbol \leq . A *residuated k -algebra*, or ra_k , is a σ_k -structure $\mathbf{A} = \langle A, \mathbf{t}, \mathbf{r}_1, \dots, \mathbf{r}_k, \leq \rangle$, where $\langle A, \leq \rangle$ is a partially ordered set and \mathbf{A} satisfies the following k -ary residuation property: for all $a_1, \dots, a_k, c \in A$ and every $j \in \{1, \dots, k\}$,

$$\mathbf{t}(a_1, \dots, a_k) \leq c \iff a_j \leq \mathbf{r}_j(a_1, \dots, a_{j-1}, c, a_{j+1}, \dots, a_k). \tag{11}$$

To make the representation of the properties of ra_k s briefer, we use the following notation: given a non-empty set A , a tuple $\mathbf{a} \in A^k$, as well as $i \in \{1, \dots, k\}$ and $c \in A$, we denote by $\mathbf{a}[i := c]$ the k -tuple obtained from \mathbf{a} by replacing its i^{th} component a_i

by c , i.e., the k -tuple $\langle a_1, \dots, a_{i-1}, c, a_{i+1}, \dots, a_k \rangle$. Then (11) can be written as

$$\mathbf{t}(\mathbf{a}) \leq c \iff a_j \leq \mathbf{r}_j(\mathbf{a}[j := c]).$$

First, we prove some properties of residuated k -algebras.

Lemma 2 Let $\mathbf{A} = \langle A, \mathbf{t}, \mathbf{r}_1, \dots, \mathbf{r}_k, \leq \rangle$ be a ra_k . Then the following hold.

(a) \mathbf{t} is monotone in each coordinate, i.e., for any $b \in A$, $\mathbf{a} \in A^k$ and $i \in \{1, \dots, k\}$,

$$b \leq a_i \implies \mathbf{t}(\mathbf{a}[i := b]) \leq \mathbf{t}(\mathbf{a}).$$

(b) For every $j \in \{1, \dots, k\}$, \mathbf{r}_j is monotone in the j^{th} coordinate, i.e., for any $b \in A$ and $\mathbf{a} \in A^k$,

$$a_j \leq b \implies \mathbf{r}_j(\mathbf{a}) \leq \mathbf{r}_j(\mathbf{a}[j := b]).$$

(c) For every $j \in \{1, \dots, k\}$ and $\mathbf{a} \in A^k$,

$$\mathbf{t}(\mathbf{a}[j := \mathbf{r}_j(\mathbf{a})]) \leq a_j.$$

(d) For every $j \in \{1, \dots, k\}$ and $i \in \{1, \dots, k\} - \{j\}$, \mathbf{r}_j is antitone in the i^{th} coordinate, i.e., for any $b \in A$ and $\mathbf{a} \in A^k$,

$$b \leq a_i \implies \mathbf{r}_j(\mathbf{a}) \leq \mathbf{r}_j(\mathbf{a}[i := b]).$$

Proof (a) Observe that $\mathbf{t}(\mathbf{a}) \leq \mathbf{t}(\mathbf{a})$ implies, by (11), $a_i \leq \mathbf{r}_i(\mathbf{a}[i := \mathbf{t}(\mathbf{a})])$, whence $b \leq \mathbf{r}_i(\mathbf{a}[i := \mathbf{t}(\mathbf{a})])$, and so, by (11), $\mathbf{t}(\mathbf{a}[i := b]) \leq \mathbf{t}(\mathbf{a})$.

(b) Observe that $\mathbf{r}_j(\mathbf{a}) \leq \mathbf{r}_j(\mathbf{a})$ implies, by (11), $\mathbf{t}(\mathbf{a}[j := \mathbf{r}_j(\mathbf{a})]) \leq a_j$, whence $\mathbf{t}(\mathbf{a}[j := \mathbf{r}_j(\mathbf{a})]) \leq b$, and so, by (11), $\mathbf{r}_j(\mathbf{a}) \leq \mathbf{r}_j(\mathbf{a}[j := b])$.

(c) Observe that $\mathbf{r}_j(\mathbf{a}) \leq \mathbf{r}_j(\mathbf{a})$ implies, by (11), $\mathbf{t}(\mathbf{a}[j := \mathbf{r}_j(\mathbf{a})]) \leq a_j$.

(d) If $b \leq a_i$ then, by (a), $\mathbf{t}(\mathbf{a}[j := \mathbf{r}_j(\mathbf{a})][i := b]) \leq \mathbf{t}(\mathbf{a}[j := \mathbf{r}_j(\mathbf{a})])$, hence, by (c), $\mathbf{t}(\mathbf{a}[j := \mathbf{r}_j(\mathbf{a})][i := b]) \leq a_j$. Thus, by (11), $\mathbf{r}_j(\mathbf{a}) \leq \mathbf{r}_j(\mathbf{a}[i := b])$. \square

Henceforth, given $i, j, \ell \in \{1, \dots, k\}$, we often write, for brevity, $i \neq j$ instead of $i \in \{1, \dots, k\} - \{j\}$ and $i \neq j, i \neq \ell$ instead of $i \in \{1, \dots, k\} - \{j, \ell\}$.

Lemma 3 Let k be an integer with $k \geq 1$ and $\mathbf{A} = \langle A, \mathbf{t}, \mathbf{r}_1, \dots, \mathbf{r}_k, \leq \rangle$ a ra_k . Then, for all $\mathbf{a}, \mathbf{c} \in A^k$ and $j, \ell \in \{1, \dots, k\}$,

- (a) $\bigwedge_{i=1}^k a_i \leq c_i \implies \mathbf{t}(\mathbf{a}) \leq \mathbf{t}(\mathbf{c});$
- (b) $\bigwedge_{i \neq j} a_i \leq c_i \ \& \ a_j \leq \mathbf{r}_j(\mathbf{c}) \implies \mathbf{t}(\mathbf{a}) \leq c_j;$
- (c) $\bigwedge_{i \neq j} a_i \leq c_i \ \& \ \mathbf{t}(\mathbf{c}) \leq a_j \implies c_j \leq \mathbf{r}_j(\mathbf{a});$
- (d) $\bigwedge_{i \neq j} a_i \leq c_i \ \& \ c_j \leq a_j \implies \mathbf{r}_j(\mathbf{c}) \leq \mathbf{r}_j(\mathbf{a});$
- (e) $\bigwedge_{i \neq j, i \neq \ell} a_i \leq c_i \ \& \ a_\ell \leq \mathbf{r}_\ell(\mathbf{c}) \ \& \ c_\ell \leq a_j \implies c_j \leq \mathbf{r}_j(\mathbf{a}).$

Proof (a) This follows by repeated applications of Lemma 2(a).
 (b) By the assumptions and Lemma 2(a), $\mathbf{t}(\mathbf{a}) \leq \mathbf{t}(\mathbf{c}[j := \mathbf{r}_j(\mathbf{c})])$. Since, by Lemma 2(c), $\mathbf{t}(\mathbf{c}[j := \mathbf{r}_j(\mathbf{c})]) \leq c_j$, it follows that $\mathbf{t}(\mathbf{a}) \leq c_j$.
 (c) By the assumptions and Lemma 2(a), $\mathbf{t}(\mathbf{a}[j := c_j]) \leq \mathbf{t}(\mathbf{c})$. Thus, by (11), $c_j \leq \mathbf{r}_j(\mathbf{a}[j := \mathbf{t}(\mathbf{c})])$. By the assumption that $\mathbf{t}(\mathbf{c}) \leq a_j$ and Lemma 2(b), $\mathbf{r}_j(\mathbf{a}[j := \mathbf{t}(\mathbf{c})]) \leq \mathbf{r}_j(\mathbf{a})$. Thus, $c_j \leq \mathbf{r}_j(\mathbf{a})$.
 (d) This follows by applications of Lemma 2(b) and Lemma 2(d).
 (e) From the assumption that $c_i \leq a_i$, for every $i \in \{1, \dots, k\} - \{j, \ell\}$, we obtain, by Lemma 2(b), $\mathbf{r}_\ell(\mathbf{c}) \leq \mathbf{r}_\ell(\mathbf{a}[j := c_j][\ell := c_\ell])$. Since, by assumption, $a_\ell \leq \mathbf{r}_\ell(\mathbf{c})$, we obtain $a_\ell \leq \mathbf{r}_\ell(\mathbf{a}[j := c_j][\ell := c_\ell])$, hence, by (11), $\mathbf{t}(\mathbf{a}[j := c_j]) \leq c_\ell$. Since, by assumption $c_\ell \leq a_j$, we obtain $\mathbf{t}(\mathbf{a}[j := c_j]) \leq a_j$, hence, by (11), $c_j \leq \mathbf{r}_j(\mathbf{a})$. \square

We extend to ra_k s the frame representation theory for $rogs$ outlined in Sect. 2. A ra_k -frame is a structure $\mathfrak{F} = \langle P, \leq, R \rangle$, where $\langle P, \leq \rangle$ is a partially ordered set and R is a $(k + 1)$ -ary relation on P monotone in the last coordinate and antitone in the other coordinates, i.e., such that, for all $f_1, f'_1, \dots, f_k, f'_k, h, h' \in P$,

$$R(f_1, \dots, f_k, h) \ \& \ \bigwedge_{i=1}^k f'_i \leq f_i \ \& \ h \leq h' \implies R(f'_1, \dots, f'_k, h'). \tag{12}$$

Let $\mathfrak{F} = \langle P, \leq, R \rangle$ be a ra_k -frame and define, for all $X_1, \dots, X_k \in U(P)$ and $j \in \{1, \dots, k\}$,

$$\mathbf{t}(X_1, \dots, X_k) := \{h \in P \mid (\exists f_1, \dots, f_k \in P) [\bigwedge_{i=1}^k f_i \in X_i \ \& \ R(f_1, \dots, f_k, h)]\};$$

and

$$\begin{aligned} \mathbf{r}_j(X_1, \dots, X_k) &:= \{f_j \in P \mid (\forall f_{i \neq j}, h \in P) \\ &\quad \& \ f_i \in X_i \ \& \ R(f_1, \dots, f_k, h) \implies h \in X_j\}. \end{aligned}$$

Since \mathfrak{F} satisfies (12), so defined $\mathbf{t}, \mathbf{r}_1, \dots, \mathbf{r}_k$ are operations on $U(P)$. The definitions of $\mathbf{t}, \mathbf{r}_1, \dots, \mathbf{r}_k$ ensure that (11) is satisfied with respect to the partial order \subseteq on $U(P)$. Thus, $\mathbf{A}_{\mathfrak{F}} = \langle U(P), \mathbf{t}, \mathbf{r}_1, \dots, \mathbf{r}_k, \subseteq \rangle$ is a ra_k , which we call the ra_k associated with \mathfrak{F} .

If $\mathbf{A} = \langle A, \mathbf{t}, \mathbf{r}_1, \dots, \mathbf{r}_k, \leq \rangle$ is a ra_k , then we may obtain a ra_k -frame as follows. Define a $(k + 1)$ -ary relation R on $U(A)$ by

$$R(f_1, \dots, f_k, h) \iff (\forall \mathbf{a} \in A^k) \left[\bigwedge_{i=1}^k a_i \in f_i \implies \mathbf{t}(\mathbf{a}) \in h \right].$$

Then $\mathfrak{F}_{\mathbf{A}} = \langle U(A), \subseteq, R \rangle$ is a ra_k -frame.

7 Partial σ_k -Structures and Partial ra_k s

Let k be an integer with $k \geq 1$. A *partial σ_k -structure* is a structure $\mathbf{B} = \langle B, \mathbf{t}^{\mathbf{B}}, \mathbf{r}_1^{\mathbf{B}}, \dots, \mathbf{r}_k^{\mathbf{B}}, \leq^{\mathbf{B}} \rangle$, where $\leq^{\mathbf{B}}$ is a binary relation on B and $\mathbf{t}^{\mathbf{B}}$ and $\mathbf{r}_j^{\mathbf{B}}$, for every $j \in \{1, \dots, k\}$, are partial k -ary operations on B , i.e., partial functions from B^k into B . The domains of $\mathbf{t}^{\mathbf{B}}$ and $\mathbf{r}_j^{\mathbf{B}}$, for every $j \in \{1, \dots, k\}$, are denoted by, respectively, $\text{dom}(\mathbf{t}^{\mathbf{B}})$ and $\text{dom}(\mathbf{r}_j^{\mathbf{B}})$. A *partial ra_k* is a partial σ_k -structure $\mathbf{B} = \langle B, \mathbf{t}^{\mathbf{B}}, \mathbf{r}_1^{\mathbf{B}}, \dots, \mathbf{r}_k^{\mathbf{B}}, \leq^{\mathbf{B}} \rangle$ that is a partial substructure of a ra_k , i.e., for which there exists a ra_k $\mathbf{A} = \langle A, \mathbf{t}^{\mathbf{A}}, \mathbf{r}_1^{\mathbf{A}}, \dots, \mathbf{r}_k^{\mathbf{A}}, \leq^{\mathbf{A}} \rangle$ such that $B \subseteq A$, $\leq^{\mathbf{B}} = \leq^{\mathbf{A}} \upharpoonright_B$ and

$$\begin{aligned} \mathbf{t}^{\mathbf{B}}(\mathbf{a}) &= \mathbf{t}^{\mathbf{A}}(\mathbf{a}) \text{ for all } \mathbf{a} \in \text{dom}(\mathbf{t}^{\mathbf{B}}); \\ \mathbf{r}_j^{\mathbf{B}}(\mathbf{a}) &= \mathbf{r}_j^{\mathbf{A}}(\mathbf{a}) \text{ for every } j \in \{1, \dots, k\} \text{ and all } \mathbf{a} \in \text{dom}(\mathbf{r}_j^{\mathbf{B}}). \end{aligned}$$

By an *embedding* of a partial σ_k -structure $\mathbf{B} = \langle B, \mathbf{t}^{\mathbf{B}}, \mathbf{r}_1^{\mathbf{B}}, \dots, \mathbf{r}_k^{\mathbf{B}}, \leq^{\mathbf{B}} \rangle$ into a σ_k -structure $\mathbf{A} = \langle A, \mathbf{t}^{\mathbf{A}}, \mathbf{r}_1^{\mathbf{A}}, \dots, \mathbf{r}_k^{\mathbf{A}}, \leq^{\mathbf{A}} \rangle$ we mean a map $\alpha : B \rightarrow A$ such that $a \leq^{\mathbf{B}} b$ if, and only if, $\alpha(a) \leq^{\mathbf{A}} \alpha(b)$, and

$$\begin{aligned} \alpha(\mathbf{t}^{\mathbf{B}}(\mathbf{a})) &= \mathbf{t}^{\mathbf{A}}(\alpha(\mathbf{a})) \text{ for all } \mathbf{a} \in \text{dom}(\mathbf{t}^{\mathbf{B}}); \\ \alpha(\mathbf{r}_j^{\mathbf{B}}(\mathbf{a})) &= \mathbf{r}_j^{\mathbf{A}}(\alpha(\mathbf{a})) \text{ for every } j \in \{1, \dots, k\} \text{ and all } \mathbf{a} \in \text{dom}(\mathbf{r}_j^{\mathbf{B}}), \\ &\text{where } \alpha(\mathbf{a}) \text{ denotes } \langle \alpha(a_1), \dots, \alpha(a_k) \rangle. \end{aligned}$$

For clarity we note that the last condition in the above definition of an embedding is necessary as it does not follow from the other conditions.

We now give an intrinsic characterization of partial ra_k s, similar to the characterization of partial $rogs$ given in Theorem 1.

Theorem 10 *Let k be an integer with $k \geq 1$. A partial σ_k -structure $\mathbf{B} = \langle B, \mathbf{t}^{\mathbf{B}}, \mathbf{r}_1^{\mathbf{B}}, \dots, \mathbf{r}_k^{\mathbf{B}}, \leq^{\mathbf{B}} \rangle$ is a partial ra_k if, and only if, it satisfies the following conditions, for all $j, \ell \in \{1, \dots, k\}$:*

- (i') $\langle B, \leq^{\mathbf{B}} \rangle$ is a partially ordered set;
- (ii') $(\forall \mathbf{a}, \mathbf{c} \in \text{dom}(\mathbf{t}^{\mathbf{B}})) [\bigwedge_{i=1}^k a_i \leq^{\mathbf{B}} c_i \implies \mathbf{t}^{\mathbf{B}}(\mathbf{a}) \leq^{\mathbf{B}} \mathbf{t}^{\mathbf{B}}(\mathbf{c})]$;
- (iii') $(\forall \mathbf{a} \in \text{dom}(\mathbf{t}^{\mathbf{B}})) (\forall \mathbf{c} \in \text{dom}(\mathbf{r}_j^{\mathbf{B}}))$
 $[\bigwedge_{i \neq j} a_i \leq^{\mathbf{B}} c_i \ \& \ a_j \leq^{\mathbf{B}} \mathbf{r}_j^{\mathbf{B}}(\mathbf{c}) \implies \mathbf{t}^{\mathbf{B}}(\mathbf{a}) \leq^{\mathbf{B}} c_j]$;
- (iv') $(\forall \mathbf{c} \in \text{dom}(\mathbf{t}^{\mathbf{B}})) (\forall \mathbf{a} \in \text{dom}(\mathbf{r}_j^{\mathbf{B}}))$
 $[\bigwedge_{i \neq j} a_i \leq^{\mathbf{B}} c_i \ \& \ \mathbf{t}^{\mathbf{B}}(\mathbf{c}) \leq^{\mathbf{B}} a_j \implies c_j \leq^{\mathbf{B}} \mathbf{r}_j^{\mathbf{B}}(\mathbf{a})]$;
- (v') $(\forall \mathbf{c} \in \text{dom}(\mathbf{r}_j^{\mathbf{B}})) (\forall \mathbf{a} \in \text{dom}(\mathbf{r}_j^{\mathbf{B}}))$
 $[\bigwedge_{i \neq j} a_i \leq^{\mathbf{B}} c_i \ \& \ c_j \leq^{\mathbf{B}} a_j \implies \mathbf{r}_j^{\mathbf{B}}(\mathbf{c}) \leq^{\mathbf{B}} \mathbf{r}_j^{\mathbf{B}}(\mathbf{a})]$;
- (vi') $(\forall \mathbf{c} \in \text{dom}(\mathbf{r}_\ell^{\mathbf{B}})) (\forall \mathbf{a} \in \text{dom}(\mathbf{r}_j^{\mathbf{B}})) (\forall \mathbf{c} \in \text{dom}(\mathbf{r}_\ell^{\mathbf{B}}))$
 $[\bigwedge_{i \neq j, i \neq \ell} a_i \leq^{\mathbf{B}} c_i \ \& \ a_\ell \leq^{\mathbf{B}} \mathbf{r}_\ell^{\mathbf{B}}(\mathbf{c}) \ \& \ c_\ell \leq^{\mathbf{B}} a_j \implies c_j \leq^{\mathbf{B}} \mathbf{r}_j^{\mathbf{B}}(\mathbf{a})]$.

Proof By Lemma 3, conditions (i'–vi') hold in every ra_k , hence also in every partial ra_k .

Conversely, suppose $\mathbf{B} = \langle B, \mathbf{r}_1^{\mathbf{B}}, \dots, \mathbf{r}_k^{\mathbf{B}}, \leq^{\mathbf{B}} \rangle$ is a partial σ_k -structure satisfying (*i'*-*vi'*). Define a $(k + 1)$ -ary relation $R^{\mathbf{B}}$ on $U(B)$ by

$$R^{\mathbf{B}}(f_1, \dots, f_k, h) \iff (\forall \mathbf{a} \in \text{dom}(\mathbf{t}^{\mathbf{B}})) [\&_{i=1}^k a_i \in f_i \implies \mathbf{t}^{\mathbf{B}}(\mathbf{a}) \in h] \& \&_{j=1}^k (\forall \mathbf{a} \in \text{dom}(\mathbf{r}_j^{\mathbf{B}})) [\&_{i \neq j} a_i \in f_i \& \mathbf{r}_j^{\mathbf{B}}(\mathbf{a}) \in f_j \implies a_j \in h].$$

It is straightforward to check that $\mathfrak{F} = \langle U(B), \subseteq, R^{\mathbf{B}} \rangle$ is a ra_k -frame. Let $\mathbf{A} = \langle U(U(B)), \mathbf{t}, \mathbf{r}_1, \dots, \mathbf{r}_k \rangle$ be the ra_k associated with \mathfrak{F} , and let $\mu : B \rightarrow U(U(B))$ be the map defined by $\mu(a) = \{f \in U(B) \mid a \in f\}$. We show that μ is an embedding of \mathbf{B} into \mathbf{A} . For brevity, for $\mathbf{a} \in B^k$, we write $\mu(\mathbf{a})$ instead of $\langle \mu(a_1), \dots, \mu(a_k) \rangle$.

The proof that $a \leq^{\mathbf{B}} b$ if, and only if, $\mu(a) \subseteq \mu(b)$ is as in the proof of Theorem 1. Let $\mathbf{a} \in \text{dom}(\mathbf{t}^{\mathbf{B}})$. We show that $\mu(\mathbf{t}^{\mathbf{B}}(\mathbf{a})) = \mathbf{t}(\mu(\mathbf{a}))$, where

$$\mu(\mathbf{t}^{\mathbf{B}}(\mathbf{a})) = \{h \in U(B) \mid \mathbf{t}^{\mathbf{B}}(\mathbf{a}) \in h\}$$

and

$$\mathbf{t}(\mu(\mathbf{a})) = \{h \in U(B) \mid (\exists f_1, \dots, f_k \in U(B)) [\&_{i=1}^k a_i \in f_i \& R^{\mathbf{B}}(f_1, \dots, f_k, h)]\}.$$

First, assume $h \in \mathbf{t}(\mu(\mathbf{a}))$. Then there exist $f_1, \dots, f_k \in U(B)$ such that $a_i \in f_i$, for every $i \in \{1, \dots, k\}$, and $R^{\mathbf{B}}(f_1, \dots, f_k, h)$. Thus, by definition of $R^{\mathbf{B}}$, we obtain $\mathbf{t}^{\mathbf{B}}(\mathbf{a}) \in h$ and so $h \in \mu(\mathbf{t}^{\mathbf{B}}(\mathbf{a}))$. Therefore, $\mathbf{t}(\mu(\mathbf{a})) \subseteq \mu(\mathbf{t}^{\mathbf{B}}(\mathbf{a}))$.

Conversely, assume $h \in \mu(\mathbf{t}^{\mathbf{B}}(\mathbf{a}))$, i.e., $\mathbf{t}^{\mathbf{B}}(\mathbf{a}) \in h$. Let $f_i := [a_i]$, for every $i \in \{1, \dots, k\}$; these sets are clearly in $U(B)$. We show that $R^{\mathbf{B}}(f_1, \dots, f_k, h)$. Suppose $\mathbf{c} \in \text{dom}(\mathbf{t}^{\mathbf{B}})$ and $c_i \in f_i$, for every $i \in \{1, \dots, k\}$. Then, $a_i \leq^{\mathbf{B}} c_i$, for every $i \in \{1, \dots, k\}$; hence, by (*ii'*), $\mathbf{t}^{\mathbf{B}}(\mathbf{a}) \leq^{\mathbf{B}} \mathbf{t}^{\mathbf{B}}(\mathbf{c})$. Since $\mathbf{t}^{\mathbf{B}}(\mathbf{a}) \in h$ and $h \in U(B)$, we obtain $\mathbf{t}^{\mathbf{B}}(\mathbf{c}) \in h$. Suppose $j \in \{1, \dots, k\}$, $\mathbf{c} \in \text{dom}(\mathbf{r}_j^{\mathbf{B}})$, as well as $c_i \in f_i$, for every $i \neq j$, and $\mathbf{r}_j^{\mathbf{B}}(\mathbf{c}) \in f_j$. Then, $a_i \leq c_i$, for every $i \neq j$, and $a_j \leq \mathbf{r}_j^{\mathbf{B}}(\mathbf{c})$, hence, by (*iii'*), $\mathbf{t}^{\mathbf{B}}(\mathbf{a}) \leq c_j$, whence $c_j \in h$. Thus, $R^{\mathbf{B}}(f_1, \dots, f_k, h)$. Since $a_i \in f_i$, for every $i \in \{1, \dots, k\}$, we obtain $h \in \mathbf{t}(\mu(\mathbf{a}))$. Therefore, $\mu(\mathbf{t}^{\mathbf{B}}(\mathbf{a})) \subseteq \mathbf{t}(\mu(\mathbf{a}))$.

Let $j \in \{1, \dots, k\}$ be fixed and let $\mathbf{a} \in \text{dom}(\mathbf{r}_j^{\mathbf{B}})$. We show that $\mu(\mathbf{r}_j^{\mathbf{B}}(\mathbf{a})) = \mathbf{r}_j(\mu(\mathbf{a}))$, where

$$\mu(\mathbf{r}_j^{\mathbf{B}}(\mathbf{a})) = \{f_j \in U(B) \mid \mathbf{r}_j^{\mathbf{B}}(\mathbf{a}) \in f_j\}$$

and

$$\mathbf{r}_j(\mu(\mathbf{a})) = \{f_j \in U(B) \mid (\forall f_{i \neq j}, h \in U(B)) [\&_{i \neq j} a_i \in f_i \& R^{\mathbf{B}}(f_1, \dots, f_k, h)] \implies a_j \in h \}.$$

First, assume $f_j \in \mu(\mathbf{r}_j^{\mathbf{B}}(\mathbf{a}))$, i.e., $\mathbf{r}_j^{\mathbf{B}}(\mathbf{a}) \in f_j$. Suppose $a_i \in f_i$, for all $i \neq j$, and $R^{\mathbf{B}}(f_1, \dots, f_k, h)$. Then, $a_j \in h$, by definition of $R^{\mathbf{B}}$, hence $f_j \in \mathbf{r}_j(\mu(\mathbf{a}))$.

Conversely, assume $f_j \notin \mu(\mathbf{r}_j^{\mathbf{B}}(\mathbf{a}))$, i.e., $\mathbf{r}_j^{\mathbf{B}}(\mathbf{a}) \notin f_j$. To prove that $f_j \notin \mathbf{r}_j(\mu(\mathbf{a}))$, we show that there exist $f_i \in U(B)$, where $i \in \{1, \dots, k\} - \{j\}$, and $h \in U(B)$ such that $a_i \in f_i$, for every $i \neq j$, as well as $R^{\mathbf{B}}(f_1, \dots, f_k, h)$ and $a_j \notin h$. Let $f_i := [a_i]$, for all $i \in \{1, \dots, k\} - \{j\}$, and let $h := \{b \in B \mid b \not\leq^{\mathbf{B}} a_j\}$; these sets are clearly in $U(B)$. It should also be clear that $a_i \in f_i$, for all $i \neq j$, and that $a_j \notin h$. We next show that $R^{\mathbf{B}}(f_1, \dots, f_k, h)$.

Suppose that $\mathbf{c} \in \text{dom}(\mathbf{t}^{\mathbf{B}})$ and $c_i \in f_i$, for every $i \in \{1, \dots, k\}$. Then, $a_i \leq^{\mathbf{B}} c_i$, for every $i \neq j$. Since, by assumption, $\mathbf{r}_j^{\mathbf{B}}(\mathbf{a}) \notin f_j$ and $f_j \in U(B)$, it follows that $c_j \not\leq^{\mathbf{B}} \mathbf{r}_j^{\mathbf{B}}(\mathbf{a})$. Then, by (iv'), $\mathbf{t}^{\mathbf{B}}(\mathbf{c}) \not\leq^{\mathbf{B}} a_j$. Thus, $\mathbf{t}^{\mathbf{B}}(\mathbf{c}) \in h$, as required.

We show, next, that the second clause of the definition of $R^{\mathbf{B}}$ holds for the j we fixed. Suppose $\mathbf{c} \in \text{dom}(\mathbf{r}_j^{\mathbf{B}})$, as well as $c_i \in f_i$, for all $i \neq j$, and $\mathbf{r}_j^{\mathbf{B}}(\mathbf{c}) \in f_j$. Then $a_i \leq^{\mathbf{B}} c_i$, for all $i \neq j$. Since $\mathbf{r}_j^{\mathbf{B}}(\mathbf{a}) \notin f_j$, we also obtain $\mathbf{r}_j^{\mathbf{B}}(\mathbf{c}) \not\leq^{\mathbf{B}} \mathbf{r}_j^{\mathbf{B}}(\mathbf{a})$; hence, by (v'), $c_j \not\leq^{\mathbf{B}} a_j$. Thus, $c_j \in h$, as required.

We show, last, that the same clause holds for every $\ell \in \{1, \dots, k\} - \{j\}$. Fix such an ℓ and suppose that $\mathbf{c} \in \text{dom}(\mathbf{r}_\ell^{\mathbf{B}})$, as well as $c_i \in f_i$, for all $i \neq \ell$, and $\mathbf{r}_\ell^{\mathbf{B}}(\mathbf{c}) \in f_\ell$. Then, $a_i \leq^{\mathbf{B}} c_i$ for every i distinct from ℓ and j , and $a_\ell \leq \mathbf{r}_\ell^{\mathbf{B}}(\mathbf{c})$. Since $\mathbf{r}_j^{\mathbf{B}}(\mathbf{a}) \notin f_j$, we also obtain $c_j \not\leq^{\mathbf{B}} \mathbf{r}_j^{\mathbf{B}}(\mathbf{a})$, hence, by (vi'), $c_\ell \not\leq^{\mathbf{B}} a_j$. Thus, $c_\ell \in h$, as required.

Thus, $R^{\mathbf{B}}(f_1, \dots, f_k, h)$, which implies that $f_j \notin \mathbf{r}_j(\mu(\mathbf{a}))$ and, hence, that $\mathbf{r}_j(\mu(\mathbf{a})) \subseteq \mu(\mathbf{r}_j^{\mathbf{B}}(\mathbf{a}))$.

Thus, μ is an embedding of \mathbf{B} into \mathbf{A} . It follows that \mathbf{B} is a partial ra_k , which completes the proof. \square

In the case $k = 2$, the conditions (i'–vi') of Theorem 10 are equivalent to conditions (i–x) of Theorem 1. In the case $k = 1$, the conditions (i'–vi') may be presented in a simpler form, as given in the following corollary.

Corollary 1 *A partial σ_1 -structure $\mathbf{B} = \langle B, \mathbf{t}^{\mathbf{B}}, \mathbf{r}^{\mathbf{B}}, \leq^{\mathbf{B}} \rangle$ is a partial ra_1 if, and only if, it satisfies the following conditions:*

- (i'') $\langle B, \leq^{\mathbf{B}} \rangle$ is a partially ordered set;
- (ii'') $(\forall a, c \in \text{dom}(\mathbf{t}^{\mathbf{B}})) [a \leq^{\mathbf{B}} c \implies \mathbf{t}^{\mathbf{B}}(a) \leq \mathbf{t}^{\mathbf{B}}(c)]$;
- (iii'') $(\forall a, c \in \text{dom}(\mathbf{r}^{\mathbf{B}})) [a \leq^{\mathbf{B}} c \implies \mathbf{r}^{\mathbf{B}}(a) \leq^{\mathbf{B}} \mathbf{r}^{\mathbf{B}}(c)]$;
- (vi'') $(\forall a \in \text{dom}(\mathbf{t}^{\mathbf{B}})) (\forall c \in \text{dom}(\mathbf{r}^{\mathbf{B}})) [\mathbf{t}^{\mathbf{B}}(a) \leq^{\mathbf{B}} c \iff a \leq^{\mathbf{B}} \mathbf{r}^{\mathbf{B}}(c)]$.

8 Complexity of the Universal Theory of Residuated Algebras

In this section we consider classes of residuated algebras over fixed signatures that may contain finitely many fusion operations and their residuals. We prove that the universal theory of each such class is coNP-complete.

Let k be an integer with $k \geq 1$. By a k -residuation tuple, we mean a tuple $\langle \mathbf{t}, \mathbf{r}_1, \dots, \mathbf{r}_k \rangle$ of $k + 1$ operation symbols, each of arity k . By a *residuation signature* we mean a signature consisting of a binary relation symbol \leq and a finite set of residuation tuples.

Let ρ be a residuation signature. A *residuated ρ -algebra*, or ra_ρ , is a ρ -structure \mathbf{A} , with universe A and binary relation $\leq^{\mathbf{A}}$ such that $\langle A, \leq^{\mathbf{A}} \rangle$ is a partially ordered set and, for each $k \geq 1$ and each k -residuation tuple $\langle \mathbf{t}, \mathbf{r}_1, \dots, \mathbf{r}_k \rangle$ in ρ , \mathbf{A} has k -ary operations $\mathbf{t}^{\mathbf{A}}, \mathbf{r}_1^{\mathbf{A}}, \dots, \mathbf{r}_k^{\mathbf{A}}$ such that the structure $\langle A, \mathbf{t}^{\mathbf{A}}, \mathbf{r}_1^{\mathbf{A}}, \dots, \mathbf{r}_k^{\mathbf{A}}, \leq^{\mathbf{A}} \rangle$ is a ra_k , that is, it satisfies (11). The class of all residuated ρ -algebras is denoted by \mathcal{RA}_ρ .

Let ρ be a residuation signature. A *partial ρ -structure* is a structure \mathbf{B} that has universe B , a binary relation $\leq^{\mathbf{B}}$ on B and, for each $k \geq 1$ and each k -residuation tuple $\langle \mathbf{t}, \mathbf{r}_1, \dots, \mathbf{r}_k \rangle$ in ρ , \mathbf{B} has corresponding partial k -ary operations $\mathbf{t}^{\mathbf{B}}, \mathbf{r}_1^{\mathbf{B}}, \dots, \mathbf{r}_k^{\mathbf{B}}$. The domain of each operation $\mathbf{f}^{\mathbf{B}}$ in \mathbf{B} is denoted by $\text{dom}(\mathbf{f}^{\mathbf{B}})$. A *partial ra_ρ* is a partial ρ -structure \mathbf{B} that is a partial substructure of a ra_ρ , that is, there exists a ra_ρ \mathbf{A} with $B \subseteq A$, $\leq^{\mathbf{B}} = \leq^{\mathbf{A}} \upharpoonright_B$ and, for each $k \geq 1$ and each k -residuation tuple $\langle \mathbf{t}, \mathbf{r}_1, \dots, \mathbf{r}_k \rangle$ in ρ ,

$$\begin{aligned} \mathbf{t}^{\mathbf{B}}(\mathbf{a}) &= \mathbf{t}^{\mathbf{A}}(\mathbf{a}) \text{ for all } \mathbf{a} \in \text{dom}(\mathbf{t}^{\mathbf{B}}); \\ \mathbf{r}_j^{\mathbf{B}}(\mathbf{a}) &= \mathbf{r}_j^{\mathbf{A}}(\mathbf{a}) \text{ for every } j \in \{1, \dots, k\} \text{ and all } \mathbf{a} \in \text{dom}(\mathbf{r}_j^{\mathbf{B}}). \end{aligned}$$

We next give an intrinsic characterization of partial ra_ρ 's.

Theorem 11 *Let ρ be a residuation signature. A partial ρ -structure \mathbf{B} is a partial ra_ρ if, and only if, condition (i') of Theorem 10 holds and, for each $k \geq 1$ and each k -residuation tuple $\langle \mathbf{t}, \mathbf{r}_1, \dots, \mathbf{r}_k \rangle$ in ρ , conditions (ii'–vi') of Theorem 10 hold.*

Proof The proof is an adaptation of the proof of Theorem 10. If \mathbf{B} is a partial ra_ρ , then the required conditions hold. Conversely, suppose that \mathbf{B} is a partial ρ -structure satisfying the conditions of the theorem. For each residuation tuple in ρ , we define a corresponding relation $R^{\mathbf{B}}$ on $U(B)$ as in the proof of Theorem 10. Next, construct ra_ρ \mathbf{A} with universe $U(U(B))$ and operations, corresponding to each relation $R^{\mathbf{B}}$, defined as in Sect. 6. Then the map $\mu : B \rightarrow U(U(B))$ defined by $\mu(a) = \{f \in U(B) \mid a \in f\}$ can be shown to be an embedding of \mathbf{B} into \mathbf{A} , completing the proof. \square

For each residuation signature ρ , quantifier-free ρ -formulas and universal ρ -sentences are defined as in Sect. 4 for the signature σ . The definition of satisfaction of quantifier-free ρ -formulas in ra_ρ s and partial ra_ρ s is analogous to that given in Sect. 4 for *rogs*. The *size* of a quantifier-free ρ -formula φ , denoted $\text{size}(\varphi)$, is defined as the number of variables in φ plus the total number of occurrences of operation symbols in φ .

The following result is an analogue of Theorem 2.

Theorem 12 *Let ρ be a residuation signature. A quantifier-free ρ -formula φ is satisfiable in \mathcal{RA}_ρ if, and only if, it is satisfiable in a partial ra_ρ whose cardinality does not exceed $\text{size}(\varphi)$.*

By the above results, following the methods used for *rogs*, we obtain the following result.

Theorem 13 *Let ρ be a residuation signature. Satisfiability of quantifier-free ρ -formulas in \mathcal{RA}_ρ is NP-complete. The universal theory of \mathcal{RA}_ρ is coNP-complete.*

9 Conclusion

We conclude by offering some future research directions related to the results presented here. Firstly, it is common in the study of substructural logics—such as NL—that various structural rules are considered alongside the existing set of rules. Structural rules typically correspond to familiar algebraic properties of the related classes of algebras. For example, the exchange rule corresponds to commutativity of the \circ operation in *rogs*, while the weakening rule corresponds to the integrality property of *irogs*. The computational complexity of the universal theory for such extensions of *rogs* may be an interesting research direction. For residuated k -algebras, with $k \geq 3$, there may be scope for investigation of structural rules in general, as well as related complexity questions. Classes of residuated 1-algebras have been investigated in the context of ordered sets with a pair of residuated unary operations, or Galois operators, where the ordered set is usually a poset, as in the case here, or a lattice. There seems to be scope for investigating computational complexity questions for various classes of such algebras.

Acknowledgements We are grateful to the anonymous reviewers for their comments, which have helped to improve the paper.

References

- Aarts, E., & Trautwein, K. (1995). Non-associative Lambek categorial grammar in polynomial time. *Mathematical Logic Quarterly*, 41, 476–484.
- Avron, A. (1996). The method of hypersequents in the proof theory of propositional non-classical logics. In M. Hyland, C. Steinhorn, C. Steinhorn, & W. Hodges (Eds.), *Logic: From foundations to applications* (pp. 1–32). Clarendon Press.
- Bezhanišvili, G., Bezhanišvili, N., & Iemhoff, R. (2016). Stable canonical rules. *The Journal of Symbolic Logic*, 81(1), 284–315.
- Blok, W. J., & Van Alten, C. J. (2002). The finite embeddability property for residuated lattices, pocirms and BCK-algebras. *Algebra Universalis*, 48(3), 253–271.
- Blok, W. J., & Van Alten, C. J. (2005). On the finite embeddability property for residuated ordered groupoids. *Transactions of AMS*, 357(10), 4141–4157.
- Blyth, T. S., & Janowitz, M. F. (1972). *Residuation theory*. Pergamon Press.
- Bulińska, M. (2009). On the complexity of nonassociative Lambek calculus with unit. *Studia Logica*, 93, 1–14.
- Buszkowski, W. (1989). *Logical foundations of Ajdukiewicz and Lambek categorial grammars*. Polish Scientific Publishers (in Polish).
- Buszkowski, W. (1998). Algebraic structures in categorial grammar. *Theoretical Computer Science*, 199, 5–24.
- Buszkowski, W. (2005). Lambek calculus with nonlogical axioms. In: Casadio, C., Scott, P. J., & Seely, R. A (Eds), *Language and Grammar: Studies in Mathematical Linguistics and Natural Language, CSLI Lecture Notes, vol 168, Center for the Study of Language and Information* (pp. 77–94).
- Buszkowski, W. (2011). Interpolation and FEP for logics of residuated algebras. *Logic Journal of the IGPL*, 19(3), 437–454.
- Chang, C. C., & Keisler, H. J. (1990). *Model theory* (3rd ed., Vol. 73). North Holland, New York: Studies in Logic and the Foundations of Mathematics.
- Cook, S. (1971). The complexity of theorem proving procedures. In *Proceedings of the third annual ACM symposium on theory of computing* (pp. 151–158).

- Dunn, J. M. (1993). Partial gaggles applied to logics with restricted structural rules. In P. Schroeder-Heister & K. Došen (Eds.), *Substructural logics, studies in logic and computation* (Vol. 2, pp. 72–108). Clarendon Press.
- Farulewski, M. (2008). Finite embeddability property for residuated groupoids. *Reports on Mathematical Logic*, 43, 25–42.
- Fuchs, L. (1963). *Partially ordered algebraic systems*. Addison-Wesley.
- Kemhoff, R. (2015). On rules. *Journal of Philosophical Logic*, 44(6), 697–711.
- Iemhoff, R. (2016). Consequence relations and admissible rules. *Journal of Philosophical Logic*, 45(3), 327–348.
- Jäger, G. (2004). Residuation, structural rules and context freeness. *Journal of Logic, Language and Information*, 13, 47–59.
- Jalali, R. (2021). Proof complexity of substructural logics. *Annals of Pure and Applied Logic*, 172(7), 102972.
- Jeřábek, E. (2009). Canonical rules. *The Journal of Symbolic Logic*, 74(4), 1171–1205.
- Kandulski, M. (1997). On generalized Ajdukiewicz and Lambek calculi and grammars. *Fundamenta Informaticae*, 30(2), 169–181.
- Kołowska-Gawiejniewicz, M. (1997). Powerset residuated algebras and generalized Lambek calculus. *Mathematical Logic Quarterly*, 43(1), 60–72.
- Lambek, J. (1958). The mathematics of sentence structure. *American Mathematical Monthly*, 65, 154–170.
- Lambek, J. (1961). On the calculus of syntactic types. In: Jacobson, R. (Ed.), *Structure of language and its mathematical aspects, proceedings of symposia in applied mathematics* (Vol. XII). American Mathematical Society.
- Lellmann, B. (2016). Hypersequent rules with restricted contexts for propositional modal logics. *Theoretical Computer Science*, 656, 76–105.
- Sambin, G., Battilotti, G., & Faggian, C. (2000). Basic logic: Reflection, symmetry, visibility. *The Journal of Symbolic Logic*, 65(3), 979–1013.
- Shkatov, D., & Van Alten, C. J. (2019). Complexity of the universal theory of bounded residuated distributive lattice-ordered groupoids. *Algebra Universalis*, 80(3), 36.
- Shkatov, D., & Van Alten, C. J. (2020). Complexity of the universal theory of modal algebras. *Studia Logica*, 108(2), 221–237.
- Shkatov, D., & Van Alten, C. J. (2021). Computational complexity for bounded distributive lattices with negation. *Annals of Pure and Applied Logic*, 172(7), 102962.
- Van Alten, C. J. (2013). Partial algebras and complexity of satisfiability and universal theory for distributive lattices, Boolean algebras and Heyting algebras. *Theoretical Computer Science*, 501, 82–92.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.