

Embeddings into almost self-centered graphs of given radius

Kexiang Xu¹ · Haiqiong Liu¹ · Kinkar Ch. Das² · Sandi Klavžar^{3,4,5}

Published online: 30 May 2018

© Springer Science+Business Media, LLC, part of Springer Nature 2018

Abstract A graph is almost self-centered (ASC) if all but two of its vertices are central. An almost self-centered graph with radius r is called an r -ASC graph. The r -ASC index $\theta_r(G)$ of a graph G is the minimum number of vertices needed to be added to G such that an r -ASC graph is obtained that contains G as an induced subgraph. It is proved that $\theta_r(G) \leq 2r$ holds for any graph G and any $r \geq 2$ which improves the earlier known bound $\theta_r(G) \leq 2r + 1$. It is further proved that $\theta_r(G) \leq 2r - 1$ holds if $r \geq 3$ and G is of order at least 2. The 3-ASC index of complete graphs is determined. It is proved that $\theta_3(G) \in \{3, 4\}$ if G has diameter 2 and for several classes of graphs of diameter 2 the exact value of the 3-ASC index is obtained. For instance, if a graph G of diameter 2 does not contain a diametrical triple, then $\theta_3(G) = 4$. The 3-ASC index of paths of order $n \geq 1$, cycles of order $n \geq 3$, and trees of order $n \geq 10$ and diameter $n - 2$ are also determined, respectively, and several open problems proposed.

✉ Kexiang Xu
kexxu1221@126.com

Haiqiong Liu
lhqstu@163.com

Kinkar Ch. Das
kinkardas2003@gmail.com

Sandi Klavžar
sandi.klavzar@fmf.uni-lj.si

¹ College of Science, Nanjing University of Aeronautics and Astronautics, Nanjing 210016, Jiangsu, People's Republic of China

² Department of Mathematics, Sungkyunkwan University, Suwon 440-746, Republic of Korea

³ Faculty of Mathematics and Physics, University of Ljubljana, Ljubljana, Slovenia

⁴ Faculty of Natural Sciences and Mathematics, University of Maribor, Maribor, Slovenia

⁵ Institute of Mathematics, Physics and Mechanics, Ljubljana, Slovenia

Keywords Eccentricity · Diameter · Almost self-centered graph · Graph of diameter 2 · ASC index

Mathematics Subject Classification 05C12 · 05C05 · 05C75

1 Introduction and preliminaries

Almost self-centered graphs (ASC graphs for short) were introduced in Klavžar et al. (2011) as the graphs which contain exactly two non-central vertices. If the radius of an ASC graph G is r , then G is more precisely called an r -ASC graph. The introduction of these graphs was in particular motivated with network designs in which two (expensive) resources that need to be far away due to interference reasons must be installed. Numerous additional situations appear in which two specific locations are desired, for instance in trees (Huang et al. 2000; Wang et al. 2009), block graphs (Cheng et al. 2014), and interval graphs (Hong and Kang 2012). While in the first two of these papers 2-centers are studied in trees, the results could be used also in the general case. For a given graph G , one could first select a suitable subtree (spanning or otherwise), then determine a 2-center, and finally lift the obtained solution to G .

Graphs dual to the almost self-centered graphs were studied in Klavžar et al. (2014) and named almost-peripheral graphs (AP graphs for short). Very recently, a measure of non-self-centrality was introduced in Xu et al. (2016), where ASC graphs and AP graphs, along with a newly defined weakly AP graphs, play a significant role as extremal graphs in studies of this new measure.

Among other results it was proved in Klavžar et al. (2011) that for any connected graph G and any $r \geq 2$ there exists an r -ASC graph which contains G as an induced subgraph. Consequently, the r -ASC index $\theta_r(G)$ of G was introduced as the minimum number of vertices needed to add to G in order to obtain an r -ASC graph that contains G as an induced subgraph. It was proved that for any connected graph G we have $\theta_2(G) \leq 2$ and that $\theta_r(G) \leq 2r + 1$ holds for any $r \geq 3$. Moreover, the 2-ASC index was determined exactly for complete bipartite graphs, paths and cycles. The study of ASC graphs was continued in Balakrishnan et al. (2012) where ASC graphs were characterized among chordal graphs and among median graphs, and in Klavžar et al. (2017) where existing constructions of ASC graphs were revisited and new constructions of such graphs were developed.

We note in passing that graph eccentricity is frequently applied in chemical graph theory via the so-called eccentricity-based topological indices. Although these indices are pure graph theory concepts, they have a substantial use in theoretical chemistry, cf. Gupta and Singh (2002), Gupta et al. (2000, 2002) and Sharma et al. (1997). The eccentricity-based topological indices include eccentric distance sum (Gupta et al. 2002; Ilić et al. 2011), Zagreb eccentricity indices (Das et al. 2013), and connective eccentricity index (Xu et al. 2016; Yu and Feng 2013; Yu et al. 2014). See also Das and Nadjafi-Arani (2017), Su et al. (2015), Morgan et al. (2011), Tomescu (2008, 2010) and Wu et al. (2013) for some recent results on the general distance-based topological indices and a recent survey (Xu et al. 2014) on extremal results on general distance-based topological indices. Moreover, the applications of eccentricity to networks and

location theory can be seen in Krnc and Škrekovski (2015) and Puerto et al. (2008), respectively.


This paper is organized as follows. In the rest of this section definitions and notations needed are given. In the following section we first prove that if G is a graph of order $n \geq 1$ and $r \geq 2$, then $\theta_r(G) \leq 2r$, thus improving the so-far best known bound $\theta_r(G) \leq 2r + 1$. In the main result of the section we further strengthen the bound by proving that $\theta_r(G) \leq 2r - 1$ holds if $r \geq 3$ and G has at least two vertices. In the rest of the paper we then consider the 3-ASC index. The obtained results and (new) techniques involved indicate that to determine the 3-ASC index of a graph is generally a much more complex task than to determine its 2-ASC index. In Sect. 3, we determine the 3-ASC index of complete graphs and investigate the index on graphs of diameter 2. [See Palacios (2015) for some recent results on the Kirchhoff index of graphs with diameter 2.] We prove that if G is such a graph, then either $\theta_3(G) = 3$ or $\theta_3(G) = 4$. For several classes of graphs of diameter 2 the exact value is determined. In Sect. 4 we turn our attention to graphs with large diameter and determine the 3-ASC index of paths, cycles, and trees of order n and diameter $n - 2$. We conclude the paper with several open problems.

We only consider finite, undirected, simple graphs throughout this paper. The *degree* $\deg_G(v)$ of a vertex v of a graph G is the cardinality of the neighborhood $N_G(v)$ of v . The closed neighborhood $N_G[v]$ of v is $N_G(v) \cup \{v\}$. The maximum and minimum degree of G are denoted by $\Delta(G)$ and $\delta(G)$, respectively. A vertex of degree 1 is called a *pendant vertex*, the edge incident with a pendant vertex is called a *pendant edge*.

The *distance* $d_G(u, v)$ between vertices u and v is the length (that is, the number of edges) of a shortest (u, v) -path. The *eccentricity* $\text{ecc}_G(u)$ of a vertex u is the maximum distance from u to other vertices in G . The *diameter* $\text{diam}(G)$ of G is the maximum eccentricity among its vertices and the *radius* $\text{rad}(G)$ is the minimum eccentricity of its vertices. A vertex v is an *eccentric vertex* of a vertex u if $d_G(u, v) = \text{ecc}_G(u)$. The *eccentric set* $\text{Ecc}_G(u)$ of u is the set of all its eccentric vertices. A vertex u of G is called a *central vertex* if $\text{ecc}_G(u) = \text{rad}(G)$ and is called a *diametrical vertex* if $\text{ecc}_G(u) = \text{diam}(G)$. The *center* $C(G)$ of G consists of all its central vertices, while the *periphery* $P(G)$ of G contains all its diametrical vertices. A graph G is *self-centered* (SC graph for short) if $C(G) = V(G)$; we refer to the survey on SC graphs (Buckley 1989) and to their recent application in Maddaloni and Zamfirescu (2016). If $\text{rad}(G) = r$, then we also say that G is an r -SC graph. Vertices u and v for which $d_G(u, v) = \text{diam}(G)$ holds will be referred to as a *diametrical pair*. Similarly, a triple of vertices that are pairwise at distance $\text{diam}(G)$ is called a *diametrical triple*. When no ambiguity occurs, the subscript G will be omitted in the notations from the last two paragraphs.

If G is a graph, then \overline{G} denotes the complement of G . The *disjoint union* of (vertex-disjoint) graphs G_1 and G_2 will be denoted with $G_1 \cup G_2$, while the *join* of G_1 and G_2 will be denoted by $G_1 \oplus G_2$. Recall that $G_1 \oplus G_2$ is obtained from $G_1 \cup G_2$ by adding an edge between any vertex of G_1 and any vertex of G_2 . If $X \subseteq V(G)$, then $G[X]$ denotes the subgraph of G induced by X . Finally, throughout this paper we use P_n , C_n , and K_n to denote the path graph, the cycle graph, and the complete graph on n vertices, respectively. In addition, for any two non-adjacent vertices $x, y \in V(G)$, we mean that an edge xy is added into G by joining x with y in G .

Fig. 1 Graph \widehat{G}_r



2 Improving upper bounds on the r -ASC index

It was proved in Klavžar et al. (2011, Corollary 4.1) that $\theta_r(G) \leq 2r + 1$ holds for any graph G . In this section we first sharpen this result as follows. Interestingly, the below construction which gives a better upper bound is simpler than the construction from Klavžar et al. (2011).

Theorem 2.1 *If G is a graph of order $n \geq 1$ and $r \geq 2$, then $\theta_r(G) \leq 2r$.*

Proof Let \widehat{G}_r be the graph obtained from G as follows. Add three new vertices w , x_1 , y_1 , and add an edge between each of them and any vertex of G , so that in this way $3n$ edges are added. Further, add vertices x_2, \dots, x_{r-1} , y_2, \dots, y_{r-1} , and w' . Finally, add edges $x_1x_2, \dots, x_{r-2}x_{r-1}, y_1y_2, \dots, y_{r-2}y_{r-1}, x_{r-1}w'$, and $y_{r-1}w'$. The construction is illustrated in Fig. 1.

It is straightforward to verify that $d_{\widehat{G}_r}(w, w') = r + 1$ and $\text{ecc}_{\widehat{G}_r}(w) = \text{ecc}_{\widehat{G}_r}(w') = r + 1$. If $a \in V(G)$, then the sequence of vertices $a, x_1, \dots, x_{r-1}, w', y_{r-1}, \dots, y_1, a$ induces an isometric cycle of length $2r$. Using this fact it follows by a simple inspection that for any vertex $x \in V(\widehat{G}_r)$, $x \neq w, w'$, we have $\text{ecc}_{\widehat{G}_r}(x) = r$. We conclude that \widehat{G}_r is an r -ASC graph and consequently $\theta_r(G) \leq 2r$. \square

Next we characterize the graphs G with $\theta_r(G) = 2r$ for any $r \geq 3$. Before doing it, we need the following result.

Lemma 2.2 *Let G be a connected graph of order $n \geq 2$ and $r \geq 3$. Then $\theta_r(G) \leq 2r - 1$.*

Proof The graph C'_{2r} , which is obtained from C_{2r} by attaching a pendant vertex to it, is an r -ASC graph. It follows that $\theta_r(K_2) \leq 2r - 1$. In the rest of the proof we may hence assume that $n \geq 3$. It suffices to construct a graph H of order at most $n + 2r - 1$ as an r -ASC embedding graph of G .

Now we construct a graph H with vertex set $V(G) \cup \{x_1, \dots, x_{r-1}, y_1, \dots, y_{r-1}, w\}$. Set $V_0 = \{x_1, \dots, x_{r-1}, y_1, \dots, y_{r-1}, w\}$. We choose two arbitrary vertices x and y with $xy \in E(G)$. In H , the adjacency relation of V_0 is $x_1 - x_2 - \dots - x_{r-1} - y_{r-1} - \dots - y_1 - w$ and x, y are joined with x_1 and y_1 , respectively. Let $N_2(x)$ and $N_2(y)$ be the set of vertices in G at distance 2 from x and y , respectively. For any vertex $v \in N(x)$, we join v with y_1 in H . And we join v with x_1 in H for any vertex $v \in N(y) \setminus N(x)$.




Fig. 2 Construction from the proof of Lemma 2.2

Set $V^*(G) = V(G) \setminus (N(x) \cup N(y))$ and note that $x, y \notin V^*(G)$. Next we consider the positions of all vertices (if any) from $V^*(G)$ in H . For any vertex $v \in V^*(G)$, if $v \in N_2(y)$, v is joined with y_1 in H ; if $v \in N_2(x) \setminus N_2(y)$, then $vy_2 \in E(H)$; while if $z \in V^*(G) \setminus (N_2(x) \cup N_2(y))$, then $zy_1, zy_3 \in E(H)$ for $r \geq 4$ or $zy_1, zx_2 \in E(H)$ for $r = 3$.

This construction is illustrated in Fig. 2, where a thick edge from a vertex to a thick set indicates that the vertex is adjacent to every vertex in the set. The dashed thick line from x_1 indicates that x_1 is adjacent to every vertex in $N(y) \setminus N(x)$, while the dashed line from y_2 says that y_2 is adjacent to every vertex in $N_2(x) \setminus N_2(y)$. The figure is drawn for the cases when $r \geq 4$. If $r = 3$ then one only needs to replace the edge zy_3 (that is, from a typical vertex z from $V^*(G) \setminus (N_2(x) \cup N_2(y))$ to y_3) with the edge zx_2 . Note that if $r = 3$ then the vertices y_3 and x_3 do not exist, so that we can imagine that in the case $r = 3$ the vertex y_3 is identified with x_2 (and x_3 with y_2) and then the edge zy_3 is just the edge zx_2 .

Clearly, w is pendant in H with $wy_1 \in E(H)$. Moreover, $d_H(w, x_{r-2}) = r + 1$. We have $\text{ecc}_H(v) = r$ for any vertex $v \in V_0 \cup \{x, y\}$ with $v \notin \{x_{r-2}, w\}$ from the fact that the set $V_0 \cup \{x, y\}$ induces an isometric subgraph C'_{2r} of H . Similarly we have $\text{ecc}_H(u) = r$ for any vertex $u \in V(G) \setminus \{x, y\}$. In particular, for the case $r = 3$, we have $\text{Ecc}_H(u) = \{x\}$ if $u \in V^*(G)$; $\text{Ecc}_H(u) = \{x_2\}$ if $u \in N(x) \cup N_2(y)$; $\text{Ecc}_H(u) = \{w, x_1\}$ if $u \in N_2(x)$ and $\text{Ecc}_H(u) = \{w\}$ if $u \in N(y)$. Therefore H is an r -ASC embedding graph of G , finishing the proof of this lemma. \square

We are now ready for the main result of this section which further sharpens Theorem 2.1 by proving that $\theta_r(G) \leq 2r - 1$ holds if $r \geq 3$ and G has at least two vertices.




Fig. 3 Case 1 from the proof of Theorem 2.3




Fig. 4 Case 2 from the proof of Theorem 2.3

Theorem 2.3 *If G is a graph of order n and $r \geq 3$, then $\theta_r(G) = 2r$ if and only if $G \cong K_1$.*

Proof Note that C'_{2r} is an r -ASC graph. If $G \cong K_1$, considering the fact that any r -ASC has order $n \geq 2r + 1$ (Klavžar et al. 2011), the result $\theta_r(G) = 2r$ holds immediately. Conversely, if G is a graph of order $n \geq 2$, in view of Theorem 2.1, we only need to prove that $\theta_r(G) \leq 2r - 1$. Equivalently, it suffices to construct a graph H of order at most $n + 2r - 1$ as an r -ASC embedding graph of G . Denote by h the number of isolated vertices in G and set $V_0 = \{x_1, \dots, x_{r-1}, y_1, \dots, y_{r-1}, w\}$. Based on the value of h , we distinguish the following three cases.

Case 1 $h \geq 2$.

In this case we may assume that x_0, y_0 are two isolated vertices in G . In H , the adjacency relation of $V_0 \cup \{x_0, y_0\}$ is $x_0 - x_1 - \dots - x_{r-1} - y_0 - y_{r-1} - \dots - y_1 - x_0$ with x_0w being a pendant edge, that is, $V_0 \cup \{x_0, y_0\}$ induces a graph C'_{2r} in H . Moreover, any other vertex v than x_0, y_0 is joined with x_1 and y_1 in H . The construction is shown in Fig. 3, where the thick edges from x_1 and y_1 indicate that x_1 and x_2 are adjacent to all the vertices in $V(G) \setminus \{x_0, y_0\}$.

It can be easily checked that $d_H(w, y_0) = r + 1$ and any other vertex in H has eccentricity r . This ensures that H is an r -ASC embedding graph of G .

Case 2 $h = 1$.

In this case let v be an isolated vertex in G . Let the adjacency relation of $V_0 \setminus \{w\}$ and v in H be $x_1 - \dots - x_{r-1} - v - y_{r-1} - \dots - y_1$. Any vertex of G different from v is joined with x_1 and y_1 in H . Choosing an arbitrary but fixed vertex $u \in V(G) \setminus \{v\}$, we join w with u . This defines H . See Fig. 4 where this construction is shown; again

thick edges from x_1 and y_1 indicate that x_1 and y_1 are adjacent to all the vertices in $V(G) \setminus \{v\}$.

Note that $d_H(w, v) = r + 1$ with w being pendant and any other vertex than w, v has eccentricity r in H . Thus H is an r -ASC embedding graph of G as desired.

Case 3 $h = 0$.

In this case, if G is a connected graph, then we are done by Lemma 2.2. If not, by choosing any component of G , we run the same operation on each component of the graph G as that in the proof of Lemma 2.2 and join any vertex v from other component(s) with y_1, y_3 for $r \geq 4$ or with y_1, x_2 for $r = 3$. In this way we obtain an r -ASC embedding graph of G , completing the proof of the theorem. \square

3 3-ASC index of graphs with diameter at most 2

In this section we first determine the 3-ASC index of graphs of diameter 1. Clearly, a graph of order $n > 1$ has diameter 1 if and only if $G \cong K_n$. Next we prove the 3-ASC index of graphs with diameter 1. For consistency we include K_1 in the following theorem.

Theorem 3.1 $\theta_3(K_1) = 6$ and if $n \geq 2$, then $\theta_3(K_n) = 5$.

Proof $\theta_3(K_1) = 6$ by Theorem 2.3. In the rest of the proof we assume that $n \geq 2$.

We first show that $\theta_3(K_n) \geq 5$. Assume that G is a graph obtained from K_n by adding at most three vertices. Then $\text{diam}(G) \leq 4$. Moreover, the equality holds if and only if either $G \cong G'$ is formed by joining a pendant vertex of a path P_3 to some but not all vertices of K_n , or $G \cong G''$ is obtained by joining a vertex of a path P_2 to some but not all vertices of K_n and joining another new vertex to some other different vertices with degree $n - 1$ in it. In both cases there is at least one vertex x in G with $\text{ecc}_G(x) = 2$. Thus G is not a 3-ASC graph which in turn implies that $\theta_3(K_n) \geq 4$.

Suppose next that $\theta_3(K_n) = 4$ and let G be a 3-ASC graph obtained by adding vertices $V_0 = \{x, y, z, w\}$ to K_n . If $G[V_0] \cong P_4$, then G contains G' or G'' as an induced graph, where G' and G'' are the graphs defined in the above paragraph. Otherwise, $\text{diam}(G) < 4$, a clear contradiction. Since $\text{diam}(G) = 4$, there exists at least one vertex v in G with $\text{ecc}_G(v) = 2$, but this contradicts the fact that G is a 3-ASC graph.

Hence $G[V_0] \cong P_4$ must hold. Let the adjacency relation of V_0 in G be $w - z - y - x$ and assume without loss of generality that w has a neighbor in K_n . Then x is a pendant vertex of G because otherwise any vertex of G would have eccentricity at most 3. If at least one of the vertices y and z , say y , is adjacent to one or more vertices of K_n , then $\text{ecc}_G(y) = 2$, which is a contradiction to the fact that G is a 3-ASC graph. Therefore, $\text{deg}_G(z) = \text{deg}_G(y) = 2$. Now, if w is adjacent to all the vertices of K_n , then $\text{ecc}_G(z) = 2$, otherwise $\text{ecc}_G(x) = 5$. Hence in both cases G is not a 3-ASC graph. We conclude that $\theta_3(K_n) \geq 5$.

It remains to prove that $\theta_3(K_n) \leq 5$ for $n \geq 2$. For this sake consider the graph G_n which is schematically shown in Fig. 5. In the graph G_n , the vertices u and v are adjacent to all the vertices of the K_{n-1} leading to two subgraphs isomorphic to K_n . The two white vertices of G_n have eccentricity 4 while all the black vertices have

Fig. 5 Graph $G_n, n \geq 1$





Fig. 6 Graph H for $G = K_{2,3}$



eccentricity 3 (hereafter the white vertices denote the diametrical vertices and black vertices denote central ones). Hence G_n is a 3-ASC graph that contains K_n . \square

The graph G_2 from the proof of Theorem 3.1 is of order seven which makes it a smallest 3-ASC graph. Now we turn to graphs of diameter 2 and first bound their 3-ASC index as follows.


Lemma 3.2 *If G is a graph of diameter 2, then $3 \leq \theta_3(G) \leq 4$.*

Proof To prove the upper bound we need to construct a graph H of order $|V(G)| + 4$ as a 3-ASC embedding graph of G . Set $V(H) = V(G) \cup \{w, x, y, z\}$ and let the adjacency relation between the new vertices be $w - z - y - x$. Let v be a vertex in G with $\text{ecc}_G(v) = 2$ and u an eccentric vertex of v in G , that is, $d_G(v, u) = 2$. Join the vertex u in G with x and join all vertices from $V(G) \setminus \{u, v\}$ with w . This defines H (see Fig. 6 for an example of H when $G = K_{2,3}$).

Note that $\text{deg}_H(y) = \text{deg}_H(z) = \text{deg}_H(x) = 2$ and that the neighborhood of v remains unchanged in H . Then we arrive at $d_H(v, y) = 4$ from the structure of H . Moreover, any vertex v' from $N_G(u)$ has eccentricity 3 in H with $\text{Ecc}_H(v') = \{y\}$. Similarly, $\text{Ecc}_H(v') = \{x, y\}$ for any vertex $v' \in \text{Ecc}_G(u) \setminus \{v\}$, since both x and y have a largest distance 3 to v' in H . It follows that $\text{ecc}_H(v') = 3$ for any vertex $v' \in \text{Ecc}_G(u) \setminus \{v\}$. In addition, we have $d_H(w, x) = 3 = d_H(z, u)$, which implies that $\text{ecc}_H(u) = \text{ecc}_H(z) = 3 = \text{ecc}_H(w) = \text{ecc}_H(x)$. Thus H is a 3-ASC embedding graph of G as desired.

Next we deal with the lower bound on $\theta_3(G)$. Note that any graph obtained by adding one more vertex to G has diameter at most 3. Thus we have $\theta_3(G) \geq 2$.

Fig. 7 Graph H'_5



Assume that G' is an embedding 3-ASC graph of G by adding two more vertices x and y . Then G' has diameter 4. Hence the adjacency relation of x, y in G' is either $y - x - z$ with y being pendant, $\deg_{G'}(x) \geq 2$ and $z \in V(G)$ with $\text{ecc}_G(z) = 2$, or $x - u$ and $y - v$ where $d_G(u, v) = 2$ with $\text{ecc}_G(u) = \text{ecc}_G(v) = 2$. Considering that G has diameter 2, in the latter case, there is at least one vertex w as a common neighbor of vertices u and v in G . Then we deduce that $\text{ecc}_{G'}(z) = 2$ in the former case or $\text{ecc}_{G'}(w) = 2$ in the latter case, either of which contradicts the fact that G' is a 3-ASC graph. So $\theta_3(G) \geq 3$, finishing the proof of this lemma. \square

It is interesting to note that if $G = P_3$, then the graph H constructed in the proof of Lemma 3.2 is the graph of order 7 mentioned before the lemma. Since the order of a smallest 3-ASC graph is 7, it follows that $\theta_3(P_3) = 4$. There is a unique 2-SC graph C_4 of order 4 with diameter 2. By Lemma 3.2, $\theta_3(C_4) \leq 4$ and one can prove that actually $\theta_3(C_4) = 4$ holds. Consider next C_5 . By Lemma 3.2, $\theta_3(C_5) \leq 4$. But, to our surprise, 4 is not the minimum number of vertices needed to be added for obtaining a 3-ASC embedding of C_5 . Denote by C'_n the graph which consists of the cycle C_n and an additional vertex adjacent to exactly two consecutive vertices in C_n . As observed in Klavžar et al. (2011), the graph C'_{2r+1} is an r -ASC graph. Let H'_5 (shown in Fig. 7) be the graph obtained from C'_5 by respectively attaching a pendant vertex to the two non-adjacent vertices of degree 2 in C'_5 . Since H'_5 is a 3-ASC graph, we conclude that $\theta_3(C_5) = 3$. This example can be generalized as follows.

Theorem 3.3 *Let G be a 2-SC graph. If G contains a diametrical pair u, v and vertices u', v' such that $u' \in N_G(u) \setminus N_G(v)$, $v' \in N_G(v) \setminus N_G(u)$ and $N_G(u) \cap N_G(v) \subseteq \text{Ecc}_G(u') \cap \text{Ecc}_G(v')$, then $\theta_3(G) = 3$.*

Proof By Lemma 3.2 it suffices to construct a 3-ASC embedding graph of G . Let H be the graph obtained from G by adding three new vertices x, y, z and edges $xu, yv, zu',$ and zv' . The vertices x and y are pendant in H and hence $d_H(x, y) = 4$. Consequently, $\text{ecc}_H(u) = \text{ecc}_H(v) = 3$ with $\text{Ecc}_H(u) = \{y\}$ and $\text{Ecc}_H(v) = \{x\}$. Since $N_G(u) \cap N_G(v) \subseteq \text{Ecc}_G(u') \cap \text{Ecc}_G(v')$, any vertex $w \in N_G(u) \cap N_G(v)$ is at distance 2 from both u' and v' in G . Thus any vertex $w \in N_G(u) \cap N_G(v)$ is an eccentric vertex of z in H with $d_H(w, z) = 3$. Then $\text{ecc}_H(z) = 3$ holds. This in turn implies that $\text{ecc}_H(w) = 3$ for any vertex $w \in N_G(u) \cap N_G(v)$. Moreover, all the vertices in $N_G(u) \setminus N_G(v)$, including u' , have eccentricity 3 in H with y as their common eccentric vertex. By symmetry, all the vertices in $N_G(v) \setminus N_G(u)$, including v' , have eccentricity 3 in H . Finally, each vertex (if any) in $\text{Ecc}_G(u) \cap \text{Ecc}_G(v)$ has eccentricity 3 in H with x and y as its eccentric vertices. We conclude that H is a 3-ASC embedding graph of G . \square

By Lemma 3.2, the 3-ASC index of a graph G of diameter 2 is either 3 or 4. In the rest of this section we make a partial corresponding classification, where we will always assume that the graph G considered is of order n and of diameter 2. We first prove:

Theorem 3.4 *If G does not contain a diametrical triple and does not satisfy the assumption of Theorem 3.3, then $\theta_3(G) = 4$.*

Proof By Lemma 3.2 it suffices to prove that $\theta_3(G) \neq 3$. Suppose on the contrary that G^* is a 3-ASC embedding graph of the graph G with $V(G^*) = V(G) \cup \{x, y, z\}$ and set $V_0 = \{x, y, z\}$.

Assume first that G has a vertex v of degree $n - 1$. Note that then there exists a vertex u with $\text{ecc}_G(u) = 2$ adjacent to v in G . We now distinguish two cases.

Case 1 $\text{ecc}_{G^*}(v) = 4$.

In this case the adjacency relation of V_0 is $x - y - z$ with one vertex, say z , being pendant. Considering that G^* has diameter 4, we infer that any vertex u with $\text{ecc}_G(u) = 2$ must be adjacent to x in G^* . But then $\text{ecc}_{G^*}(x) = 2$, which is impossible because of the 3-ASC property of G^* .

Case 2 $\text{ecc}_{G^*}(v) = 3$.

In this case, clearly, we have $\text{Ecc}_{G^*}(v) \subseteq V_0$. Since $\text{ecc}_{G^*}(v) = \text{ecc}_G(v) + 2$, we have $|\text{Ecc}_{G^*}(v)| \leq 2$. If $|\text{Ecc}_{G^*}(v)| = 2$, we may assume that $\text{Ecc}_{G^*}(v) = \{y, z\}$. Then $x \in N_{G^*}(y) \cap N_{G^*}(z)$ and there exists a vertex w with $wx \in E(G^*)$. But now we get $\text{ecc}_{G^*}(w) = 2$, which is a contradiction. When $|\text{Ecc}_{G^*}(v)| = 1$, we may assume that $\text{Ecc}_{G^*}(v) = \{z\}$ and $v - w - y - z$ is a (v, z) -path of length 3. Note that $\text{ecc}_{G^*}(w) = 3$. Now we consider the position of x in G^* . If $xv \in E(G^*)$, then $\text{ecc}_{G^*}(w) = 2$, a contradiction again. If $d_{G^*}(x, v) = 2$, we assume that $w' \in N_{G^*}(x) \cap N_{G^*}(v)$. If $ww' \in E(G)$, we get the same result $\text{ecc}_{G^*}(w) = 2$. We observe that x is not adjacent to any vertex from $N_{G^*}(w)$. If not, we have $\text{ecc}_{G^*}(w) = 2$ which is impossible. In particular, $xy \notin E(G^*)$. Now we only need to deal with the subcase when $d_G(w, w') = 2$. Let w'' be a common neighbor of w and w' in G . If $xz \in E(G^*)$, then $xw'vwyzx$ is an induced cycle C_6 . Considering that G contains no diametrical triple, then G^* is a 3-SC graph. This contradicts the 3-ASC property of G^* . Therefore $xz \notin E(G^*)$. Combining $d_{G^*}(v, z) = 3$ with $\deg_G(v) = n - 1$, we find that z is pendant in G^* . Obviously, $\text{ecc}_{G^*}(z) = \text{ecc}_{G^*}(y) + 1$. It follows that $\text{ecc}_{G^*}(z) = 4$ from the 3-ASC property of G^* . Then $2 \leq d_{G^*}(y, x) \leq 3$. If $d_{G^*}(y, x) = 3$, then there must be a (y, x) -path, say $y - y_1 - x_1 - x$, of length 3 where $y_1, x_1 \in V(G) \setminus \{v\}$. This deduces $\text{ecc}_{G^*}(y_1) = 2$, a contradiction again. While if $d_{G^*}(y, x) = 2$, there must be a neighbor v' of v such that v' is a common neighbor of x and y in G^* . Then $\text{ecc}_{G^*}(v') = 2$. Both these possibilities are impossible from the 3-ASC property of G^* .

We have thus proved that no vertex of G can be of degree $n - 1$, that is, G must be a 2-SC graph. Suppose first that the adjacency relation of V_0 in G^* is $x - y - z$. We will show that x or z must be a pendant vertex of G^* . Otherwise, each vertex from V_0 has degree at least 2 in G^* . Let x' and z' from G be neighbors of x and z , respectively, in G^* . If $\text{deg}_{G^*}(y) > 2$, then any vertex of G^* has eccentricity at most 3, a contradiction since G^* is a 3-ASC graph. Hence $\text{deg}_{G^*}(y) = 2$. Clearly $\text{ecc}_{G^*}(x) = \text{ecc}_{G^*}(z) = 3$, therefore there must exist $u \in V(G)$ such that $d_{G^*}(y, u) = 4$. Now $d_{G^*}(y, x') =$

$d_{G^*}(y, z') = 2$ and we have $d_{G^*}(u, x') = d_{G^*}(u, z') = 2$. If $d_{G^*}(x', z') = 2$, then we have a contradiction with x', z' and u being a diametrical triple of G . Hence $d_{G^*}(x', z') = 1$ or $x' = z'$. In both cases, we have that $\text{ecc}_{G^*}(x') = \text{ecc}_{G^*}(z') = 2$, again a contradiction with G^* being a 3-ASC graph. Therefore both x and z cannot have a neighbor in G and suppose that z is a pendant vertex of G^* and that x' is a neighbor of x in G . Every vertex $u \in \text{Ecc}_G(x')$ must be adjacent to x or to y , since otherwise we have $d_{G^*}(z, u) = 5$ as a contradiction. But now we have $\text{ecc}_{G^*}(x) = 2$ a contradiction again with G^* being a 3-ASC graph.

Suppose now that $G^*[V_0] = P_2 \cup K_1$. Without loss of generality we may assume that $P_2 = xy$. If one of x and y , say y , is a pendant vertex, then $\text{ecc}_{G^*}(y) = 4$ and let $x' \in V(G)$ be a neighbor of x in G on the diametrical path of length 4 that starts in y . Observe that $\text{ecc}_{G^*}(x') = 2$, a contradiction with G^* being a 3-ASC graph. So, we may assume that x and y are not pendant and that x' and y' from G are neighbors of x and y , respectively, in G^* . Every vertex from G is at distance at most 3 to each vertex of V_0 and this implies that z is one endvertex of a path of length 4 and either x or y , say x , is the other. But now $\text{ecc}_{G^*}(y') = 2$, the same contradiction again.

Finally, let V_0 be an independent set in G^* . Then V_0 contains two diametrical vertices, say x and y , that is, $d_{G^*}(x, y) = 4$. Let $x - x' - w - y' - y$ be a diametrical path in G^* . Clearly, $w \in N_G(x') \cap N_G(y')$. We now claim that $\text{ecc}_{G^*}(w) = 2$. To prove it, it suffices to show that $d_{G^*}(w, z) = 2$. Otherwise, we have $d_{G^*}(w, z) = 3$. Since x' and y' do not lie in any diametrical triple, then there are two vertices $x_0, y_0 \in V(G)$ with $x_0 \in N_G(x') \setminus N_G(y'), y_0 \in N_G(y') \setminus N_G(x')$ and $w \in \text{Ecc}_G(x_0) \cap \text{Ecc}_G(y_0)$ with $zx_0, zy_0 \in E(G^*)$. But now G just satisfies the assumption of Theorem 3.3, which is impossible. Therefore we get the contradiction $\text{ecc}_{G^*}(w) = 2$ to the 3-ASC property of G . This completes the proof of the theorem. □

By Theorem 3.4 it remains to consider those graphs G of diameter 2 that contain diametrical triples. We next give two sufficient conditions that guarantee that $\theta_3(G) = 3$.

Theorem 3.5 *If a graph G contains vertices u and v of a diametrical triple of G such that $\delta(G[\text{Ecc}_G(u) \cap \text{Ecc}_G(v)]) = 0$, then $\theta_3(G) = 3$.*

Proof By Lemma 3.2 we only need to prove that $\theta_3(G) \leq 3$.

Since u and v belong to a diametrical triple, there exists a vertex $w \in \text{Ecc}_G(u) \cap \text{Ecc}_G(v)$, $w \neq u, v$. By the theorem’s assumption we may assume that w is an isolated vertex in $G[\text{Ecc}_G(u) \cap \text{Ecc}_G(v)]$. Now we construct a graph H as follows. Let $V(H) = V(G) \cup \{x, y, z\}$ and $E(G) \cup \{ux, vz, xy, yz\} \subseteq E(H)$. In addition, we join x to all (if any) vertices from $\text{Ecc}_G(u) \cap \text{Ecc}_G(v)$ different from w . This construction is presented in Fig. 8.

Since w is an isolated vertex in $G[\text{Ecc}_G(u) \cap \text{Ecc}_G(v)]$, we have $d_G(w, w') = 2$ for any vertex $w' \in (\text{Ecc}_G(u) \cap \text{Ecc}_G(v)) \setminus \{w\}$. Clearly, $d_G(w, u) = 2 = d_G(w, v)$. So $d_H(w, y) = 4$. Any vertex from $N_G(u) \cap N_G(v)$ has eccentricity 3 since y is its unique eccentric vertex with distance 3 to it in H . In addition, $\text{ecc}_H(u) = 3$ from the fact that $\text{Ecc}_H(u) = \{z\}$ with $d_H(u, z) = 3$. Similarly, we also have $\text{ecc}_H(v) = 3$. For any vertex $w' \in (\text{Ecc}_G(u) \cap \text{Ecc}_G(v)) \setminus \{w\}$, considering that $w'x \in E(H)$, we




Fig. 8 Construction from the proof of Theorem 3.5




Fig. 9 Construction from the proof of Theorem 3.6

conclude that $\text{ecc}_H(w') = 3$ since z is the only one vertex with the largest distance 3 to w' in H . For any vertex $v' \in N_G(u) \setminus N_G(v)$, we have $\text{Ecc}_H(v') = \{y, z\}$ with $d_H(v', y) = 3 = d_H(v', z)$ being largest. Then it follows that $\text{ecc}_H(v') = 3$. Moreover, $\text{ecc}_H(v') = 3$ with $\text{Ecc}_H(v') = \{y, x\}$ for any vertex $v' \in N_G(v) \setminus N_G(u)$ by an analogous reasoning. We conclude that H is a 3-ASC embedding graph of G . \square

Theorem 3.6 *If a graph G contains vertices u and v of a diametrical triple of G such that $G[\text{Ecc}_G(u) \cap \text{Ecc}_G(v)]$ contains P_3 as an induced subgraph, then $\theta_3(G) = 3$.*

Proof Applying Lemma 3.2 again we only need to prove that $\theta_3(G) \leq 3$.

Let $w_1w_2w_3$ be an induced path in $G[\text{Ecc}_G(u) \cap \text{Ecc}_G(v)]$. We construct a graph H obtained from G by adding vertices x, y, z and edges xy, yz, ux, vz, w_3x . Moreover, we join all vertices (if any) from $(\text{Ecc}_G(u) \cap \text{Ecc}_G(v)) \setminus N_G[w_1]$ with the vertex x in H . See Fig. 9 for an illustration of the construction.

Thus the neighborhood of any vertex from $N_G[w_1]$ or not in $(\text{Ecc}_G(u) \cap \text{Ecc}_G(v)) \cup \{u, v\}$ remains unchanged in H . From the construction of H we first infer that $d_H(w_1, y) = 4$ since $w_1 w_2 w_3 x y$ is a shortest path between them. Any vertex from $\text{Ecc}_G(u) \cap \text{Ecc}_G(v)$, except w_2 and w_1 , has eccentricity 3 in H because z is its unique eccentric vertex and is at distance 3. Also we have $\text{ecc}_H(w_2) = 3$ from the fact that $\text{Ecc}_H(w_2) = \{y, z\}$ with $d_H(w_2, y) = 3 = d_H(w_2, z)$. Any vertex from $N_G(u) \cap N_G(v)$ has eccentricity 3 in H because y is its unique eccentric vertex and is at distance 3. Furthermore, $\text{ecc}_H(w) = 3$ with $\text{Ecc}_H(w) = \{y, z\}$ for any vertex $w \in N_G(u) \setminus N_G(v)$. By symmetry, we get $\text{ecc}_H(w) = 3$ for any vertex $w \in N_G(v) \setminus N_G(u)$. Also we claim that $\text{ecc}_H(x) = 3 = \text{ecc}_H(z)$ because $w_1 \in \text{Ecc}_H(x)$ and $w_2 \in \text{Ecc}_H(z)$, respectively, in H . Obviously we have $\text{ecc}_H(u) = 3 = \text{ecc}_H(v)$ from $\text{Ecc}_H(u) = \{z\}$ and $\text{Ecc}_H(v) = \{x\}$, respectively. Therefore H is a 3-ASC embedding graph of G . \square

Note that the Petersen graph PG fulfills the assumption of Theorem 3.3, hence $\theta_3(PG) = 3$. But PG does not fulfill the condition of Theorem 3.5. Further, $\theta_3(K_{1,3}) = 3$ holds by Theorem 3.5. However, $K_{1,3}$ does not fulfill the assumption of Theorem 3.3 or that of Theorem 3.6. Therefore, none of Theorems 3.3, 3.5, and 3.6 gives a necessary condition for $\theta_3(G) = 3$.

In view of Theorems 3.5 and 3.6 one might wonder whether $\theta_3(G) = 3$ holds as soon as G contains a diametrical triple. This is not the case as it can be verified on the graph $K_1 \oplus 3P_2$. More generally, we have the following result.

Theorem 3.7 *If for any non-adjacent vertices u, v the subgraph induced by $\text{Ecc}_G(u) \cap \text{Ecc}_G(v)$ is a disjoint union of complete graphs on at least two vertices such that any outer neighbor of each vertex from any of these complete subgraphs belongs to $N_G(u) \cap N_G(v)$ and G does not satisfy the assumption of Theorem 3.3, then $\theta_3(G) = 4$.*

Proof By Lemma 3.2 it suffices to prove that $\theta_3(G) > 3$. Suppose on the contrary that $\theta_3(G) = 3$ and let H be a 3-ASC graph obtained from G by adding vertices $V_0 = \{x, y, z\}$ and some edges (while keeping G to be induced in H). We distinguish two cases.

Case 1 No vertex from V_0 is pendant in H .

We first claim that if $H[V_0]$ is complete or disconnected, then we can find some vertex with eccentricity 2 in H , contradicting the 3-ASC property of H .

Suppose first that $H[V_0]$ is complete. Then considering that H has exactly two diametrical vertices with eccentricity 4, we observe that one vertex, say x , from V_0 has eccentricity 4 in H . Let $w \in V(G)$ be the other diametrical vertex in H . Then there is a vertex $w' \in V(G)$ with $d_H(w', x) = d_H(w', w) = 2$. It follows that $\text{ecc}_H(w') = 2$.

Assume next that $H[V_0]$ is disconnected. Then $H[V_0] \cong K_1 \cup P_2$ or consists of three isolated vertices. In the former case, we can assume that x is isolated in $H[V_0]$. Then $2 \leq \min\{d_H(x, y), d_H(x, z)\} \leq 3$ since H is a 3-ASC graph. Without loss of generality assume that $\min\{d_H(x, y), d_H(x, z)\} = d_H(x, y)$. If $d_H(x, y) = 2$, then there is a vertex $w \in V(G)$ as a common neighbor of x and y in H . Thus we get

$\text{ecc}_H(w) = 2$. When $d_H(x, y) = 3$, we assume that $xx', yy' \in E(H)$ where $x', y' \in V(G)$. Then $d_G(x', y') = 1$ and $\text{ecc}_H(y') = 2$. In the latter case, we conclude that there is a diametrical vertex in V_0 of H , say $\text{ecc}_H(x) = 4$. Then the other diametrical vertex of H must be also in V_0 . Otherwise, assume that $w \in V(G)$ has eccentricity 4 in H . Then the distance from w to any neighbor of x in G is 3, which is impossible because G has diameter 2. Assume that $\text{ecc}_H(y) = 4$. Let z' be an arbitrary neighbor of z in H and x', y' the arbitrary neighbors of x and y , respectively, in H . First we claim that neither of x and y belongs to $\text{Ecc}_H(z)$. If not, without loss of generality let $x \in \text{Ecc}_H(z)$. Considering that $\text{ecc}_H(z') = 3$, we have $d_H(z', x) = 3$. Then $d_H(z, x) = 4$, which is a clear contradiction. Thus we have $d_H(x, z) = 2 = d_H(y, z)$ from $\text{ecc}_H(z) = 3$. Now we conclude that $zx', zy' \in E(H)$. It follows that $\text{ecc}_H(w) = 2$ for any vertex $w \in N_G(x') \cap N_G(y')$. This proves the claim.


So we are left with the situation when $H[V_0] \cong P_3$. Assume that the adjacency relation of V_0 in H is $x - y - z$. By the case assumption, there exists edges $xu, zv \in E(H)$. Note first that $u \neq v$, for otherwise we would have $\text{ecc}_H(u) = 2$. The same conclusion holds if $d_H(u, v) = 1$.

Hence $d_H(u, v) = 2$ must hold. By the theorem’s assumption, $\text{Ecc}_G(u) \cap \text{Ecc}_G(v)$ induces a disjoint union of complete graphs $K_{n_k}, n_k \geq 2$. Note first that w is not adjacent to any vertex from V_0 for any vertex $w \in N_G(u) \cap N_G(v)$, for otherwise $\text{ecc}_H(w) = 2$ would hold. Recall that G has diameter 2. If there is at least one vertex in each K_{n_k} adjacent to some vertex from V_0 , then the distance is at most 3 from any vertex from $G[\text{Ecc}_G(u) \cap \text{Ecc}_G(v)]$ to any vertex in V_0 of H . Thus any vertex in H has eccentricity at most 3, a contradiction to the fact that H is a 3-ASC graph. Hence there must be some complete component K_{n_i} , from $G[\text{Ecc}_G(u) \cap \text{Ecc}_G(v)]$, such that none of vertices of K_{n_i} is adjacent to any vertex from V_0 in H . Let w_1, w_2 be arbitrary vertices of K_{n_i} . Recalling that any vertex from V_0 is not adjacent to any vertex from $N_G(u) \cap N_G(v)$ in H , from the theorem’s condition that any outer neighbor of any vertex of these complete subgraphs belongs to $N_G(u) \cap N_G(v)$, we find that $d_H(w_1, y) = d_H(w_2, y) = 4$. We conclude that H is not a 3-ASC graph.

Case 2 V_0 contains at least one pendant vertex of H .

Denote by h the number of pendant vertices from V_0 in H . If $h = 1$, we may assume that $\text{deg}_H(z) = 1$. Considering that H is 3-ASC graph, we observe that $\text{ecc}_H(z) = 4$. Assume that $z - z' - w_0 - w' - w$ is a diametrical path in H . Then $d_H(z', w) = 3$, which implies that at most one vertex from $\{w', z\}$ belongs to $V(G)$. If $w_0 \in V(G)$, we claim that $\text{ecc}_H(w_0) = 2$ as a contradiction. In fact, clearly, $\text{ecc}_H(w_0) = 2$ if $\{z, w'\} \subseteq V_0$. When $|\{z', w\} \cap V_0| = 1$, without loss of generality, assume that $z' = y \in V_0$. Now it suffices to prove that $d_H(w_0, x) = 2$. Otherwise, we have $d_H(w_0, x) = 3$, that is, there exists a vertex $x' \in \text{Ecc}_G(w_0)$ such that $xx' \in E(H)$. Note that $\text{ecc}_H(x) = 3$. Then $d_H(x, z) \in \{2, 3\}$. If $d_H(x, z) = 2$, then $xz' \in E(H)$, which implies $d_H(w_0, x) = 2$, contradicting the fact that $d_H(w_0, x) = 3$. Thus $d_H(x, z) = 3$ holds. We get $d_H(x', z') = 3$, i.e., $d_H(x', z) = 4$ as another contradiction. While $w_0 \notin V(G)$, without loss of generality, we may assume that $w_0 = x$. Next we divide into the following two subcases. For $z' \in V(G)$, recalling that z', w cannot belong to $V(G)$ simultaneously, we have $w \notin V(G)$, i.e., $w = y$. In this subcase, we claim that z' and w' forms a diametrical pair in G . Let $w^* \in N_G(w') \cap N_G(z')$. Then the distance from w^* to any vertex in V_0 is at most 2 in H . Recalling that G has diameter 2. Then

Fig. 10 Graph G with $\theta_3(G) = 3$



$\text{ecc}_H(w^*) = 2$, which is also impossible. If $z' \notin V(G)$, then $z' = y$. So the adjacency relation of V_0 in H is $x - y - z$ with z being pendant. If $\text{deg}_G(w') = n - 1$, then $\text{ecc}_H(x) = 2$ clearly. If $\text{ecc}_G(w') = 2$, we consider the vertices from $\text{Ecc}_G(w')$. Now we conclude that w_1 is adjacent to x or y in H for any vertex $w_1 \in \text{Ecc}_G(w')$. Otherwise, $d_H(w_1, z) > 4$, contradicting to the 3-ASC property of H . Therefore, we get $\text{ecc}_H(x) = 2$, a clear contradiction comes again.

If $h = 2$, without loss of generality, we assume that x, y are both pendant in H . Clearly, $\text{ecc}_H(x) = 4 = \text{ecc}_H(y)$. Let $xx', yy' \in E(H)$. Since H is 3-ASC graph, there is a diametrical path $x - x' - w - y' - y$ with $w \in V(G) \cup \{z\}$. Note that $d_G(x', y') = 2$. Without loss of generality, assume that $w \in V(G)$. Since G does not satisfy the assumption of Theorem 3.3, we have $d_H(w, z) \leq 2$ (if G satisfies the condition of Theorem 3.3 and $z \in N_2(x') \cap N_2(y')$ in H , then $d_H(w, z) = 3$ holds possibly), which implies that $\text{ecc}_H(w) = 2$ as an obvious contradiction. If $h = 3$, then each of x, y and z has eccentricity 4 in H (if one of them has eccentricity 3, then its unique neighbor has eccentricity 2, a contradiction), contradicting the definition of 3-ASC graph. □

Consider the graph G shown in Fig. 10. It is of diameter 2 and fulfils the first condition of Theorem 3.7. For the selected vertices u and v we have $\text{Ecc}_G(u) \cap \text{Ecc}_G(v) = \{x, y\}$. Note that each of these vertices has an outer neighbor that does not belong to $N_G(u) \cap N_G(v)$. Hence the second condition of the theorem is not fulfilled. In the figure an embedding of G into a 3-ASC graph is also shown, therefore $\theta_3(G) = 3$. In summary, the technical condition about the outer neighbors from Theorem 3.7 cannot be avoided.

Recall that the well-known cocktail party graph $CP(n)$ is a complete graph on $2n$ vertices with a 1-factor removed. By Theorem 3.4, $\theta_3(CP(n)) = 4$ but $CP(n)$ does not fulfill the condition of Theorem 3.7. In addition, from Theorem 3.7 we get that $\theta_3(K_1 \oplus tP_2) = 4$ holds for any $t \geq 3$, but $K_1 \oplus tP_2, t \geq 3$, does not fulfill the condition of Theorem 3.4. Therefore, Theorems 3.4 and 3.7 give sufficient but not necessary conditions for G having $\theta_3(G) = 4$.





Fig. 11 The graphs H_1 and H_2

Fig. 12 Embedding of P_9 into a 3-ASC graph of order 11



4 3-ASC index of some graphs with large diameter

In this section we will determine the 3-ASC index of paths of order $n \geq 1$, cycles of order $n \geq 3$, and trees of order $n \geq 10$ and diameter $n - 2$.

4.1 Paths

In this subsection we will always assume that $V(P_n) = \{v_1, \dots, v_n\}$ and $E(P_n) = \{v_i v_{i+1} : i = 1, \dots, n - 1\}$. We will also use the graph C_{2k}^* , $k \geq 2$, which is obtained from C_{2k} by attaching a pendant vertex to one of its vertices. As observed in Klavžar et al. (2011), the graph C_{2r}^* is an r -ASC graph. First we deal with the cases when n is small.

The cases $n = 1$ and $n = 2$ are covered by Theorem 3.1. We have already observed that a smallest 3-ASC graph is of order 7. Since each of the paths P_3, P_4, P_5 , and P_6 is an induced subgraph of C_6^* , it follows that $\theta_3(P_3) = 4, \theta_3(P_4) = 3, \theta_3(P_5) = 2$, and $\theta_3(P_6) = 1$. For $n = 7$ and $n = 8$, consider the graphs H_1 and H_2 from Fig. 11. By a straightforward checking we find that both H_1 and H_2 are 3-ASC graphs. Therefore $\theta_3(P_7) = \theta_3(P_8) = 1$.

In the rest we may thus assume that $n \geq 9$ and first prove:

Lemma 4.1 *If $n \geq 9$, then $\theta_3(P_n) \leq 2$.*

Proof For $n = 9$ the assertion follows from the embedding presented in Fig. 12.

For $n \geq 10$ we construct a graph G_n schematically shown in Fig. 13, where all the vertices v_{10}, \dots, v_n are adjacent to both x and y .

It is straightforward to check that G_n is a 3-ASC embedding graph of P_n . □

Lemma 4.2 *If $n \geq 9$, then $\theta_3(P_n) \geq 2$.*

Proof Since paths P_n are not 3-ASC graphs we have $\theta_3(P_n) > 0$ for all $n \geq 9$. In the following we prove that $\theta_3(P_n) > 1$ for $n \geq 9$. Assume on the contrary that there




Fig. 13 Graphs G_n

exists a 3-ASC graph G with $V(G) = V(P_n) \cup \{x\}$ that contains P_n as an induced subgraph.

First assume that x is a diametrical vertex in G and let v_i be its diametrical pair, so that $d_G(x, v_i) = 4$. If $i = 1$, then clearly $xv_1, xv_2, xv_3 \notin E(G)$ and $xv_4 \in E(G)$. Since P_n is an induced subgraph of G , we conclude that $d_G(v_1, v_j) \geq 4$ for $j \geq 5$, which contradicts the fact that G is a 3-ASC graph. Similarly we get a contradiction if $i = 2$. By symmetry, $i \neq n - 1$ and $i \neq n$. Finally, if $2 < i < n - 2$, then x is not adjacent to any vertex of $v_{i-2}, v_{i-1}, v_{i+1}$ and v_{i+2} . Thus $d_G(v_{i-2}, v_{i+2}) = 4$, another contradiction.

It follows that the two diametrical vertices of G are on P_n , say v_i and v_j , where $i < j$. Then $d_G(v_i, v_j) = 4$. Clearly, xv_i and xv_j cannot belong to $E(G)$ simultaneously. We distinguish the following two cases.

Case 1 Either $xv_i \in E(G)$ or $xv_j \in E(G)$.

We may without loss of generality assume that $xv_i \in E(G)$. If $i > 1$, then $d_G(v_{i-1}, v_j) \geq 4$, which implies that G has at least three diametrical vertices. Hence $i = 1$ must hold. Since $xv_1 \in E(G)$ and $d_G(v_1, v_j) = 4$, neither of v_j and v_{j-1} is adjacent to x in G . If $j < n$, then the same conclusion holds also for v_{j+1} . In any case, $d_G(v_{j-4}, v_j) = 4$. This is only possible if $j - 4 = i = 1$, that is, if $j = 5$. As already argued above, $xv_4, xv_5, xv_6 \notin E(G)$, for otherwise $d_G(v_1, v_5) \leq 3$ would hold. But then it follows that $d_G(v_9, v_5) = 4$, so we would again have at least three diametrical vertices. Case 1 hence cannot happen.

Case 2 Neither of v_i and v_j is adjacent to x in G .

Suppose first that $i \geq 2$ and $j \leq n - 1$. Then x must be adjacent to each vertex from $V_1 = \{v_k : 1 \leq k < i\}$ because $d_G(v_k, v_j) \geq 4$ holds if $v_kx \notin E(G)$. Analogously, x must be adjacent to each vertex from $V_2 = \{v_t : j < t \leq n\}$. Since $d_G(v_i, v_j) = 4$ we clearly have that $j - i \geq 4$. If $j - i = 4$, then $xv_{i+2} \notin E(G)$, for otherwise $\text{ecc}_G(x) = 2$. Considering that $d_G(v_{i+3}, v_k) \leq 3$ for any vertex $v_k \in V_1$, we have $xv_{i+3} \in E(G)$. By symmetry, we get $xv_{i+1} \in E(G)$. Hence $\text{ecc}_G(x) = 2$ holds again, a contradiction. If $j - i > 4$, then from $\text{ecc}_G(x) = 3$ we find out that there are three consecutive vertices v_s, v_{s+1}, v_{s+2} where $i \leq s < s + 2 \leq j$ such that each of them has degree 2 in G . But now there must be a vertex $v_p \in V_1 \cup V_2$ such that $d_G(v_{s+1}, v_p) = 4$.

Let next $i = 1$ and $j < n$. Then x must be adjacent to all vertices $v_t, t > j$, because otherwise $d_G(v_t, v_1) \geq 4$ would hold for such a vertex v_t . Assume that $xv_2 \notin E(G)$. Then $j = 5$ because otherwise $d_G(v_2, v_j) > 3$. But then $d_G(v_1, v_6) > 3$, a contradiction. Hence necessary $xv_2 \in E(G)$. As above we now find consecutive vertices v_s, v_{s+1}, v_{s+2} such that none of them is adjacent to x in G where $3 \leq s <$

$s + 2 \leq j$, for otherwise $\text{ecc}_G(x) = 2$. But then $d_G(v_{s+1}, v_1) > 3$ for $s > 3$, or $d_G(v_{s+1}, v_p) > 3$ where $p > j$ for $s = 3$, another contradiction.

The case when $i > 1$ and $j = n$ is symmetric to the last case. Hence we are left with the situation when $i = 1$ and $j = n$. If $xv_2 \notin E(G)$, then we have $d_G(v_1, v_{n-1}) > 3$. So $xv_2 \in E(G)$ and by a parallel argument also $xv_{n-1} \in E(G)$. Now, as above, we have a triple of vertices v_s, v_{s+1}, v_{s+2} none of which is adjacent to x in G . But then $d_G(v_{s+1}, v_1) > 3$ or $d_G(v_{s+1}, v_n) > 3$, a final contradiction. \square

Combining Lemmas 4.1 and 4.2 with the arguments given before then we have the following result.

Theorem 4.3 *The 3-ASC index of paths is:*

$$\theta_3(P_n) = \begin{cases} 7 - n, & 1 \leq n \leq 5; \\ 1, & 6 \leq n \leq 8; \\ 2, & n \geq 9. \end{cases}$$

4.2 Cycles

Next we determine the 3-ASC index of cycles. The triangle C_3 is a complete graph, so $\theta_3(C_3) = 5$ by Theorem 3.1. We have already observed that $\theta_3(C_4) = 4$ and $\theta_3(C_5) = 3$. Recall that C_6^* (the graph obtained by attaching a pendant vertex to one of the vertices of C_6) is a 3-ASC graph, hence $\theta_3(C_6) = 1$. Recall further that C'_{2r+1} is the graph which consists of a cycle C_{2r+1} and a vertex adjacent to exactly two consecutive vertices in C_{2r+1} and that C'_{2r+1} is an r -ASC graph. Thus C'_7 is 3-ASC graph and $\theta_3(C_7) = 1$ holds accordingly. Let in addition C''_8 be the graph obtained by joining five consecutive vertices of C_8 with a new vertex. It can be easily checked that C''_8 is a 3-ASC graph and thus $\theta_3(C_8) = 1$. The result for all cycles reads as follows.

Theorem 4.4 *The 3-ASC index of cycles is:*

$$\theta_3(C_n) = \begin{cases} 8 - n, & 3 \leq n \leq 5; \\ 1, & 6 \leq n \leq 8; \\ 2, & n \geq 9. \end{cases}$$

Proof The result for $n \leq 8$ has been established above, hence in the following we will assume that $n \geq 9$. Let $V(C_n) = \{v_1, \dots, v_n\}$, $E(C_n) = \{v_i v_{i+1} : i = 1, \dots, n - 1\} \cup \{v_n v_1\}$, and for a positive integer k , $1 \leq k \leq n$, we set $|k|_n = \min\{k, n - k\}$.

We first prove that $\theta_3(C_n) \leq 2$. For $n = 9$ consider the graph G shown in Fig. 14. It can be routinely checked that G is a 3-ASC graph (recall that two vertices filled white are diametrical vertices as mentioned before).

When $n \geq 10$, we construct the graph H_n as shown in Fig. 15. In H_n both x and y are adjacent to any vertex from $\{v_k : 10 \leq k \leq n\}$. The graph H_n is a 3-ASC embedding graph of C_n , completing the proof of upper bound on $\theta_3(C_n)$.

To complete the proof we need to show that $\theta_3(C_n) \geq 2$. As C_n , $n \geq 9$, is not a 3-ASC graph, we only need to prove that $\theta_3(C_n) > 1$. Suppose on the contrary that a graph G with vertex set $V(C_n) \cup \{x\}$ is a 3-ASC embedding graph of C_n .

Fig. 14 The graph G





Fig. 15 The graph H_n



If $\text{ecc}_G(x) = 4$, then there exists an induced internal (v_i, v_j) -path P (that is, a path whose internal vertices are of degree 2) of length 6 in G such that $xv_i, xv_j \in E(G)$ and $d_G(x, v_t) = 4$ where $v_t \in V(P)$ with $|t - i|_n = |t - j|_n = 3$. Thus we conclude that $d_G(v_t, v_s) \geq 4$ for $v_s \in V(C_n) \setminus V(P)$. This is an apparent contradiction to the 3-ASC property of G . If $\text{ecc}_G(x) = 3$, then, considering a vertex being at distance 3 from x , we infer that there is an induced internal (v_i, v_j) -path P with $|j - i|_n = 4$ or 5 and $v_k \in V(P)$ such that both v_i and v_j are adjacent to x in G and that neither of v_i and v_j is adjacent to v_k . Now we consider the eccentricity of v_k in G . If $\text{ecc}_G(v_k) = 3$, then any vertex from $V(C_n) \setminus V(P)$ is adjacent to v_i or v_j . However, this is impossible when $n \geq 9$. Then we get $\text{ecc}_G(v_k) = 4$. Without loss of generality, assume that $i < j$ and v_i has a neighbor $v_{i-1} \in V(C_n) \setminus V(P)$ and v_j has a neighbor $v_{j+1} \in V(C_n) \setminus V(P)$. Let $V_0 = V(C_n) \setminus (V(P) \cup \{v_{i-1}, v_{j+1}\})$. Since $n \geq 9$, we have $|V_0| \geq 2$ for $|j - i|_n = 4$ or $|V_0| \geq 1$ for $|j - i|_n = 5$. Therefore any vertex in V_0 must have eccentricity at least 4 in G . Moreover, when $|j - i|_n = 5$, v_{i-1} or v_{j+1} has eccentricity 4 in G . This contradicts the fact that G is a 3-ASC graph and we are done. \square

4.3 Trees of order n and diameter $n - 2$

Now we turn to $\theta_3(T)$ for trees T of order n and diameter $n - 2$. Any such tree can be obtained by attaching a pendant vertex to any non-pendant vertex of the path P_{n-1} . Throughout this subsection let $V(T) = \{v_1, \dots, v_n\}$ and $E(T) = \{v_i v_{i+1} : i = 1, \dots, n - 2\} \cup \{v_k v_n\}$, where k is a fixed integer from $\{2, \dots, n - 2\}$.

Lemma 4.5 *If T is a tree of order $n \geq 10$ and diameter $n - 2$, then $\theta_3(T) \leq 2$.*

Proof Based on the value of k introduced above we distinguish two cases.

Case 1 $k = 5$.

Let G be the graph obtained from $T \cup P_2$, where $P_2 = xy$, by adding edges xv_i , $i \in \{1, 2\} \cup \{7, \dots, n - 1\}$ and yv_j , $j \in \{1, 4\} \cup \{10, \dots, n - 1, n\}$. Then it is straightforward to verify that G is a 3-ASC embedding graph of T where v_5, v_9 are two diametrical vertices in G .

Case 2 $k \neq 5$.

In this case we can construct a similar graph H as in Case 1, but now one more edge xv_n is added to H (except that $yv_n \notin E(G)$ for $k = 6$ otherwise $\text{ecc}_G(y) = 2$). It is not difficult to check that H is a 3-ASC embedding graph of T with v_5, v_9 being its unique diametrical vertices. \square

Lemma 4.6 *If T is a tree of order $n \geq 10$ and diameter $n - 2$, then $\theta_3(T) \geq 2$.*

Proof Since T is clearly not a 3-ASC graph, it suffices to prove that $\theta_3(T) > 1$. Suppose on the contrary that there exists a 3-ASC embedding graph G of T with $V(G) = V(T) \cup \{x\}$. We distinguish the following two cases.

Case 1 $xv_n \notin E(G)$.

In this case v_n is pendant in G with $v_k v_n \in E(G)$. Note that G is a 3-ASC graph. Then $\text{ecc}_G(v_n) = 4$ and $\text{ecc}_G(v_k) = 3$.

Suppose first that $xv_k \in E(G)$. Let v_i be the vertex with $d(v_n, v_i) = 4$. Clearly, $d_T(v_k, v_i) \geq 3$ and $xv_i \notin E(G)$. If $v_j \in V(T) \setminus \{v_n, v_i\}$, then $d_G(v_n, v_j) \leq 3$ and $d_G(v_i, v_j) \leq 3$ and consequently $d_G(v_k, v_j) \leq 2$. Obviously, any vertex v_j ($j \neq n, i$) with $d_T(v_j, v_k) \geq 3$ must be adjacent to x in G . If $d_T(v_i, v_k) > 3$, then we get $\text{ecc}_G(x) = 2$, which is impossible because of the 3-ASC property of G . Hence $d_T(v_k, v_i) = 3$ and without loss of generality we may assume that $k < i$. First we claim that $xv_s \in E(G)$ for any vertex v_s with $s < k$. Otherwise, we have $d_G(v_s, v_i) = 4$. Then $\text{ecc}_G(v_s) \geq 4$, contradicting to the 3-ASC property of G . Since $\text{ecc}_G(x) = 3$, any vertex in the (v_i, v_k) -path except v_k has degree 2 in G . Then any vertex v_p with $d_T(v_p, v_{k+2}) > 3$ has the property that $d_G(v_{k+2}, v_p) = 4$. Clearly, $\text{ecc}_G(v_{k+2})$ and $\text{ecc}_G(v_p)$ are at least 4, contradicting the 3-ASC property of G , again.

Now we consider the subcase when $xv_k \notin E(G)$. From $\text{ecc}_G(v_n) = 4$, we find that there is at most one vertex v_p with $d_G(v_n, v_p) = 4$ and $d_G(v_n, v_j) \leq 3$ for any vertex v_j with $j \neq p$. Considering that v_n is pendant in G , all vertices except v_p in T are at distance at most 2 from v_k in G . But, combining the fact that $xv_k \notin E(G)$, we have $d_G(v_j, v_k) \geq 3$ for any vertex v_j with $d_T(v_j, v_k) \geq 3$ with $j \neq p$. A contradiction occurs.

Case 2 $xv_n \in E(G)$.

In this case we first assume that $\text{ecc}_G(v_n) = 4$. Then $\text{ecc}_G(v_k) = 3$ and there exists a vertex v_i with $d_G(v_n, v_i) = 4$, i.e., $d_G(v_k, v_i) = 3$ and $d_G(x, v_i) = 3$. If $i = 1$, we find that $d_G(v_i, v_j) \geq 4$ for any vertex v_j with $k + 1 \leq j \leq n - 1$, which is impossible since G is a 3-ASC graph. By symmetry, $i \neq n - 1$. It follows that v_i has two neighbors v_{i-1} and v_{i+1} in T . Moreover, neither of v_{i-1} and v_{i+1} is adjacent to x in G .

Now we further consider the position of the vertex v_i in T . Clearly, $d_G(v_i, v_k) = 3 \leq d_T(v_i, v_k)$. If $d_T(v_k, v_i) > 3$, then, from $d_G(x, v_i) = 3$, we get $d_G(v_i, v_k) \geq 4$, which is a clear contradiction. Thus $d_T(v_k, v_i) = 3$. If $xv_k \notin E(G)$, then, assuming without loss of generality that $k < i$, we find that $d_G(v_k, v_{i+1}) = 4$, contradicting the

3-ASC property of G . Next we deal with the subcase when $xv_k \in E(G)$. Note that $n \geq 10$. There is at least one vertex v_j in T such that $d_T(v_i, v_j) \geq 4$. Then in this subcase we have $d_G(v_i, v_j) \geq 4$ whenever $v_jx \in E(G)$ or not. A contradiction comes again.

We have thus proved that $\text{ecc}_G(v_n) = 3$ must hold. Next we distinguish the following two subcases.

Subcase 2.1 $xv_k \notin E(G)$.

In this subcase, we first prove $\text{ecc}_G(x) = 3$. If not, then we have $\text{ecc}_G(x) = 4$. Then x must lie on an induced cycle C_8 of G where $v_i \in V(C_8)$ with $d_G(x, v_i) = 4$, or on an induced (x, v_i) -path with $d_G(x, v_i) = 4$ and v_i being pendant in G . In the former case, we can find at least one vertex $v_j \in V(C_8)$ such that $d_G(v_n, v_j) \geq 4$. This is a clear contradiction. In the latter case, we claim that v_i is the other diametrical vertex in G , that is, $d_G(v_i, v_s) \leq 3$ for any vertex v_s with $1 \leq s \leq n$. But this cannot be obeyed for any vertex v_s with $d_T(v_s, v_i) > 3$ for $n \geq 10$.

Next we claim that $\text{ecc}_G(v_k) = 3$ and assume that on the contrary $\text{ecc}_G(v_k) = 4$. Then there is exactly one vertex, say v_i , with $d_G(v_i, v_k) = 4$. Without loss of generality, assume that $i > k$. Evidently, we have $d_T(v_i, v_k) \geq 4$ and $xv_i \notin E(G)$. Considering that $\text{ecc}_G(v_n) = \text{ecc}_G(x) = 3$, from the structure of G , we observe that $d_G(x, v_i) = 2$. From the fact that $d_G(v_i, v_j) \leq 3$ and $d_G(v_k, v_j) \leq 3$ for any vertex $v_j \in V(T) \setminus \{v_i, v_k\}$, we find that any vertex v_j with $j \in \{m : m \leq k-1, \text{ or } m \geq i+1\}$ must be adjacent to x in G .

If $d_T(v_i, v_k) = 4$, then $i = k + 4$. Clearly, $xv_{k+2} \notin E(G)$. If not, we have $\text{ecc}_G(x) = 2$, which is impossible. Since $d_G(v_n, v_{k+3}) \leq 3$, we get $xv_{k+3} \in E(G)$ from the fact that neither of v_{k+2} and v_{k+4} is adjacent to x in G . Moreover, we have $xv_{k+1} \notin E(G)$. Otherwise, we get $\text{ecc}_G(x) = 2$, which is impossible. Then v_{k+1} is just the eccentric vertex of x in G . Thus there must be a vertex v_j ($j \neq k+3, k+4$) with $d_T(v_k, v_j) \geq 3$ such that $d_G(v_{k+1}, v_j) = 4$, a contradiction.

If $d_T(v_i, v_k) > 4$, we have $xv_j \in E(G)$ for any vertex v_j with $k+3 < j < i$, since $d_G(v_k, v_j) \leq 3$. Moreover, neither of v_{k+1} and v_{k+2} is adjacent to x in G , for otherwise $\text{ecc}_G(x) = 2$ would hold. But then $d_G(v_i, v_{k+1}) \geq 4$, a contradiction.

We have thus proved that $\text{ecc}_G(x) = \text{ecc}_G(v_k) = 3$. From $\text{ecc}_G(v_k) = 3$ we find that $xv_j \in E(G)$ for any vertex v_j with $d_T(v_j, v_k) \geq 4$. Because $\text{ecc}_G(x) = 3$, we infer that there exist three consecutive vertices, say v_{p-1}, v_p, v_{p+1} , such that they are at distance at most 3 to v_k and that none of them is adjacent to x in G . It implies that $d_G(x, v_p) = 3$. Since $n \geq 10$, there are at least two vertices v_j with $d_G(v_j, v_p) = 4$, contradicting to the 3-ASC property of G .

Subcase 2.2 $xv_k \in E(G)$.

In this subcase we first assume that $\text{ecc}_G(x) = 3$. Then there exists a vertex v_i with $d_G(x, v_i) = 3$. We may again assume without loss of generality that $i > k$. If v_i is pendant in G , then there are at least two vertices at distance at least 4 from v_i in G , contradicting to the 3-ASC property of G . If v_i is non-pendant in G , then there are three consecutive vertices v_{i-1}, v_i, v_{i+1} , such that none of them is adjacent to x in G and $d_G(x, v_i) = 3$ with $d_T(v_i, v_k) \geq 2$. If $d_T(v_k, v_i) \geq 3$, we find that $d_G(v_n, v_i) = 4$, which is impossible because of $\text{ecc}_G(v_n) = 3$. While $d_T(v_i, v_k) = 2$, for $n \geq 10$, there exist at least two vertices v_p with $d_T(v_i, v_p) \geq 4$ such that $d_G(v_i, v_p) \geq 4$. This contradicts the 3-ASC property of G . So it follows that $\text{ecc}_G(x) = 4$.

If $\text{ecc}_G(x) = 4$, then there is a vertex v_p with $d_G(x, v_p) = 4$. Obviously, $d_T(v_k, v_p) \geq 3$. Considering that v_n, v_k, x form a triangle in G , we get $d_G(v_n, v_p) \geq 4$, the final contradiction. \square

Therefore Lemmas 4.5 and 4.6 immediately imply:

Theorem 4.7 *If T is a tree of order $n \geq 10$ and diameter $n - 2$, then $\theta_3(T) = 2$.*

In Theorem 4.7 we did not consider trees of small order. Note that $K_{1,3}$ is the unique tree of the smallest order of interest (that is, of order 4). For $n = 5$ there is also a unique tree of interest, while for $n = 6$ there are two non-isomorphic trees of diameter 4. Any of these trees is an induced subgraph of C_6^* which is in turn a smallest 3-ASC graph, hence we have the 3-ASC index for $4 \leq n \leq 6$. However, for $7 \leq n \leq 9$ the variety of the trees of interest becomes larger, hence the determination of the 3-ASC index for all of them would be too extensive to be done here.

5 Three open problems

We conclude the paper with three open problems. First, in view of Theorems 2.1 and 2.3 we pose the following very ambitious task:

Problem 5.1 *For any given integer $k \in [1, 5]$ characterize the graphs with $\theta_3(G) = k$.*

Next, in view of Lemma 3.2 and Theorems 3.3–3.7, it is natural to pose:

Problem 5.2 *Characterize the graphs of diameter 2 with the 3-ASC index equal to 3 (equivalently, with the 3-ASC index equal to 4).*

A classical result from random graph theory asserts that almost any graph has diameter 2, cf. Diestel (2006, p. 312, Exercise 7). Since by Lemma 3.2 the 3-ASC index of such a graph is either 3 or 4, our third problem reads as follows.

Problem 5.3 *Is it true that the 3-ASC index of almost every graphs is 3 (resp. 4)?*

Acknowledgements Kexiang Xu is supported by NNSF of China (No. 11671202), China Postdoctoral Science Foundation (2013M530253, 2014T70512). Kinkar Ch. Das is supported by National Research Foundation funded by the Korean government with the Grant No. 2017R1D1A1B03028642. Sandi Klavžar acknowledges the financial support from the Slovenian Research Agency (research core funding No. P1-0297). We are much grateful to two anonymous referees for their careful reading and helpful comments on our paper which have greatly improved the original version of this paper.

References

- Balakrishnan K, Brešar B, Changat M, Klavžar S, Peterin I, Subhamathi AR (2012) Almost self-centered median and chordal graphs. *Taiwan J Math* 16:1911–1922
- Buckley F (1989) Self-centered graphs, *Graph Theory and Its Applications: East and West*. Ann N Y Acad Sci 576:71–78
- Cheng YK, Kang LY, Yan H (2014) The backup 2-median problem on block graphs. *Acta Math Appl Sin Engl Ser* 30:309–320

- Das KC, Nadjafi-Arani MJ (2017) On maximum Wiener index of trees and graphs with given radius. *J Comb Optim* 34:574–587
- Das KC, Lee DW, Graovac A (2013) Some properties of the Zagreb eccentricity indices. *ARS Math Contemp* 6:117–125
- Diestel R (2006) *Graph theory*. Springer, Berlin
- Gupta S, Singh M (2002) Application of graph theory: relationship of eccentric connectivity index and Wiener's index with anti-inflammatory activity. *J Math Anal Appl* 266:259–268
- Gupta S, Singh M, Madan AK (2000) Connective eccentricity index: a novel topological descriptor for predicting biological activity. *J Mol Graph Model* 18:18–25
- Gupta S, Singh M, Madan AK (2002) Eccentric distance sum: a novel graph invariant for predicting biological and physical properties. *J Math Anal Appl* 275:386–401
- Hong YM, Kang LY (2012) Backup 2-center on interval graphs. *Theor Comput Sci* 445:25–35
- Huang TC, Lin JC, Chen HJ (2000) A self-stabilizing algorithm which finds a 2-center of a tree. *Comput Math Appl* 40:607–624
- Ilić A, Yu G, Feng L (2011) On the eccentric distance sum of graphs. *J Math Anal Appl* 381:590–600
- Klavžar S, Narayankar KP, Walikar HB (2011) Almost self-centered graphs. *Acta Math Sin (Engl Ser)* 27:2343–2350
- Klavžar S, Narayankar KP, Walikar HB, Lokesh SB (2014) Almost-peripheral graphs. *Taiwan J Math* 18:463–471
- Klavžar S, Liu H, Singh P, Xu K (2017) Constructing almost peripheral and almost self-centered graphs revisited. *Taiwan J Math* 21:705–717
- Krnc M, Škrekovski R (2015) Group centralization of network indices. *Discrete Appl Math* 186:147–157
- Maddaloni A, Zamfirescu CT (2016) A cut locus for finite graphs and the farthest point mapping. *Discrete Math* 339:354–364
- Morgan MJ, Mukwambi S, Swart HC (2011) On the eccentric connectivity index of a graph. *Discrete Math* 311:1229–1234
- Palacios JL (2015) On the Kirchhoff index of graphs with diameter 2. *Discrete Appl Math* 184:196–201
- Puerto J, Tamir A, Mesa JA, Pérez-Brito D (2008) Center location problems on tree graphs with subtree-shaped customers. *Discrete Appl Math* 156:2890–2910
- Sharma V, Goswami R, Madan AK (1997) Eccentric connectivity index: a novel highly discriminating topological descriptor for structure-property and structure-activity studies. *J Chem Inf Comput Sci* 37:273–282
- Su G, Xiong L, Su X, Chen X (2015) Some results on the reciprocal sum-degree distance of graphs. *J Comb Optim* 30:435–446
- Tomescu I (2008) Properties of connected graphs having minimum degree distance. *Discrete Math* 309:2745–2748
- Tomescu I (2010) Ordering connected graphs having small degree distances. *Discrete Appl Math* 158:1714–1717
- Wang HL, Wu B, Chao KM (2009) The backup 2-center and backup 2-median problems on trees. *Networks* 53:39–49
- Wu B, An X, Liu G, Yan G, Liu X (2013) Minimum degree, edge-connectivity and radius. *J Comb Optim* 26:585–591
- Xu K, Liu M, Das KC, Gutman I, Furtula B (2014) A survey on graphs extremal with respect to distance-based topological indices. *MATCH Commun Math Comput Chem* 71:461–508
- Xu K, Das KC, Liu H (2016) Some extremal results on the connective eccentricity index of graphs. *J Math Anal Appl* 433:803–817
- Xu K, Das KC, Maden AD (2016) On a novel eccentricity-based invariant of a graph. *Acta Math Sin (Engl Ser)* 32:1477–1493
- Yu G, Feng L (2013) On the connective eccentricity index of graphs. *MATCH Commun Math Comput Chem* 69:611–628
- Yu G, Qu H, Tang L, Feng L (2014) On the connective eccentricity index of trees and unicyclic graphs with given diameter. *J Math Anal Appl* 420:1776–1786