



Fast Reaction Limits via Γ -Convergence of the Flux Rate Functional

Mark A. Peletier¹ · D. R. Michiel Renger²

Received: 30 September 2020 / Revised: 30 March 2021 / Accepted: 5 June 2021 /
Published online: 22 July 2021
© The Author(s) 2021, corrected publication 2021

Abstract

We study the convergence of a sequence of evolution equations for measures supported on the nodes of a graph. The evolution equations themselves can be interpreted as the forward Kolmogorov equations of Markov jump processes, or equivalently as the equations for the concentrations in a network of linear reactions. The jump rates or reaction rates are divided in two classes; ‘slow’ rates are constant, and ‘fast’ rates are scaled as $1/\epsilon$, and we prove the convergence in the fast-reaction limit $\epsilon \rightarrow 0$. We establish a Γ -convergence result for the rate functional in terms of both the concentration at each node and the flux over each edge (the level-2.5 rate function). The limiting system is again described by a functional, and characterises both fast and slow fluxes in the system. This method of proof has three advantages. First, no condition of detailed balance is required. Secondly, the formulation in terms of concentration and flux leads to a short and simple proof of the Γ -convergence; the price to pay is a more involved compactness proof. Finally, the method of proof deals with approximate solutions, for which the functional is not zero but small, without any changes.

Keywords Quasi-steady state approximation · Γ -Convergence · Linear network · Rate functional · Fast reaction limit

1 Introduction

The aim of this paper is to prove a fast-reaction limit for a sequence of evolution equations on a graph. We first specify the system.

Let $\mathcal{G} = (\mathcal{V}, \mathcal{R})$ be a finite directed disconnected graph with weights $\kappa^\epsilon : \mathcal{R} \rightarrow [0, \infty)$. For each edge $r \in \mathcal{R}$ we denote $r = (r_-, r_+)$, with $r_-, r_+ \in \mathcal{V}$ the corresponding source and target nodes. We consider the classical problem of deriving effective equations for the flow

✉ D. R. Michiel Renger
renger@wias-berlin.de

¹ Department of Mathematics and Computer Science and Institute for Complex Molecular Systems (ICMS), Eindhoven University of Technology, Eindhoven, The Netherlands

² Weierstrass-Institut für Angewandte Analysis und Stochastik, Berlin, Germany

on $(\mathcal{V}, \mathcal{R})$ with two different rates:

$$\dot{\rho}^\epsilon(t) = -\operatorname{div}(\kappa^\epsilon \otimes \rho^\epsilon(t)), \quad \rho^\epsilon(0) \text{ fixed.} \tag{1.1}$$

with discrete divergence $(\operatorname{div} A)_x := \sum_{r_-=x} A_r - \sum_{r_+=x} A_r$, product $(\kappa^\epsilon \otimes \rho)_{r \in \mathcal{R}} := \kappa_r^\epsilon \rho_{r_-}$, and $t \in [0, T], T > 0$. We assume that the space of edges is a disjoint union $\mathcal{R} = \mathcal{R}_{\text{slow}} \cup \mathcal{R}_{\text{fast}}$ so that

$$\kappa_r^\epsilon = \begin{cases} \kappa_r, & r \in \mathcal{R}_{\text{slow}}, \\ \frac{1}{\epsilon} \kappa_r. & r \in \mathcal{R}_{\text{fast}}. \end{cases} \tag{1.2}$$

We are interested in the limiting behaviour as $\epsilon \rightarrow 0$, where the fast edges equilibrate instantaneously onto a slow manifold. Such limits, also known as ‘Quasi-Steady-State Approximations’, have a long history in the literature, see for example [41,43].

1.1 Γ -Convergence of the Large-Deviations Rate

Often, one is not only interested in convergence of the dynamics, but also in convergence of some variational structure such as a gradient structure, or more generally an ‘action’ functional that is minimised by the dynamics (1.1). Of course this convergence is particularly relevant if this action has a physical meaning. The functional that we study in this paper can be interpreted as an action functional in the following way.

Consider a microscopic system of n independent particles $X_i^\epsilon(t) \in \mathcal{V}, i = 1, \dots, n$ that randomly jump from state $X_i^\epsilon(t_-) = r_-$ to a new state $X_i^\epsilon(t) = r_+$ with Markov intensity κ_r^ϵ . This is a typical microscopic model for a (bio)chemical system of unimolecular reactions with multiple time scales. The concentration of particles in state x is then $\rho_x^{n,\epsilon}(t) := n^{-1} \sum_{i=1}^n \mathbb{1}_{\{X_i^\epsilon(t)=x\}}$, and the vector of random concentrations $\rho^{n,\epsilon}(t)$ converges to the deterministic solution $\rho^\epsilon(t)$ of (1.1) by Kurtz’ classical result [21]. For large but finite particle numbers n , there is a small probability that $\rho^{n,\epsilon}(t)$ deviates significantly from $\rho^\epsilon(t)$. These small probabilities are best understood through a large deviations principle [1,14,22]:

$$-n^{-1} \log \operatorname{Prob} (\rho^{n,\epsilon} \approx \rho) \stackrel{n \rightarrow \infty}{\sim} \mathcal{I}_0^\epsilon(\rho(0)) + \mathcal{I}^\epsilon(\rho), \quad \text{where} \tag{1.3a}$$

$$\mathcal{I}^\epsilon(\rho) := \inf_{\substack{j \in L^1([0,T]; \mathbb{R}^{\mathcal{R}}): \\ \dot{\rho} = -\operatorname{div} j}} \sum_{r \in \mathcal{R}} \int_{[0,T]} s(j_r(t) \mid \kappa_r^\epsilon \rho_{r_-}(t)) dt, \tag{1.3b}$$

$$s(a \mid b) := \begin{cases} a \log \frac{a}{b} - a + b & \text{for } a, b > 0, \\ b & \text{for } a = 0, b \geq 0, \\ \infty & \text{otherwise.} \end{cases} \tag{1.3c}$$

and \mathcal{I}_0^ϵ reflects whatever randomness is taken for the initial concentration $\rho^{n,\epsilon}(0)$. We stress that this formula is typical for Markov jump processes; choosing a different microscopic model for the dynamics could lead to different functionals.

If the network satisfies detailed balance, then the rate functional (1.3b) can be related to a gradient flow [30,31,34,35]. We shall revisit the detailed balance condition in Sect. 1.9. For a similar interpretation in terms of an action without the detailed balance condition, see [3,38].

Note that \mathcal{I}^ϵ is indeed minimised by solutions ρ^ϵ of (1.1). This implies that we can consider the equation $\mathcal{I}^\epsilon = 0$ as a variational formulation of the Eq. (1.1); this is the point of view known as ‘curves of maximal slope’ [2] or the ‘energy-dissipation principle’ [27]. An important advantage of this choice of formulation is that Γ -convergence of \mathcal{I}^ϵ implies converge of the minimising dynamics (see [11, Cor. 7.24] and [27]); in other words, one can

prove convergence of the solutions by proving Γ -convergence of the functionals. This is also the method that we adopt in this paper.

1.2 Γ -Convergence of the Flux Large-Deviations Rate

One difficulty in proving Γ -convergence of the functional $\mathcal{I}_0^\epsilon + \mathcal{I}^\epsilon$, however, is that \mathcal{I}^ϵ is implicitly defined by a constrained minimisation problem. The constrained infimum of the sum in (1.3b) is an infimal convolution (see [30, Sec. 3.4]). This shows that the evolution of the concentrations in different nodes are strongly intertwined, which considerably complicates the mathematical analysis. For example, the related work [10] requires an orthogonality assumption to decouple the concentrations.

We can however avoid this difficulty by considering a different functional instead. Observe that the variable $j_r(t)$ in (1.3b) has the interpretation of a flux: it measures how much mass is transported through edge r at time t . Naturally, one can rephrase (1.1) in terms of this flux as the coupled system

$$\dot{\rho}^\epsilon(t) = -\operatorname{div} j^\epsilon(t) \quad \text{and} \quad j^\epsilon(t) = \kappa^\epsilon \otimes \rho^\epsilon(t), \quad \rho^\epsilon(0) \text{ fixed.} \tag{1.4}$$

On the level of the microscopic particle system one can also define the random particle flux $J^{n,\epsilon}$, which yields the large-deviation principle [4,37,38]:

$$-n^{-1} \log \operatorname{Prob}((\rho^{n,\epsilon}, J^{n,\epsilon}) \approx (\rho, j)) \stackrel{n \rightarrow \infty}{\sim} \mathcal{I}_0^\epsilon(\rho(0)) + \mathcal{J}^\epsilon(\rho, j), \quad \text{where} \tag{1.5}$$

$$\mathcal{J}^\epsilon(\rho, j) := \begin{cases} \sum_{r \in \mathcal{R}} \int_{[0, T]} s(j_r(t) \mid \kappa_r^\epsilon \rho_{r-}(t)) dt, & \text{if } \rho \in W^{1,1}([0, T]; \mathbb{R}^{\mathcal{V}}), j \in L^1([0, T]; \mathbb{R}^{\mathcal{R}}), \\ & \text{and } \dot{\rho} = -\operatorname{div} j, \\ \infty, & \text{otherwise.} \end{cases} \tag{1.6}$$

Indeed, the functional \mathcal{J}^ϵ is related to (1.3b) by $\mathcal{I}^\epsilon(\rho) = \inf_{\dot{\rho} = -\operatorname{div} j} \mathcal{J}^\epsilon(\rho, j)$, which is consistent with the ‘contraction principle’ in large-deviations theory. Its minimiser (1.4) follows the same evolution as the minimiser (1.1), but provides with more information: the flux. From a physics perspective, this additional information is important to understand non-equilibrium thermodynamics; see for example [3,30] and [38, Sec. 4]. From a mathematical perspective, we will use the property that the flux functional \mathcal{J}^ϵ is a sum over edges to decompose networks into separate components.

The goal of this paper is to prove convergence of the functional $\mathcal{I}_0^\epsilon + \mathcal{J}^\epsilon$ to a limit functional, whose minimiser describes the effective dynamics for (1.4). As a consequence, we obtain Γ -convergence of the functional $\mathcal{I}_0^\epsilon + \mathcal{I}^\epsilon$, convergence of solutions of the flux ODE (1.4), and convergence of solutions of the ODE (1.1).

In order to track diverging fluxes and vanishing concentrations, we shall introduce a number of rescalings before taking the Γ -limit, as we explain in the next section.

1.3 Network Decomposition: Nodes

We decompose the network into different components according to their scaling behaviour. To explain the main ideas, consider the example of Fig. 1. Recall from (1.2) that we assume that $\mathcal{R} = \mathcal{R}_{\text{slow}} \cup \mathcal{R}_{\text{fast}}$, where the slow edges have rates of order 1, and the fast edges of order $1/\epsilon$.




Fig. 1 An example of a network with slow and fast edges

The first step in the decomposition is to categorise the nodes. In the example, node 5 is expected to have low concentration, since any mass at node 5 will be quickly transported to node 4. We make this statement precise by considering the equilibrium concentration. Since we assume the network to be disconnected, there exists a unique equilibrium concentration $0 < \pi^\epsilon \in \mathbb{R}^{\mathcal{V}}$ for the dynamics (1.1); we will always assume that π^ϵ is normalised, i.e. $\sum_{x \in \mathcal{V}} \pi_x^\epsilon = 1$. We use the equilibrium concentrations to subdivide the nodes into two classes, $\mathcal{V} = \mathcal{V}_0 \cup \mathcal{V}_1$, where

$$\mathcal{V}_0 := \left\{ x \in \mathcal{V} : \pi_x^\epsilon \xrightarrow{\epsilon \rightarrow 0} \pi_x > 0 \right\}, \quad \text{and} \quad \mathcal{V}_1 := \left\{ x \in \mathcal{V} : \frac{1}{\epsilon} \pi_x^\epsilon \xrightarrow{\epsilon \rightarrow 0} \tilde{\pi}_x > 0 \right\}, \quad (1.7)$$

and the tilde is used to stress that the quantity is rescaled. This decomposition implies that π_x^ϵ is either of order 1 or of order ϵ . In fact, one can construct networks with $\mathcal{R} = \mathcal{R}_{\text{slow}} \cup \mathcal{R}_{\text{fast}}$ with stationary states π_x^ϵ of order ϵ^2, ϵ^3 , or higher, but in this paper such networks will be ruled out by our assumption that there are no “leak” fluxes (see below). We introduce a further subdivision of the nodes after categorising the fluxes.

1.4 Network Decomposition: Fluxes

We expect that j_r^ϵ is comparable to $\kappa_r^\epsilon \rho_{r_-}^\epsilon$, which in turn we expect to be comparable to $\kappa_r^\epsilon \pi_{r_-}^\epsilon$. Hence the flux or amount of mass being transported through an edge r not only depends on the order of κ_r^ϵ , but also on the amount of available mass in the source node r_- , of order $\pi_{r_-}^\epsilon$. Therefore the scaling behaviour of the flux falls into one of the following four different categories:

j_r^ϵ	$r_- \in \mathcal{V}_0$		$r_- \in \mathcal{V}_1$	
$r \in \mathcal{R}_{\text{slow}}$	$\mathcal{O}(1)$	“slow”	$\mathcal{O}(\epsilon)$	“leak”
$r \in \mathcal{R}_{\text{fast}}$	$\mathcal{O}(1/\epsilon)$	“fast cycle”	$\mathcal{O}(1)$	“damped”

In this paper we rule out “leak” fluxes by assumption, so that $\mathcal{R} = \mathcal{R}_{\text{slow}} \cup \mathcal{R}_{\text{fctc}} \cup \mathcal{R}_{\text{damp}}$, with

$$\mathcal{R}_{\text{fctc}} := \{ r \in \mathcal{R}_{\text{fast}} : r_- \in \mathcal{V}_0 \} \quad \text{and} \quad \mathcal{R}_{\text{damp}} := \{ r \in \mathcal{R}_{\text{fast}} : r_- \in \mathcal{V}_1 \}.$$

Let us now explain these four categories in more detail by considering the example network of Fig. 1, which can now be redrawn as Fig. 2.




Fig. 2 The example from Fig. 1, redrawn using the categorisation of nodes and fluxes (left); the final reduction to a two-node network in the limit $\epsilon \rightarrow 0$ (right)

1. What we shall call the *slow fluxes* are fluxes through a slow edge that start at a node in \mathcal{V}_0 . Typically, these slow fluxes will be of order $\mathcal{O}(1)$, and they depend on ϵ only indirectly through dependence on the other fluxes.
2. For the fast edges however, there is a fundamental difference between the fluxes $1 \rightarrow 2 \rightarrow 3 \rightarrow 1$ and the flux $5 \rightarrow 4$. The three fluxes $1 \rightarrow 2 \rightarrow 3 \rightarrow 1$ constitute a cycle of fast edges, with fluxes of order $\mathcal{O}(1/\epsilon)$. Therefore mass will rotate very fast through this cycle, and in the limit $\epsilon \rightarrow 0$, the mass present in the cycle will instantaneously equilibrate over these three edges. Moreover, any mass inserted into this cycle through the slow flux $4 \rightarrow 1$ will also instantaneously equilibrate over the nodes in the cycle, and any mass removed from the cycle through the slow flux $2 \rightarrow 5$ may be withdrawn from any node in the cycle. Practically this means that in the limit the cycle/diconnected component $1 \rightarrow 2 \rightarrow 3 \rightarrow 1$ acts as one node $c := \{1, 2, 3\}$. We shall see in Lemma 3.2 that all edges with $r \in \mathcal{R}_{\text{fast}}$ and $r_- \in \mathcal{V}_0$ are indeed part of a cycle, which justifies the name *fast cycle*.
3. By contrast, the fast edge $5 \rightarrow 4$ is not part of a fast diconnected component. One does expect mass in node 5 to be transported very fast into node 4, but since there is no fast inflow, the mass in node 5 will be strongly depleted after the initial time. After this, the amount of mass that will be actually transported through edge $5 \rightarrow 4$ is fully subject to the amount of inflow of mass into node 5 by the slow fluxes $2 \rightarrow 5$ and $4 \rightarrow 5$, and will therefore be of $\mathcal{O}(1)$. We shall call the flux $5 \rightarrow 4$ a *damped flux*; its corresponding edge is fast, but the flux is damped by the fact that there is not enough mass available in the source node 5. In the limit, any mass that is inserted into node 5 from node 2 or 4 will be immediately pushed into node 4.
4. Now imagine a flux $5 \rightarrow 1$, not drawn in the picture. Since there is a damped flux going out of node 5, almost all mass from node 5 will follow that flux into node 4, whereas very little mass from node 5 would leak away into node 1. We shall call such fluxes *leak fluxes*. Since they contribute little to the behaviour of the whole network we rule out this possibility by assumption. As mentioned above, this also rules out the possibility of higher orders of π_x^ϵ , see Lemma 3.1.

An even further subdivision of $\mathcal{R}_{\text{damp}}$ will be discussed in Sect. 1.8, but this will not be needed in the general discussion.

1.5 Network Decomposition: Connected Components

After categorising the fluxes, we now further subdivide the nodes of \mathcal{V}_0 into $\mathcal{V}_0 = \mathcal{V}_{0\text{fcyc}} \cup \mathcal{V}_{0\text{slow}}$, consisting of nodes that are part of a fast cycle and the remainder:

$$\mathcal{V}_{0\text{fcyc}} := \{x \in \mathcal{V}_0 : \exists r \in \mathcal{R}_{\text{fcyc}}, r_- = x\}, \quad \text{and} \quad \mathcal{V}_{0\text{slow}} := \mathcal{V}_0 \setminus \mathcal{V}_{0\text{fcyc}}.$$

The notation reflects the expectation that the concentration in the nodes in $\mathcal{V}_{0\text{fcyc}}$ will instantaneously equilibrate over the disconnected components of the graph $(\mathcal{V}_0, \mathcal{R}_{\text{fcyc}})$. We collect these components in the set

$$\mathfrak{C} := \left\{ \mathfrak{c} \subset \mathcal{V}_0 : \forall x, y \in \mathfrak{c}, \exists (r^k)_{k=1}^K \subset \mathcal{R}_{\text{fcyc}}, r_-^1 = x, r_+^k = r_-^{k+1} \text{ for } k = 1, \dots, K-1, r_+^K = y \right\}.$$

To each $\mathfrak{c} \in \mathfrak{C}$ corresponds the equilibrium mass

$$\pi_{\mathfrak{c}}^\epsilon := \sum_{x \in \mathfrak{c}} \pi_x^\epsilon, \quad \mathfrak{c} \in \mathfrak{C}. \tag{1.8}$$

We will see in Lemma 3.2 that a component $\mathfrak{c} \in \mathfrak{C}$ can be written as a union of cycles in the graph $(\mathcal{V}_{0\text{fcyc}}, \mathcal{R}_{\text{fcyc}})$. Consequently, if there exists a fast-cycle path from x to y then there also exists a fast-cycle path from y to x . This remark also implies that each fast component \mathfrak{c} is a subset of $\mathcal{V}_{0\text{fcyc}}$.

Observe that, as suggested in Fig. 2, certain combinations of nodes become slaved to each other, in the sense that their concentrations move in unison. This is for instance the case for all nodes in a fast cycle. It is therefore common in the literature to pass to the limit by a coarse-grained description that neglects the difference between those nodes; see for instance [5,16,33] or more generally [18, Ch. 4] or [20, Ch. 2]. By contrast, we preserve the information about the separate nodes and we keep track of the fast cycle as well as the fluxes between these nodes. This is motivated by our Theorem 1.1, which yields sufficient compactness in the \mathcal{V}_1 -concentrations, damped fluxes and fast cycle fluxes. The Γ -convergence of the coarse-grained functional follows directly from our result, see Remark 6.2.

1.6 Rescaled Flux and Initial Functionals

In Sects. 1.3 and 1.4 we categorised the nodes and fluxes by their typical scaling behaviour. We shall prove that the scaling behaviour of these categories is not only typical for the effective dynamics but actually for any dynamics with finite large-deviation cost. In order to do so we rescale all concentrations and fluxes according to their respective scalings.

We expect concentrations ρ_x^ϵ to follow π_x^ϵ , and therefore to be of order order 1 on \mathcal{V}_0 and of order ϵ on \mathcal{V}_1 . This motivates the rescaling the concentrations by working with the densities u^ϵ , defined by

$$u_x^\epsilon(t) := \frac{\rho_x^\epsilon(t)}{\pi_x^\epsilon(t)},$$

where $x \in \mathcal{V} \cup \mathfrak{C}$, using (1.8). Although $\mathcal{V}_{0\text{fcyc}} = \bigcup \mathfrak{C}$, we study $u_x^\epsilon(t)$ for $x \in \mathcal{V}_{0\text{slow}} \cup \mathcal{V}_{0\text{fcyc}} \cup \mathfrak{C}$, assuming that $u_{\mathfrak{c}}^\epsilon$ and $u_x^\epsilon, x \in \mathfrak{c}$ are related by

$$\pi_{\mathfrak{c}}^\epsilon u_{\mathfrak{c}}^\epsilon(t) := \sum_{x \in \mathfrak{c}} \pi_x^\epsilon u_x^\epsilon(t), \tag{1.9}$$

which we consider as a special continuity equation, additional to $\rho^\epsilon = -\operatorname{div} j^\epsilon$. The distinction between u_x^ϵ and u_c^ϵ allows for two different notions of compactness: a weaker compactness for u_x^ϵ with $x \in \mathcal{V}_{0\text{fcyc}}$, and a stronger compactness for u_c^ϵ for any $c \in \mathcal{C}$.

As explained in Sect. 1.4, the fluxes are expected to scale as $j_r^\epsilon(t) = \mathcal{O}(\kappa_r^\epsilon \pi_{r-}^\epsilon)$. The slow and damped fluxes are of order 1 and therefore need not be rescaled. For fast cycle fluxes, of order $1/\epsilon$, we introduce the rescaled flux \tilde{j}_r^ϵ , defined by

$$j_r^\epsilon(t) =: \kappa_r^\epsilon \rho_{r-}^\epsilon(t) + \frac{1}{\sqrt{\epsilon}} \tilde{j}_r^\epsilon(t) = \frac{1}{\epsilon} \kappa_r \pi_{r-}^\epsilon(t) u_{r-}^\epsilon(t) + \frac{1}{\sqrt{\epsilon}} \tilde{j}_r^\epsilon(t), \quad r \in \mathcal{R}_{\text{fcyc}}.$$

It turns out that this deviation from $\kappa_r^\epsilon \rho_{r-}^\epsilon$ of order $1/\sqrt{\epsilon}$ is the right choice for sequences along which $\mathcal{I}_0^\epsilon + \mathcal{J}^\epsilon$ is bounded, since this scaling is natural in the context of the compactness and Γ -limit results that we prove below.

To shorten the expressions we shall write

$$\begin{aligned} u_{\mathcal{V}_{0\text{slow}}} &:= (u_x)_{x \in \mathcal{V}_{0\text{slow}}}, & u_{\mathcal{V}_{0\text{fcyc}}} &:= (u_x)_{x \in \mathcal{V}_{0\text{fcyc}}}, & u_{\mathcal{C}} &:= (u_c)_{c \in \mathcal{C}}, & u_{\mathcal{V}_1} &:= (u_x)_{x \in \mathcal{V}_1}, \\ \tilde{j}_{\mathcal{R}_{\text{slow}}} &:= (\tilde{j}_r)_{r \in \mathcal{R}_{\text{slow}}}, & \tilde{j}_{\mathcal{R}_{\text{damp}}} &:= (\tilde{j}_r)_{r \in \mathcal{R}_{\text{damp}}}, & \tilde{j}_{\mathcal{R}_{\text{fcyc}}} &:= (\tilde{j}_r)_{r \in \mathcal{R}_{\text{fcyc}}}, \end{aligned}$$

and finally by a slight abuse of notation $(u, j) := (u_{\mathcal{V}_{0\text{slow}}}, u_{\mathcal{V}_{0\text{fcyc}}}, u_{\mathcal{C}}, u_{\mathcal{V}_1}, \tilde{j}_{\mathcal{R}_{\text{slow}}}, \tilde{j}_{\mathcal{R}_{\text{damp}}}, \tilde{j}_{\mathcal{R}_{\text{fcyc}}})$. With these rescalings and notation we now rewrite the large-deviations rate functional (1.6) as:

$$\begin{aligned} \tilde{\mathcal{J}}^\epsilon(u, j) &= \tilde{\mathcal{J}}^\epsilon(u_{\mathcal{V}_{0\text{slow}}}, u_{\mathcal{V}_{0\text{fcyc}}}, u_{\mathcal{C}}, u_{\mathcal{V}_1}, \tilde{j}_{\mathcal{R}_{\text{slow}}}, \tilde{j}_{\mathcal{R}_{\text{damp}}}, \tilde{j}_{\mathcal{R}_{\text{fcyc}}}) \\ &:= \mathcal{J}^\epsilon(\pi^\epsilon u, (j_{\mathcal{R}_{\text{slow}}}, j_{\mathcal{R}_{\text{damp}}}, \epsilon^{-1} \kappa \otimes \pi^\epsilon u^\epsilon + \epsilon^{-1/2} \tilde{j}_{\mathcal{R}_{\text{fcyc}}})) \\ &= \sum_{r \in \mathcal{R}_{\text{slow}}} \underbrace{\int_{[0, T]} s(j_r(t) \mid \kappa_r \pi_{r-}^\epsilon u_{r-}(t)) dt}_{=: \tilde{\mathcal{J}}_{\text{slow}}^\epsilon(u_{\mathcal{V}_{0\text{slow}}}, u_{\mathcal{V}_{0\text{fcyc}}})} \\ &\quad + \sum_{r \in \mathcal{R}_{\text{damp}}} \underbrace{\int_{[0, T]} s(j_r(t) \mid \frac{1}{\epsilon} \kappa_r \pi_{r-}^\epsilon u_{r-}(t)) dt}_{=: \tilde{\mathcal{J}}_{\text{damp}}^\epsilon(u_{\mathcal{V}_1}, \tilde{j}_{\mathcal{R}_{\text{damp}}})} \\ &\quad + \sum_{r \in \mathcal{R}_{\text{fcyc}}} \underbrace{\int_{[0, T]} s\left(\frac{1}{\epsilon} \kappa_r \pi_{r-}^\epsilon u_{r-}(t) + \frac{1}{\sqrt{\epsilon}} \tilde{j}_r(t) \mid \frac{1}{\epsilon} \kappa_r \pi_{r-}^\epsilon u_{r-}(t)\right) dt}_{=: \tilde{\mathcal{J}}_{\text{fcyc}}^\epsilon(u_{\mathcal{V}_{0\text{fcyc}}}, \tilde{j}_{\mathcal{R}_{\text{fcyc}}})}, \end{aligned} \tag{1.10}$$

where $\tilde{\mathcal{J}}^\epsilon = \infty$ if the finiteness condition of (1.6) or $\tilde{\mathcal{J}}$ condition (1.9) is violated. Recall that $\pi_{r-}^\epsilon \approx \pi_{r-}$ for $r \in \mathcal{R}_{\text{slow}}$ and $\pi_{r-}^\epsilon \approx \epsilon \tilde{\pi}_{r-}$ for $r \in \mathcal{R}_{\text{damp}}$, so that the two functionals $\tilde{\mathcal{J}}_{\text{slow}}^\epsilon$ and $\tilde{\mathcal{J}}_{\text{damp}}^\epsilon$ are very similar.

In order to control the initial condition we include the initial large-deviation rate function \mathcal{I}_0^ϵ in the analysis. As mentioned in Sect. 1.1, this function depends on the choice of the initial probability. As is common, we choose the random dynamics to start independently at the invariant measure. Since linear reactions correspond to independent copies of the process, the particles modelled by the invariant measure are also independent, and hence $\mathcal{I}_0^\epsilon(\rho(0)) = \sum_{x \in \mathcal{V}} s(\rho_x(0) \mid \pi^\epsilon)$ by Sanov’s Theorem [12, Th. 6.2.10]. We again rescale

this functional to work with densities instead:

$$\begin{aligned} \tilde{\mathcal{I}}_0^\epsilon(u(0)) &= \tilde{\mathcal{I}}_0^\epsilon(u_{\mathcal{V}_{0\text{slow}}}(0), u_{\mathcal{V}_{0\text{fcyc}}}(0), u_{\mathcal{C}}(0), u_{\mathcal{V}_1}(0)) := \mathcal{I}_0^\epsilon(\pi^\epsilon \otimes u(0)) \\ &:= \sum_{x \in \mathcal{V}_{0\text{slow}}} s(\pi_x^\epsilon u_x(0) \mid \pi_x^\epsilon) + \sum_{x \in \mathcal{V}_{0\text{fcyc}}} s(\pi_x^\epsilon u_x(0) \mid \pi_x^\epsilon) + \sum_{x \in \mathcal{V}_1} s(\pi_x^\epsilon u_x(0) \mid \pi_x^\epsilon). \end{aligned} \tag{1.11}$$

The minimiser of $\tilde{\mathcal{I}}_0$ is the vector $u(0) \equiv 1$.

1.7 Main Results: Compactness and Γ -Convergence

We now focus on the Γ -limit of the rescaled functional $\tilde{\mathcal{I}}_0^\epsilon + \tilde{\mathcal{J}}^\epsilon$, in the space

$$\begin{aligned} \Theta &:= C([0, T]; \mathbb{R}^{\mathcal{V}_{0\text{slow}}}) \times L^\infty([0, T]; \mathbb{R}^{\mathcal{V}_{0\text{fcyc}}}) \times C([0, T]; \mathbb{R}^{\mathcal{C}}) \times \mathcal{M}([0, T]; \mathbb{R}^{\mathcal{V}_1}) \\ &\quad \times L^{\mathcal{C}}([0, T]; \mathbb{R}^{\mathcal{R}_{\text{slow}}}) \times \mathcal{M}([0, T]; \mathbb{R}^{\mathcal{R}_{\text{damp}}}) \times L^{\mathcal{C}}([0, T]; \mathbb{R}^{\mathcal{R}_{\text{fcyc}}}), \end{aligned}$$

where C is the space of continuous functions, \mathcal{M} denotes spaces of bounded measures, and $L^{\mathcal{C}}$ denote Orlicz spaces corresponding to the nice Young function (see Sect. 2.2):

$$\mathcal{C}(a) := \inf_{p-q=a} s(p \mid 1) + s(q \mid 1).$$

We always make the implicit assumption that $u_{\mathcal{C}}$ and $u_{\mathcal{V}_{0\text{fcyc}}}$ are connected by (1.9).

We make Θ into a topological space by equipping each space C with the uniform topology, each L^∞ and $L^{\mathcal{C}}$ with their weak-* topologies and each measure space \mathcal{M} with the narrow topology (defined by duality with continuous functions).

Of course Γ -Convergence properties strongly depend on the chosen topology. In fact, it is known that different topologies may lead to different Γ -limits [11, Ch. 6], [28, Sec. 2.6]. The choice of this particular topological space Θ is motivated by our first main result:

Theorem 1.1 (Equicoercivity) *Let $(u^\epsilon, j^\epsilon)_{\epsilon>0} \subset \Theta$ such that*

$$\tilde{\mathcal{I}}_0^\epsilon(u^\epsilon(0)) + \tilde{\mathcal{J}}^\epsilon(u^\epsilon, j^\epsilon) \leq C \quad \text{for some } C > 0.$$

Then there exists a Θ -convergent subsequence.

This equicoercivity identifies a topology that is generated by the sequence of functionals itself, and therefore natural for the Γ -convergence. Note that the topologies for $u_{\mathcal{V}_0}$ and $u_{\mathcal{C}}$ are much stronger than the other ones. This will be needed to interchange limits $\lim_{\epsilon \rightarrow 0} \lim_{t \downarrow 0} u_{\mathcal{V}_{0\text{slow}}}^\epsilon(t)$ and $\lim_{t \downarrow 0} \lim_{\epsilon \rightarrow 0} u_{\mathcal{V}_{0\text{slow}}}^\epsilon(t)$ in order to converge in the continuity equation later on. By contrast, such strong compactness is not to be expected for $u_{\mathcal{V}_1}^\epsilon$, nor is it needed, since the $u_{\mathcal{V}_1}^\epsilon(0)$ will not play a role in the limit due to instantaneous equilibration.

Our second main result is the Γ -convergence:

Theorem 1.2 *In the topological space Θ :*

$$\tilde{\mathcal{I}}_0^\epsilon + \tilde{\mathcal{J}}_{\text{slow}}^\epsilon + \tilde{\mathcal{J}}_{\text{fcyc}}^\epsilon + \tilde{\mathcal{J}}_{\text{damp}}^\epsilon =: \tilde{\mathcal{I}}_0^\epsilon + \tilde{\mathcal{J}}^\epsilon \xrightarrow{\epsilon \rightarrow 0} \tilde{\mathcal{I}}_0^0 + \tilde{\mathcal{J}}^0 := \tilde{\mathcal{I}}_0^0 + \tilde{\mathcal{J}}_{\text{slow}}^0 + \tilde{\mathcal{J}}_{\text{fcyc}}^0 + \tilde{\mathcal{J}}_{\text{damp}}^0,$$

where, setting $u_{r-} := u_c$ for any $r_- \in \mathfrak{c}$,

$$\begin{aligned} \tilde{\mathcal{I}}_0^0(u(0)) &:= \sum_{x \in \mathcal{V}_{0slow}} s(\pi_x u_x(0) \mid \pi_x) + \sum_{\mathfrak{c} \in \mathfrak{C}} s(\pi_{\mathfrak{c}} u_{\mathfrak{c}}(0) \mid \pi_{\mathfrak{c}}), \\ \tilde{\mathcal{J}}_{slow}^0(u_{\mathcal{V}_{0slow}}, u_{\mathcal{V}_{0fyc}}, j_{\mathcal{R}_{slow}}) &:= \sum_{r \in \mathcal{R}_{slow}} \int_{[0, T]} s(j_r(t) \mid \kappa_r \pi_{r-} u_{r-}(t)) dt, \\ \tilde{\mathcal{J}}_{damp}^0(u_{\mathcal{V}_1}, j_{\mathcal{R}_{damp}}) &:= \sum_{r \in \mathcal{R}_{damp}} \int_{[0, T]} s(j_r \mid \kappa_r \tilde{\pi}_{r-} u_{r-})(dt), \\ \tilde{\mathcal{J}}_{fyc}^0(u_{\mathcal{V}_{0fyc}}, \tilde{\mathcal{J}}_{\mathcal{R}_{fyc}}) &:= \frac{1}{2} \sum_{r \in \mathcal{R}_{fyc}} \int_{[0, T]} \frac{\tilde{j}_r(t)^2}{\kappa_r \pi_{r-} u_{r-}(t)} dt, \end{aligned}$$

and we set $\tilde{\mathcal{J}}^0 = \infty$ if the limit continuity equations (3.14) are violated.

The explicit form (3.14) of the limit continuity equations will be derived in Lemma 3.14, after the required notions are introduced and the required results about the network and continuity equations are proven. In our third main result, explained in the next section, we show that both the densities $u_{\mathcal{V}_1}$ and the damped fluxes $j_{\mathcal{R}_{damp}}$ may become measure-valued in time; therefore we use a slight generalisation of the function s to measure-valued trajectories, i.e.:

$$\int_{[0, T]} s(j_r \mid \kappa_r \tilde{\pi}_{r-} u_{r-})(dt) := \begin{cases} \int_{[0, T]} s\left(\frac{dj_r}{\kappa_r \tilde{\pi}_{r-} du_{r-}}(t) \mid 1\right) \kappa_r \tilde{\pi}_{r-} u_{r-}(dt), & \text{if } j_r \ll u_{r-}, \\ \infty, & \text{otherwise.} \end{cases} \quad (1.12)$$

Comparing Theorem 1.2 with Fig. 1, we see that the limit functional contains additional information about the \mathcal{V}_1 nodes that contract to a single node in the limit, and about all slow, fast cycle and damped fluxes. Due to this additional information, the proof of the Γ -convergence is relatively straightforward, e.g. without the need of unfolding techniques. This illustrates our ‘philosophical’ message that the mathematics becomes easier if one takes fluxes into account, which was also observed in [37] where the large-deviation principle (1.6) was proven.

1.8 Main Result: The Development of Spikes

The equicoercivity of $u_{\mathcal{V}_1}^\epsilon$ and j_{damp}^ϵ will be derived by uniform L^1 -bounds in Lemmas 3.5 and 3.7. From these bounds one can only extract compactness as measures, in the narrow sense, so that $u_{\mathcal{V}_1}^\epsilon$ and j_{damp}^ϵ may develop measure-valued singularities or *spikes* in time.

For the densities $u_{\mathcal{V}_1}^\epsilon$, such spikes can not be ruled out, regardless of the network structure. This is easy to see from the fact that these densities become fully uncoupled in the limit continuity equation (3.14d). From (1.12) one sees that one may choose large u_r for $r \in \mathcal{V}_1$, provided $j_r \ll u_{r-}$.

For the fluxes j_{damp}^ϵ , the occurrence of spikes is related to the presence of *damped cycles*, i.e. cycles of damped fluxes. The example of Figs. 1 and 2 has no such damped cycles, but Fig. 3 illustrates the concept.




Fig. 3 An example of a network with a cycle of damped fluxes. On the left the network with the distinction between fast and slow edges; on the right the redrawn network with the node and flux classification as in Fig. 2

To study this we further subdivide $\mathcal{R}_{\text{damp}}$ into damped cycles and the rest, $\mathcal{R}_{\text{damp}} = \mathcal{R}_{\text{dcyc}} \cup \mathcal{R}_{\text{dnocyc}}$, where

$$\mathcal{R}_{\text{dcyc}} := \{r^0 \in \mathcal{R}_{\text{damp}} : \exists (r^k)_{k=1}^K \subset \mathcal{R}_{\text{damp}}, r_+^k = r_-^{k-1} \text{ for } k = 1, \dots, K - 1, r_+^K = r_-^0\},$$

$$\mathcal{R}_{\text{dnocyc}} := \mathcal{R}_{\text{damp}} \setminus \mathcal{R}_{\text{dcyc}}.$$

The relation between damped cycles and spikes in the damped fluxes is summarised in our third main result:

Theorem 1.3 (i) For any sequence $(u^\epsilon, j^\epsilon)_{\epsilon>0} \subset \Theta$ such that $\tilde{\mathcal{I}}_0^\epsilon(u^\epsilon(0)) + \tilde{\mathcal{J}}^\epsilon(u^\epsilon, j^\epsilon) \leq C$ for some $C > 0$ and $(u^\epsilon, j^\epsilon) \xrightarrow{\Theta} (u, j)$, we have $j_{\mathcal{R}_{\text{dnocyc}}} \in L^{\mathcal{C}}([0, T]; \mathbb{R}^{\mathcal{R}_{\text{dnocyc}}})$.

(ii) If $\mathcal{R}_{\text{dcyc}} \neq \emptyset$ then there exists a sequence $(u^\epsilon, j^\epsilon)_{\epsilon>0} \subset \Theta$ with $\tilde{\mathcal{I}}_0^\epsilon(u^\epsilon(0)) + \tilde{\mathcal{J}}^\epsilon(u^\epsilon, j^\epsilon) \leq C$ for some $C > 0$ and $(u^\epsilon, j^\epsilon) \xrightarrow{\Theta} (u, j)$ such that

$$j_{\mathcal{R}_{\text{dcyc}}} \in \mathcal{M}([0, T]; \mathbb{R}^{\mathcal{R}_{\text{dcyc}}}) \setminus L^1([0, T]; \mathbb{R}^{\mathcal{R}_{\text{dcyc}}}).$$

As a consequence of Theorems 1.2 and 1.3(ii), if $\mathcal{R}_{\text{dcyc}} \neq \emptyset$ then there is a $(u, j) \in \Theta$ with $j_{\mathcal{R}_{\text{dcyc}}} \in \mathcal{M}([0, T]; \mathbb{R}^{\mathcal{R}_{\text{dcyc}}}) \setminus L^1([0, T]; \mathbb{R}^{\mathcal{R}_{\text{dcyc}}})$ for which $\tilde{\mathcal{I}}_0^\epsilon(u(0)) + \tilde{\mathcal{J}}^\epsilon(u, j) \leq \infty$.

1.9 Related Literature

As mentioned in the introduction, this work is related to classical quasi-steady state approximation theory; see e.g. [18, Sec. 4.2] or [20, Sec. 3.1]. We mention two recent works [10,33] that study fast-reaction limits in connection with another underlying structure, namely a gradient structure. A gradient structure consists of an energy $\frac{1}{2}\mathcal{I}_0^\epsilon(c)$ and a non-negative convex dissipation potential $\Psi_c^\epsilon(\xi)$ such that the evolution equation (1.1) can be rewritten as $D\Psi_{c(t)}^\epsilon(\dot{c}(t)) = -D\frac{1}{2}\mathcal{I}_0^\epsilon(c(t))$. Both studies work on the level of concentrations rather than fluxes, under the assumption that the ϵ -dependent evolution equation (1.1) satisfies detailed balance, and under the assumption that damped fluxes do not occur. The detailed balance condition is needed for the ϵ -dependent equation to have a gradient structure, and the absence of damped fluxes guarantees that the gradient structure is not destroyed in the limit.

Disser et al. [10] study general, possibly non-linear reaction networks with mass-action kinetics. Under the detailed balance assumption such equations have a gradient structure with quadratic dissipation potential, as discovered in [24,26]. The authors show the convergence of that gradient structure by the notion of E-convergence as defined in [28]. In order to do so they assume linearly independent stoichiometric coefficients, which can be seen as a decoupling or orthogonality between the slow and the fast reactions. In this paper we do not need such an assumption because the flux setting automatically decouples the reactions.

Mielke and Stephan [33] study the linear setting, similarly to the current paper. Contrary to Disser et al., they use the gradient structure that is related to the large-deviation principle (1.3b) in the sense of [31], again under the detailed balance assumption. They prove the convergence of that gradient structure, using the stronger notion of tilted EDP-convergence; see [23, 26,29]. This result implies convergence of the large-deviation rate functions \mathcal{I}^ϵ , under the more restrictive assumptions mentioned above, but also for a wide range of tilted energies simultaneously. In the recent follow-up paper [32], this setting is generalised to the case of nonlinear systems, modelled on the class of chemical reactions with mass-action kinetics, still under the detailed balance condition.

1.10 Overview

Section 2 contains preliminaries that are needed throughout the paper. In Sect. 3, we study properties of the network, the continuity equations, and their limits, and we derive equicoercivity in Θ . In Sect. 4 we prove our main Γ -convergence result, Theorem 1.2. In Sect. 5 we prove the relation between spikes and damped cycles, Theorem 1.3. Finally, in Sect. 6 we derive implications for Γ -convergence of the density large deviations, and for convergence of solutions to the effective dynamics.

2 Preliminaries

We first provide a list of basic facts that will be used throughout the paper. After this we introduce the Orlicz space $L^\mathcal{C}$. Next we recall a FIR inequality that bounds the free energy and Fisher information by the rate functional which will be needed to derive compactness of densities later on. Finally, we state a number of convex dual formulations of a number of relevant functionals.

2.1 Basic Properties

We will use the following properties of the functions $s(\cdot|\cdot)$ and \mathcal{C} . For any $a, b \geq 0$ and $p \in \mathbb{R}$, we have:

$$s(a | b) := a \log \frac{a}{b} - a + b \geq s(\alpha b | b) + \partial_a s(\alpha b | b) = (1 - \alpha)b + a \log \alpha \quad \forall \alpha > 0, \tag{2.1}$$

$$s(a | b) \leq \frac{a^2}{b} - 2a + b \quad (\text{using } \log x \leq x - 1), \tag{2.2}$$

$$\begin{aligned} \mathcal{C}(a) &:= \inf_{p-q=a} s(p | 1) + s(q | 1) \\ &= s\left(\frac{1}{2}a + \sqrt{1 + a^2/4} \mid 1\right) + s\left(-\frac{1}{2}a + \sqrt{1 + a^2/4} \mid 1\right) \\ &= \int_0^a \sinh^{-1}(\hat{a}/2) \, d\hat{a} = 2(\cosh^*(a/2) + 1), \end{aligned}$$

$$\mathcal{C}^*(p) := \sup_{a \in \mathbb{R}} pa - \mathcal{C}(a) \tag{2.3}$$

$$= 2(\cosh(p) - 1), \tag{2.4}$$

$$s(a | b) = b(s(a/b | 1) + s(1 | 1)) \geq b \mathcal{C}(\frac{a-b}{b}), \tag{2.5}$$

$$\mathcal{C}(\delta p) = \delta \int_0^P \sinh^{-1}(\delta q/2) dq \stackrel{\text{(concave)}}{\geq} \delta^2 \int_0^P \sinh^{-1}(q/2) dq = \delta^2 \mathcal{C}(p) \quad \forall \delta \in [0, 1], \tag{2.6}$$

$$\mathcal{C}(\delta p) \stackrel{\text{(convex)}}{\leq} \delta \mathcal{C}(p) \quad \forall \delta \in [0, 1]. \tag{2.7}$$

2.2 Orlicz Space

The functions $\mathcal{C}, \mathcal{C}^*$ defined above form a convex dual pair of N-functions (“nice Young functions” [39, Sec. 1.3]). The primal function \mathcal{C} satisfies the Δ_2 property: $\mathcal{C}(2p) \leq 4\mathcal{C}(p)$ (but \mathcal{C}^* does not). We shall use the corresponding Orlicz space (see [39, Th. 3.3.13]):

$$L^{\mathcal{C}}([0, T]; \mathbb{R}^{\mathcal{R}}) := \left\{ j : [0, T] \rightarrow \mathbb{R}^{\mathcal{R}}, \exists a > 0 \text{ such that } \sum_{r \in \mathcal{R}} \int_{[0, T]} \mathcal{C}(\frac{1}{a} j_r(t)) dt < \infty \right\},$$

$$\|j\|_{L^{\mathcal{C}}} := \sup_{\substack{\zeta \in L^{\mathcal{C}^*}([0, T]; \mathbb{R}^{\mathcal{R}}): \\ \sum_{r \in \mathcal{R}} \int_{[0, T]} \mathcal{C}^*(|\zeta_r(t)|) dt \leq 1}} \int_{[0, T]} |j(t) \cdot \zeta(t)| dt = \inf_{a > 0} \frac{1}{a} \left(1 + \sum_{r \in \mathcal{R}} \int_{[0, T]} \mathcal{C}(a j_r(t)) dt \right). \tag{2.8}$$

The final characterisation above implies that

$$a \|j\|_{L^{\mathcal{C}}} \leq 1 + \sum_r \int_{[0, T]} \mathcal{C}(a j_r(t)) dt \quad \text{for all } a > 0. \tag{2.9}$$

We also introduce the space (see [39, Prop. 3.4.3])

$$M^{\mathcal{C}^*}([0, T]; \mathbb{R}^{\mathcal{R}}) := \left\{ \zeta : [0, T] \rightarrow \mathbb{R}^{\mathcal{R}}, \forall a > 0 \text{ there holds } \sum_{r \in \mathcal{R}} \int_{[0, T]} \mathcal{C}^*(\frac{1}{a} \zeta_r(t)) dt < \infty \right\}$$

$$= \overline{\text{span} \left\{ \text{step functions } \zeta \in L^{\mathcal{C}^*}([0, T]; \mathbb{R}^{\mathcal{R}}) \right\}}^{L^{\mathcal{C}^*}} \subsetneq L^{\mathcal{C}^*}([0, T]; \mathbb{R}^{\mathcal{R}}).$$

Then $(M^{\mathcal{C}^*}([0, T]; \mathbb{R}^{\mathcal{R}}))^* \simeq L^{\mathcal{C}}([0, T]; \mathbb{R}^{\mathcal{R}})$ [39, Thms 4.1.6 & 4.1.7], and, since \mathcal{C} satisfies the Δ_2 -property, also $(L^{\mathcal{C}}([0, T]; \mathbb{R}^{\mathcal{R}}))^* \simeq (L^{\mathcal{C}^*}([0, T]; \mathbb{R}^{\mathcal{R}}))^*$ [39, Cor. 4.1.9]. In particular, the first of these isomorphisms defines the weak-* topology on $L^{\mathcal{C}}([0, T]; \mathbb{R}^{\mathcal{R}})$.

2.3 An FIR Inequality

There are various related notions of Fisher information for discrete systems in the literature [7,15,25]. The notion that we use is:

$$FI^\epsilon(u) := \frac{1}{2} \sum_{r \in \mathcal{R}} \int_{[0, T]} \kappa_r^\epsilon \pi_{r_-}^\epsilon \left(\sqrt{u_{r_-}(t)} - \sqrt{u_{r_+}(t)} \right)^2 dt, \tag{2.10}$$

where $\kappa_r^\epsilon \pi_{r_-}^\epsilon u_{r_+}(t) = \kappa_r^\epsilon \frac{\pi_{r_-}^\epsilon}{\pi_{r_+}^\epsilon} \rho_{r_+}(t)$ appears as the backward jump rate for the time-reversed process. This expression can be recognised as a weighted version of the Hellinger distance between two measures u_{r_-} and u_{r_+} on the set of edges \mathcal{R} .

Recall the definitions of $\tilde{\mathcal{I}}_0$ from (1.11) and $\tilde{\mathcal{J}}^\epsilon$ from (1.10). Using arguments from Macroscopic Fluctuation Theory, one can show the following inequality, that is sometimes known as the FIR inequality in the literature [17,19,40]:

Lemma 2.1 (FIR inequality) *Let $(u_{\mathcal{V}_0}^\epsilon, u_{\mathcal{V}_1}^\epsilon, j_{\mathcal{R}_{slow}}^\epsilon, j_{\mathcal{R}_{damp}}^\epsilon, \tilde{j}_{\mathcal{R}_{ficyc}}^\epsilon) \in \Theta$ be such that $\tilde{\mathcal{I}}_0^\epsilon(u^\epsilon(0)) + \mathcal{J}^\epsilon(u^\epsilon \pi^\epsilon, j^\epsilon) < \infty$. Then*

$$\sup_{0 \leq t \leq T} \frac{1}{2} \tilde{\mathcal{I}}_0^\epsilon(u^\epsilon(t)) + FI^\epsilon(u^\epsilon) \leq \frac{1}{2} \tilde{\mathcal{I}}_0^\epsilon(u^\epsilon(0)) + \mathcal{J}^\epsilon(u^\epsilon \pi^\epsilon, j^\epsilon). \tag{2.11}$$

The proof is a simple rewriting of the results of [17], [19, Cor. 4] and [40], and we omit it. From the boundedness of $\tilde{\mathcal{I}}_0^\epsilon(u^\epsilon(0)) + \mathcal{J}^\epsilon(u^\epsilon \pi^\epsilon, j^\epsilon)$ assumed above, the inequality (2.11) implies boundedness of both $\tilde{\mathcal{I}}_0^\epsilon(u^\epsilon(T))$ and $FI^\epsilon(u^\epsilon \pi^\epsilon)$; this will be important in deducing compactness for the densities $u_{\mathcal{V}_1}^\epsilon$.

2.4 Dual Formulations

We recall convex dual formulations for the entropic and quadratic functionals and the Fisher information.

Lemma 2.2 ([37, Prop. 3.5], [2, Lemma 9.4.4]) *If $u \in L^1([0, T])$,*

$$\sup_{\zeta \in C([0, T])} \int_{[0, T]} [\zeta(t)j(t) - u(t)(e^{\zeta(t)} - 1)] dt = \begin{cases} \int_{[0, T]} s(j(t) \mid u(t)) dt, & \text{if } j \in L^1([0, T]), j \ll u, \\ \infty, & \text{otherwise,} \end{cases}$$

and if $u \in \mathcal{M}([0, T])$,

$$\sup_{\zeta \in C([0, T])} \int_{[0, T]} [\zeta(t)j(dt) - u(dt)(e^{\zeta(t)} - 1)] = \begin{cases} \int_{[0, T]} s(j \mid u)(dt), & \text{if } j \in \mathcal{M}([0, T]), j \ll u, \\ \infty, & \text{otherwise,} \end{cases}$$

using the notation (1.12).

Lemma 2.3 ([2, Lemma 9.4.4]) *If $u \in L^1([0, T])$, $u \geq 0$, then*

$$\sup_{\zeta \in C([0, T])} \int_{[0, T]} \left[\zeta(t)\tilde{j}(t) - \frac{1}{2}u(t)\zeta(t)^2 \right] dt = \begin{cases} \frac{1}{2} \int_{[0, T]} \left(\frac{\tilde{j}(t)}{u(t)} \right)^2 u(t) dt, & \text{if } \tilde{j} \ll u, \\ \infty, & \text{otherwise.} \end{cases}$$

Proposition 2.4 *For $u \in L^1([0, T]; \mathbb{R}^{\mathcal{V}})$,*

$$FI^\epsilon(u) = \sup_{\substack{p \in C([0, T]; \mathbb{R}^{2\mathcal{R}}): \\ p_{r_-} < 1, p_{r_+} < 1, \\ (p_{r_-} - 1)(p_{r_+} - 1) > 1}} \sum_{r \in \mathcal{R}} \kappa_{r_-}^\epsilon \pi_{r_-}^\epsilon \int_{[0, T]} \kappa_{r_-}^\epsilon \pi_{r_-}^\epsilon [p_{r_-}(t)u_{r_-}(t) + p_{r_+}(t)u_{r_+}(t)] dt.$$

Proof The proof follows from a simple duality argument, using the property

$$(\sqrt{a} - \sqrt{b})^2 = \sup_{p < 1, q < 1, (p-1)(q-1) > 1} ap + bq.$$

□

Remark 2.5 The duality formulation above gives an interpretation of FI^ϵ for measure-valued u . In fact, the supremum is finite for any finite measure u :

$$\begin{aligned} & \sup_{\substack{p \in C([0, T]; \mathbb{R}^{2\mathcal{R}}): \\ p_{r_-} < 1, p_{r_+} < 1, \\ (p_{r_-} - 1)(p_{r_+} - 1) > 1}} \sum_{r \in \mathcal{R}} \kappa_{r_-}^\epsilon \pi_{r_-}^\epsilon \int_{[0, T]} [p_{r_-}(t)u_{r_-}(dt) + p_{r_+}(t)u_{r_+}(dt)] \\ &= \sup_{\substack{q \in C([0, T]; \mathbb{R}^{2\mathcal{R}}): \\ q_{r_-} > 0, q_{r_+} > 0, \\ q_{r_-} q_{r_+} > 1}} \sum_{r \in \mathcal{R}} \kappa_{r_-}^\epsilon \pi_{r_-}^\epsilon \int_{[0, T]} [(1 - q_{r_-}(t))u_{r_-}(dt) + (1 - q_{r_+}(t))u_{r_+}(dt)] \\ &\leq \sum_{r \in \mathcal{R}} \kappa_{r_-}^\epsilon \pi_{r_-}^\epsilon (\|u_{r_-}\|_{TV} + \|u_{r_+}\|_{TV}). \end{aligned}$$

This shows that a uniformly bounded Fisher information does not rule out the development of singularities in the densities, as explained in Sect. 1.8.

3 Network Properties and Compactness

In this section we study the network decomposition introduced in Sects. 1.5, 1.4, 1.3, and in particular the implications for the continuity equation. We derive estimates for sublevel sets of the rate functional and deduce compactness of these sublevel sets in the topological space Θ as defined in Sect. 1.7. We then use that topology to derive the limiting continuity equations. In addition, we show that any sequence of bounded cost will equilibrate over the fast cycle components, and then prove a stronger equilibration result that will be needed in the construction of the recovery sequence in Sect. 4.

3.1 Network Properties and the Continuity Equations

As mentioned in Sect. 1.3, the absence of leak fluxes, through which the non-equilibrium steady state flux is of order ϵ , implies that $\mathcal{V} = \mathcal{V}_0 \cup \mathcal{V}_1$. We first prove this claim.

Lemma 3.1 *Let π^ϵ be the unique invariant measure of (1.1). Let us repeat the assumptions from Sect. 1:*

1. π^ϵ is normalised, i.e. $\sum_{x \in \mathcal{V}} \pi_x^\epsilon = 1$;
2. The graph $(\mathcal{V}, \mathcal{R})$ is disconnected;
3. There are no ‘leak’ fluxes: if $r \in \mathcal{R}_{slow}$ then $r_- \in \mathcal{V}_0$.

Then $\mathcal{V} = \mathcal{V}_0 \cup \mathcal{V}_1$, i.e. for each $x \in \mathcal{V}$, either π_x^ϵ or $\epsilon^{-1}\pi_x^\epsilon$ converges to a positive limit.

Proof The invariant measure π^ϵ satisfies the equation

$$\left(\sum_{r \in \mathcal{R}: r_- = x} \kappa_r^\epsilon \right) \pi_x^\epsilon = \sum_{r \in \mathcal{R}: r_+ = x} \kappa_r^\epsilon \pi_{r_-}^\epsilon \quad \text{for each } x \in \mathcal{V}. \tag{3.1}$$

To force a contradiction, assume that $\tilde{\mathcal{V}} := \mathcal{V} \setminus (\mathcal{V}_0 \cup \mathcal{V}_1)$ is non-empty. By summing (3.1) over $x \in \mathcal{V}_0 \cup \mathcal{V}_1$ and cancelling terms that appear on both sides we deduce a ‘macro’ version of (3.1),

$$\sum_{\substack{r \in \mathcal{R}: r_+ \in \tilde{\mathcal{V}}, \\ r_- \in \mathcal{V}_0 \cup \mathcal{V}_1}} \kappa_r^\epsilon \pi_{r_-}^\epsilon = \sum_{\substack{r \in \mathcal{R}: r_- \in \tilde{\mathcal{V}}, \\ r_+ \in \mathcal{V}_0 \cup \mathcal{V}_1}} \kappa_r^\epsilon \pi_{r_-}^\epsilon. \tag{3.2}$$

Pick edges with leading order fluxes:

$$r^{\text{in}} \in \operatorname{argmax} \{ \kappa_r^\epsilon \pi_{r_-}^\epsilon : r \in \mathcal{R}, r_- \in \mathcal{V}_0 \cup \mathcal{V}_1, r_+ \in \tilde{\mathcal{V}} \},$$

$$r^{\text{out}} \in \operatorname{argmax} \{ \kappa_r^\epsilon \pi_{r_-}^\epsilon : r \in \mathcal{R}, r_- \in \tilde{\mathcal{V}}, r_+ \in \mathcal{V}_0 \cup \mathcal{V}_1 \}.$$

It follows from (3.2) that

$$\frac{1}{|\mathcal{R}|} \frac{\kappa_{r^{\text{in}}}^\epsilon \pi_{r^{\text{in}}}^\epsilon}{\kappa_{r^{\text{out}}}^\epsilon} \leq \pi_{r^{\text{out}}}^\epsilon \leq |\mathcal{R}| \frac{\kappa_{r^{\text{in}}}^\epsilon \pi_{r^{\text{in}}}^\epsilon}{\kappa_{r^{\text{out}}}^\epsilon}.$$

As leak fluxes do not occur, $\kappa_{r^{\text{in}}}^\epsilon \pi_{r^{\text{in}}}^\epsilon$ is of either order 1 or $1/\epsilon$. Therefore $\pi_{r^{\text{out}}}^\epsilon$ must be either of order 1, ϵ or $1/\epsilon$, the latter being ruled out by the normalisation assumption. We conclude that $r^{\text{out}} \in \mathcal{V}_0 \cup \mathcal{V}_1$ which contradicts $r^{\text{out}} \in \tilde{\mathcal{V}}$. □

We further decomposed \mathcal{V}_0 into $\mathcal{V}_{0\text{fcyc}}$ and $\mathcal{V}_{0\text{slow}}$, where $\mathcal{V}_{0\text{fcyc}}$ is defined as all nodes $x \in \mathcal{V}_0$ such there is at least one fast reaction that leaves x . The name $\mathcal{V}_{0\text{fcyc}}$ (‘fast cycle’) reflects the fact that all nodes in this set belong to a cycle of fast fluxes, as the following simple lemma shows:

Lemma 3.2 *The subgraph $(\mathcal{V}_{0\text{fcyc}}, \mathcal{R}_{\text{fcyc}})$ consists purely of cycles. More explicitly, let $x^1 \in \mathcal{V}_{0\text{fcyc}}$. Then there exists a cycle $(r^k)_{k=1}^K \subset \mathcal{R}_{\text{fcyc}}, r_+^k = r_-^{k+1}$ for $k = 1, \dots, K - 1, r_-^1 = x^1 = r_+^K$. Similarly any $r \in \mathcal{R}_{\text{fcyc}}$ is part of such a fast cycle.*

Proof Let $r^1 \in \mathcal{R}_{\text{fcyc}}$ with $r_-^1 = x^1$, which exists by assumption $x^0 \in \mathcal{V}_{0\text{fcyc}}$, and let $x^2 := r_+^1$. The equilibrium equation in x^2 reads:

$$\pi_{x^2}^\epsilon \sum_{r \in \mathcal{R}: r_- = x^2} \kappa_r^\epsilon = \sum_{r \in \mathcal{R}: r_+ = x^2} \kappa_r^\epsilon \pi_{r_-}^\epsilon \geq \kappa_{r^1}^\epsilon \pi_{x^1}^\epsilon.$$

The right-hand side is of order $1/\epsilon$, and so for the left-hand side $\pi_{x^2}^\epsilon$ must be order 1 (or higher, which is ruled out by assumption), and the sum contains at least one $r^2 := r \in \mathcal{R}_{\text{fast}}$. It follows that $x^2 \in \mathcal{V}_{0\text{fcyc}}$ and $r^2 \in \mathcal{R}_{\text{fcyc}}$. We then repeat the same argument, which only terminates when $x^{K+1} = x^1$. The second claim is true by the same argument. □

We can then enumerate all possible edges from and to $\mathcal{V}_{0\text{slow}}, \mathcal{V}_{0\text{fcyc}}$, and \mathcal{V}_1 .

- Lemma 3.3** (i) *If $x \in \mathcal{V}_{0\text{slow}}$, then all incoming edges $r \in \mathcal{R}, r_+ = x$ are either in $\mathcal{R}_{\text{slow}}$ or in $\mathcal{R}_{\text{damp}}$, and all outgoing edges $r \in \mathcal{R}, r_+ = x$ are in $\mathcal{R}_{\text{slow}}$.*
 (ii) *If $x \in \mathcal{V}_{0\text{fcyc}}$, then the incoming edges could be of any type, and all outgoing edges $r \in \mathcal{R}, r_- = x$ are either in $\mathcal{R}_{\text{slow}}$ or in $\mathcal{R}_{\text{fcyc}}$.*
 (iii) *If $x \in \mathcal{V}_1$, then all incoming fluxes $r \in \mathcal{R}, r_+ = x$ are either in $\mathcal{R}_{\text{slow}}$ or $\mathcal{R}_{\text{damp}}$, and all outgoing fluxes $r \in \mathcal{R}, r_- = x$ are in $\mathcal{R}_{\text{damp}}$.*

Proof For $x \in \mathcal{V}_{0\text{slow}}$ or $\mathcal{V}_{0\text{fcyc}}$, the statement follows immediately from the definitions of $\mathcal{R}_{\text{slow}}, \mathcal{R}_{\text{damp}}$ and $\mathcal{R}_{\text{fcyc}}$. For $x \in \mathcal{V}_1$ any slow outgoing edge will be of leak type that we ruled out by assumption and any fast outgoing edge is damped. Since all outgoing edges are of order 1, an incoming fast cycle edge of order $1/\epsilon$ would imply that π_x^ϵ is of order $1/\epsilon$, which is ruled out by the conservation of mass. □

We can now write down the rescaled continuity equations. Although for any $\epsilon > 0$ all densities u and fluxes j and \tilde{j} have $W^{1,1}$ and L^1 regularity respectively, provided the rate

functional (1.6) is finite, some of this regularity is lost in the regime $\epsilon \rightarrow 0$. Therefore it will be useful to write the continuity equations in a different form. In the following we will say that

$$\pi_x^\epsilon \dot{u}_x^\epsilon = \sum_r j_r^\epsilon \quad \text{in the weak sense,}$$

whenever

$$-\int_{[0,T]} \dot{\phi}(t) \pi_x^\epsilon u_x^\epsilon(dt) = \sum_r \int_{[0,T]} \phi(t) j_r^\epsilon(dt) \quad \text{for all } \phi \in C_0^1([0, T]), \quad (3.3)$$

where we identify $u_x(dt) = u_x(t) dt$ and $j_r(dt) = j_r(t) dt$ wherever possible. If for a fixed $\epsilon > 0$ we have $\tilde{\mathcal{I}}_0^\epsilon + \tilde{\mathcal{J}}^\epsilon < \infty$, then by (1.6) we know that all densities are absolutely continuous and all fluxes have L^1 -densities. We will then say that

$$\pi_x^\epsilon \dot{u}_x^\epsilon = \sum_r j_r^\epsilon \quad \text{in the mild sense,}$$

whenever for all $0 \leq t_0 \leq t_1 \leq T$,

$$\pi_x^\epsilon u_x^\epsilon(t_1) - \pi_x^\epsilon u_x^\epsilon(t_0) = \sum_r \int_{t_0}^{t_1} j_r^\epsilon(t) dt. \quad (3.4)$$

Corollary 3.4 *After rescaling, the continuity equations (1.4) and (1.9) are, for fixed $\epsilon > 0$, in the weak sense,*

$$\pi_c^\epsilon u_c^\epsilon = \sum_{x \in \mathfrak{C}} \pi_x^\epsilon u_x^\epsilon \quad \text{for } c \in \mathfrak{C}, \quad (3.5a)$$

$$\pi_x^\epsilon \dot{u}_x^\epsilon = \sum_{\substack{r \in \mathcal{R}_{slow}: \\ r_+ = x}} j_r^\epsilon + \sum_{\substack{r \in \mathcal{R}_{damp}: \\ r_+ = x}} j_r^\epsilon - \sum_{\substack{r \in \mathcal{R}_{slow}: \\ r_- = x}} j_r^\epsilon \quad \text{for } x \in \mathcal{V}_{0slow}, \quad (3.5b)$$

$$\begin{aligned} \pi_x^\epsilon \dot{u}_x^\epsilon &= \sum_{\substack{r \in \mathcal{R}_{slow}: \\ r_+ = x}} j_r^\epsilon + \sum_{\substack{r \in \mathcal{R}_{damp}: \\ r_+ = x}} j_r^\epsilon - \sum_{\substack{r \in \mathcal{R}_{slow}: \\ r_- = x}} j_r^\epsilon \\ &+ \sum_{\substack{r \in \mathcal{R}_{fctc}: \\ r_+ = x}} \left(\frac{1}{\epsilon} k_r \pi_{r_-}^\epsilon u_{r_-}^\epsilon + \frac{1}{\sqrt{\epsilon}} j_r^\epsilon \right) - \sum_{\substack{r \in \mathcal{R}_{fctc}: \\ r_- = x}} \left(\frac{1}{\epsilon} k_r \pi_{r_-}^\epsilon u_{r_-}^\epsilon + \frac{1}{\sqrt{\epsilon}} j_r^\epsilon \right), \quad \text{for } x \in \mathcal{V}_{0fctc}, \end{aligned} \quad (3.5c)$$

$$\pi_x^\epsilon \dot{u}_x^\epsilon = \sum_{\substack{r \in \mathcal{R}_{slow}: \\ r_+ = x}} j_r^\epsilon + \sum_{\substack{r \in \mathcal{R}_{damp}: \\ r_+ = x}} j_r^\epsilon - \sum_{\substack{r \in \mathcal{R}_{damp}: \\ r_- = x}} j_r^\epsilon, \quad \text{for } x \in \mathcal{V}_1. \quad (3.5d)$$

If in addition $\tilde{\mathcal{I}}_0^\epsilon(u^\epsilon(0)) + \tilde{\mathcal{J}}^\epsilon(u^\epsilon, j^\epsilon) < \infty$, then these equations also hold in the mild sense of (3.4).

3.2 Boundedness of Densities and Fluxes

The aim of this section is to prove uniform bounds that are needed to derive the equicoercivity Theorem 1.1 later on.

Lemma 3.5 (Boundedness of densities) *Let $(u^\epsilon, j^\epsilon)_{\epsilon > 0} \subset \Theta$ such that $\tilde{\mathcal{I}}_0^\epsilon(u^\epsilon(0)) + \tilde{\mathcal{J}}^\epsilon(u^\epsilon, j^\epsilon) \leq C$ for some $C > 0$. Then*

1. $(u_{\mathcal{V}_{0slow}}^\epsilon, u_{\mathfrak{C}}^\epsilon)$ and $u_{\mathcal{V}_{0fctc}}^\epsilon$ are uniformly bounded in $C([0, T]; \mathbb{R}^{\mathcal{V}_{0slow} \cup \mathfrak{C}})$ and $L^\infty([0, T]; \mathbb{R}^{\mathcal{V}_{0fctc}})$;
2. $(u_{\mathcal{V}_{0slow}}^\epsilon, u_{\mathcal{V}_{0fctc}}^\epsilon, u_{\mathcal{V}_1}^\epsilon)$ is uniformly bounded in $L^1([0, T]; \mathbb{R}^{\mathcal{V}_{0slow} \cup \mathcal{V}_{0fctc} \cup \mathcal{V}_1})$;

3. $\|u_x^\epsilon\|_{C([0,T])} \rightarrow 0$ for all $x \in \mathcal{V}_1$ as $\epsilon \rightarrow 0$.

Proof From (2.1) and mass conservation we derive a uniform bound on the total mass for each $t \in [0, T]$:

$$\sum_{x \in \mathcal{V}} \pi_x^\epsilon u_x^\epsilon(t) = \sum_{x \in \mathcal{V}} \pi_x^\epsilon u_x^\epsilon(0) \leq \tilde{\mathcal{I}}_0^\epsilon(u^\epsilon(0)) + (e - 1) \sum_x \pi_x^\epsilon \leq C + e - 1. \tag{3.6}$$

This implies the C -bounds on $u_{\mathcal{V}_{0\text{slow}}}^\epsilon$, $u_{\mathcal{C}}^\epsilon$, and the L^∞ bound on $u_{\mathcal{V}_{0\text{ficyc}}}^\epsilon$.

From the FIR inequality (2.11) we deduce that

$$\begin{aligned} C &\geq \frac{1}{2} \tilde{\mathcal{I}}_0^\epsilon(u^\epsilon(0)) + \tilde{\mathcal{J}}^\epsilon(u_{\mathcal{V}_0}^\epsilon, u_{\mathcal{V}_1}^\epsilon, j_{\mathcal{R}_{\text{slow}}}^\epsilon, j_{\mathcal{R}_{\text{damp}}}^\epsilon, \tilde{j}_{\mathcal{R}_{\text{ficyc}}}^\epsilon) \\ &\geq \text{FI}^\epsilon(u^\epsilon) = \frac{1}{2} \sum_{r \in \mathcal{R}} \kappa_r^\epsilon \pi_{r_-}^\epsilon \int_{[0,T]} \left(\sqrt{u_{r_-}^\epsilon(t)} - \sqrt{u_{r_+}^\epsilon(t)} \right)^2 dt. \end{aligned} \tag{3.7}$$

Hence by (1.7), for ϵ sufficiently small and any $r \in \mathcal{R}$:

$$\int_{[0,T]} \left(\sqrt{u_{r_-}^\epsilon(t)} - \sqrt{u_{r_+}^\epsilon(t)} \right)^2 dt \leq \begin{cases} \frac{4C}{\kappa_r \pi_{r_-}}, & r \in \mathcal{R}_{\text{slow}}, \\ \frac{4C}{\kappa_r \pi_{r_-}}, & r \in \mathcal{R}_{\text{damp}}, \\ \frac{4C}{\kappa_r \pi_{r_-}} \epsilon, & r \in \mathcal{R}_{\text{ficyc}}. \end{cases}$$

Since \mathcal{V} is finite, \mathcal{V}_0 cannot be empty, since otherwise the total mass in the system would vanish. Take an arbitrary $x^0 \in \mathcal{V}_0$; by (3.6) we have $\|u_{x^0}^\epsilon\|_{L^\infty(0,T)} \leq 2(C + e - 1)/\pi_{x^0}$ for sufficiently small ϵ . Now take an arbitrary $y \in \mathcal{V}$. By irreducibility of the graph $(\mathcal{V}, \mathcal{R})$ there exists a sequence of edges $x^0 \xrightarrow{r^{01}} x^1 \xrightarrow{r^{12}} \dots \rightarrow x^n = y$. For the first edge we find, using the inequality $a \leq 2(\sqrt{a} - \sqrt{b})^2 + 2b$,

$$\begin{aligned} \int_{[0,T]} u_{x^1}^\epsilon(t) dt &\leq 2 \int_{[0,T]} \left(\sqrt{u_{x^0}^\epsilon(t)} - \sqrt{u_{x^1}^\epsilon(t)} \right)^2 dt \\ &+ 2 \int_{[0,T]} u_{x^0}^\epsilon(t) dt \leq \frac{8C}{\kappa_{01} \pi_{x^0}} + 4T \frac{C + e - 1}{\pi_{x^0}}. \end{aligned}$$

Repeating this procedure for all edges yields that u_y^ϵ is uniformly bounded in $L^1(0, T)$.

Finally we prove the vanishing of $\epsilon u_{\mathcal{V}_1}^\epsilon$. We also deduce from (2.11) that for all $0 \leq t \leq T$,

$$\begin{aligned} C &\geq \frac{1}{2} \tilde{\mathcal{I}}_0^\epsilon(u^\epsilon(t)) \geq \frac{1}{2} \sum_{x \in \mathcal{V}_1} s(\epsilon u_x^\epsilon(t) \tilde{\pi}_x^\epsilon | \epsilon \tilde{\pi}_x^\epsilon) = \frac{1}{2} \sum_{x \in \mathcal{V}_1} \epsilon \tilde{\pi}_x^\epsilon s(u_x^\epsilon(t) | 1) \geq \sum_{x \in \mathcal{V}_1} \epsilon \tilde{\pi}_x^\epsilon \eta(u_x^\epsilon(t)), \\ &\text{with } \eta(\tau) := \begin{cases} \frac{1}{2} [\tau \log \tau - \tau + 1] & \text{if } \tau \geq 1, \\ 0 & \text{if } 0 \leq \tau \leq 1. \end{cases} \end{aligned}$$

Since the $\tilde{\pi}_x^\epsilon$ are bounded away from zero, we find that

$$\eta(u_x^\epsilon(t)) \leq \frac{C}{\epsilon} \implies 0 \leq u_x^\epsilon(t) \leq \eta^{-1}\left(\frac{C}{\epsilon}\right),$$

where η^{-1} is the right-continuous generalised inverse of η . Since η is superlinear at infinity, $\epsilon \eta^{-1}(C/\epsilon) \rightarrow 0$ as $\epsilon \rightarrow 0$, and we find that $\|u_{\mathcal{V}_1}^\epsilon\|_{C([0,T])} \rightarrow 0$ as $\epsilon \rightarrow 0$. \square

Lemma 3.6 (Boundedness of slow fluxes) *Let $(u^\epsilon, j^\epsilon)_{\epsilon>0} \subset \Theta$ such that $\tilde{\mathcal{I}}_0^\epsilon(u^\epsilon(0)) + \tilde{\mathcal{J}}^\epsilon(u^\epsilon, j^\epsilon) \leq C$ for some $C > 0$. Then the slow fluxes $j_{\mathcal{R}_{\text{slow}}}^\epsilon$ are uniformly bounded in*

$L^\mathcal{C}([0, T]; \mathbb{R}^{\mathcal{R}_{\text{slow}}})$. It follows that there is a non-decreasing function $\omega : [0, \infty) \rightarrow [0, \infty)$ with $\lim_{\sigma \downarrow 0} \omega(\sigma) = 0$ such that for all $0 \leq t_0 \leq t_1 \leq T$,

$$\sup_{\epsilon > 0} \sum_{r \in \mathcal{R}_{\text{slow}}} \int_{t_0}^{t_1} j_r^\epsilon(t) dt \leq \omega(t_1 - t_0). \tag{3.8}$$

Proof Again by (1.6) we know that $j_{\mathcal{R}_{\text{slow}}}^\epsilon$ and $\rho_{\mathcal{V}_0}^\epsilon$ both have L^1 -densities. Writing $Z := (C + e - 1) \sum_{r \in \mathcal{R}_{\text{slow}}} \kappa_r$,

$$\begin{aligned} C &\stackrel{(1.10)}{\geq} \sum_{r \in \mathcal{R}_{\text{slow}}} \int_{[0, T]} \underbrace{\left[s(j_r^\epsilon(t) \mid \pi_{r_-}^\epsilon u_{r_-}^\epsilon \kappa_r) \right]}_{\text{non-increasing in } \pi_{r_-}^\epsilon u_{r_-}^\epsilon(t) \kappa_r} - \underbrace{\pi_{r_-}^\epsilon u_{r_-}^\epsilon(t) \kappa_r}_{\geq 0} + \pi_{r_-}^\epsilon u_{r_-}^\epsilon(t) \kappa_r dt \\ &\stackrel{(3.6)}{\geq} \sum_{r \in \mathcal{R}_{\text{slow}}} \int_{[0, T]} [s(j_r^\epsilon(t) \mid Z) - Z] dt \\ &\stackrel{(2.5)}{\geq} Z \sum_{r \in \mathcal{R}_{\text{slow}}} \int_{[0, T]} \mathcal{C} \left(\frac{j_r^\epsilon(t) - Z}{Z} \right) dt - Z |\mathcal{R}_{\text{slow}}| T \\ &\stackrel{(2.9)}{\geq} \|j^\epsilon - Z\|_{L^\mathcal{C}([0, T]; \mathbb{R}^{\mathcal{R}_{\text{slow}}})} - Z(|\mathcal{R}_{\text{slow}}| T + 1). \end{aligned}$$

The proof of estimate (3.8) follows from the definition (2.8) of the Orlicz norm and the superlinearity of \mathcal{C} . Define the function

$$\omega : [0, \infty) \rightarrow [0, \infty), \quad \omega(\sigma) := \inf_{\beta > 0} \left\{ \frac{\tilde{C}}{\beta} : |\mathcal{R}_{\text{slow}}| \mathcal{C}^*(\beta) \leq \frac{1}{\sigma} \right\},$$

where \tilde{C} is the bound on $j_{\mathcal{R}_{\text{slow}}}^\epsilon$ in $L^\mathcal{C}([0, T]; \mathbb{R}_+^{\mathcal{R}_{\text{slow}}})$. The function ω is non-decreasing by construction, and $\lim_{\sigma \downarrow 0} \omega(\sigma) = 0$ because \mathcal{C}^* is finite on all of \mathbb{R} .

Fix $0 \leq t_0 \leq t_1 \leq T$ and take $\beta > 0$ such that $(t_1 - t_0) |\mathcal{R}_{\text{slow}}| \mathcal{C}^*(\beta) \leq 1$. Set

$$\zeta_r(t) := \beta \mathbb{1}_{[t_0, t_1]}(t) \mathbb{1}_{\mathcal{R}_{\text{slow}}}(r).$$

Then $\sum_{r \in \mathcal{R}} \int_{[0, T]} \mathcal{C}^*(\zeta_r(t)) dt \leq 1$ and so $\zeta \in L^{\mathcal{C}^*}([0, T]; \mathbb{R}^{\mathcal{R}})$. Plugging this function ζ in (2.8) produces the estimate

$$\sum_{r \in \mathcal{R}_{\text{slow}}} \int_{t_0}^{t_1} j_r^\epsilon(t) dt = \sum_{r \in \mathcal{R}_{\text{slow}}} \frac{1}{\beta} \int_{[0, T]} j_r^\epsilon(t) \zeta(t) dt \leq \frac{1}{\beta} \|j_{\mathcal{R}_{\text{slow}}}^\epsilon\|_{L^\mathcal{C}([0, T]; \mathbb{R}^{\mathcal{R}_{\text{slow}}})} \leq \frac{\tilde{C}}{\beta}.$$

The estimate (3.8) follows from taking the infimum over β . □

Although the form of the rate functional is almost the same for the slow and damped fluxes, the damped fluxes lack a $C([0, T])$ -bound on the corresponding densities. Therefore we obtain a weaker bound on the damped fluxes:

Lemma 3.7 (Boundedness of damped fluxes) *Let $(u^\epsilon, j^\epsilon)_{\epsilon > 0} \subset \Theta$ such that $\tilde{\mathcal{I}}_0^\epsilon(u^\epsilon(0)) + \tilde{\mathcal{J}}^\epsilon(u^\epsilon, j^\epsilon) \leq C$ for some $C > 0$. Then the damped fluxes $j_{\mathcal{R}_{\text{damp}}}^\epsilon$ are uniformly bounded in $L^1([0, T]; \mathbb{R}^{\mathcal{R}_{\text{damp}}})$. In addition, for all $\sigma > 0$,*

$$\limsup_{\epsilon \rightarrow 0} \sup_{\substack{0 \leq t_0 \leq t_1 \leq T: \\ |t_1 - t_0| < \sigma}} \sum_{\substack{r \in \mathcal{R}_{\text{damp}}: \\ r_+ \in \mathcal{V}_0}} \int_{t_0}^{t_1} j_r^\epsilon(t) dt \leq \omega(\sigma), \tag{3.9}$$

where ω is the modulus of continuity of Lemma 3.6.

Proof Again by (1.10) we can assume that $u_{\mathcal{V}_1}^\epsilon$ and $j_{\mathcal{R}_{\text{damp}}}^\epsilon$ have L^1 -densities, at least for $\epsilon > 0$. This allows us to write

$$\begin{aligned} C &\stackrel{(1.10)}{\geq} \sum_{r \in \mathcal{R}_{\text{damp}}} \int_{[0, T]} s(j_r^\epsilon(t) \mid \frac{1}{\epsilon} \kappa_r \pi_{r_-}^\epsilon u_{r_-}^\epsilon(t)) dt \\ &\stackrel{(2.1)}{\geq} \sum_{r \in \mathcal{R}_{\text{damp}}} \int_{[0, T]} ((1 - e) \frac{1}{\epsilon} \kappa_r \pi_{r_-}^\epsilon u_{r_-}^\epsilon + j_r^\epsilon(t)) dt, \end{aligned}$$

and so $\|j_{\mathcal{R}_{\text{damp}}}^\epsilon\|_{L^1([0, T]; \mathcal{R}_{\text{damp}})} \leq C + (e - 1) \|u_{\mathcal{V}_1}^\epsilon\|_{L^1([0, T]; \mathbb{R}^{\mathcal{V}_1})} \sup_{\epsilon > 0, r \in \mathcal{R}_{\text{damp}}} \frac{1}{\epsilon} \kappa_r \pi_{r_-}^\epsilon$, which is uniformly bounded by Lemma 3.5 and the assumption $\frac{1}{\epsilon} \pi_{r_-}^\epsilon \rightarrow \tilde{\pi}_{r_-}$.

Next we prove the estimate (3.9) by summing the mild formulation of the continuity equations (3.5d) over all $x \in \mathcal{V}_1$, for arbitrary $0 \leq t_0 \leq t_1 \leq T$:

$$\sum_{\substack{r \in \mathcal{R}_{\text{damp}}: \\ r_- \in \mathcal{V}_1}} \int_{t_0}^{t_1} j_r^\epsilon(t) dt - \sum_{\substack{r \in \mathcal{R}_{\text{damp}}: \\ r_+ \in \mathcal{V}_1}} \int_{t_0}^{t_1} j_r^\epsilon(t) dt = \sum_{\substack{r \in \mathcal{R}_{\text{slow}}: \\ r_+ \in \mathcal{V}_1}} \int_{t_0}^{t_1} j_r^\epsilon(t) dt - \sum_{x \in \mathcal{V}_1} \pi_x^\epsilon (u_x^\epsilon(t_1) - u_x^\epsilon(t_0)).$$

Since the first two sums have common terms corresponding to $r_-, r_+ \in \mathcal{V}_1$, we can remove them to find

$$\sum_{\substack{r \in \mathcal{R}_{\text{damp}}: \\ r_- \in \mathcal{V}_1 \\ r_+ \in \mathcal{V}_0}} \int_{t_0}^{t_1} j_r^\epsilon(t) dt - \sum_{\substack{r \in \mathcal{R}_{\text{damp}}: \\ r_- \in \mathcal{V}_0 \\ r_+ \in \mathcal{V}_1}} \int_{t_0}^{t_1} j_r^\epsilon(t) dt = \sum_{\substack{r \in \mathcal{R}_{\text{slow}}: \\ r_+ \in \mathcal{V}_1}} \int_{t_0}^{t_1} j_r^\epsilon(t) dt - \sum_{x \in \mathcal{V}_1} \pi_x^\epsilon (u_x^\epsilon(t_1) - u_x^\epsilon(t_0)).$$

The second sum is a sum over the empty set, and applying the estimate (3.8) we find

$$\sum_{\substack{r \in \mathcal{R}_{\text{damp}}: \\ r_- \in \mathcal{V}_1 \\ r_+ \in \mathcal{V}_0}} \int_{t_0}^{t_1} j_r^\epsilon(t) dt \leq \omega(t_1 - t_0) + \sum_{x \in \mathcal{V}_1} \pi_x^\epsilon \|u_x^\epsilon\|_{C([0, T])}.$$

The estimate (3.9) then follows from part 3 of Lemma 3.5 together with $\frac{1}{\epsilon} \pi_{r_-}^\epsilon \rightarrow \tilde{\pi}_{r_-}$. \square

Lemma 3.8 (Boundedness of fast fluxes) *Let $(u^\epsilon, j^\epsilon)_{\epsilon > 0} \subset \Theta$ such that $\tilde{T}_0^\epsilon(u^\epsilon(0)) + \tilde{J}^\epsilon(u^\epsilon, j^\epsilon) \leq C$ for some $C > 0$. Then the fast cycle fluxes $\tilde{j}_{\mathcal{R}_{\text{fcyc}}}^\epsilon$ are uniformly bounded in $L^\infty([0, T]; \mathbb{R}^{\mathcal{R}_{\text{fcyc}}})$.*

Proof Similar to the proof of Lemma 3.6 we write $Z := (C + e - 1) \sum_{r \in \mathcal{R}_{\text{fcyc}}} \kappa_r$, so that due to the total mass estimate (3.6), for each $r \in \mathcal{R}_{\text{fcyc}}$:

$$\kappa_r \pi_{r_-}^\epsilon u_{r_-}^\epsilon(t) / Z \leq 1. \tag{3.10}$$

Again using the existence of L^1 -densities:

$$\begin{aligned}
 C &\stackrel{(1.10)}{\geq} \sum_{r \in \mathcal{R}_{\text{ficyc}}} \int_{[0, T]} s \left(\frac{1}{\epsilon} \kappa_r \pi_{r-}^\epsilon u_{r-}^\epsilon(t) + \frac{1}{\sqrt{\epsilon}} \tilde{j}_r^\epsilon(t) \mid \frac{1}{\epsilon} \kappa_r \pi_{r-}^\epsilon u_{r-}^\epsilon(t) \right) dt \\
 &\stackrel{(2.5)}{\geq} \sum_{r \in \mathcal{R}_{\text{ficyc}}} \int_{[0, T]} \frac{1}{\epsilon} \kappa_r \pi_{r-}^\epsilon u_{r-}^\epsilon(t) \mathcal{C} \left(\frac{\frac{1}{\sqrt{\epsilon}} \tilde{j}_r^\epsilon(t)}{\frac{1}{\epsilon} \kappa_r \pi_{r-}^\epsilon u_{r-}^\epsilon(t)} \right) dt \\
 &\stackrel{(2.6)}{\geq} \sum_{r \in \mathcal{R}_{\text{ficyc}}} \int_{[0, T]} Z \frac{\kappa_r \pi_{r-}^\epsilon u_{r-}^\epsilon(t)}{Z} \mathcal{C} \left(\frac{\tilde{j}_r^\epsilon(t)/Z}{\kappa_r \pi_{r-}^\epsilon u_{r-}^\epsilon(t)/Z} \right) dt \\
 &\stackrel{(2.7), (3.10)}{\geq} \sum_{r \in \mathcal{R}_{\text{ficyc}}} \int_{[0, T]} Z \mathcal{C} \left(\frac{\tilde{j}_r^\epsilon(t)}{Z} \right) dt \\
 &\stackrel{(2.9)}{\geq} \|\tilde{j}^\epsilon\|_{L^\mathcal{C}([0, T]; \mathbb{R}^{\mathcal{R}_{\text{ficyc}}})} - Z.
 \end{aligned}$$

□

Lemma 3.9 (Equicontinuity of $u_{\mathcal{V}_{0\text{slow}}}^\epsilon$ and $u_{\mathcal{C}}^\epsilon$.)

Let $(u_{\mathcal{V}_0}^\epsilon, j^\epsilon)_{\epsilon > 0} \subset \Theta$ such that $\tilde{I}_0^\epsilon(u^\epsilon(0)) + \tilde{J}^\epsilon(u^\epsilon, j^\epsilon) \leq C$ for some $C > 0$. Then there exists a continuous non-decreasing function $\bar{\omega} : [0, \infty) \rightarrow [0, \infty)$ with $\lim_{\sigma \downarrow 0} \bar{\omega}(\sigma) = 0$ such that for all $0 \leq t_0 \leq t_1 \leq T$,

$$\limsup_{\epsilon \rightarrow 0} \left\{ \sum_{\mathfrak{c} \in \mathcal{C}} |u_{\mathfrak{c}}^\epsilon(t_1) - u_{\mathfrak{c}}^\epsilon(t_0)| + \sum_{x \in \mathcal{V}_{0\text{slow}}} |u_x^\epsilon(t_1) - u_x^\epsilon(t_0)| \right\} \leq \bar{\omega}(t_1 - t_0). \tag{3.11}$$

Proof Fix $0 \leq t_0 \leq t_1 \leq T$. Take $x \in \mathcal{V}_{0\text{slow}}$ and note that by (3.5b) and (3.8)

$$\pi_x^\epsilon(u_x^\epsilon(t_1) - u_x^\epsilon(t_0)) \geq - \sum_{\substack{r \in \mathcal{R}_{\text{slow}} \\ r_+ = x}} \int_{t_0}^{t_1} j_r^\epsilon(t) dt \geq -\omega(t_1 - t_0),$$

where we again used the mild formulation of the continuity equations. To estimate the difference from the other side we write

$$\begin{aligned}
 \pi_x^\epsilon(u_x^\epsilon(t_1) - u_x^\epsilon(t_0)) &\leq \sum_{\substack{r \in \mathcal{R}_{\text{slow}} \\ r_+ = x}} \int_{t_0}^{t_1} j_r^\epsilon(t) dt + \sum_{\substack{r \in \mathcal{R}_{\text{damp}} \\ r_+ = x}} \int_{t_0}^{t_1} j_r^\epsilon(t) dt \\
 &\leq \omega(t - s) + \sum_{\substack{r \in \mathcal{R}_{\text{damp}} \\ r_+ = x}} \int_{t_0}^{t_1} j_r^\epsilon(t) dt,
 \end{aligned}$$

and by (3.9) one part of (3.11) follows.

The same line of reasoning leads to a corresponding statement about $|u_{\mathfrak{c}}^\epsilon(t_1) - u_{\mathfrak{c}}^\epsilon(t_0)|$ for any $\mathfrak{c} \in \mathcal{C}$, after one sums the continuity equations (3.5c) over all $x \in \mathfrak{c}$ to find

$$\sum_{x \in \mathfrak{c}} \pi_x^\epsilon(u_x^\epsilon(t_1) - u_x^\epsilon(t_0)) = \sum_{\substack{r \in \mathcal{R}_{\text{slow}} \\ r_+ \in \mathfrak{c}}} \int_{t_0}^{t_1} j_r^\epsilon(t) dt + \sum_{\substack{r \in \mathcal{R}_{\text{damp}} \\ r_+ \in \mathfrak{c}}} \int_{t_0}^{t_1} j_r^\epsilon(t) dt - \sum_{\substack{r \in \mathcal{R}_{\text{slow}} \\ r_- \in \mathfrak{c}}} \int_{t_0}^{t_1} j_r^\epsilon(t) dt.$$

We omit the details.

□

3.3 Compactness of Densities and Fluxes

In this brief section we derive the compactness of level sets, and hence the equicoercivity of Theorem 1.1.

Corollary 3.10 *Let $(u^\epsilon, j^\epsilon)_{\epsilon>0} \subset \Theta$ such that $\tilde{I}_0^\epsilon(u^\epsilon(0)) + \tilde{J}^\epsilon(u^\epsilon, j^\epsilon) \leq C$ for some $C > 0$. Then one can choose a sequence $\epsilon_n \rightarrow 0$ and a limit point $(u, j) \in \Theta$ such that*

$$u_{\mathcal{V}_{0slow}}^{\epsilon_n} \longrightarrow u_{\mathcal{V}_{0slow}} \quad \text{in } C([0, T]; \mathbb{R}^{\mathcal{V}_{0slow}}), \tag{3.12a}$$

$$u_{\mathcal{V}_{0fcyc}}^{\epsilon_n} \xrightarrow{*} u_{\mathcal{V}_{0fcyc}} \quad \text{in } L^\infty([0, T]; \mathbb{R}^{\mathcal{V}_{0fcyc}}), \tag{3.12b}$$

$$u_{\mathcal{C}}^{\epsilon_n} \longrightarrow u_{\mathcal{C}} \quad \text{in } C([0, T]; \mathbb{R}^{\mathcal{C}}), \tag{3.12c}$$

$$u_{\mathcal{V}_1}^{\epsilon_n} \xrightarrow{\text{narrow}} u_{\mathcal{V}_1} \quad \text{in } \mathcal{M}([0, T]; \mathbb{R}^{\mathcal{V}_1}), \tag{3.12d}$$

$$\epsilon u_{\mathcal{V}_1}^{\epsilon} \longrightarrow 0 \quad \text{in } C([0, T]; \mathbb{R}^{\mathcal{V}_1}), \tag{3.12e}$$

$$j_{\mathcal{R}_{slow}}^{\epsilon_n} \xrightarrow{*} j_{\mathcal{R}_{slow}} \quad \text{in } L^{\mathcal{C}}([0, T]; \mathbb{R}^{\mathcal{R}_{slow}}), \tag{3.12f}$$

$$j_{\mathcal{R}_{damp}}^{\epsilon_n} \xrightarrow{\text{narrow}} j_{\mathcal{R}_{damp}} \quad \text{in } \mathcal{M}([0, T]; \mathbb{R}^{\mathcal{R}_{damp}}), \tag{3.12g}$$

$$\tilde{j}_{\mathcal{R}_{fcyc}}^{\epsilon_n} \xrightarrow{*} \tilde{j}_{\mathcal{R}_{fcyc}} \quad \text{in } L^{\mathcal{C}}([0, T]; \mathbb{R}^{\mathcal{R}_{fcyc}}). \tag{3.12h}$$

It follows that $u_{\mathcal{V}_{0slow}}$ and $u_{\mathcal{C}}$ are continuous.

Proof The boundedness given by Lemmas 3.5, 3.6, and 3.7 immediately implies the weak-* and narrow compactness of (3.12b), (3.12d), (3.12e), (3.12f), (3.12g), and (3.12h); we extract a subsequence that converges in this sense.

The additional uniform convergences of (3.12a) and (3.12c) follow from an alternative version of the classical Arzelà-Ascoli theorem, which we state and prove in the appendix. This version applies to sequences that are uniformly bounded and *asymptotically uniformly equicontinuous*. The uniform boundedness of $u_{\mathcal{V}_{0slow}}^{\epsilon_n}$ and $u_{\mathcal{C}}^{\epsilon_n}$ follow by Lemma 3.5, and the asymptotic uniform equicontinuity is the statement of Lemma 3.9. The uniform convergences (3.12a) and (3.12c) then follow by Theorem A.1 (up to extraction of a subsequence). \square

Remark 3.11 A similar coercivity result is proven in [33, Prop. 5.9] for a system in detailed balance and with $\mathcal{R}_{damp} = \emptyset = \mathcal{V}_1$. More precisely, they obtain uniform compactness of $(u_{\mathcal{V}_{0slow}}^\epsilon, u_{\mathcal{C}}^\epsilon)$ as we do, and strong L^2 -compactness of $u_{\mathcal{V}_{0fcyc}}^\epsilon$. Our strengthened equilibration Lemma 3.17 will imply strong L^p -compactness of $u_{\mathcal{V}_{0fcyc}}^\epsilon$ for any $1 \leq p < \infty$.

From now on we shall consider sequences that converge in the sense of (3.12).

3.4 Equilibration on Fast Cycle Components

In this section we prove that all mass on fast cycles will instantaneously spread over each node in the fast cycle component.

Lemma 3.12 *Let $(u^\epsilon, j^\epsilon)_{\epsilon>0} \subset \Theta$ such that $\tilde{I}_0^\epsilon(u^\epsilon(0)) + \tilde{J}^\epsilon(u^\epsilon, j^\epsilon) \leq C$ converge to (u, j) in Θ in the sense of (3.12). Then $u_x(t) \equiv u_c(t)$ for almost every $t \in [0, T]$ and each component $c \in \mathcal{C}$, and $\text{div } \tilde{j}_{\mathcal{R}_{fcyc}}(t) \equiv 0$ for almost every $t \in [0, T]$.*

Proof For any $x \in c \in \mathfrak{C}$ and $t \in [0, T]$, the mild formulation of the continuity equation is:

$$\begin{aligned} \pi_x^\epsilon u_x^\epsilon(t) - \pi_x^\epsilon u_x^\epsilon(0) &= \sum_{\substack{r \in \mathcal{R}_{\text{slow}}: \\ r_+ = x}} \int_0^t j_r^\epsilon(s) ds + \sum_{\substack{r \in \mathcal{R}_{\text{damp}}: \\ r_+ = x}} \int_0^t j_r^\epsilon(s) ds - \sum_{\substack{r \in \mathcal{R}_{\text{slow}}: \\ r_- = x}} \int_0^t j_r^\epsilon(s) ds \\ &+ \sum_{\substack{r \in \mathcal{R}_{\text{ficyc}}: \\ r_+ = x}} \frac{1}{\epsilon} \kappa_r \pi_{r_-}^\epsilon \int_0^t u_{r_-}^\epsilon(s) ds + \frac{1}{\sqrt{\epsilon}} \int_0^t \tilde{j}_r^\epsilon(s) ds \\ &- \sum_{\substack{r \in \mathcal{R}_{\text{ficyc}}: \\ r_- = x}} \frac{1}{\epsilon} \kappa_r \pi_{r_-}^\epsilon \int_0^t u_{r_-}^\epsilon(s) ds + \frac{1}{\sqrt{\epsilon}} \int_0^t \tilde{j}_r^\epsilon(s) ds. \end{aligned} \tag{3.13}$$

All terms in the first line are uniformly bounded in $L^1(0, T)$, and the same holds for the u^ϵ and \tilde{j} in the second and third lines. First multiplying the equation by ϵ , and then letting $\epsilon \rightarrow 0$ thus yields:

$$\sum_{\substack{r \in \mathcal{R}_{\text{ficyc}}: \\ r_+ = x}} \kappa_r \pi_{r_-} u_{r_-}(t) = \sum_{\substack{r \in \mathcal{R}_{\text{ficyc}}: \\ r_- = x}} \kappa_r \pi_{r_-} u_{r_-}(t).$$

Without the $u_{r_-}(t)$ factors, this is exactly the equation for the steady state π for a network consisting only of the fast edges. Since the component c containing x is disconnected, this equation has a unique solution up to a multiplicative constant, i.e. $u_x(t) \equiv a_c(t)$ on c , for some $a_c \in L^\infty([0, T])$. To identify a_c , use (3.5a) together with the convergences (3.12b) and (3.12c) to find for the limit

$$\pi_c u_c \leftarrow \pi_c^\epsilon u_c^\epsilon = \sum_{x \in c} \pi_x^\epsilon u_x^\epsilon \rightarrow \sum_{x \in c} \pi_x u_x = \pi_c a_c,$$

so that indeed $u_x \equiv a_c = u_c$.

The same argument, multiplying (3.13) by $\sqrt{\epsilon}$ and letting $\epsilon \rightarrow 0$, shows that

$$\sum_{\substack{r \in \mathcal{R}_{\text{ficyc}}: \\ r_+ = x}} \tilde{j}_r(t) = \sum_{\substack{r \in \mathcal{R}_{\text{ficyc}}: \\ r_- = x}} \tilde{j}_r(t).$$

□

Remark 3.13 Alternatively, the fact that u is constant on c can also be seen from the FIR inequality (3.7) together with the lower semicontinuity that follows from Proposition 2.4. The FIR inequality can however not be used to make a similar statement about divergence-free fast fluxes.

3.5 The Limiting Continuity Equations

We again place ourselves in the setting of Sect. 3.3 and derive the continuity equations satisfied in the limit.

Lemma 3.14 *Let $(u^\epsilon, j^\epsilon)_{\epsilon>0} \subset \Theta$ be such that $\tilde{T}_0^\epsilon(u^\epsilon(0)) + \tilde{J}^\epsilon(u^\epsilon, j^\epsilon) \leq C$, and assume that (u^ϵ, j^ϵ) converges to (u, j) in the sense of (3.12). Then the limit satisfies the continuity*

equations

$$\pi_x \dot{u}_x = \sum_{\substack{r \in \mathcal{R}_{slow}: \\ r_+ = x}} j_r + \sum_{\substack{r \in \mathcal{R}_{damp}: \\ r_+ = x}} j_r - \sum_{\substack{r \in \mathcal{R}_{slow}: \\ r_- = x}} j_r, \quad \text{for } x \in \mathcal{V}_{0slow}, \quad (3.14a)$$

$$\pi_c \dot{u}_c = \sum_{\substack{r \in \mathcal{R}_{slow}: \\ r_+ \in c}} j_r + \sum_{\substack{r \in \mathcal{R}_{damp}: \\ r_+ \in c}} j_r - \sum_{\substack{r \in \mathcal{R}_{slow}: \\ r_- \in c}} j_r, \quad \text{for } c \in \mathcal{C}, \quad (3.14b)$$

$$u_x = u_c \quad \text{and} \quad \sum_{\substack{r \in \mathcal{R}_{ficyc}: \\ r_+ = x}} \tilde{j}_r = \sum_{\substack{r \in \mathcal{R}_{ficyc}: \\ r_- = x}} \tilde{j}_r, \quad \text{for } x \in c \in \mathcal{C}, \quad (3.14c)$$

$$0 = \sum_{\substack{r \in \mathcal{R}_{slow}: \\ r_+ = x}} j_r + \sum_{\substack{r \in \mathcal{R}_{damp}: \\ r_+ = x}} j_r - \sum_{\substack{r \in \mathcal{R}_{damp}: \\ r_- = x}} j_r, \quad \text{for } x \in \mathcal{V}_1. \quad (3.14d)$$

These equations hold on $[0, T]$ in the weak sense of (3.3).

Proof Equation (3.14a) follows directly from Eq. (3.5b) by the convergence properties of Corollary 3.10. For fixed $c \in \mathcal{C}$ we sum equation (3.5c) over all $x \in c$ to find

$$\sum_{x \in c} \pi_x^\epsilon \dot{u}_x^\epsilon = \sum_{\substack{r \in \mathcal{R}_{slow}: \\ r_+ \in c}} j_r^\epsilon + \sum_{\substack{r \in \mathcal{R}_{damp}: \\ r_+ \in c}} j_r^\epsilon - \sum_{\substack{r \in \mathcal{R}_{slow}: \\ r_- \in c}} j_r^\epsilon.$$

Note that the final two sums in (3.5c) cancel by Lemma 3.2. The left-hand side equals $\pi_c^\epsilon \dot{u}_c^\epsilon$ and converges in distributional sense by (3.12c); the remaining terms also converge by (3.12f) and (3.12g). The limit equation is (3.14b).

Equation (3.14c) is the content of Lemma 3.12. Finally, to prove (3.14d) we write (3.5d) for $x \in \mathcal{V}_1$ as

$$\epsilon \tilde{\pi}_x^\epsilon \dot{u}_x^\epsilon = \sum_{\substack{r \in \mathcal{R}_{slow}: \\ r_+ = x}} j_r^\epsilon + \sum_{\substack{r \in \mathcal{R}_{damp}: \\ r_+ = x}} j_r^\epsilon - \sum_{\substack{r \in \mathcal{R}_{damp}: \\ r_- = x}} j_r^\epsilon.$$

The left-hand side converges to zero in distributional sense by (3.12e), and the right-hand side again converges by (3.12f) and (3.12g). \square

As an immediate consequence, the Γ -limit $\tilde{\mathcal{I}}_0^0 + \tilde{\mathcal{J}}^0$ from Theorem 1.2 can only be finite if these limit continuity equations (3.14) hold.

Note that although the densities $u_{\mathcal{V}_1}$ do appear in the limit rate functional $\tilde{\mathcal{J}}_{damp}^0$, they become decoupled from the other variables in the sense that they have vanished completely from the continuity equations. Furthermore, if one does not take fluxes into account, the mass flowing into a \mathcal{V}_1 node will be instantaneously distributed over the next nodes, which would lead to a contracted network as drawn on the right of Fig. 2. At the level of fluxes this is contraction is reflected in (3.14d).

Remark 3.15 Note that $L^\infty([0, T]) \ni u_x \stackrel{\text{a.e.}}{=} u_c \in C([0, T])$, so that in general $u_x(0) \neq u_c(0)$; the mass that is initially present will be spread out over the component c at every positive time $t > 0$, but not at $t = 0$. The same principle can be seen in the strengthened equilibration in the next section, which only holds in the time interval $[t_0, T]$ for any $t_0 > 0$.

Remark 3.16 If there are no damped cycles, as in Sect. 1.8, then Lemmas 3.7 and 3.14 show that $u_{\mathcal{V}_{0slow}} \in W^{1,\mathcal{C}}([0, T]; \mathbb{R}^{\mathcal{V}_{0slow}})$ and similarly $u_c \in W^{1,\mathcal{C}}([0, T]; \mathbb{R}^{\mathcal{C}})$.

3.6 Strengthened Equilibration on Fast Cycle Components

In the previous sections we derived that for a sequence with uniformly bounded cost $\tilde{J}_0^\epsilon + \tilde{J}^\epsilon$, concentrations u_x^ϵ in a fast cycle $x \in \mathfrak{c} \in \mathfrak{C}$ converge weakly-* in $L^\infty([0, T])$, whereas the weighted sum $u_\mathfrak{c}^\epsilon$ converges uniformly in $C([0, T])$. We now show that the convergence of u_x^ϵ can be strengthened to uniform convergence as well, as long as one does not include time 0 in the interval. This result will be needed later on for the construction of the recovery sequence, see Sect. 4.2.

Recall from Sect. 3.2 that sequences with bounded cost have uniformly bounded fluxes in L^1 . Together with the continuity equations, this will be the only requirement of the following result.

Lemma 3.17 *Let $(u^\epsilon, j^\epsilon)_{\epsilon>0}$ in Θ such that each (u^ϵ, j^ϵ) satisfy the continuity equations (3.5), and assume that all fluxes $j_{\mathcal{R}_{slow}}^\epsilon, j_{\mathcal{R}_{damp}}^\epsilon, \tilde{j}_{\mathcal{R}_{ficyc}}^\epsilon$ are L^1 -valued and uniformly bounded in $L^1(0, T; \mathbb{R}^{\mathcal{R}_{slow}})$, $L^1(0, T; \mathbb{R}^{\mathcal{R}_{damp}})$ and $L^1(0, T; \mathbb{R}^{\mathcal{R}_{ficyc}})$, and that $u_\mathfrak{c}^\epsilon \rightarrow u_\mathfrak{c}$ in $C([0, T]; \mathbb{R}^\mathfrak{C})$. Then for any $t_0 \in (0, T]$,*

$$u_x^\epsilon \rightarrow u_\mathfrak{c} \quad \text{strongly in } L^\infty([t_0, T]) \quad \text{for each } x \in \mathfrak{c} \in \mathfrak{C}. \tag{3.15}$$

If in addition,

$$-(\operatorname{div} j^\epsilon(t))_x := \sum_{\substack{r \in \mathcal{R}_{slow}: \\ r_+ = x}} j_r^\epsilon(t) + \sum_{\substack{r \in \mathcal{R}_{damp}: \\ r_+ = x}} j_r^\epsilon(t) - \sum_{\substack{r \in \mathcal{R}_{slow}: \\ r_- = x}} j_r^\epsilon(t),$$

and

$$-\frac{1}{\sqrt{\epsilon}}(\operatorname{div} \tilde{j}^\epsilon(t))_x := \frac{1}{\sqrt{\epsilon}} \sum_{\substack{r \in \mathcal{R}_{ficyc}: \\ r_+ = x}} \tilde{j}_r^\epsilon(t) - \frac{1}{\sqrt{\epsilon}} \sum_{\substack{r \in \mathcal{R}_{ficyc}: \\ r_- = x}} \tilde{j}_r^\epsilon(t)$$

are both uniformly bounded in $L^\infty(0, T; \mathbb{R}^{\mathcal{V}_{oficyc}})$ and $u_x^\epsilon(0) = u_\mathfrak{c}^\epsilon(0)$ for each $x \in \mathfrak{c} \in \mathfrak{C}$, then

$$u_x^\epsilon \rightarrow u_\mathfrak{c} \quad \text{strongly in } L^\infty([0, T]) \quad \text{for each } x \in \mathfrak{c} \in \mathfrak{C}. \tag{3.16}$$

Proof We prove the result for one fast cycle $\mathfrak{c} \in \mathfrak{C}$; for the length of this proof we therefore restrict the parameter x to the elements of \mathfrak{c} . To exploit the stochastic structure we temporarily write $\rho_x^\epsilon(t) := \pi_x^\mathfrak{c} u_x^\epsilon(t)$, and

$$(A^\top \rho^\epsilon(t))_x := \sum_{\substack{r \in \mathcal{R}_{ficyc}: \\ r_+ = x}} \kappa_r \rho_{r_-}^\epsilon(t) - \rho_x^\epsilon(t) \sum_{\substack{r \in \mathcal{R}_{ficyc}: \\ r_- = x}} \kappa_r,$$

so that A is the generator matrix of the Markov chain that consists of the irreducible fast cycle \mathfrak{c} , which does not depend on ϵ . Note that A has one-dimensional kernel spanned by the all-ones vector $\mathbf{1} := (1, \dots, 1)$, and A^\top has one-dimensional kernel spanned by π^ϵ . Recall from (3.5c) that for each $x \in \mathfrak{c}$:

$$\dot{\rho}_x^\epsilon(t) = -(\operatorname{div} j^\epsilon(t))_x - \frac{1}{\sqrt{\epsilon}}(\operatorname{div} \tilde{j}^\epsilon(t))_x + \frac{1}{\epsilon}(A^\top \rho^\epsilon(t))_x =: f_x^\epsilon(t) + \frac{1}{\epsilon}(A^\top \rho^\epsilon(t))_x.$$

Writing $\mu_x^\epsilon(t) := \rho_x^\epsilon(t) - (\pi_x^\epsilon/\pi_c^\epsilon) \sum_{y \in c} \rho_y^\epsilon(t)$, we calculate, using the characterisation of the kernel of A , that

$$\dot{\mu}_x^\epsilon(t) = f_x^\epsilon(t) - \underbrace{\frac{\pi_x^\epsilon}{\pi_c^\epsilon} \sum_{y \in c} f_y^\epsilon(t)}_{g_x^\epsilon(t)} + \frac{1}{\epsilon} (A^\top \mu^\epsilon(t))_x \quad \text{for } x \in c.$$

This equation can be integrated to

$$\mu^\epsilon(t) = e^{A^\top t/\epsilon} \mu^\epsilon(0) + \int_0^t e^{A^\top(t-s)/\epsilon} g^\epsilon(s) ds.$$

Recall that A is irreducible so that it has only one zero eigenvalue and no other eigenvalues on the imaginary axis (e.g. [13, Lemma V.4.2]). We restrict the operator A^\top to the orthogonal complement of the zero eigenspace by defining

$$\mathbf{1}^\perp := \{v \in \mathbb{R}^c : \mathbf{1}^\top v = 0\}, \quad \text{and} \quad B := A^\top|_{\mathbf{1}^\perp},$$

and equip $\mathbf{1}^\perp$ with a norm $\|\cdot\|$.

Since $\mathbf{1}^\top A^\top = 0$, B is well defined, and B has the same eigenvalues and eigenvectors as A^\top except for the zero eigenvalue-eigenvector pair. Therefore $\alpha(B) := \max \operatorname{Re} \sigma(B) < 0$. Since the δ -pseudospectrum $\sigma_\delta(B)$ converges to the spectrum $\sigma(B)$ as $\delta \rightarrow 0$, there exists a $\delta > 0$ such that $\lambda := -\sup \operatorname{Re} \sigma_\delta(B) > 0$. By [42, Th. 15.2] we find that there exists $M > 0$ such that

$$\|e^{Bt}\| \leq M e^{-\lambda t}.$$

Because $(\mu^\epsilon(t))_t, (g^\epsilon(t))_t \subset \mathbf{1}^\top$ we can estimate,

$$\begin{aligned} \|\mu^\epsilon(t)\| &\leq \|e^{A^\top t/\epsilon} \mu^\epsilon(0)\| + \int_0^t \|e^{A^\top(t-s)/\epsilon} g^\epsilon(s)\| ds \\ &= \|e^{Bt/\epsilon} \mu^\epsilon(0)\| + \int_0^t \|e^{B(t-s)/\epsilon} g^\epsilon(s)\| ds \\ &\leq M e^{-\lambda t/\epsilon} \|\mu^\epsilon(0)\| + M \int_0^t e^{-\lambda(t-s)/\epsilon} \|g^\epsilon(s)\| ds. \end{aligned} \tag{3.17}$$

The fact that the $L^1(0, T)$ -norms of g^ϵ are bounded uniformly in ϵ implies that $\mu^\epsilon(t)$ converges to zero uniformly in $t \in [t_0, T]$ for any $t_0 > 0$. This proves (3.15).

To prove the stronger statement of (3.16), note that the strengthened assumptions imply that $\|\mu^\epsilon(0)\|$ and $\|g^\epsilon\|_{L^\infty(0,T)}$ converge to zero as $\epsilon \rightarrow 0$. In combination with (3.17) this proves (3.15). □

4 Γ -Convergence

This section is devoted to the proof of the main Γ -convergence Theorem 1.2, which consists of the lower bound, Proposition 4.1, and the existence of a recovery sequence in Proposition 4.5.

4.1 Γ -Lower Bounds

The Γ -lower bound is summarised in the following.

Proposition 4.1 (Γ -lower bound) *For any sequence $(u^\epsilon, j^\epsilon) \rightarrow (u, j)$ in Θ ,*

$$\liminf_{\epsilon \rightarrow 0} \tilde{\mathcal{I}}_0^\epsilon(u^\epsilon(0)) + \tilde{\mathcal{J}}^\epsilon(u^\epsilon, j^\epsilon) \geq \tilde{\mathcal{I}}_0^0(u(0)) + \tilde{\mathcal{J}}^0(u, j).$$

Proof We treat each functional $\tilde{\mathcal{I}}_0^\epsilon, \tilde{\mathcal{J}}_{\text{slow}}^\epsilon, \tilde{\mathcal{J}}_{\text{damp}}^\epsilon$ and $\tilde{\mathcal{J}}_{\text{fcyc}}^\epsilon$ separately, and without loss of generality we may always assume that $\tilde{\mathcal{I}}_0^\epsilon + \tilde{\mathcal{J}}^\epsilon \leq C$ for some $C \geq 0$ and hence the continuity equations (3.5) hold; otherwise the lower bound is trivial. This is carried out in the next Lemmas 4.2, 4.3 and 4.4. \square

For the initial condition, recall the definitions of $\tilde{\mathcal{I}}_0^\epsilon$ and $\tilde{\mathcal{I}}_0$ from (1.11) and Sect. 1.7, and observe that the first one depends on $u_{\mathcal{V}_{0\text{fcyc}}}(0)$ whereas the second depends on $u_{\mathcal{C}}(0)$, which may be different, see Remark 3.15. Hence the Γ -convergence of $\tilde{\mathcal{I}}_0^\epsilon$ to $\tilde{\mathcal{I}}_0$ does not hold in $\mathbb{R}^{\mathcal{V}_{0\text{slow}}} \times \mathbb{R}^{\mathcal{V}_{0\text{fcyc}}} \times \mathbb{R}^{\mathcal{C}} \times \mathbb{R}^{\mathcal{V}_1}$, but only in the path-space convergence of (3.12).

Lemma 4.2 (Γ -lower bound for the initial condition) *Let $(u^\epsilon, j^\epsilon)_{\epsilon>0} \subset \Theta$ be a sequence such that $\tilde{\mathcal{I}}_0^\epsilon(u^\epsilon(0)) + \tilde{\mathcal{J}}^\epsilon(u^\epsilon, j^\epsilon) \leq C$ and $(u^\epsilon, j^\epsilon) \rightarrow (u, j)$ in Θ (i.e. in the sense of (3.12)). Then:*

$$\liminf_{\epsilon \rightarrow 0} \tilde{\mathcal{I}}_0^\epsilon(u^\epsilon(0)) \geq \tilde{\mathcal{I}}_0^0(u(0)).$$

Proof By uniform convergence, $u_{\mathcal{V}_{0\text{slow}}}^\epsilon(0) \rightarrow u_{\mathcal{V}_{0\text{slow}}}(0)$ so that

$$\sum_{x \in \mathcal{V}_{0\text{slow}}} s(\pi_x^\epsilon u_x^\epsilon(0) \mid \pi_x^\epsilon) \rightarrow \sum_{x \in \mathcal{V}_{0\text{slow}}} s(\pi_x u_x(0) \mid \pi_x),$$

and clearly

$$\sum_{x \in \mathcal{V}_1} s(\pi_x^\epsilon u_x^\epsilon(0) \mid \pi_x^\epsilon) \geq 0.$$

Lemma 3.2 shows that every $x \in \mathcal{V}_{0\text{fcyc}}$ is part of exactly one component $c \in \mathcal{C}$. From (1.8), Jensen’s inequality and the continuity equation (3.5a),

$$\begin{aligned} \sum_{x \in \mathcal{V}_{0\text{fcyc}}} s(\pi_x^\epsilon u_x^\epsilon(0) \mid \pi_x^\epsilon) &= \sum_{c \in \mathcal{C}} \pi_c^\epsilon \sum_{x \in c} s(u_x^\epsilon(0) \mid 1) \frac{\pi_x^\epsilon}{\pi_c^\epsilon} \\ &\geq \sum_{c \in \mathcal{C}} \pi_c^\epsilon s(\sum_{x \in c} u_x^\epsilon(0) \frac{\pi_x^\epsilon}{\pi_c^\epsilon} \mid 1) \\ &= \sum_{c \in \mathcal{C}} s(\pi_c^\epsilon u_c^\epsilon(0) \mid \pi_c^\epsilon) \rightarrow \sum_{c \in \mathcal{C}} s(\pi_c u_c(0) \mid \pi_c), \end{aligned}$$

again by uniform convergence of u_c^ϵ . \square

Lemma 4.3 (Γ -lower bound for the slow and damped fluxes)

Let $(u^\epsilon, j^\epsilon)_{\epsilon>0} \subset \Theta$ be a sequence such that $\tilde{\mathcal{I}}_0^\epsilon(u^\epsilon(0)) + \tilde{\mathcal{J}}^\epsilon(u^\epsilon, j^\epsilon) \leq C$ and $(u^\epsilon, j^\epsilon) \rightarrow (u, j)$ in Θ . Then:

$$\liminf_{\epsilon \rightarrow 0} \tilde{\mathcal{J}}_{\text{slow}}^\epsilon(u_{\mathcal{V}_{0\text{slow}}}^\epsilon, u_{\mathcal{V}_{0\text{fcyc}}}^\epsilon, j_{\mathcal{R}_{\text{slow}}}^\epsilon) \geq \tilde{\mathcal{J}}_{\text{slow}}^0(u_{\mathcal{V}_{0\text{slow}}}, u_{\mathcal{V}_{0\text{fcyc}}}, j_{\mathcal{R}_{\text{slow}}}),$$

and

$$\liminf_{\epsilon \rightarrow 0} \tilde{J}_{damp}^\epsilon(u_{\nu_1}^\epsilon, j_{\mathcal{R}_{damp}}^\epsilon) \geq \tilde{J}_{damp}^0(u_{\nu_1}, j_{\mathcal{R}_{damp}}).$$

Proof Recall the uniform L^1 -bounds on the slow and damped fluxes from Lemmas 3.6 and 3.7. The statement for slow fluxes follows directly from rewriting

$$\tilde{J}_{slow}^0(u_{\nu_0}^\epsilon, j_{\mathcal{R}_{slow}}^\epsilon) = \sum_{r \in \mathcal{R}_{slow}} \int_{[0, T]} \left[s \left(j_r^\epsilon(t) \mid \kappa_r \pi_{r-} u_{r-}^\epsilon(t) \right) + j_r^\epsilon(t) \log \frac{\pi_{r-}}{\pi_{r-}^\epsilon} - \pi_{r-}^\epsilon + \pi_{r-} \right] dt, \tag{4.1}$$

together with the joint lower semicontinuity from Lemma 2.2, and $\pi^\epsilon \rightarrow \pi > 0$. The argument for the damped fluxes is the same after generalising to possible measure-valued trajectories in time. \square

Lemma 4.4 (Γ -lower bound for the fast cycle fluxes) *Let $(u^\epsilon, j^\epsilon)_{\epsilon > 0} \subset \Theta$ be a sequence such that $\tilde{I}_0^\epsilon(u^\epsilon(0)) + \tilde{J}^\epsilon(u^\epsilon, j^\epsilon) \leq C$ and $(u^\epsilon, j^\epsilon) \rightarrow (u, j)$ in Θ . Then:*

$$\liminf_{\epsilon \rightarrow 0} \tilde{J}_{fyc}^\epsilon(u_{\nu_{0fyc}}^\epsilon, \tilde{j}_{\mathcal{R}_{fyc}}^\epsilon) \geq \tilde{J}_{fyc}^0(u_{\nu_{0fyc}}, \tilde{j}_{\mathcal{R}_{fyc}}).$$

A similar statement is proven in [6, Th. 2].

Proof To simplify notation we prove the statement for one arbitrary $r \in \mathcal{R}_{fyc}$. We first note that

$$\pi_{r-}^\epsilon u_{r-}^\epsilon \rightharpoonup \pi_{r-} u_{r-} \quad \text{in } L^1([0, T]) \quad \text{and} \quad \sup_{\epsilon > 0} \|\pi_{r-}^\epsilon u_{r-}^\epsilon\|_{L^1} < \infty,$$

and that for any test function $\zeta \in C([0, T])$,

$$\frac{1}{\epsilon} e^{\sqrt{\epsilon} \zeta} - \frac{1}{\epsilon} - \frac{1}{\sqrt{\epsilon}} \zeta \xrightarrow[\epsilon \rightarrow 0]{L^\infty} \frac{1}{2} \zeta^2.$$

It then follows that the following integral converges:

$$\int_{[0, T]} \kappa_r \pi_{r-}^\epsilon u_{r-}^\epsilon(t) \left(\frac{1}{\epsilon} e^{\sqrt{\epsilon} \zeta(t)} - \frac{1}{\epsilon} - \frac{1}{\sqrt{\epsilon}} \zeta(t) \right) dt \rightarrow \frac{1}{2} \int_{[0, T]} \kappa_r \pi_{r-} u_{r-}(t) \zeta(t)^2 dt.$$

Using the dual formulations of Lemmas 2.2 and 2.3,

$$\begin{aligned} & \liminf_{\epsilon \rightarrow 0} \int_{[0, T]} s \left(\frac{1}{\epsilon} \kappa_r \pi_{r-}^\epsilon u_{r-}^\epsilon(t) + \frac{1}{\sqrt{\epsilon}} \tilde{j}_r^\epsilon(t) \mid \frac{1}{\epsilon} \kappa_r \pi_{r-}^\epsilon u_{r-}^\epsilon(t) \right) \\ & \geq \sup_{\zeta \in C([0, T])} \liminf_{\epsilon \rightarrow 0} \int_{[0, T]} \left[\zeta(t) \tilde{j}_r^\epsilon(t) - \kappa_r \pi_{r-}^\epsilon u_{r-}^\epsilon(t) \left(\frac{1}{\epsilon} e^{\sqrt{\epsilon} \zeta(t)} - \frac{1}{\epsilon} - \frac{1}{\sqrt{\epsilon}} \zeta(t) \right) \right] dt \\ & = \sup_{\zeta \in C([0, T])} \int_{[0, T]} \left[\zeta(t) \tilde{j}_r(t) - \frac{1}{2} \kappa_r \pi_{r-} u_{r-}(t) \zeta(t)^2 \right] dt \\ & = \begin{cases} \frac{1}{2} \int_{[0, T]} \left(\frac{\tilde{j}_r(t)}{u_{r-}(t)} \right)^2 \kappa_r \pi_{r-} u_{r-}(t) dt, & \text{if } \tilde{j}_r \ll u_{r-}, \\ \infty, & \text{otherwise.} \end{cases} \end{aligned}$$

\square

4.2 Γ -Recovery Sequence

For each of the four functionals separately, convergence is easily shown using a constant sequence $(u^\epsilon, j^\epsilon) \equiv (u, j)$. However, such a constant sequence is not a valid recovery sequence as it violates the continuity equations (3.5). The construction of the recovery sequence is summarised in the following proposition.

Proposition 4.5 (Γ -recovery sequence) *For any (u, j) in Θ there exists a sequence $(u^\epsilon, j^\epsilon)_\epsilon \subset \Theta$ such that $(u^\epsilon, j^\epsilon) \rightarrow (u, j)$ in Θ and*

$$\lim_{\epsilon \rightarrow 0} \tilde{I}_0^\epsilon(u^\epsilon(0)) + \tilde{J}^\epsilon(u^\epsilon, j^\epsilon) = \tilde{I}_0^0(u(0)) + \tilde{J}^0(u, j).$$

Proof In Lemma 4.7 we first show that (u, j) can be approximated by a regularised (u^δ, j^δ) such that the limit functional converges, i.e. such that $\tilde{I}^0(u^\delta(0)) + \tilde{J}^0(u^\delta, j^\delta) \rightarrow \tilde{I}^0(u(0)) + \tilde{J}^0(u, j)$ as $\delta \rightarrow 0$. In Lemma 4.9 we construct a recovery sequence (u^ϵ, j^ϵ) corresponding to such regularised (u^δ, j^δ) , and then use a diagonal argument to construct a recovery sequence for arbitrary (u, j) , see for example [9, Prop. 6.2]. □

Remark 4.6 So far, we only assumed $\sum_{x \in \mathcal{V}} \pi_x^\epsilon = 1$, whereas the total mass $\sum_{x \in \mathcal{V}} \pi_x^\epsilon u_x^\epsilon(t)$ is only bounded above by (3.6). All arguments in this paper can be extended to the case where the total mass is fixed. In that case the construction of the recovery sequence becomes slightly more involved, since adding mass to certain nodes must be balanced by subtracting mass from other nodes.

Lemma 4.7 (Approximation of the limit functional) *Let $(u, j) \in \Theta$ such that $\tilde{I}_0^0(u(0)) + \tilde{J}^0(u, j) < \infty$, so (u, j) satisfies the limit continuity equations (3.14). Then there exists a sequence $(u^\delta, j^\delta)_{\delta > 0} \in \Theta$ such that for each $\delta > 0$,*

1. $(u^\delta, j^\delta) \in C_b^\infty([0, T]; \mathbb{R}^{\mathcal{V}_{0slow}} \times \mathbb{R}^{\mathcal{V}_{0fycyc}} \times \mathbb{R}^{\mathcal{C}} \times \mathbb{R}^{\mathcal{V}_1} \times \mathbb{R}^{\mathcal{R}_{slow}} \times \mathbb{R}^{\mathcal{R}_{damp}} \times \mathbb{R}^{\mathcal{R}_{fycyc}})$,
2. (u^δ, j^δ) satisfies the limit continuity equations (3.14),
3. $\inf_{t \in [0, T]} u_x^\delta(t) > 0$ for all $x \in \mathcal{V}_{0slow} \cup \mathcal{V}_{0fycyc} \cup \mathcal{C} \cup \mathcal{V}_1$,
4. $j_r^\delta \geq \sum_{x \in \mathcal{V}_1} \delta \tilde{\pi}_x \|\dot{u}_x^\delta\|_{L^\infty}$ for all $r \in \mathcal{R}_{slow} \cup \mathcal{R}_{damp}$, and as $\delta \rightarrow 0$,
5. $(u^\delta, j^\delta) \rightarrow (u, j)$ in Θ ,
6. $\tilde{I}_0^0(u^\delta(0)) + \tilde{J}^0(u^\delta, j^\delta) \rightarrow \tilde{I}_0^0(u(0)) + \tilde{J}^0(u, j)$.

Proof We construct the approximation in three steps.

Step 1: convolution. Note that for each $x \in \mathcal{V}_0$ the concentration $t \mapsto u_x(t)$ is continuous; for $x \in \mathcal{V}_{0slow}$ this follows from the definition of Θ , and for $x \in \mathcal{V}_{0fycyc}$ this follows from the continuity of $t \mapsto u_{\mathcal{C}}(t)$ in Θ and the continuity equation (3.14c). We first extend $u_{\mathcal{V}_0}$ beyond $[0, T]$ by constants, and $u_{\mathcal{V}_1}$ and j by zero. Observe that with this extension the pair (u, j) satisfies the continuity equation (3.14) in the sense of distributions on the whole time interval \mathbb{R} (which is a stronger statement than the usual interpretation (3.3)). We then approximate (u, j) by convoluting with the heat kernel: $(u^\delta, j^\delta) := (u * \theta^\delta, j * \theta^\delta)$, where $\theta^\delta(t) := (4\pi\delta)^{-1/2} e^{-t^2/(4\delta)}$. Since (u, j) satisfies the linear continuity equations (3.14) in the sense of distributions on \mathbb{R} , they are also satisfied for the convolution (u^δ, j^δ) .

It is easily checked that $(u^\delta, j^\delta)|_{[0, T]} \rightarrow (u, j)|_{[0, T]}$ in Θ . The initial conditions $u_x^\delta(0)$ converge for $x \in \mathcal{V}_{0slow} \cup \mathcal{C}$ and so by continuity $\tilde{I}_0^0(u^\delta(0)) \rightarrow \tilde{I}_0^0(u(0))$. The bound $\liminf_{\delta \rightarrow 0} \tilde{J}^0(u^\delta, j^\delta) \geq \tilde{J}^0(u, j)$ is for free because of lower semicontinuity (see Sect. 2.4).

The bound in the other direction is obtained by exploiting the joint convexity of $(u, j) \mapsto \tilde{\mathcal{J}}^0(u, j)$ and applying Jensen’s inequality to the probability measure θ^δ ; see [38, Lem. 3.12]. **Step 2: add constants to the densities.** For the next step we further approximate the sequence (u^δ, j^δ) , but to reduce clutter we now assume that the procedure above is already applied so that we are given a smooth and bounded (u, j) . We make all densities positive by adding a constant $\delta > 0$, i.e.

$$u_x^\delta(t) := u_x(t) + \delta \quad \text{for } 0 \leq t \leq T, x \in \mathcal{V}.$$

It follows automatically that $u_c^\delta = u_c + \delta$. We leave the fluxes j invariant, and the resulting pair (u^δ, j^δ) again satisfies the limiting continuity equations (3.14). The following lemma shows that the limit functional $\tilde{\mathcal{I}}_0 + \tilde{\mathcal{J}}^0$ converges along the sequence (u^δ, j^δ) .

Lemma 4.8 *Recall the definition (1.12) of s for measures. Let $a, b \in \mathcal{M}_{\geq 0}([0, T])$ satisfy*

$$\int_{[0, T]} s(a|b)(dt) < \infty.$$

Then setting $b^\delta(dt) := b(dt) + \delta dt$ we have

$$\lim_{\delta \rightarrow 0} \int_{[0, T]} s(a|b^\delta)(dt) = \int_{[0, T]} s(a|b)(dt).$$

Proof of Lemma 4.8 We write

$$s(a|b^\delta)(dt) = a(dt) \log \frac{da}{db^\delta}(t) - a(dt) + b^\delta(dt).$$

After integration over $[0, T]$ the final term $b^\delta([0, T])$ converges to $b([0, T])$ as $\delta \rightarrow 0$; in the first term the argument of the logarithm is decreasing in δ , and therefore the first term converges by the Monotone Convergence Theorem. \square

Step 3: add constant fluxes. Again to reduce clutter we may assume that we are given an (u, j) satisfying properties 1, 2, and 3 of the Lemma. By irreducibility of the network there exists a cycle $(r^k)_{k=1}^K \subset \mathcal{R}_{\text{ficyc}}, r_+^k = r_-^{k+1}$ for $k = 1, \dots, K - 1, r_-^1 = r_+^1 = r_+^K$, such that each damped flux $r \in \mathcal{R}_{\text{damp}}$ is contained in the cycle at least once. Note that some fluxes r may occur multiple times, namely $n(r) := \#\{k = 1, \dots, K : r^k = r\}$ times in the cycle. For each $k = 1, \dots, K$ we define the new approximation:

$$\begin{cases} j_{r^k}^\delta := j_{r^k} + n(r^k) \sum_{y \in \mathcal{V}_1} \delta \tilde{\pi}_y \|\dot{u}_y\|_{L^\infty}, & r^k \in \mathcal{R}_{\text{damp}} \cup \mathcal{R}_{\text{slow}}, \\ \tilde{j}_{r^k}^\delta := \tilde{j}_{r^k} + \sqrt{\epsilon} n(r^k) \sum_{y \in \mathcal{V}_1} \delta \tilde{\pi}_y \|\dot{u}_y\|_{L^\infty}, & r^k \in \mathcal{R}_{\text{ficyc}}. \end{cases}$$

Substituting these modified fluxes into the limit continuity equations (3.14) shows that the concentrations are left unchanged, since some extra mass is being pushed around in cycles. Since the fluxes are only changed by adding a constant, it is easily checked that $(u, j^\delta) \rightarrow (u, j)$ in Θ , and by Lemma 4.8 we find $\tilde{\mathcal{J}}^0(u, j^\delta) \rightarrow \tilde{\mathcal{J}}^0(u, j)$ as $\delta \rightarrow 0$. \square

We now construct a recovery sequence (u^ϵ, j^ϵ) for $(u, j) \in \Theta$ that is regularised by Lemma 4.7. The difficulty is to construct the sequence such that the continuity equations hold in the \mathcal{V}_1 and $\mathcal{V}_{0\text{ficyc}}$ nodes. The problem with the \mathcal{V}_1 nodes is that the continuity equations (3.5d) and (3.14d) are different, but $u_{\mathcal{V}_1}^\epsilon$ needs to converge to $u_{\mathcal{V}_1}$. This will be done by transporting exactly the right amount of mass from certain \mathcal{V}_0 -nodes to the \mathcal{V}_1 -nodes. To satisfy the continuity equations in the $\mathcal{V}_{0\text{ficyc}}$ nodes, we define $u_{\mathcal{V}_{0\text{ficyc}}}^\epsilon$ through the continuity equations, and use the strengthened convergence result of Sect. 3.6 to pass to the limit. \square

Lemma 4.9 (Recovery sequence for regularised paths) *Let $(u, j) \in \Theta$ satisfy properties 1, 2, 3 and 4 of Lemma 4.7. Then there exists a sequence $(u^\epsilon, j^\epsilon) \in \Theta$ such that:*

1. (u^ϵ, j^ϵ) satisfies the ϵ -dependent continuity equations (3.5);
2. $(u^\epsilon_{\mathcal{V}_{0\text{slow}}}, u^\epsilon_{\mathcal{R}_{\text{fcyc}}}, u^\epsilon_{\mathfrak{c}}, u^\epsilon_{\mathcal{V}_1}, j^\epsilon_{\mathcal{R}_{\text{slow}}}, j^\epsilon_{\mathcal{R}_{\text{damp}}}, \tilde{j}^\epsilon_{\mathcal{R}_{\text{fcyc}}}) \rightarrow (u_{\mathcal{V}_{0\text{slow}}}, u_{\mathcal{R}_{\text{fcyc}}}, u_{\mathfrak{c}}, u_{\mathcal{V}_1}, j_{\mathcal{R}_{\text{slow}}}, j_{\mathcal{R}_{\text{damp}}}, \tilde{j}_{\mathcal{R}_{\text{fcyc}}})$ uniformly on $[0, T]$;
3. $\tilde{I}_0^\epsilon(u^\epsilon(0)) \rightarrow \tilde{I}_0^0(u(0))$ and $\tilde{J}^\epsilon(u^\epsilon, j^\epsilon) \rightarrow \tilde{J}^0(u, j)$.

Proof For ease of notation we pick only one node $\hat{x} \in \mathcal{V}_{0\text{slow}}$, whose density is bounded from below by assumption. We will approximate all fluxes such that a little mass is transported from node \hat{x} to all \mathcal{V}_1 -nodes, as follows. Since the network is irreducible, there exists, for each $y \in \mathcal{V}_1$, a connecting chain $Q(\hat{x}, y) := (r^{k,y})_{k=1}^{K_y} \subset \mathcal{R}$, $r_-^{1,y} = \hat{y}$, $r_+^{k,y} = r_-^{k+1,y}$ for $k = 1, \dots, K_y - 1$ and $r^{K_y} = y$. For these connecting chains we may assume without loss of generality that no $r \in \mathcal{R}$ occurs multiple times in a chain $Q(\hat{x}, y)$. Define for all $r \in \mathcal{R}$:

$$\begin{cases} j_r^\epsilon := j_r + \sum_{y \in \mathcal{V}_1: r \in Q(\hat{x}, y)} \pi_y^\epsilon \dot{u}_y, & r \in \mathcal{R}_{\text{slow}} \cup \mathcal{R}_{\text{damp}}, \\ \tilde{j}_r^\epsilon := \tilde{j}_r + \frac{1}{\sqrt{\epsilon}} \sum_{y \in \mathcal{V}_1: r \in Q(\hat{x}, y)} \pi_y^\epsilon \dot{u}_y, & r \in \mathcal{R}_{\text{fcyc}}. \end{cases}$$

Note that by the assumed properties 1 and 3 of Lemma 4.7 together with $\pi_{\mathcal{V}_1}^\epsilon \rightarrow 0$, all approximated fluxes $j_r^\epsilon, \tilde{j}_r^\epsilon$ are non-negative for ϵ small enough. Clearly all fluxes j^ϵ converge uniformly to j , since $\pi_{\mathcal{V}_1}^\epsilon / \sqrt{\epsilon} \rightarrow 0$. For the initial conditions, set

$$\begin{aligned} u_x^\epsilon(0) &:= \frac{\pi_x^\epsilon}{\pi_x} u_x(0), && \text{for all } x \in \mathcal{V}_{0\text{slow}}, \\ u_{\mathfrak{c}}^\epsilon(0) &= u_x^\epsilon(0) := u_x(0) = u_{\mathfrak{c}}(0), && \text{for all } x \in \mathfrak{c} \in \mathfrak{C}, \\ u_x^\epsilon(0) &:= u_x(0), && \text{for all } x \in \mathcal{V}_1, \end{aligned} \tag{4.2}$$

and define the paths u^ϵ by the continuity equations (3.5).

More precisely, by construction for $x \in \mathcal{V}_{0\text{slow}}$:

$$\begin{aligned} \pi_x^\epsilon u_x^\epsilon(t) &\stackrel{(3.5b)}{:=} \pi_x u_x(0) + \sum_{\substack{r \in \mathcal{R}_{\text{slow}}: \\ r_+ = x}} \int_0^t j_r^\epsilon(s) ds + \sum_{\substack{r \in \mathcal{R}_{\text{damp}}: \\ r_+ = x}} \int_0^t j_r^\epsilon(s) ds - \sum_{\substack{r \in \mathcal{R}_{\text{slow}}: \\ r_- = x}} \int_0^t j_r^\epsilon(s) ds \\ &= \pi_x u_x(t) - \mathbb{1}_{\{x = \hat{x}\}} \sum_{y \in \mathcal{V}_1} \pi_y^\epsilon u_y(t), \end{aligned}$$

which is bounded away from zero (for ϵ small enough) by the assumed properties 1 and 3 of Lemma 4.7 together with $\pi_{\mathcal{V}_1}^\epsilon \rightarrow 0$. Clearly $u_x^\epsilon \rightarrow u_x$ uniformly.

For $x \in \mathcal{V}_1$, the densities will be constant in ϵ , since:

$$\begin{aligned} \pi_x^\epsilon u_x^\epsilon(t) &\stackrel{(3.5d)}{:=} \pi_x^\epsilon u_x(0) + \sum_{\substack{r \in \mathcal{R}_{\text{slow}}: \\ r_+ = x}} \int_0^t j_r^\epsilon(s) ds + \sum_{\substack{r \in \mathcal{R}_{\text{damp}}: \\ r_+ = x}} \int_0^t j_r^\epsilon(s) ds \\ &\quad - \sum_{\substack{r \in \mathcal{R}_{\text{damp}}: \\ r_- = x}} \int_0^t j_r^\epsilon(s) ds = \pi_x^\epsilon u_x(t). \end{aligned}$$

For $x \in \mathfrak{c} \in \mathfrak{C}$, the density $u_x^\epsilon(t)$ is defined as the solution of the coupled equations:

$$\begin{aligned} \pi_x^\epsilon \dot{u}_x^\epsilon & \stackrel{(3.5c)}{:=} \sum_{\substack{r \in \mathcal{R}_{\text{slow}}: \\ r_+ = x}} j_r^\epsilon + \sum_{\substack{r \in \mathcal{R}_{\text{damp}}: \\ r_+ = x}} j_r^\epsilon - \sum_{\substack{r \in \mathcal{R}_{\text{slow}}: \\ r_- = x}} j_r^\epsilon \\ & + \sum_{\substack{r \in \mathcal{R}_{\text{fcyc}}: \\ r_+ = x}} \left(\frac{1}{\epsilon} \kappa_r \pi_{r_-}^\epsilon u_{r_-}^\epsilon + \frac{1}{\sqrt{\epsilon}} \tilde{j}_r^\epsilon \right) - \sum_{\substack{r \in \mathcal{R}_{\text{fcyc}}: \\ r_- = x}} \left(\frac{1}{\epsilon} \kappa_r \pi_{r_-}^\epsilon u_{r_-}^\epsilon + \frac{1}{\sqrt{\epsilon}} \tilde{j}_r^\epsilon \right) \end{aligned} \tag{4.3}$$

with initial condition (4.2). Summing over $x \in \mathfrak{c}$ yields:

$$\begin{aligned} \pi_{\mathfrak{c}}^\epsilon \dot{u}_{\mathfrak{c}}^\epsilon & \stackrel{(3.5a)}{:=} \sum_{x \in \mathfrak{c}} \pi_x^\epsilon \dot{u}_x^\epsilon = \sum_{\substack{r \in \mathcal{R}_{\text{slow}}: \\ r_+ \in \mathfrak{c}}} j_r^\epsilon + \sum_{\substack{r \in \mathcal{R}_{\text{damp}}: \\ r_+ \in \mathfrak{c}}} j_r^\epsilon - \sum_{\substack{r \in \mathcal{R}_{\text{slow}}: \\ r_- \in \mathfrak{c}}} j_r^\epsilon \\ & + \sum_{\substack{r \in \mathcal{R}_{\text{fcyc}}: \\ r_+ \in \mathfrak{c}}} \left(\frac{1}{\epsilon} \kappa_r \pi_{r_-}^\epsilon u_{r_-}^\epsilon + \frac{1}{\sqrt{\epsilon}} \tilde{j}_r^\epsilon \right) - \sum_{\substack{r \in \mathcal{R}_{\text{fcyc}}: \\ r_- \in \mathfrak{c}}} \left(\frac{1}{\epsilon} \kappa_r \pi_{r_-}^\epsilon u_{r_-}^\epsilon + \frac{1}{\sqrt{\epsilon}} \tilde{j}_r^\epsilon \right) \\ & \stackrel{(3.14b), (3.14c)}{=} \pi_{\mathfrak{c}} \dot{u}_{\mathfrak{c}}. \end{aligned}$$

Together with the initial condition (4.2) this shows that $u_{\mathfrak{c}}^\epsilon \rightarrow u_{\mathfrak{c}}$ uniformly. Since all fluxes are uniformly bounded (and actually $\text{div } \tilde{j}^\epsilon \equiv 0$) and $u_x^\epsilon(0) = u_{\mathfrak{c}}^\epsilon(0)$ for $x \in \mathfrak{c} \in \mathfrak{C}$ we can apply Lemma 3.17 to (4.3) to derive that $u_x^\epsilon \rightarrow u_{\mathfrak{c}}$ uniformly on $[0, T]$ for all $x \in \mathfrak{c}$. Thus indeed all variables $(u_{\mathcal{V}_{0\text{slow}}}^\epsilon, u_{\mathcal{R}_{\text{fcyc}}}^\epsilon, u_{\mathfrak{C}}^\epsilon, u_{\mathcal{V}_1}^\epsilon, j_{\mathcal{R}_{\text{slow}}}^\epsilon, j_{\mathcal{R}_{\text{damp}}}^\epsilon, \tilde{j}_{\mathcal{R}_{\text{fcyc}}}^\epsilon) \rightarrow (u_{\mathcal{V}_{0\text{slow}}}, u_{\mathcal{R}_{\text{fcyc}}}, u_{\mathfrak{C}}, u_{\mathcal{V}_1}, j_{\mathcal{R}_{\text{slow}}}, j_{\mathcal{R}_{\text{damp}}}, \tilde{j}_{\mathcal{R}_{\text{fcyc}}})$ uniformly, which was to be shown.

To show convergence of $\tilde{\mathcal{I}}^\epsilon(u^\epsilon(0))$,

$$\begin{aligned} \tilde{\mathcal{I}}_0^\epsilon(u^\epsilon(0)) & = \sum_{x \in \mathcal{V}_{0\text{slow}}} \pi_x^\epsilon s\left(\frac{\pi_x^\epsilon}{\pi_x} u_x(0) \mid 1\right) + \sum_{\mathfrak{c} \in \mathfrak{C}} \sum_{x \in \mathfrak{c}} \pi_x^\epsilon s(u_{\mathfrak{c}}(0) \mid 1) + \sum_{x \in \mathcal{V}_1} \pi_x^\epsilon s(u_x(0) \mid 1) \\ & \rightarrow \sum_{x \in \mathcal{V}_{0\text{slow}}} \pi_x s(u_x(0) \mid 1) + \sum_{\mathfrak{c} \in \mathfrak{C}} \pi_{\mathfrak{c}} s(u_{\mathfrak{c}}(0) \mid 1) = \tilde{\mathcal{I}}_0^0(u(0)). \end{aligned}$$

To show convergence of $\tilde{\mathcal{J}}^\epsilon(u^\epsilon, j^\epsilon)$, we use the fact that all fluxes and densities are uniformly bounded, that is for ϵ sufficiently small and all $t \in [0, T]$,

$$\begin{aligned} 0 \leq j_r^\epsilon(t) & \leq 2 \|j_r\|_{L^\infty} < \infty, & r \in \mathcal{R}_{\text{slow}} \cup \mathcal{R}_{\text{damp}}, \\ 0 \leq \tilde{j}_r^\epsilon(t) & \leq 2 \|\tilde{j}_r\|_{L^\infty} < \infty, & r \in \mathcal{R}_{\text{fcyc}}, \\ 0 < \frac{1}{2} \inf_{t \in [0, T]} u_x(t) & \leq u_x^\epsilon(t) \leq 2 \|u_x\|_{L^\infty}, & x \in \mathcal{V}_{0\text{slow}} \cup \mathcal{V}_{0\text{fcyc}} \cup \mathfrak{C} \cup \mathcal{V}_1. \end{aligned}$$

The convergence of the integrals for $r \in \mathcal{R}_{\text{slow}}$ and $r \in \mathcal{R}_{\text{damp}}$ then follows by dominated convergence:

$$\begin{aligned} \int_{[0, T]} s(j_r^\epsilon(t) \mid \kappa_r \pi_{r_-}^\epsilon u_{r_-}^\epsilon(t)) dt & \rightarrow \int_{[0, T]} s(j_r(t) \mid \kappa_r \pi_{r_-} u_{r_-}(t)) dt, & r \in \mathcal{R}_{\text{slow}}, \\ \int_{[0, T]} s(j_r^\epsilon(t) \mid \frac{1}{\epsilon} \kappa_r \pi_{r_-}^\epsilon u_{r_-}^\epsilon(t)) dt & \rightarrow \int_{[0, T]} s(j_r(t) \mid \kappa_r \tilde{\pi}_{r_-} u_{r_-}(t)) dt, & r \in \mathcal{R}_{\text{damp}}. \end{aligned}$$

Similarly for $r \in \mathcal{R}_{\text{fyc}}$, by dominated convergence,

$$\int_{[0,T]} s \left(\frac{1}{\epsilon} \kappa_r \pi_{r-}^\epsilon u_{r-}^\epsilon(t) + \frac{1}{\sqrt{\epsilon}} \tilde{j}_r^\epsilon(t) \mid \frac{1}{\epsilon} \kappa_r \pi_{r-}^\epsilon u_{r-}^\epsilon(t) \right) dt$$

$$\stackrel{(2.2)}{\leq} \int_{[0,T]} \frac{\tilde{j}_r^\epsilon(t)^2}{\kappa_r \pi_{r-}^\epsilon u_{r-}^\epsilon(t)} dt \rightarrow \int_{[0,T]} \frac{\tilde{j}_r(t)^2}{\kappa_r \pi_{r-}^\epsilon u_{r-}^\epsilon(t)} dt.$$

The inequality in the other direction follows from Lemma 4.4. □

5 Spikes and Damped Cycles

As explained in Sect. 1.8, the uniform L^1 -bounds on the damped fluxes $j_{\mathcal{R}_{\text{damp}}}^\epsilon$ and small concentrations $u_{\mathcal{V}_1}^\epsilon$, can not prevent limits from becoming measure-valued in time, that is, both may develop atomic or Cantor parts. The question when these spikes in damped fluxes may occur is answered in our Theorem 1.3; this section is devoted to the proof of both statements in that theorem. The first part of Theorem 1.3 rules out spikes for damped fluxes that are not chained in a cycle. The second part shows that spikes may occur in damped flux cycles.

Recall the subdivision $\mathcal{R}_{\text{damp}} = \mathcal{R}_{\text{dcyc}} \cup \mathcal{R}_{\text{dnocyc}}$ from Sect. 1.8

5.1 No Spikes in Damped Fluxes Outside Cycles

Proof of Theorem 1.3(i) Take an arbitrary $r^0 \in \mathcal{R}_{\text{dnocyc}}$ coming out of node $r_-^0 =: x \in \mathcal{V}_1$. By Lemma 3.3, all fluxes flowing out of node x are damped, and all fluxes flowing into node x are either slow or damped. The weak formulation of the continuity equation in x now reads:

$$\int_{[0,T]} \phi(t) \left[\sum_{\substack{r^1 \in \mathcal{R}_{\text{damp}}: \\ r_+^1 = x}} j_{r^1}^\epsilon(dt) - \sum_{\substack{r^1 \in \mathcal{R}_{\text{slow}}: \\ r_+^1 = x}} j_{r^1}^\epsilon(t) dt - \sum_{\substack{r^1 \in \mathcal{R}_{\text{damp}}: \\ r_+^1 = x}} j_{r^1}^\epsilon(dt) \right]$$

$$= - \int_{[0,T]} \dot{\phi}(t) \pi_x^\epsilon u_x^\epsilon(t) dt,$$

for all $\phi \in C_0^1([0, T])$. By the uniform L^1 -bounds on u^ϵ and the vanishing π_x^ϵ , the right-hand side above converges to zero, and so (for arbitrary measurable sets $dt \subset [0, T]$):

$$0 \leq j_{r^0}(dt) \leq \sum_{\substack{r^1 \in \mathcal{R}_{\text{damp}}: \\ r_+^1 = r_-^0 (=x)}} j_{r^1}(dt) = \sum_{\substack{r^1 \in \mathcal{R}_{\text{slow}}: \\ r_+^1 = r_-^0}} j_{r^1}(t) dt + \sum_{\substack{r^1 \in \mathcal{R}_{\text{damp}}: \\ r_+^1 = r_-^0}} j_{r^1}(dt).$$

Applying the same inequality for each damped flux $r_+^1 = r_-^0$, we get:

$$0 \leq j_{r^0}(dt) \leq \sum_{\substack{r^1 \in \mathcal{R}_{\text{slow}}: \\ r_+^1 = r_-^0}} j_{r^1}(t) dt + \sum_{\substack{r^1 \in \mathcal{R}_{\text{damp}}: \\ r_+^1 = r_-^0}} \left[\sum_{\substack{r^2 \in \mathcal{R}_{\text{slow}}: \\ r_+^2 = r_-^1}} j_{r^2}(t) dt + \sum_{\substack{r^2 \in \mathcal{R}_{\text{damp}}: \\ r_+^2 = r_-^1}} j_{r^2}(dt) \right].$$

We now apply this procedure recursively until the right-hand side contains slow fluxes only. This is possible because by assumption any damped flux that already appeared during this procedure can not reappear in the inequality. Exploiting that eventually the right-hand side is a sum over slow fluxes that are in $L^\mathcal{C}(0, T)$, by the Radon-Nikodym Lemma the left-hand j_{r^0} also has a $L^\mathcal{C}$ -density. □

5.2 Finite-Cost Spikes in Damped Flux Cycles

We now prove that fluxes in $\mathcal{R}_{\text{dcyc}}$ may actually develop singularities.

Proof of Theorem 1.3(ii) If $\mathcal{R}_{\text{dcyc}} \neq \emptyset$ then there exists a disconnected damped component $\mathfrak{d} \subset \mathcal{V}_1$ such that $\forall x, y \in \mathfrak{d} \exists (r^k)_{k=1}^K \subset \mathcal{R}_{\text{dcyc}}, r_-^1 = x, r_+^k = r_-^{k+1}$ for $k = 1, \dots, K - 1, r_+^K = y$ (cf. Sect. 1.5). By irreducibility and mass conservation there exists at least one $r^{\text{in}} \in \mathcal{R}_{\text{slow}} \cup \mathcal{R}_{\text{dnocyc}}$ with $r_+^{\text{in}} \in \mathfrak{d}$ and at least one $r^{\text{out}} \in \mathcal{R}_{\text{dnocyc}}$ with $r_-^{\text{out}} \in \mathfrak{d}$. We first assume:

1. that all edges in \mathfrak{d} are chained in a cycle, i.e. $\mathfrak{d} := (x^k)_{k=1}^K, \mathcal{R}_{\text{dcyc}} \cap \{r_- \in \mathfrak{d}\} = (r^k)_{k=1}^K$ with $r_-^k = x^k$ for $k = 1, \dots, K, r_+^k = x^{k+1}$ for $k = 1, \dots, K - 1$ and $r_+^K = x^1,$
2. that $r_+^{\text{in}} = x^1$ and $r_-^{\text{out}} = x^l,$ and
3. that $x^0 := r_-^{\text{in}}$ and $x^{K+1} := r_+^{\text{out}}$ both lie in $\mathcal{V}_0,$ see Fig. 4.

Initially we concentrate all mass in $x^0,$ i.e. $u_{x^0}^\epsilon(0) := 1/\pi_{x^0}^\epsilon,$ and $u_x^\epsilon(0) = 0$ for all other nodes $x \in \mathcal{V}.$ The rate functional of the initial condition is indeed uniformly bounded:

$$\tilde{\mathcal{I}}_0^\epsilon(u^\epsilon(0)) = s(1 \mid \underbrace{\pi_{x^0}^\epsilon}_{\mathcal{O}(1)}) + \underbrace{\sum_{x \neq x^0} \pi_x^\epsilon}_{\leq 1}.$$

Define the continuous piecewise affine triangle function:

$$\Delta_T^\epsilon(t) := \begin{cases} 0, & 0 \leq t \leq \frac{1}{2}T - \frac{1}{2}\sqrt{\epsilon}, \\ t - \frac{1}{2}T + \frac{1}{2}\sqrt{\epsilon}, & \frac{1}{2}T - \frac{1}{2}\sqrt{\epsilon} < t \leq \frac{1}{2}T, \\ \frac{1}{2}T + \frac{1}{2}\sqrt{\epsilon} - t, & \frac{1}{2}T < t \leq \frac{1}{2}T + \frac{1}{2}\sqrt{\epsilon}, \\ 0, & \frac{1}{2}T + \frac{1}{2}\sqrt{\epsilon} < t \leq T. \end{cases}$$

Note that $\int_0^T \Delta_T^\epsilon(t) dt = \epsilon/4.$ For the dynamics, we will first transport a little bit of mass from x^0 into each node of the cycle $\mathfrak{d},$ then develop a spike at $t = T/2,$ and then release all mass from the cycle through $r^{\text{out}}.$

$$\begin{aligned} j_{r^{\text{in}}}^\epsilon(t) &:= K \mathbb{1}_{(T/2 - \sqrt{\epsilon}/2, T/2)}(t), \\ j_{r^k}^\epsilon(t) &:= a_k \mathbb{1}_{(T/2 - \sqrt{\epsilon}/2, T/2)}(t) + \frac{1}{\epsilon} \Delta_T^\epsilon(t) + b_k \mathbb{1}_{(T/2, T/2 + \sqrt{\epsilon}/2)}(t), \quad k = 1, \dots, K \\ j_{r^{\text{out}}}^\epsilon(t) &:= K \mathbb{1}_{(T/2, T/2 + \sqrt{\epsilon}/2)}(t), \end{aligned}$$

where $a_k := K - k$ and $b_k := k - l + K \mathbb{1}_{\{k < l\}}.$ We set all other fluxes $j_r^\epsilon, \tilde{j}_r^\epsilon$ in the network to 0. By construction, $j_r^\epsilon \xrightarrow{\text{narrow}} \frac{1}{4} \delta_{T/2},$ which is singular as was to be shown.

We now show that the functional $\tilde{\mathcal{J}}^\epsilon$ is uniformly bounded. To calculate the densities, note that the $\frac{1}{\epsilon} \Delta_T^\epsilon$ terms in $j_{r^k}^\epsilon$ are divergence free. The mild formulation of the continuity equation (3.5d) thus yields for all $k = 1, \dots, K$ and $t \in [0, T],$

$$\begin{aligned} \pi_{x^0}^\epsilon u_{x^0}^\epsilon(t) - 1 &= \begin{cases} 0, & 0 \leq t < T/2 - \sqrt{\epsilon}/2, \\ -K(t - \frac{1}{2}T + \frac{1}{2}\sqrt{\epsilon}), & T/2 - \sqrt{\epsilon}/2 < t \leq T/2, \\ -\frac{1}{2}K\sqrt{\epsilon}, & t \geq T/2, \end{cases} \\ \pi_{x^k}^\epsilon u_{x^k}^\epsilon(t) &= \mathbb{1}_{\{k=1\}} j_{r^{\text{in}}}^\epsilon[0, t] + j_{r^{k-1}}^\epsilon[0, t] - j_{r^k}^\epsilon[0, t] - \mathbb{1}_{\{k=l\}} j_{r^{\text{out}}}^\epsilon[0, t] = \Delta_T^\epsilon(t), \\ \pi_{x^{K+1}}^\epsilon u_{x^{K+1}}^\epsilon(t) &= \begin{cases} 0, & 0 \leq t < T/2 - \sqrt{\epsilon}/2, \\ K(t - \frac{1}{2}T + \frac{1}{2}\sqrt{\epsilon}), & T/2 - \sqrt{\epsilon}/2 < t \leq T/2, \\ \frac{1}{2}K\sqrt{\epsilon}, & t \geq T/2. \end{cases} \end{aligned}$$

The dynamic part of the rate functional is:

$$\begin{aligned}
 \tilde{J}^\epsilon(u_{\mathcal{V}_0}^\epsilon, u_{\mathcal{V}_1}^\epsilon, j_{\mathcal{R}_{\text{slow}}}^\epsilon, j_{\mathcal{R}_{\text{damp}}}^\epsilon, \tilde{j}_{\mathcal{R}_{\text{fcyc}}}^\epsilon) &= \underbrace{\int_{[0,T]} s(j_{r,\text{in}}^\epsilon(t) \mid \kappa_{r,\text{in}} \pi_{x^0}^\epsilon u_{x^0}^\epsilon(t)) dt}_{(I)} \\
 &+ \sum_{k=1}^K \underbrace{\int_{[0,T]} s(j_{r,k}^\epsilon(t) \mid \frac{1}{\epsilon} \kappa_{r,k} \pi_{x^k}^\epsilon u_{x^k}^\epsilon(t)) dt}_{(II)} \\
 &+ \underbrace{\int_{[0,T]} s(j_{r,\text{out}}^\epsilon(t) \mid \frac{1}{\epsilon} \kappa_{r,\text{out}} \pi_{x^l}^\epsilon u_{x^l}^\epsilon(t)) dt}_{(III)} \\
 &+ \sum_{r^{\text{in}} \neq r \in \mathcal{R}_{\text{slow}}: r_- = x^0} \underbrace{\int_{[0,T]} s(0 \mid \kappa_{r_-} \pi_{x^0}^\epsilon u_{x^0}^\epsilon(t)) dt}_{(IV)} \\
 &+ \sum_{r^{\text{out}} \neq r \in \mathcal{R}_{\text{slow}}: r_- = x^{K+1}} \underbrace{\int_{[0,T]} s(0 \mid \kappa_{r_-} \pi_{x^{K+1}}^\epsilon u_{x^{K+1}}^\epsilon(t)) dt}_{(V)}.
 \end{aligned}$$

By a long but simple calculation, these integrals can be calculated explicitly:

$$\begin{aligned}
 (I) &= \kappa_{r,\text{in}}(T - \sqrt{\epsilon}(\frac{1}{2} + \frac{1}{4}KT)) + \frac{1}{2}\sqrt{\epsilon}(K \log \frac{K}{\kappa_{r,\text{in}}} - K) + s(1 - \frac{1}{2}K\sqrt{\epsilon} \mid 1) + \frac{1}{8}\kappa_{r,\text{in}}K\epsilon, \\
 (II) &= \int_0^{\sqrt{\epsilon}/2} s(a_k + \frac{1}{\epsilon}t \mid \frac{1}{\epsilon}\kappa_{r,k}\pi_{x^k}^\epsilon t) dt + \int_0^{\sqrt{\epsilon}/2} s(b_k + \frac{1}{\epsilon}t \mid \frac{1}{\epsilon}\kappa_{r,k}\pi_{x^k}^\epsilon t) dt \\
 &= \frac{1}{2}\epsilon a_k^2 \log\left(\frac{\epsilon a_k + \frac{1}{2}\sqrt{\epsilon}}{\epsilon a_k}\right) + (\frac{1}{2}\sqrt{\epsilon}a_k + \frac{1}{8}) \log\left(\frac{2\sqrt{\epsilon}a_k + 1}{\kappa_{r,k}}\right) - \frac{1}{4}\sqrt{\epsilon}a_k + \frac{1}{8}\kappa_{r,k} - \frac{1}{8} \\
 &\quad + \frac{1}{2}\epsilon b_k^2 \log\left(\frac{\epsilon b_k + \frac{1}{2}\sqrt{\epsilon}}{\epsilon b_k}\right) + (\frac{1}{2}\sqrt{\epsilon}b_k + \frac{1}{8}) \log\left(\frac{2\sqrt{\epsilon}b_k + 1}{\kappa_{r,k}}\right) - \frac{1}{4}\sqrt{\epsilon}b_k + \frac{1}{8}\kappa_{r,k} - \frac{1}{8}, \\
 (III) &= \int_0^{\sqrt{\epsilon}/2} s(0 \mid \frac{1}{\epsilon}\kappa_{r,\text{out}}t) dt + \int_0^{\sqrt{\epsilon}/2} s(K \mid \frac{1}{\epsilon}\kappa_{r,\text{out}}t) dt = \frac{1}{2}K\sqrt{\epsilon} \log\left(\frac{2K\sqrt{\epsilon}}{\kappa_{r,\text{out}}}\right) + \frac{1}{4}\kappa_{r,\text{out}}, \\
 (IV) &= \kappa_{r_-} \underbrace{\int_{[0,T]} \pi_{x^0}^\epsilon u_{x^0}^\epsilon(t) dt}_{\leq 1} \quad \text{and} \quad (V) = \kappa_{r_-} \underbrace{\int_{[0,T]} \pi_{x^{K+1}}^\epsilon u_{x^{K+1}}^\epsilon(t) dt}_{\leq 1}.
 \end{aligned}$$

It thus follows that $\tilde{I}_0^\epsilon + \tilde{J}^\epsilon$ is uniformly bounded as claimed.

Recall the three assumptions we made in the beginning of the proof. The second assumption is just notational. The first assumption, that all edges in \mathfrak{d} are chained in a cycle, can easily be relaxed by fixing additional concentrations and damped fluxes to 0, which keeps the rate functional finite. The third assumption would be violated if there were a chain of damped fluxes between a \mathcal{V}_0 -node x^0 and x^1 or between a \mathcal{V}_0 -node x^l and x^{K+1} ; in that case we can again set these fluxes equal to $j_{r,\text{in}}^\epsilon$ and $j_{r,\text{out}}^\epsilon$ respectively, without having the rate functional blowing up, which relaxes the last assumption. \square

6 Implications for Large Deviations and the Effective Dynamics

We now prove two consequences: the Γ -convergence of the density large deviations, and the convergence of ϵ -level solutions to the solution of the effective dynamics.




Fig. 4 A diconnected component \mathfrak{d} of damped fluxes, chained in a cycle

6.1 Γ -Convergence of the Density Large Deviatons

As a consequence of our main Γ -convergence result, we obtain the Γ -convergence for the density large-deviation rate functional $\tilde{\mathcal{I}}_0^\epsilon + \tilde{\mathcal{I}}^\epsilon$ given by

$$\tilde{\mathcal{I}}^\epsilon(u_{\mathcal{V}_{0\text{slow}}}, u_{\mathcal{V}_{0\text{fycyc}}}, u_{\mathcal{C}}, u_{\mathcal{V}_1}) := \inf_{\substack{j_{\mathcal{R}_{\text{slow}}} \in L^\mathcal{C}([0, T]; \mathbb{R}^{\mathcal{R}_{\text{slow}}}), \\ j_{\mathcal{R}_{\text{damp}}} \in \mathcal{M}([0, T]; \mathbb{R}^{\mathcal{R}_{\text{damp}}}), \\ \tilde{j}_{\mathcal{R}_{\text{fycyc}}} \in L^\mathcal{C}([0, T]; \mathbb{R}^{\mathcal{R}_{\text{fycyc}}})}} \tilde{\mathcal{J}}^\epsilon(u_{\mathcal{V}_{0\text{slow}}}, u_{\mathcal{V}_{0\text{fycyc}}}, u_{\mathcal{C}}, u_{\mathcal{V}_1}, j_{\mathcal{R}_{\text{slow}}}, j_{\mathcal{R}_{\text{damp}}}, \tilde{j}_{\mathcal{R}_{\text{fycyc}}}).$$

Corollary 6.1 *In $C([0, T]; \mathbb{R}^{\mathcal{V}_{0\text{slow}}}) \times L^\infty([0, T]; \mathbb{R}^{\mathcal{V}_{0\text{fycyc}}}) \times C([0, T]; \mathbb{R}^{\mathcal{C}}) \times \mathcal{M}([0, T]; \mathbb{R}^{\mathcal{V}_1})$ (equipped with the uniform, weak-*, uniform, and narrow topologies),*

$$\tilde{\mathcal{I}}_0^\epsilon + \tilde{\mathcal{I}}^\epsilon \xrightarrow{\Gamma} \tilde{\mathcal{I}}_0^0 + \tilde{\mathcal{I}}^0,$$

where

$$\tilde{\mathcal{I}}^0(u_{\mathcal{V}_0}, u_{\mathcal{V}_1}) := \inf_{\substack{j_{\mathcal{R}_{\text{slow}}} \in L^\mathcal{C}([0, T]; \mathbb{R}^{\mathcal{R}_{\text{slow}}}), \\ j_{\mathcal{R}_{\text{damp}}} \in \mathcal{M}([0, T]; \mathbb{R}^{\mathcal{R}_{\text{damp}}}), \\ \tilde{j}_{\mathcal{R}_{\text{fycyc}}} \in L^\mathcal{C}([0, T]; \mathbb{R}^{\mathcal{R}_{\text{fycyc}}})}} \tilde{\mathcal{J}}^0(u_{\mathcal{V}_0}, u_{\mathcal{V}_1}, j_{\mathcal{R}_{\text{slow}}}, j_{\mathcal{R}_{\text{damp}}}, \tilde{j}_{\mathcal{R}_{\text{fycyc}}}).$$

Proof The proof is more-or-less classic but we include it here for completeness. For brevity we write $u = (u_{\mathcal{V}_{0\text{slow}}}, u_{\mathcal{V}_{0\text{fycyc}}}, u_{\mathcal{C}}, u_{\mathcal{V}_1})$ and $j = (j_{\mathcal{R}_{\text{slow}}}, j_{\mathcal{R}_{\text{damp}}}, \tilde{j}_{\mathcal{R}_{\text{fycyc}}})$.

To prove the Γ -lower bound, take an arbitrary convergent sequence $u_{\mathcal{V}_0}^\epsilon \rightarrow u_{\mathcal{V}_0}$, $u_{\mathcal{V}_1}^\epsilon \xrightarrow{\text{narrow}} u_{\mathcal{V}_1}$, and choose a corresponding sequence j^ϵ that satisfies for each $\epsilon > 0$ the inequality

$$\tilde{\mathcal{J}}^\epsilon(u^\epsilon, j^\epsilon) \leq \inf_j \tilde{\mathcal{J}}^\epsilon(u^\epsilon, j) + \epsilon.$$

Without loss of generality we assume that $\sup_{\epsilon > 0} \tilde{\mathcal{I}}_0^\epsilon(u^\epsilon(0)) + \tilde{\mathcal{J}}^\epsilon(u^\epsilon, j^\epsilon) < \infty$. Hence by Corollary 3.10 there exists a subsequence (u^ϵ, j^ϵ) (without changing notation) that converges in the sense of (3.12) to a limit (u, j) . From the Γ -lower bound Lemmas 4.2, 4.3 and 4.4, we find that:

$$\liminf_{\epsilon \rightarrow 0} \tilde{\mathcal{I}}_0^\epsilon(u^\epsilon(0)) \geq \tilde{\mathcal{I}}_0^0(u(0)), \quad \text{and}$$

$$\liminf_{\epsilon \rightarrow 0} \inf_j \tilde{\mathcal{J}}^\epsilon(u^\epsilon, j) \geq \liminf_{\epsilon \rightarrow 0} \tilde{\mathcal{J}}^\epsilon(u^\epsilon, j^\epsilon) - \epsilon \geq \tilde{\mathcal{J}}^0(u, j) \geq \inf_j \tilde{\mathcal{J}}^0(u, j).$$

This proves the lower bound

$$\liminf_{\epsilon \rightarrow 0} \tilde{I}_0^\epsilon(u^\epsilon(0)) + \tilde{I}^\epsilon(u^\epsilon) \geq \tilde{I}_0^0(u(0)) + \tilde{I}^0(u).$$

For the recovery property, take an arbitrary u with $\tilde{I}_0^0(u(0)) + \tilde{I}^0(u) < \infty$, and for an arbitrary $\delta > 0$, a flux j^δ such that

$$\tilde{J}^0(u, j^\delta) \leq \inf_j \tilde{J}^0(u, j) + \delta = \tilde{I}^0(u) + \delta.$$

Proposition 4.5 provides a recovery sequence (u^ϵ, j^ϵ) for (u, j^δ) and the sequences $(\tilde{I}_0^\epsilon)_\epsilon$ and $(\tilde{J}^\epsilon)_\epsilon$, hence:

$$\begin{aligned} \limsup_{\epsilon \rightarrow 0} \tilde{I}_0^\epsilon(u^\epsilon(0)) &\leq \tilde{I}_0^0(u(0)), & \text{and} \\ \limsup_{\epsilon \rightarrow 0} \tilde{I}^\epsilon(u^\epsilon) &\leq \limsup_{\epsilon \rightarrow 0} \tilde{J}^\epsilon(u^\epsilon, j^\epsilon) \leq \tilde{J}^0(u, j^\delta) \leq \tilde{I}^0(u) + \delta. \end{aligned}$$

Since $\delta > 0$ is arbitrary, the recovery property follows. □

Remark 6.2 By the same argument one may also contract further to obtain Γ -convergence of the functional

$$\begin{aligned} (u_{\mathcal{V}_{0\text{slow}}}, u_{\mathcal{C}}) &\mapsto \inf \tilde{I}_0^\epsilon(u_{\mathcal{V}_{0\text{slow}}}(0), u_{\mathcal{V}_{0\text{fcyc}}}(0), u_{\mathcal{C}}(0), u_{\mathcal{V}_1}(0)) \\ &+ \tilde{J}^\epsilon(u_{\mathcal{V}_{0\text{slow}}}, u_{\mathcal{V}_{0\text{fcyc}}}, u_{\mathcal{C}}, u_{\mathcal{V}_1}, j_{\mathcal{R}_{\text{slow}}}, j_{\mathcal{R}_{\text{damp}}}, \tilde{j}_{\mathcal{R}_{\text{fcyc}}}) \\ &\text{over all } u_{\mathcal{V}_{0\text{fcyc}}}, u_{\mathcal{V}_1}, j_{\mathcal{R}_{\text{slow}}}, j_{\mathcal{R}_{\text{damp}}}, \text{ and } \tilde{j}_{\mathcal{R}_{\text{fcyc}}}. \end{aligned}$$

This is the coarse-grained description for which Γ -convergence is studied in [33].

6.2 Convergence to the Effective Equations

For any pair (u, j) at which the limiting functional \tilde{J}^0 vanishes, the densities satisfy the following set of equations in the weak sense of (3.3):

$$\pi_x \dot{u}_x = \sum_{\substack{r \in \mathcal{R}_{\text{slow}}: \\ r_+ = x}} \kappa_r \pi_{r-} u_{r-} + \sum_{\substack{r \in \mathcal{R}_{\text{damp}}: \\ r_+ = x}} \kappa_r \tilde{\pi}_{r-} u_{r-} - \sum_{\substack{r \in \mathcal{R}_{\text{slow}}: \\ r_- = x}} \kappa_r \pi_x u_x \quad \text{for } x \in \mathcal{V}_{0\text{slow}}, \tag{6.1a}$$

$$\pi_{\mathcal{C}} \dot{u}_{\mathcal{C}} = \sum_{\substack{r \in \mathcal{R}_{\text{slow}}: \\ r_+ \in \mathcal{C}}} \kappa_r \pi_{r-} u_{r-} + \sum_{\substack{r \in \mathcal{R}_{\text{damp}}: \\ r_+ \in \mathcal{C}}} \kappa_r \tilde{\pi}_{r-} u_{r-} - \sum_{\substack{r \in \mathcal{R}_{\text{slow}}: \\ r_- \in \mathcal{C}}} \kappa_r \pi_{r-} u_{r-} \quad \text{for } \mathcal{C} \in \mathcal{C}, \tag{6.1b}$$

$$u_x = u_{\mathcal{C}} \quad \text{for any } x \in \mathcal{C} \in \mathcal{C}, \tag{6.1c}$$

$$0 = \sum_{\substack{r \in \mathcal{R}_{\text{slow}}: \\ r_+ = x}} \kappa_r \pi_{r-} u_{r-} + \sum_{\substack{r \in \mathcal{R}_{\text{damp}}: \\ r_+ = x}} \kappa_r \tilde{\pi}_{r-} u_{r-} - \sum_{\substack{r \in \mathcal{R}_{\text{damp}}: \\ r_- = x}} \kappa_r \tilde{\pi}_x u_x \quad \text{for } x \in \mathcal{V}_1. \tag{6.1d}$$

We first prove existence and uniqueness for these equations.

Lemma 6.3 Fix an initial condition $u(0) \in \mathbb{R}^{\mathcal{V}}$ that is well-prepared, which means that

1. Whenever x, y are in the same connected component $\mathcal{C} \in \mathcal{C}$, we have $u_x = u_y$; we denote the common value by $u_{\mathcal{C}}$;

2. $u(0)$ satisfies the condition (6.1d).

Then the system of Eq. (6.1) has a unique solution $u \in C^\infty([0, \infty); \mathbb{R}^{\mathcal{V}})$ with initial value $u(0)$.

Proof Since $\dot{u}_x = \dot{u}_c$ whenever $x \in c \in \mathcal{C}$, Eq. (6.1b) can be rewritten as

$$\pi_c \dot{u}_x = \sum_{\substack{r \in \mathcal{R}_{\text{slow}}: \\ r_+ \in c}} \kappa_r \pi_r u_{r_-} + \sum_{\substack{r \in \mathcal{R}_{\text{damp}}: \\ r_+ \in c}} \kappa_r \tilde{\pi}_r u_{r_-} - \sum_{\substack{r \in \mathcal{R}_{\text{slow}}: \\ r_- \in c}} \kappa_r \pi_r u_{r_-} \quad \text{for all } x \in c \in \mathcal{C}, \tag{6.2}$$

The right-hand side does not depend on the choice of x within the same $c \in \mathcal{C}$; therefore, under the assumption that $u_x(0) = u_c(0)$ for all $x \in c \in \mathcal{C}$, the system (6.1) is equivalent to the set of Eqs. (6.1a)–(6.2).

This implies that the system (6.1) can be written as a differential-algebraic equation:

$$\dot{u}_{\mathcal{V}_0} = A_{\mathcal{V}_0 \rightarrow \mathcal{V}_0} u_{\mathcal{V}_0} + A_{\mathcal{V}_1 \rightarrow \mathcal{V}_0} u_{\mathcal{V}_1}, \tag{6.3a}$$

$$0 = A_{\mathcal{V}_0 \rightarrow \mathcal{V}_1} u_{\mathcal{V}_0} + A_{\mathcal{V}_1 \rightarrow \mathcal{V}_1} u_{\mathcal{V}_1}, \tag{6.3b}$$

where for $x \in \mathcal{V}_1$,

$$(A_{\mathcal{V}_1 \rightarrow \mathcal{V}_1} u_{\mathcal{V}_1})_x := \sum_{\substack{r \in \mathcal{R}_{\text{damp}} \\ r_+ = x}} \kappa_r \frac{\tilde{\pi}_r}{\tilde{\pi}_x} u_{r_-} - u_x \sum_{\substack{r \in \mathcal{R}_{\text{damp}} \\ r_- = x}} \kappa_r.$$

By Lemma 6.4 below the matrix $A_{\mathcal{V}_1 \rightarrow \mathcal{V}_1}$ is invertible, and therefore (6.3) can be cast in the form of a linear ordinary differential equation for $u_{\mathcal{V}_0}$. This equation has unique solutions with C^∞ regularity, and by transforming back we find that $u_{\mathcal{V}_1}$ has the same regularity as $u_{\mathcal{V}_0}$. □

Lemma 6.4 *Under the conditions of the previous lemma, the matrix $A_{\mathcal{V}_1 \rightarrow \mathcal{V}_1}$ is invertible.*

Proof We first note that the matrix $A_{\mathcal{V}_1 \rightarrow \mathcal{V}_1}$ can be written as

$$A_{\mathcal{V}_1 \rightarrow \mathcal{V}_1} = \text{diag}(\tilde{\pi})^{-1} (A^{\text{int}} - \text{diag}(E)) \text{diag}(\tilde{\pi}),$$

with for $x, y \in \mathcal{V}_1$,

$$A_{xy}^{\text{int}} = \kappa_{y \rightarrow x} - \delta_{xy} \sum_{y' \in \mathcal{V}_1} \kappa_{x \rightarrow y'}, \quad E_x = \sum_{y' \in \mathcal{V}_0} \kappa_{x \rightarrow y'}.$$

Since $\text{diag}(\tilde{\pi})$ is invertible, it is sufficient to show that $A^{\text{int}} - \text{diag}(E)$ is invertible.

To do this we construct a new graph $\hat{\mathcal{G}} := (\mathcal{V}_1 \cup \{\mathbf{o}\}, \mathcal{R}_{\text{int}} \cup \mathcal{R}_{\mathbf{o}})$, consisting of the nodes of \mathcal{V}_1 and a single ‘graveyard’ node \mathbf{o} ; the graveyard collects all elements of \mathcal{V}_0 into one new node. The graph $\hat{\mathcal{G}}$ has edges

$$\mathcal{R}_{\text{int}} := \left\{ (x \rightarrow y) \in \mathcal{V}_1 \times \mathcal{V}_1 \setminus \{z \rightarrow z\} : \exists r \in \mathcal{R}_{\text{damp}} \text{ such that } r_- = x \text{ and } r_+ = y \right\}$$

$$\mathcal{R}_{\mathbf{o}} := \left\{ (x \rightarrow \mathbf{o}) : x \in \mathcal{V}_1, \exists r \in \mathcal{R}_{\text{damp}} \text{ such that } r_- = x \text{ and } r_+ \in \mathcal{V}_0 \right\}.$$

Note that there are no fluxes out of \mathbf{o} .

We define a new Markov jump process $Z(t)$ on this graph $\tilde{\mathcal{G}}$, by specifying jump rates $\hat{\kappa}_{x \rightarrow y}$ for each edge $(x \rightarrow y)$ in $\tilde{\mathcal{G}}$:

$$\hat{\kappa}_{x \rightarrow y} := \begin{cases} \sum_{r \in \mathcal{R}_{\text{damp}}: r_- = x, r_+ = y} \kappa_r, & \text{for } (x \rightarrow y) \in \mathcal{R}_{\text{int}}, \\ \sum_{r' \in \mathcal{R}_{\text{damp}}: r'_- = x, r'_+ \in \mathcal{V}_0} \kappa_{r'}, & \text{for } (x \rightarrow y) \in \mathcal{R}_\bullet. \end{cases}$$

The generator for this jump process is the matrix L given by

$$L_{xy} := \begin{cases} \hat{\kappa}_{x \rightarrow y} & \text{if } (x \rightarrow y) \in \mathcal{R}_{\text{int}} \cup \mathcal{R}_\bullet \text{ (which implies } x \neq y) \\ -\sum_{y' \in \mathcal{V}_1 \cup \{\mathbf{o}\}} \hat{\kappa}_{x \rightarrow y'} & \text{if } x = y \in \mathcal{V}_1 \\ 0 & \text{otherwise.} \end{cases}$$

By construction the transpose L^\top of this generator has the following structure in terms of the splitting $\mathcal{V}_1 \cup \{\mathbf{o}\}$:

$$L^\top = \left(\begin{array}{c|c} A^{\text{int}} - \text{diag}(E) & 0 \\ \hline E^\top & 0 \end{array} \right),$$

Since the original graph \mathcal{G} is disconnected, there exists for each $x \in \mathcal{V}_1$ a path $x = x_0 \rightarrow x_1 \rightarrow \dots \rightarrow x_k$ in \mathcal{G} leading to some $x_k \in \mathcal{V}_0$; without loss of generality we assume that $x_0, x_1, \dots, x_{k-1} \in \mathcal{V}_1$. All fluxes out of nodes in \mathcal{V}_1 are damped, and so the fluxes $(x_0 \rightarrow x_1), \dots, (x_{k-1} \rightarrow x_k)$ are all in $\mathcal{R}_{\text{damp}}$. Because these fluxes also exist as fluxes \mathcal{R}_{int} in the graph $\tilde{\mathcal{G}}$, the path $x_0 \rightarrow x_1 \rightarrow \dots \rightarrow x_{k-1}$ also is a path in $\tilde{\mathcal{G}}$. By construction, \mathcal{R}_\bullet contains an edge $r = (x_{k-1} \rightarrow \mathbf{o})$ with positive rate $\hat{\kappa}_r$.

It follows that if the process $Z(t)$ starts at any $x \in \mathcal{V}_1$, then at each positive time $t > 0$ there is a positive probability that $Z(t) = \mathbf{o}$. Since the graveyard \mathbf{o} has no outgoing fluxes, the only invariant measure for the process $Z(t)$ is $\mathbf{1}_\bullet := (0, 0, \dots, 0, 1)$, and so the kernel of L^\top coincides with the span of $\mathbf{1}_\bullet$. Explicitly written, this means that for all vectors $v = (v_{\mathcal{V}_1}, v_\bullet)$:

$$(A^{\text{int}} - \text{diag}(E))v_{\mathcal{V}_1} + E^\top v_\bullet = 0 \implies (v_{\mathcal{V}_1}, v_\bullet) = \text{const} \times \mathbf{1}_\bullet \implies v_{\mathcal{V}_1} = 0.$$

Consequently the matrix $A^{\text{int}} - \text{diag}(E)$ is invertible. □

We finally derive convergence to the full effective equations.

Corollary 6.5 *For each $\epsilon > 0$ let $(u_{\mathcal{V}_0}^\epsilon, u_{\mathcal{V}_1}^\epsilon, j_{\mathcal{R}_{\text{slow}}}^\epsilon, j_{\mathcal{R}_{\text{damp}}}^\epsilon, \tilde{j}_{\mathcal{R}_{\text{fcyc}}}^\epsilon)$ in Θ solve the system of equations:*

$$\begin{cases} j_r^\epsilon(t) = \kappa_r \pi_r^\epsilon u_{r_-}^\epsilon(t), & r \in \mathcal{R}_{\text{slow}}, \\ j_r^\epsilon(t) = \frac{1}{\epsilon} \kappa_r \pi_r^\epsilon u_{r_-}^\epsilon(t), & r \in \mathcal{R}_{\text{damp}}, \\ \tilde{j}_r^\epsilon(t) = 0, & r \in \mathcal{R}_{\text{fcyc}}, \\ \pi^\epsilon \dot{u}^\epsilon(t) = -\text{div } j^\epsilon(t), & \text{in the weak sense of (3.5),} \\ u^\epsilon(0) = u^{\epsilon,0}, \end{cases}$$

where $u^{\epsilon,0}$ is given. Assume that $u_{\mathcal{V}_{\text{slow}}}^{\epsilon,0}, u_{\mathcal{C}}^{\epsilon,0}$ converge to some $u_{\mathcal{V}_{\text{slow}}}^{0,0}, u_{\mathcal{C}}^{0,0} > 0$, that $u^{0,0}$ is well-prepared in the sense of Lemma 6.3. In addition, assume that for each $x \in \mathcal{V}_1$, $\log u_x^{\epsilon,0}$ remains bounded. Then $(u_{\mathcal{V}_0}^\epsilon, u_{\mathcal{V}_1}^\epsilon, j_{\mathcal{R}_{\text{slow}}}^\epsilon, j_{\mathcal{R}_{\text{damp}}}^\epsilon, \tilde{j}_{\mathcal{R}_{\text{fcyc}}}^\epsilon)$ converges in Θ

to $(u_{\mathcal{V}_0}, u_{\mathcal{V}_1}, j_{\mathcal{R}_{slow}}, j_{\mathcal{R}_{damp}}, \tilde{j}_{\mathcal{R}_{ficyc}})$, which is the unique solution to

$$\begin{cases} j_r(t) = \kappa_r \pi_r u_{r-}(t), & r \in \mathcal{R}_{slow}, \\ j_r(t) = \kappa_r \tilde{\pi}_r u_{r-}(t), & r \in \mathcal{R}_{damp}, \\ \tilde{j}_r(t) = 0, & r \in \mathcal{R}_{ficyc}, \\ \pi \dot{u}(t) = -\operatorname{div} j(t), & \text{in the weak sense of (3.14),} \\ u(0) = u^{0,0}. \end{cases} \tag{6.4}$$

Proof Set:

$$\begin{aligned} \tilde{\mathcal{I}}_0^\epsilon(u(0)) &:= \sum_{x \in \mathcal{V}} s(\pi_x^\epsilon u_x(0) \mid \pi_x^\epsilon u_x^{\epsilon,0}) \text{ and } F^\epsilon(u(0)) \\ &:= \sum_{x \in \mathcal{V}} \pi_x^\epsilon \left(-u_x(0) \log u_x^{\epsilon,0} - 1 + u_x^{\epsilon,0} \right). \end{aligned}$$

Then for each $\epsilon > 0$, the solution $(u_{\mathcal{V}_0}^\epsilon, u_{\mathcal{V}_1}^\epsilon, j_{\mathcal{R}_{slow}}^\epsilon, j_{\mathcal{R}_{damp}}^\epsilon, \tilde{j}_{\mathcal{R}_{ficyc}}^\epsilon)$ minimises the modified functional $\tilde{\mathcal{I}}_0^\epsilon + \tilde{\mathcal{J}}^\epsilon = \tilde{\mathcal{I}}_0^\epsilon + F^\epsilon + \tilde{\mathcal{J}}^\epsilon : \Theta \rightarrow [0, \infty]$ at value zero. In particular this means that:

$$\sup_{\epsilon > 0} \tilde{\mathcal{I}}_0^\epsilon(u^\epsilon(0)) + \tilde{\mathcal{J}}^\epsilon(u^\epsilon, j^\epsilon) = \sup_{\epsilon > 0} -F^\epsilon(u^\epsilon(0)) = \sup_{\epsilon > 0} \sum_{x \in \mathcal{V}} \pi_x^\epsilon (u_x^{\epsilon,0} \log u_x^{\epsilon,0} + 1 - u_x^{\epsilon,0}) < \infty.$$

By Corollary 3.10, the sequence (u^ϵ, j^ϵ) has a subsequence that converges in the sense of (3.12) to a limit (u, j) . By the assumptions on $u^{\epsilon,0}$, the functional F^ϵ converges along the sequence (u^ϵ, j^ϵ) to the limit F^0 , where

$$F^0(v(0)) := \sum_{x \in \mathcal{V}_{0slow}} \pi_x \left(-v_x(0) \log u_x^{0,0} - 1 + u_x^{0,0} \right) + \sum_{c \in \mathcal{C}} \pi_c \left(-v_c(0) \log u_c^{0,0} - 1 + u_c^{0,0} \right).$$

With the Γ -lower bound of Proposition 4.1 it follows that

$$\begin{aligned} 0 &= \liminf_{\epsilon \rightarrow 0} \tilde{\mathcal{I}}_0^\epsilon(u^\epsilon(0)) + F^\epsilon(u^\epsilon(0)) + \tilde{\mathcal{J}}^\epsilon(u^\epsilon, j^\epsilon) \\ &\geq \tilde{\mathcal{I}}_0^0(u(0)) + F^0(u(0)) + \tilde{\mathcal{J}}^0(u, j) = \tilde{\mathcal{I}}_0^0(u(0)) + \tilde{\mathcal{J}}^0(u, j). \end{aligned}$$

Here

$$\tilde{\mathcal{I}}_0^0(v(0)) := \sum_{x \in \mathcal{V}_{0slow}} s(\pi_x u_x(0) \mid \pi_x u_x^{0,0}) + \sum_{c \in \mathcal{C}} s(\pi_c u_c(0) \mid \pi_c u_c^{0,0}).$$

It follows that the limit (u, j) is a solution of the problem $\tilde{\mathcal{I}}_0^0 + \tilde{\mathcal{J}}^0 = 0$, which coincides with (6.4). □

Acknowledgements This research has been funded by the Deutsche Forschungsgemeinschaft (DFG) through grant CRC 1114 “Scaling Cascades in Complex Systems”, Project C08. We thank Robert Patterson for the useful discussions.

Funding Open Access funding enabled and organized by Projekt Deal.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory

regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

A The Arzelà-Ascoli Theorem for Asymptotic Uniformly Equicontinuous Sequences

The classical Arzelà-Ascoli theorem asserts that a set of continuous functions on a compact set is precompact in the supremum norm if and only if it is uniformly bounded and uniformly equicontinuous. For countable sets such as sequences the uniform equicontinuity is equivalent to *asymptotic* uniform equicontinuity, and this observation leads to the alternative version below. This is mentioned in various places in the literature (e.g. [36, Rem. 2.3 (ii)] or [8, Ex. 5.27]) but since we could not find a clear statement we state and prove it here.

Theorem A.1 *Let $(f_n)_{n \geq 1}$ be a sequence of continuous real-valued functions on $[0, T]$ that satisfies*

1. $\sup_{n \geq 1} \|f_n\|_\infty < \infty$;
2. *There exists $\omega : [0, \infty) \rightarrow [0, \infty)$, non-decreasing, with $\lim_{\sigma \downarrow 0} \omega(\sigma) = 0$, such that,*

$$\limsup_{n \rightarrow \infty} \sup_{\substack{0 \leq t, s \leq T: \\ |t-s| < \sigma}} |f_n(t) - f_n(s)| \leq \omega(\sigma).$$

Then there exists a subsequence f_{n_k} that converges uniformly on $[0, T]$.

Proof We prove the result by showing that the sequence $(f_n)_n$ also is uniformly equicontinuous in the usual sense. Fix $\epsilon > 0$. Choose $N \geq 1$ and $\sigma_0 > 0$ such that

$$\forall n \geq N \quad \forall |t - s| < \sigma_0 : \quad |f_n(t) - f_n(s)| < \epsilon.$$

Next, choose $\sigma_1 > 0$ such that

$$\forall 1 \leq n < N \quad \forall |t - s| < \sigma_1 : \quad |f_n(t) - f_n(s)| < \epsilon.$$

Then for all $n \geq 1$ and $|t - s| < \sigma_0 \wedge \sigma_1$ we have $|f_n(t) - f_n(s)| < \epsilon$. This proves that $(f_n)_n$ is uniformly equicontinuous, and therefore the result follows from the classical Arzelà-Ascoli theorem. □

References

1. Agazzi, A., Dembo, A., Eckmann, J.-P.: Large deviations theory for Markov jump models of chemical reaction networks. *Annals of Applied Probability* **28**(3), 1821–1855 (2018)
2. Ambrosio, L., Gigli, N., Savaré, G.: Gradient flows in metric spaces and in the space of probability measures, 2nd edn. *Lectures in Mathematics*. ETH Zürich. Birkhauser, Basel, Switzerland (2008)
3. Bertini, L., De Sole, A., Gabrielli, D., Jona-Lasinio, G., Landim, C.: Macroscopic fluctuation theory. *Rev. Mod. Phys.* **87**(2) (2015)
4. Baiesi, M., Maes, C., Netočný, K.: Computation of current cumulants for small nonequilibrium systems. *Journal of Statistical Physics* **135**(1), 57–75 (2009)
5. Bothe, D.: Instantaneous limits of reversible chemical reactions in presence of macroscopic convection. *Journal of Differential Equations* **193**(1), 27–48 (2003)
6. Bonaschi, G.A., Peletier, M.A.: Quadratic and rate-independent limits for a large-deviations functional. *Continuum Mech. Thermodyn.* **28**(4), 1191–1219 (2016)

7. Bobkov, S.G., Tetali, P.: Modified logarithmic Sobolev inequalities in discrete settings. *Journal of Theoretical Probability* 19(2), 289–336 (2006)
8. Davidson, J.: *Stochastic Limit Theory: An Introduction for Econometricians*. OUP, Oxford, 1994
9. Duong, M. H., Laschos, V., Renger, M.: Wasserstein gradient flows from large deviations of many-particle limits. *ESAIM Control Optim. Calculus Var.* 19(4):1166–1188, (2013)
10. Disser, K., Liero, M., Zinsl, J.: Evolutionary Γ -convergence of gradient systems modeling slow and fast chemical reactions. *Nonlinearity* 31(8), 3689 (2018)
11. Dal Maso, G.: *Introduction to Γ -convergence*. Birkhäuser, Boston, MA, USA (1993)
12. Dembo, A., Zeitouni, O.: *Large deviations techniques and applications*. In: *Stochastic Modelling and Applied Probability*, vol. 38, 2nd edn. Springer, New York (1987)
13. Engel, K.-J., Nagel, R.: *A short course on operator semigroups*. Springer (2006)
14. Feng, S.: Large deviations for empirical process of mean-field interacting particle system with unbounded jumps. *The Annals of Probability* 22(4), 1679–2274 (1994)
15. Fathi, M., Shu, Y.: Curvature and transport inequalities for Markov chains in discrete spaces. *Bernoulli* 24(1), 672–698 (2018)
16. Goeke, A., Walcher, S., Zerz, E.: Quasi-steady state-intuition, perturbation theory and algorithmic algebra. In: *International Workshop on Computer Algebra in Scientific Computing*, Springer, pp. 135–151 (2015)
17. Hilder, B., Peletier, M.A., Sharma, U., Tse, O.: An inequality connecting entropy distance, Fisher information and large deviations. *Stochastic Processes and their Applications* 130(5), 2596–2638 (2020)
18. Heinrich, R., Schuster, S.: *The Regulation of Cellular Systems*. Chapman & Hall, New York (1996)
19. Kaiser, M., Jack, R.L., Zimmer, J.: Canonical structure and orthogonality of forces and currents in irreversible Markov chains. *Journal of Statistical Physics* 170(6), 1019–1050 (2018)
20. Kuehn, C.: *Multiple Time Scale Dynamics*. Springer, New York, NY, USA (2015)
21. Kurtz, T.G.: Solutions of ordinary differential equations as limits of pure jump processes. *Journal of Applied Probability* 7(1), 49–58 (1970)
22. Léonard, C.: Large deviations for long range interacting particle systems with jumps. *Annales de l’Institut Henri Poincaré, section B* 31(2), 289–323 (1995)
23. Liero, M., Mielke, A., Peletier, M.A., Renger, D.R.M.: On microscopic origins of generalized gradient structures. *Discrete Contin. Dyn. Syst. Ser. S* 10(1) (2017)
24. Maas, J.: Gradient flows of the entropy for finite Markov chains. *Functional analysis* 261(8), 2250–2292 (2011)
25. Maas, J.: *Entropic Ricci Curvature for Discrete Spaces*, Springer, Cham, pp. 159–174 (2017)
26. Mielke, A.: Geodesic convexity of the relative entropy in reversible Markov chains. *Calculus of Variations and Partial Differential Equations* 48(1), 1–31 (2013)
27. Mielke, A.: Deriving effective models for multiscale systems via evolutionary Γ -convergence. In: *Control of Self-Organizing Nonlinear Systems*, Springer, pp. 235–251 (2016)
28. Mielke, A.: *On Evolutionary Γ -Convergence for Gradient Systems*, vol. 3, Springer, Cham, pp. 187–249 (2016)
29. Mielke, A., Montefusco, A., Peletier, M.A.: Exploring families of energy-dissipation landscapes via tilting: three types of EDP convergence. *Continu. Mech. Thermodyn.* (2021)
30. Mielke, A., Patterson, R., Peletier, M.A., Renger, D.R.M.: Non-equilibrium thermodynamical principles for chemical reactions with mass-action kinetics. *SIAM Journal on Applied Mathematics* 77(4), 1562–1585 (2017)
31. Mielke, A., Peletier, M.A., Renger, D.R.M.: On the relation between gradient flows and the large-deviation principle, with applications to Markov chains and diffusion. *Potential Anal.* 41(4) (2014)
32. Mielke, A., Peletier, M.A., Stephan, A.: EDP-convergence for nonlinear fast-slow reaction systems with detailed balance. [arXiv: 2010.08458](https://arxiv.org/abs/2010.08458) (2020)
33. Mielke, A., Stephan, A.: Coarse-graining via EDP-convergence for linear fast-slow reaction systems. *Mathematical Models and Methods in Applied Sciences* 30(9), 1765–1807 (2020)
34. Onsager, L., Machlup, S.: Fluctuations and irreversible processes. *Phys. Rev.* 91(6), 1505–1512 (1953)
35. Onsager, L.: Reciprocal relations in irreversible processes I. *Phys. Rev.* 37(4), 405–426 (1931)
36. Pötscher, B.M., Prucha, I.R.: Generic uniform convergence and equicontinuity concepts for random functions: An exploration of the basic structure. *Journal of Econometrics* 60(1–2), 23–63 (1994)
37. Patterson, R.I.A., Renger, D.R.M.: Large deviations of jump process fluxes. *Mathematical Physics, Analysis and Geometry* 22(3), 21 (2019)
38. Renger, D.R.M.: Flux large deviations of independent and reacting particle systems, with implications for macroscopic fluctuation theory. *J. Stat. Phys.* 172(5) (2018)
39. Rao, M.M., Ren, Z.D.: *Theory of Orlicz Spaces*. U.S.A, Marcel Dekker, New York, N.Y. (1991)
40. Renger, D.R.M., Zimmer, J.: Orthogonality of fluxes in general nonlinear reaction networks. *Discrete & Continuous Dynamical Systems - S* 14(1), 205–217 (2021)

41. Stiefenhofer, M.: Quasi-steady-state approximation for chemical reaction networks. *Journal of Mathematical Biology* 36(6), 593–609 (1998)
42. Trefethen, L.N., Embree, M.: *Spectra And Pseudospectra: The Behavior Of Nonnormal Matrices And Operators*. Princeton University Press (2005)
43. Tikhonov, A.N.: Systems of differential equations containing small parameters in the derivatives. *Matematicheskii Sbornik* 31(73), 575–586 (1952).

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.