




Clinical Features of Gambling Disorder Patients with and Without Food Addiction: Gender-Related Considerations

Mikel Etxandi¹ · Isabel Baenas^{1,2} · Lucero Munguía^{1,3} · Gemma Mestre-Bach^{1,4} · Roser Granero^{2,5} · Mónica Gómez-Peña¹ · Laura Moragas¹ · Amparo del Pino-Gutiérrez^{2,6} · Ester Codina¹ · Bernat Mora-Maltas¹ · Eduardo Valenciano-Mendoza¹ · Marc N. Potenza^{7,8,9,10} · Ashley N. Gearhardt¹¹ · Fernando Fernández-Aranda^{1,2,3} · Susana Jiménez-Murcia^{1,2,3} 

Accepted: 3 September 2021 / Published online: 28 September 2021

© The Author(s), under exclusive licence to Springer Science+Business Media, LLC, part of Springer Nature 2021

Abstract

Although food addiction (FA) is a debated condition and it is not currently recognized as a formal diagnosis, it shares features with other addictions, such as gambling disorder (GD). However, the prevalence of FA in GD and the clinical correlates are incompletely understood, especially within women versus men. To investigate FA in patients presenting with GD. The sample included 867 patients diagnosed with GD (798 males and 69 females) attending a specialized behavioral addictions unit. FA was observed in 8.3% of GD patients (18.8% of women, 7.4% of men). More psychopathology and harm avoidance, greater body mass indices and less self-directedness and cooperativeness were associated with FA. In women, FA was associated with a longer GD duration. In men, FA was associated with earlier GD onset, greater GD and problematic alcohol use severities. Among patients with GD, FA was associated with more psychopathology and gambling patterns suggestive of more protracted or severe GD. Screening for and addressing FA condition in patients with GD may help optimize preventive and therapeutic approaches. Future studies should consider testing guidelines to improve healthy eating habits, increase physical exercise and better manage stress and other negative emotions in order to target FA in GD.

Keywords Addictive behaviors · Food addiction · Gambling disorder · Alcohol · Gender

✉ Susana Jiménez-Murcia
sjimenez@bellvitgehospital.cat

Extended author information available on the last page of the article

Introduction

Gambling disorder (GD) is considered a behavioral addiction and defined as a persistent and recurrent maladaptive pattern of gambling behavior associated with impaired functioning in personal, social, and occupational domains (APA, 2013; Fauth-Bühler et al., 2017; Potenza et al., 2019). GD frequently co-occurs with other psychiatric disorders including substance use, mood, anxiety and personality disorders (Grant & Chamberlain, 2015; Karlsson et al., 2019; Kim et al., 2018; Petry et al., 2005; Tackett et al., 2017). Although food addiction (FA) is not a formal diagnostic entity, a possible association between GD and FA has been suggested (Jiménez-Murcia et al., 2017). The prevalence of FA has been estimated at 7.8% in individuals with GD, and an association between body mass index (BMI) and FA symptomatology has been described for patients with GD (Granero, Jiménez-Murcia, et al., 2018).

At least two frameworks have been proposed when conceptualizing FA: addictive-like eating and substance-based addiction models (Fernandez-Aranda et al., 2018; Hebebranda et al., 2014; Schulte et al., 2017). This lack of consensus may in part contribute to FA not being considered as a formal mental disorder in the current editions of the Diagnostic and Statistical Manual (DSM-5) or the International Classification of Diseases (ICD-11) (APA, 2013; The World Health Organisation, 2019). The substance-based addiction framework purports that some foods, especially palatable ones with large amounts of processed sugars and fats, may promote both overeating and addictive-like behaviors by activating brain reward systems (Schulte et al., 2015a). Therefore, FA may show similarities with other addictions, such as substance use disorders (SUDs) (Gearhardt et al., 2011; Schulte et al., 2015b). Proposed similarities include a preoccupation with obtaining the desired substance, the development of tolerance, abstinence, excessive use, difficulties reducing consumption despite negative physical and psychological consequences and similar neural processes (Carter & Davis, 2010; Gearhardt et al., 2011; Volkow et al., 2011; Fletcher & Kenny, 2018; Gordon et al., 2018; Pursey et al., 2019).

Given GD's classification as a behavioral addiction, common features between FA and GD may exist (Jiménez-Murcia et al., 2017). Both are associated with difficulties in controlling behavior (Hardy et al., 2018; Saunders et al., 2017), and impaired executive functions (Mestre-Bach, Fernández-Aranda, et al., 2020a; Steward et al., 2018). As with GD, psychopathology has been associated with FA (Burrows et al., 2018; Jiménez-Murcia et al., 2019). As well, GD and FA may involve maladaptive emotional regulation, engaging in gambling or eating for negative reinforcement motivations; i.e., to alleviate negative emotions (Innamorati et al., 2017; Mestre-Bach, Fernández-Aranda, et al., 2020b). Finally, impulsivity is another common feature between both disorders (Kandeger et al., 2019; Mestre-Bach, Steward et al., 2020), that has been highly associated with the development and maintenance of behavioral addictions (Lee et al., 2019), considering the tendency to respond with little forethought with the maladaptive behaviors, in spite of the negative later repercussions (Grant & Chamberlain, 2020). Same characteristics has been found in binge-related impulsive behaviors presents in Bulimia Nervosa and Binge eating disorders (Treasure et al., 2020), that has also been related with FA (Jiménez-Murcia, Agüera, Paslakis & Munguia, 2019; Penzenstadler et al., 2018).

Differences between GD and FA have also been described. For example, GD is more frequently observed among men, although the gender gap appears to be narrowing (Abbott et al., 2018; Subramaniam et al., 2015). Gender-related differences in types of gambling, age of onset and other features have been reported in GD (Zakiniæiz et al., 2017). In

contrast, FA is more frequently observed among women (Purseley et al., 2014). Women as compared with men are also more likely tendencies to engage in addictive behaviors and eating for negative reinforcement motivations that include coping with stress, depression and anxiety (Zakiniæiz & Potenza, 2018). Thus, there is a need to consider gender in understanding how FA may relate to clinical features in GD patients.

Although these two clinical entities have begun to be studied together, gender, personality traits and psychopathology have been scarcely investigated in GD individuals with and without FA. In the study from Jimenez-Murcia et al. (2017), the comorbidity between FA and treatment-seeking patients with GD was related to poorer emotional and psychological states. Being female and younger were two condition more associated with the co-occurrence of FA and GD. Regarding personality traits, high scores in harm avoidance and self-transcendence, and low scores in cooperativeness were associated with a higher risk of presenting FA among subjects with GD. However, the authors did not find significant differences in age of onset, duration of gambling problems or drug use among others, between participants with GD and those with comorbid FA.

Therefore, the present study aimed to assess the clinical correlates of FA among GD patients, considering personality features, psychological state or substance consumption, examined in men and women separately, anticipating potential gender-related differences as the literature points out. Our main hypothesis is that the presence of FA among GD would be associated with worse clinical profiles, greater psychopathology and more severe GD. We also anticipated more frequent FA among women (versus men) with GD.

Methods

Participants

The study sample consisted of 867 GD patients including 798 men and 69 women. Participants were referred for treatment to the Pathological Gambling Unit in the Psychiatry Department at a General University Hospital, in Spain, between May of 2016 and June of 2020. This hospital is part of the Spanish Public Assistance Network and of a Public University Campus, having its own research institute. It is the local reference center for a population of around 343,000 inhabitants and the territorial tertiary reference center for more than 2 million people. Participants were diagnosed according to the DSM-5 (APA, 2013) criteria for GD by clinical psychologists and psychiatrists with more than 20 years of experience in the field of behavioral addictions and eating disorders. None met DSM-5 diagnostic criteria for any current eating disorder, being a comorbid diagnostic of ED an exclusion criteria for the present study.

Measures

The South Oaks Gambling Screen (SOGS) (Lesieur & Blume, 1987) is a 20-item screen for problem gambling behaviors and consequences during the prior twelve months. The total score reflects problem gambling severity, with a score of 4 or more indicating problem gambling. The Spanish validation of the scale achieved very good psychometric results in the adaptation study (test-retest reliability $R=0.98$, internal consistency $\alpha=0.94$ and convergent validity $R=0.92$) (Echeburúa et al., 1994). The internal consistency for this scale in the study sample was adequate ($\alpha=0.74$).

The Diagnostic Questionnaire for Pathological Gambling (Stinchfield, 2003) is a self-report questionnaire with 19 items coded in a binary fashion (yes-no) that permits assessing DSM-IV (American Psychiatric Association, 2010) and DSM-5 (APA, 2013) diagnostic criteria for GD. It was used in the present study for assessing GD. Based on the DSM-5 taxonomy, several GD-related measures may be generated: the presence/absence of each DSM inclusion criterion, the presence/absence of GD diagnosis, a dimensional measure of problem gambling severity (total number of DSM criteria, obtained as the sum of the individual criteria), and GD severity grouped in four levels [non-problem gambling (0 criteria), problem gambling (for 1–3 criteria), mild GD (4–5 criteria), moderate GD (6–7 criteria) and severe GD (8–9 criteria)]. The Spanish adaptation of the questionnaire obtained satisfactory psychometric properties: internal consistency with a Cronbach's alpha equal to 0.95 for the combined sample, satisfactory convergent validity (moderate to large correlations with other measures of problem gambling severity), and high discriminative capacity (sensitivity = 0.92 and specificity = 0.99) (Jiménez-Murcia et al., 2009). The internal consistency for this scale in the study sample was good ($\alpha = 0.80$).

The Symptom Checklist-90-Revised (SCL-90-R) (Derogatis, 1994) is a 90-item self-report questionnaire measured on an ordinal 3-point scale that evaluates a broad range of psychological problems and symptoms of psychopathology by measuring nine primary symptom dimensions: Somatization, Obsession-Compulsion, Interpersonal Sensitivity, Depression, Anxiety, Hostility, Phobic Anxiety, Paranoid Ideation, and Psychoticism. It includes three global ratings, named global severity index (overall psychological distress), positive symptom distress index (the intensity of symptoms), and positive symptom total (self-reported symptoms). The global severity index can be used as a summary of the test. The validation of the scale in a Spanish population (Derogatis, 2002) obtained a mean internal consistency of 0.75 (coefficient alpha). The internal consistency in the study was excellent for the global ratings ($\alpha = 0.98$) and the one for the subscales range from adequate to excellent ($\alpha = 0.80$ for Paranoid Ideation to $\alpha = 0.92$ for Depressive).

The Temperament and Character Inventory-Revised (TCI-R) (Cloninger, 1999) is a questionnaire with 240 items scored on a 5-point Likert scale, which measures personality factors related to three character dimensions (Self-Directedness, Cooperativeness, and Self-Transcendence) and four temperament dimensions (Harm Avoidance, Novelty Seeking, Reward Dependence and Persistence). Evaluation of the Spanish revised version (Gutiérrez-Zotes et al., 2004) had an internal consistency of 0.87 (coefficient alpha). The internal consistency in this study for the subscales ranged from adequate to very good ($\alpha = 0.70$ for Novelty Seeking to $\alpha = 0.87$ for Self-Directedness).

The Yale Food Addiction Scale 2.0 (YFAS-2) (Gearhardt et al., 2016) is a 35-item self-report questionnaire for measuring FA during the prior year. The original YFAS was based on the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV-TR) (APA, 2010) criteria for substance dependence and was adapted to the context of food consumption. The YFAS-2 is based on DSM-5 criteria for SUDs (APA, 2013) and evaluates 11 symptoms. Two measurements may be calculated: (a) a continuous symptom count score reflecting the number of fulfilled criteria (ranging from 0 to 11), and (b) a FA threshold based on the number of symptoms (at least 2) and self-reported clinically significant impairment or distress. This final measurement allows for the binary classification of FA (present versus absent). Based on the revised DSM-5 taxonomy, it is possible to establish severity cut-offs: mild (2–3 symptoms), moderate (4–5 symptoms), and severe (6–11 symptoms). The Spanish validation of the YFAS-2 (Granero, Jiménez-Murcia, et al., 2018) generated an internal consistency of 0.94 (coefficient alpha). In this study, internal consistency for the total score was excellent ($\alpha = 0.96$).

The Alcohol Use Disorders Identification Test (AUDIT) (Saunders et al., 1993) is a 10-item screening questionnaire for hazardous alcohol consumption. It includes questions about the amount and frequency of drinking, alcohol dependence, and problems caused by alcohol. A score of 8 or more is considered to indicate harmful alcohol use and a score of 12 or more in women (15 or more in men) is likely to indicate alcohol dependence. This questionnaire has shown adequate validity in Spanish samples (Delgado, 1996).

The Drug Use Disorders Identification Test (DUDIT) (Berman et al., 2003) is an 11-item self-administered instrument to identify non-alcohol drug use patterns and related problems in individuals likely to meet criteria for a substance dependence diagnosis. The total score can range from 0 to 44 (as a result of the sum of the 11 items scored from 0 to 4); higher scores reflect more severe drug use problem. The first nine items are scored on a 5-point Likert scale ranging from 0 to 4, and the last two are scored on 3-point scales (values of 0, 2, and 4).

Other variables, including sociodemographic variables and other gambling-related measures (e.g. age of onset and duration), were obtained via face-to-face clinical interviews. The interviews also assessed participants' educational attainment. Socioeconomic status was assessed using the Hollingshead Four Factor Index of Socioeconomic Status (Hollingshead, 2011).

Procedure

All participants in our sample voluntarily sought treatment for GD. Experienced psychologists and psychiatrists conducted two face-to-face clinical interviews, before and after completing study instruments. The first visit was a clinical interview, in which the medical/psychiatric history was assessed, and the subsequent visit consisted of an interview to report the results of the tests and information about the treatment, according to their clinical characteristics and factors associated with GD.

The participants included in this study met criteria for GD and did not present other comorbid behavioral addictions and did not receive any compensation for participating in the study.

Statistical Analyses

Statistical analysis was conducted with Stata16 for Windows (Stata-Corp, 2019). Between-group comparisons for personality (TCI-R), psychopathology (SCL-90R), substance use and BMI measures were based on analysis of variance adjusted for age (ANCOVA) (gender was not considered as a covariate because results stratified for men and women were obtained). Effect sizes for mean differences were estimated through Cohen's *d* coefficient, poor-low for $|d| > 0.20$, mild-moderate for $|d| > 0.50$ and large-high for $|d| > 0.80$ (Kelley & Preacher, 2012). Between-group comparisons for categorical variables were based on logistic regression also adjusted for age. Type-I error related to multiple comparisons was controlled using the Finner's method, a family-wise procedure with higher power than a Bonferroni correction (Finner & Roters, 2001).

Path analysis was used to assess the set of relationships between gender, age, the personality domains of self-directedness and harm avoidance, FA severity, global psychopathological distress and gambling preference. Path analysis constitutes an extension of multiple regression modeling, used with the aim of estimating magnitudes and significance of simultaneous associations in a set of variables, including direct and indirect effects

(mediational links) (Kline, 2005). This technique is currently used for both exploratory and confirmatory purposes, and therefore it contributes to theory testing and theory development (MacCallum & Austin, 2000). This study modeled path analyses through structural equation modeling (SEM), with the maximum-likelihood estimation method. Goodness of fit was assessed with standard indexes, and it was considered adequate fitting (Barrett, 2007) for: non-significant result in the chi-square test (χ^2), root mean square error of approximation RMSEA < 0.08, Bentler's Comparative Fit Index CFI > 0.90, Tucker-Lewis Index TLI > 0.90, and standardized root mean square residual SRMR < 0.10. Global predictive capacity was estimated with the coefficient of determination (CD).

Ethics

The present study was conducted in accordance with the latest version of the Declaration of Helsinki. The General University Hospital Clinical Research Ethics Committee approved the study, and signed informed consent was obtained from all participants (Ref: PR329/19).

Results

Sample Characteristics

Most patients were single ($n=454$, 52.4%) or married ($n=313$, 36.1%), achieved primary ($n=460$, 53.1%) or secondary ($n=331$, 38.2%) education levels, were of medium-low or low socioeconomic status ($n=723$, 83.4%) and were employed ($n=527$, 60.8%). Mean age was 40.5 years ($SD=13.8$). Most patients reported only non-strategic gambling ($n=449$; 51.8%), and 26.1% ($n=226$) and 22.1% ($n=192$) reported only strategic gambling and both gambling forms, respectively. Strategic gambling activities are those where the gambler can use knowledge of the game to influence or predict the outcome, while nonstrategic gambling implies little or no possibility of influencing the outcome (Potenza et al., 2001). The gambling activity reported as the main reason for seeking treatment in the sample was slot-machines (60.1%), followed by casino or gambling saloons (24.7%), sports-betting (16.0%), and lotteries (14.4%).

Comparison Between Patients with Positive Versus Negative FA Screening

Seventy-two participants (8.3%) screened positive for FA. FA was observed among 18.8% of women and 7.4% of men. Table 1 contains the comparison for individuals with negative versus positive FA screening, stratified by gender (separate comparisons were performed for women and men due to the potential moderator role of gender). Among women, FA was associated with harm avoidance, less self-directedness and cooperativeness, greater psychopathology and higher BMI. Among men, FA was associated with harm avoidance and self-transcendence, less self-directedness and cooperativeness, greater psychopathology, greater alcohol use problem severity and higher BMI.

Table 2 includes the prevalence of patients outside the normative ranges for the main psychological variables of the study (estimates represent proportions of participants in subclinical or clinical severity levels). Among women, FA was associated with obesity, greater psychopathology and more self-directedness. Among men, FA was associated

Table 1 Associations between FA and clinical variables with the FA measures

	Women				Men					
	FA – (n=56)		FA + (n=13)		FA – (n=739)		FA + (n=59)			
	Mean	SD	Mean	SD	Mean	SD	Mean	SD		
TCI-R Novelty seeking	108.3	12.2	106.4	8.8	110.9	13.0	111.8	12.5	.371	0.07
TCI-R Harm avoidance	108.2	17.4	118.3	17.2	98.4	15.8	107.4	15.4	.001*	0.58†
TCI-R Reward depend	101.1	14.7	99.8	10.6	97.3	13.8	95.4	16.1	.324	0.12
TCI-R Persistence	105.5	18.1	103.4	21.7	108.2	18.5	105.1	22.7	.228	0.15
TCI-R Self-directedness	128.4	20.6	108.8	18.7	130.4	21.2	117.4	18.8	.001*	0.65†
TCI-R Cooperativeness	137.7	12.2	129.3	13.8	129.6	15.6	122.9	18.8	.002*	0.39
TCI-R Transcendence	62.7	13.7	67.4	11.2	61.2	14.1	67.7	14.7	.001*	0.46
SCL-90R Somatization	1.40	0.90	2.34	0.89	0.89	0.76	1.54	0.99	.001*	0.75†
SCL-90R Obsessive/comp	1.32	0.87	2.16	0.99	1.11	0.77	1.93	0.93	.001*	0.97†
SCL-90R Interp. sensitive	1.21	0.85	2.32	0.84	0.97	0.80	1.74	0.95	.001*	0.87†
SCL-90R Depressive	1.87	1.00	2.74	0.82	1.50	0.92	2.20	0.93	.001*	0.75†
SCL-90R Anxiety	1.20	0.84	2.06	0.72	0.97	0.78	1.59	0.98	.001*	0.69†
SCL-90R Hostility	0.90	0.78	1.67	0.73	0.94	0.84	1.53	1.10	.001*	0.61†
SCL-90R Phobic anxiety	0.54	0.51	1.25	1.09	0.42	0.61	0.97	0.90	.001*	0.73†
SCL-90R Paranoid	1.09	0.84	2.21	0.86	0.94	0.80	1.60	0.99	.001*	0.74†
SCL-90R Psychotic	0.98	0.82	1.68	0.70	0.89	0.75	1.57	0.95	.001*	0.80†
SCL-90R GSI score	1.28	0.72	2.13	0.65	1.03	0.68	1.72	0.83	.001*	0.91†
SCL-90R PST score	51.20	20.16	65.91	15.96	46.42	21.96	62.03	18.86	.001*	0.76†
SCL-90R PSDI score	2.08	0.60	2.89	0.61	1.85	0.56	2.37	0.71	.001*	0.82†
Tobacco (cigarettes/day)	9.91	11.51	7.55	13.77	9.45	10.90	9.31	13.12	.925	0.01
Alcohol (AUDIT total)	2.41	4.01	1.06	1.87	5.20	5.68	7.25	8.31	.011*	0.29
Drugs (DUDIT total)	1.25	4.02	0.10	0.00	2.75	6.61	3.80	8.18	.247	0.14

Table 1 (continued)

	Women				Men				
	FA – (n = 56)		FA + (n = 13)		FA – (n = 739)		FA + (n = 59)		
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	
Body mass index (kg/m ²)	27.43	5.96	33.43	6.89	26.08	4.35	30.34	6.87	.001* 0.74[†]

SD: standard deviation; FA – : food addiction negative screening; FA + : food addiction positive screening

TCI-R: Temperament and Character Inventory – Revised

SCL-90R: Symptom Checklist-90-Revised

AUDIT: Alcohol Use Disorders Identification Test

DUDIT: Drug Use Disorders Identification Test

*Bold: significant parameter (.05 level)

[†]Bold: effect size into the range mild-moderate ($|d| > 0.50$) to large-high ($|d| > 0.80$)

Results adjusted by the covariate age

Table 2 Comparison of clinical measures by gender and FA status

	Women					Men						
	FA – (n = 56)		FA + (n = 13)			FA – (n = 739)		FA + (n = 59)				
	n	%	n	%	<i>p</i>	<i>ldl</i>	n	%	n	%	<i>p</i>	<i>ldl</i>
GD (8–9 DSM-5 criteria)	30	53.6	8	61.5	.603	0.16	354	47.9	42	71.2	.001*	0.51[†]
Obesity (BMI > 30)	18	32.1	9	69.2	.014*	0.76[†]	129	17.5	32	54.2	.001*	0.79[†]
SCL-90R Somatization	22	39.3	10	76.9	.013*	0.78[†]	288	39.0	41	69.5	.001*	0.62[†]
SCL-90R Obsessive/comp	24	42.9	10	76.9	.027*	0.71[†]	304	41.1	46	78.0	.001*	0.77[†]
SCL-90R Interp. sensitive	30	53.6	11	84.6	.040*	0.69[†]	323	43.7	48	81.4	.001*	0.80[†]
SCL-90R Depressive	30	53.6	12	92.3	.010*	0.94[†]	439	59.4	47	79.7	.001*	0.50[†]
SCL-90R Anxiety	17	30.4	10	76.9	.002*	0.97[†]	321	43.4	41	69.5	.001*	0.53[†]
SCL-90R Hostility	14	25.0	10	76.9	.001*	1.09[†]	207	28.0	31	52.5	.001*	0.51[†]
SCL-90R Phobic anxiety	13	23.2	7	53.8	.028*	0.64[†]	196	26.5	37	62.7	.001*	0.75[†]
SCL-90R Paranoid	23	41.1	11	84.6	.005*	0.94[†]	219	29.6	36	61.0	.001*	0.64[†]
SCL-90R Psychotic	30	53.6	12	92.3	.010*	0.94[†]	363	49.1	45	76.3	.001*	0.57[†]
SCL-90R GSI score	31	55.4	13	100.0	.003*	1.46[†]	419	56.7	50	84.7	.001*	0.63[†]
SCL-90R PST score	35	62.5	11	84.6	.128	0.51[†]	420	56.8	48	81.4	.001*	0.54[†]
SCL-90R PSDI score	8	14.3	8	61.5	.001*	1.03[†]	160	21.7	31	52.5	.001*	0.65[†]
TCI-R Novelty seeking	30	53.6	4	30.8	.138	0.47	324	43.8	22	37.3	.328	0.13
TCI-R Harm avoidance	28	50.0	7	53.8	.803	0.08	263	35.6	33	55.9	.002*	0.41
TCI-R Reward depend	21	37.5	6	46.2	.565	0.18	260	35.2	25	42.4	.267	0.15
TCI-R Persistence	16	28.6	6	46.2	.220	0.37	267	36.1	25	42.4	.338	0.13
TCI-R Self-directedness	41	73.2	12	92.3	.142	0.53[†]	465	62.9	47	79.7	.010*	0.37
TCI-R Cooperativeness	16	28.6	6	46.2	.220	0.37	277	37.5	37	62.7	.001*	0.51[†]
TCI-R Transcendence	16	28.6	4	30.8	.875	0.05	286	38.7	19	32.2	.323	0.14

SD: standard deviation; GD: gambling disorder

FA –: food addiction negative screening, FA +: food addiction positive screening. BMI: Body Mass Index

SCL-90R: Symptom Checklist-90-Revised

TCI-R: Temperament and Character Inventory-Revised

*Bold: significant parameter (.05 level)

[†]Bold: effect size into the range mild-moderate (*ldl*>0.50) to large-high (*ldl*>0.80)

Results adjusted by the covariate age

with higher GD severity, obesity, greater psychopathology, more self-directedness and harm avoidance and less cooperativeness.

Table S1 (supplementary) contains the comparison between patients with negative versus positive FA screening for socio-demographics, gambling profile, and problem gambling severity measured as the total number of DSM-5 criteria for GD and SOGS scores. Among women, FA was associated with lower socioeconomic status, unemployment, non-strategic gambling preference, and longer durations of gambling problems. Among men, FA was associated with higher socioeconomic status, greater GD severity and earlier age of onset of gambling.




Fig. 1 Radar chart (n = 867)




Fig. 2 Path diagram with the results of the structural equation model (n = 867). Note Only significant coefficients were retained in the model

Figure 1 displays a radar chart with the profiles related to FA, separately for women and men. Standardized scores are plotted in this graphic to allow easy interpretation as original scales had different ranges.

Path Analysis

Figure 2 displays a path diagram with standardized coefficients obtained in the path analysis (Table S2 includes the complete results for this model, including tests for direct, indirect and total effects). Only significant coefficients were retained in the final model. Only two personality domains were retained (self-directedness and harm avoidance) because the other TCI-R scales did not achieve significant effects. Adequate goodness-of-fit was achieved: $\chi^2=10.71$ ($p=0.296$), RMSEA=0.015 (95% confidence interval: 0.001 to 0.043), CFI=0.999, TLI=0.997, and SRMR=0.019. Global predictive capacity was also good (CD=0.299).

FA severity level (defined as the YFAS-2 total score) was higher for women and participants with lower self-directedness scores and higher harm avoidance level. Both personality domains also showed mediational links between gender and age with FA severity: being female was associated with decreased self-directedness and increased harm avoidance, while older age was associated with increased harm avoidance. Path analysis also suggested that FA levels led to higher psychopathology, which also was directly related to

lower self-directedness and higher harm avoidance. Gambling preference was also directly related to age, with older age linked to preferences for non-strategic gambling.

Discussion

The present work studied among GD patients, clinical features associated with FA in women and men. The prevalence of FA in the total sample was 8.3% (18.8% among women, 7.4% among men), similar than the 7.8% reported in other similar studies (Granero, Jiménez-Murcia, et al., 2018). Besides, this percentage is lower than the one corresponding to the presence of FA in non-clinical population (Gearhardt et al., 2016), the study of the common features between both disorders responds to the similar aspects involved in the addictive process, already mentioned in the literature (Innamorati et al., 2017; Jiménez-Murcia et al., 2017; Mestre-Bach, Steward, et al., 2020).

Although no relation between FA and age was found, significant differences in socioeconomic level and employment state were evidenced only among women. In this vein, women with FA had lower socioeconomic status and unemployed. Higher scores in general psychopathology and certain personality traits, namely higher harm avoidance and lower self-directedness and cooperativeness were found in both genders. However, men with FA presented higher self-transcendence, although a numerical difference in the same direction with a similar effect size was observed in women, suggesting a similar relationship in the smaller female sample. Regardless of gender, FA was associated with a higher BMI and obesity. Association between FA and types of gambling were largely negligible. Implications are discussed below.

Although a higher prevalence of GD is described in men than women (Blanco et al., 2006; Husky et al., 2015), previous studies have reported that women are more likely to engage in FA behaviors (Schulte & Gearhardt, 2018) and experience abnormal eating and weight disorders (Jiménez-Murcia et al., 2019; Romero et al., 2019; Schulte & Gearhardt, 2018), with initial studies suggesting similar relationships in patients with GD (Granero et al., 2018; Jiménez-Murcia et al., 2017). Lower socioeconomic status has been associated with GD severity among women (Jiménez-Murcia et al., 2020). Higher psychopathology, particularly depressive and anxiety symptoms, and some specific personality features, such as low self-directedness and perseverance or high reward dependence, have been linked to FA in both medical and mental conditions (Brunault et al., 2018; Imperatori, 2014; Wolz et al., 2016). FA may impact global health, with a special influence on BMI and overweight/obesity (Murphy et al., 2014; Meseri et al., 2020). Our findings among GD patients suggest a poor socioeconomic context and work difficulties (e.g., unemployment), higher emotional vulnerability (e.g., anxiety and depressive symptoms, particularly for women) and poor adaptive coping strategies to deal with stress (e.g., higher harm avoidance and lower self-directedness and cooperativeness) may characterize GD patients with FA.

Regarding clinical features, women with GD and FA had higher duration of GD in comparison with those who did not fulfill FA criteria. Male patients with both disorders were characterized by lower age of onset and higher severity of the GD. These results differ from previous studies were no differences in age of onset and duration of GD were found among patients with GD and FA (Jiménez-Murcia et al., 2017). Moreover, no significant differences were found in gambling modality associated with the presence of FA in neither gender group. Thus, our results could suggest that the presence of FA in patients with GD might be associated with a more severe GD clinical profile in both genders.

Both men and women with GD and FA appear to fit the emotionally vulnerable subtype from the pathway model proposed by Blaszczynski & Nower (2002), and also within the cluster 2–3 (moderate-functional clusters) described among FA individuals with obesity by Jimenez-Murcia (2019). In this group of patients, gambling and eating behaviors might represent potential elements of addiction as these behaviors are seen as strategies to relieve emotional discomfort they may experience in daily situations, usually perceived as threatening and stressful, or which they may not have the proper skills to manage adequately. Therefore, the identification of a vulnerable clinical profile at baseline among GD patients with FA suggests a need for early systematic identification of FA by clinicians. Although both genders have similarities, this subtype may be especially important among women, who appear more likely to experience FA, overweight/obesity and fewer socio-economic resources. Moreover, the creation and/or optimization of specific and individualized social and therapeutic approaches are needed, taking into account that the lack of social support, economic difficulties or physical limitations (e.g., related to obesity) may result in decreased access to medical services. As suggested in previous studies (Jiménez-Murcia et al., 2019), some approaches may target better nutritional and weight management, but also emotional regulation and problem-solving strategies to cope with stress may also be needed.

To the best of our knowledge, this is the first study that analyzes clinical correlates of FA in GD patients from a gender-informed perspective. The results suggest a negative influence of FA on GD, in terms of a higher presence of poor prognosis factors for GD in patients with comorbid FA, that are both similar across genders and differ by gender. With respect to the latter, a longer duration of the GD was associated with FA in women, and in men, FA was associated with an earlier age of onset and higher severities of GD and problematic alcohol use. Of note, some of these differences (particularly with respect to relationships with GD severity) may also pertain to women, as evidenced by largely similar effect sizes and the smaller sample of women.

The clinical characteristics associated with FA have been reported as having deleterious impacts on the course of GD (del Pino-Gutiérrez et al., 2017; Jiménez-Murcia et al., 2016; Valero-Solís et al., 2018). Low socioeconomic status or co-occurring psychopathology (e.g. depression, anxiety) may be particularly relevant to FA in women with GD. These factors could delay medical consultation for GD, resulting in a longer duration of GD when receiving a GD diagnosis. In the case of men, early age of gambling onset and problematic alcohol use appear particularly relevant. Previous studies have reported a relationship between FA and tobacco use among a male predominant sample with GD (Jiménez-Murcia et al., 2017). Apart from gambling and food, men with GD and FA may also use alcohol as a mechanism to cope with stress and negative emotions.

FA was more prevalent among women (versus men) with GD. These results suggest that assessing for FA in treatment-seeking individuals with GD may be particularly helpful for women's health and useful in general for identifying vulnerable patients presenting with the potential for worse courses of GD. The detection and simultaneous treatment of co-occurring psychiatric concerns among GD patients is important. Our findings suggest that the design of specific protocols to detect FA is needed, as is the testing of treatment approaches used in other care settings for addressing FA in GD populations.

In considering the pathway analyses, relationships between gender and personality features with appear mediated by FA. Therefore, the results reinforcement the interpretations of a specific vulnerable group of GD patients (i.e., women with dysfunctional personality features) in whom FA may represent a maladaptive way of dealing with higher levels of psychological distress. Similar conclusions may extend to GD, as has been also proposed

previously (Di Trani et al., 2017; Jiménez-Murcia et al., 2017), with poor emotional regulation contributing importantly to behavioral addictions (Estévez et al., 2017; Mestre-Bach, Fernández-Aranda, et al., 2020b). In this sense, the detection of FA among GD patients could lead to a greater emphasis on approaches to improve emotional regulation (e.g., with mindfulness-based or stress-reduction treatments) within GD treatment. The acquisition of adaptive emotional skills may be translated into a better stress management and reduced gambling and eating to manage such negative states.

A relationship between age and gambling preferences was found, with preferences for non-strategic gambling with older age, consistent with prior findings (Potenza, Steinberg, et al., 2006). Among patients with FA the relationship with GD subtype was only indirect, and specifically in females. This result is consistent with previous works regarding GD samples (Assanangkornchai et al., 2016; Odlaug et al., 2011). Men (particularly younger ones) tend to prefer and have problems with strategic gambling, whereas women (particularly older ones) tend to prefer and have problems with non-strategic gambling (Moragas et al., 2015; Odlaug et al., 2011; Potenza et al., 2001; Potenza et al., 2006; Stevens & Young, 2010).

Strength and Limits

Strengths of this study, such as the large clinical sample, should be mentioned. To date, this is the first study to consider FA in GD patients in a gender-sensitive manner. As the sample consisted of patients treated at a specific unit, assessments were consistently conducted. However, study limitations should also be mentioned. For instance, the treatment-seeking sample was from a region of Spain; as such, the findings may not generalize to non-clinical samples or those from other jurisdictions. The use of self-report assessments for psychiatric conditions may decrease reliability. Future studies should consider alternate assessments (e.g., structured clinical interviews). However, as FA remains a debated construct, its evaluation with a self-report measure is presently most reasonable. Another limitation of the study was the small sample size for the group of patients who met criteria for FA: the underpowered statistical accuracy reduces the chance of detecting a true effect, reproducibility of the results and generalization capacity. In addition, the cross-sectional nature of the study did not allow the study of the progression of the concurrent condition of GD with FA, nor examining the underlying mechanisms contributing to the course of this complex clinical condition.

Conclusions

In conclusion, this study characterizes the clinical profile of GD patients with and without FA from a gender-informed perspective. GD patients, especially women, with specific personality characteristics and psychopathology may be particularly prone to FA. Further, a lower socioeconomic status may also be relevant for women with GD to experience FA. Speculatively, the findings taken together suggest that GD patients may engage in FA behaviors to manage high levels of psychological distress. Alcohol and consumption, early age of onset, and severity of GD may contribute to a worse GD prognosis in the presence of FA, particularly in men. In women, a longer duration of GD was linked to FA, suggesting the need for enhancing early intervention efforts to

improve women's health. Therefore, these results support the existence of a specific vulnerable group of GD patients and suggest the relevance of designing specific screening and treatment protocol to address FA in GD patients. Further studies are necessary to increase knowledge about FA and its influence not only in GD, but also in other addictive disorders and mental health conditions. Addressing areas such as underlying mechanisms and neurobiological factors related to FA could be helpful for a better understanding of this condition and its clinical relevance.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s10899-021-10071-w>.

Authors' Contribution ME, IB, LM, GM-B and SJ-M contributed to the development of the study concept and design. RG performed the statistical analysis. ME, IB, MG-P, LM, AP-G, EC, BM-M and EV-M aided with data collection. ME, IB, LM, GM-B and SJ-M aided with interpretation of data and the writing of the manuscript. MNP, ANG, FF-A, and SJ-M revised the manuscript and provided substantial comments. FF-A and SJ-M obtained funding.

Funding First, we thank CERCA Programme / Generalitat de Catalunya for institutional support. This work was additionally supported by grants from the Ministerio de Ciencia, Innovación y Universidades (grant RTI2018-101837-B-100) and the Delegación del Gobierno para el Plan Nacional sobre Drogas (2017I067 and 2019I47), Instituto de Salud Carlos III (ISCIII) (PI17/01167) and co-funded by FEDER funds/European Regional Development Fund (ERDF), a way to build Europe. CIBERObn is an initiative of ISCIII. IB was partially supported by a Post-Residency Grant from the Research Committee of the University Hospital of Bellvitge (HUB; Barcelona, Spain) 2019–2020. LM is supported by a postdoctoral grant of the mexican institution Consejo Nacional de Ciencia y Tecnología (CONACYT). GMB was supported by a postdoctoral grant from FUNCIVA. The funders had no role in the study design, data collection and analysis, decision to publish or preparation of the manuscript. Dr. Potenza involvement was supported by the Connecticut Department of Mental Health and Addiction Services, the Connecticut Council on Problem Gambling and the National Center for Responsible Gaming.

Data Availability and Material (Data Transparency) Individuals may inquire with Dr. Jiménez-Murcia regarding availability of the data as there exist ongoing studies using the data. In order to avoid overlapping research efforts, Dr. Jiménez-Murcia will consider request on a case-by-case bases.

Code Availability Not applicable.

Declarations

Conflict of interest Dr. Potenza has consulted for and advised Game Day Data, the Addiction Policy Forum, AXA, Idorsia, and Opiant Therapeutics; has received research support from the Veteran's Administration, Mohegan Sun Casino, and the National Center for Responsible Gaming (no the International Center for Responsible Gambling); has participated in surveys, mailings, or telephone consultations related to addictions, impulse-control disorders or other health topics; has consulted for law offices and the federal public defender's office in issues related to impulse-control and addictive disorders; has provided clinical care in the Connecticut Department of Mental Health and Addiction Services Problem Gambling Services Program; has performed grant reviews for the National Institutes of Health and other agencies; has edited journals and journal sections; has given academic lectures in grand rounds, CME events and other clinical/scientific venues; and has generated books or chapters for publishers of mental health texts. Other authors report no disclosures. Dr. Fernandez-Aranda received consultancy honoraria from Novo Nordisk and editorial honoraria as EIC from Wiley. The rest of the authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

Ethical Approval The present study was carried out in accordance with the latest version of the Declaration of Helsinki. The Bellvitge University Hospital Clinical Research Ethics Committee approved the study, and signed informed consent was previously obtained from all participants (Ref: PR329/19).

Consent to Participate All subjects were informed about the study and all provided a written informed consent.

Consent for Publication Consent for publication of this article has been obtained from all authors.

References

- Abbott, M., Romild, U., & Volberg, R. (2018). The prevalence, incidence, and gender and age-specific incidence of problem gambling: Results of the Swedish longitudinal gambling study (Swelogs). *Addiction*. <https://doi.org/10.1111/add.14083>
- American Psychiatric Association. (2010). *Diagnostic and statistical manual of mental disorders* (4th ed.). American Psychiatric Association
- APA. American Psychiatric Association, (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). In *American Journal of Psychiatry*.
- Assanangkornchai, S., McNeil, E. B., Tantirangsee, N., & Kittirattanapaiboon, P. (2016). Gambling disorders, gambling type preferences, and psychiatric comorbidity among the Thai general population: Results of the 2013 national mental health survey. *Journal of Behavioral Addictions*, *5*(3), 410–418. <https://doi.org/10.1556/2006.5.2016.066>.
- Barrett, P. (2007). Structural equation modelling: Adjudging model fit. *Personality and Individual Differences*, *42*(5), 815–824. <https://doi.org/10.1016/j.paid.2006.09.018>.
- Berman, A. H., Bergman, H., Palmstierna, T., & Schlyter, F. (2003). *The Drug Use Disorders Identification Test (DUDIT) Manual*. Karolinska Institutet
- Blanco, C., Hasin, D. S., Petry, N., Stinson, F. S., & Grant, B. F. (2006). Sex differences in subclinical and DSM-IV pathological gambling: Results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Psychological Medicine*, *36*(7), 943–953
- Blaszczyński, A., & Nower, L. (2002). A pathways model of problem and pathological gambling. *Addiction*, *97*(5), 487–499. <https://doi.org/10.1046/j.1360-0443.2002.00015.x>
- Brunault, P., Ducluzeau, P. H., Courtois, R., Bourbao-Tournois, C., Delbachian, I., Réveillère, C., & Balon, N. (2018). Food addiction is associated with higher neuroticism, Lower conscientiousness, higher impulsivity, but lower extraversion in obese patient candidates for bariatric surgery. In *Substance Use and Misuse*. <https://doi.org/10.1080/10826084.2018.1433212>
- Burrows, T., Kay-Lambkin, F., Pursey, K., Skinner, J., & Dayas, C. (2018). Food addiction and associations with mental health symptoms: A systematic review with meta-analysis. *Journal of Human Nutrition and Dietetics*, *31*(4), 544–572. <https://doi.org/10.1111/jhn.12532>.
- Carter, J. C., & Davis, C. (2010). An addiction model of binge eating disorder. In *Addiction Medicine* (pp. 633–647). Springer.
- Cloninger, C. R. (1999). *The temperament and character inventory—Revised*. Center for Psychobiology of Personality, Washington University
- del Pino-Gutiérrez, A., Fernández-Aranda, F., Granero, R., Tárrega, S., Valdepérez, A., Agüera, Z., Håkansson, A., Sauvaget, A., Aymamí, N., Gómez-Peña, M., Moragas, L., & Jiménez-Murcia, S. (2017). Impact of alcohol consumption on clinical aspects of gambling disorder. *International Journal of Mental Health Nursing*, *26*(2), 121–128. <https://doi.org/10.1111/inm.12221>
- Delgado, J. M. M. (1996). *Validación de los cuestionarios breves: AUDIT, CAGE y CBA para la detección precoz del síndrome de dependencia de alcohol en Atención Primaria*. Universidad de Cadiz
- Derogatis, L. R. (1994). *SCL-90-R: Symptom Checklist-90-R. Administration, Scoring and Procedures Manual—II for the Revised Version*. Clinical Psychometric Research
- Derogatis, L. R. (2002). *SCL-90-R. Cuestionario de 90 Síntomas-Manual* (TEA (ed.)).
- Di Trani, M., Renzi, A., Vari, C., Zavattini, G. C., & Solano, L. (2017). Gambling disorder and affect regulation: The role of alexithymia and attachment style. *Journal of Gambling Studies*. <https://doi.org/10.1007/s10899-016-9637-3>.
- Echeburúa, E., Baez, C., Fernández-Montalvo, J., & Páez, D. (1994). Cuestionario de Juego Patológico de South Oaks (SOGS): Validación española. *Análisis y Modificación de La Conducta*, *20*(74), 769–791
- Estévez, A., Jáuregui, P., Sánchez-Marcos, I., López-González, H., & Griffiths, M. D. (2017). Attachment and emotion regulation in substance addictions and behavioral addictions. *Journal of Behavioral Addictions*, *6*(4), 534–544. <https://doi.org/10.1556/2006.6.2017.086>

- Fauth-Bühler, M., Mann, K., & Potenza, M. N. (2017). Pathological gambling: A review of the neurobiological evidence relevant for its classification as an addictive disorder. In *Addiction Biology*. <https://doi.org/10.1111/adb.12378>
- Fernandez-Aranda, F., Karwautz, A., & Treasure, J. (2018). Food addiction: A transdiagnostic construct of increasing interest. *European Eating Disorders Review*, 26(6), 536–540. <https://doi.org/10.1002/erv.2645>
- Finner, H., & Roters, M. (2001). On the false discovery rate and expected type I errors. *Journal of the American Statistical Association*, 88, 920–923. <https://doi.org/10.1002/1521-4036>
- Fletcher, P. C., & Kenny, P. J. (2018). Food addiction: A valid concept? *Neuropsychopharmacology*, 43(13), 2506–2513. <https://doi.org/10.1038/s41386-018-0203-9>
- Gearhardt, A., Corbin, W., & Brownell, K. (2016). Development of the Yale Food Addiction Scale Version 2.0. *Psychology of Addictive Behaviors*, 30, 113–121. <https://doi.org/10.1037/adb0000136>
- Gearhardt, A. N., Davis, C., Kuschner, R., & Brownell, K. D. (2011). The addiction potential of hyperpalatable foods. *Current Drug Abuse Reviews*, 4(3), 140–145. <https://doi.org/10.2174/1874473711104030140>
- Gordon, E. L., Ariel-Donges, A. H., Bauman, V., & Merlo, L. J. (2018). What is the evidence for “food addiction?” A systematic review. In *Nutrients*. <https://doi.org/10.3390/nu10040477>
- Granero, R., Fernández-Aranda, F., Mestre-Bach, G., Steward, T., García-Caro, B., Prever, Gaviel-Fried, B., del Pino-Gutiérrez, A., Moragas, L., Aymamí, N., Gómez-Peña, M., & Jiménez-Murcia, S., J. M., &. (2018). Clustering of treatment-seeking women with gambling disorder. *Journal of Behavioral Addictions*, 7(3), 770–780. <https://doi.org/10.1556/2006.7.2018.93>
- Granero, R., Jiménez-Murcia, S., Gerhardt, A. N., Agüera, Z., Aymamí, N., Gómez-Peña, M., & Fernández-Aranda, F. (2018). Validation of the Spanish version of the Yale Food Addiction Scale 2.0 (YFAS 2.0) and clinical correlates in a sample of eating disorder, gambling disorder, and healthy control participants. *Frontiers in Psychiatry*. <https://doi.org/10.3389/fpsy.2018.00208>
- Grant, J. E., & Chamberlain, S. R. (2015). Gambling disorder and its relationship with substance use disorders: Implications for nosological revisions and treatment. *American Journal on Addictions*. <https://doi.org/10.1111/ajad.12112>
- Grant, J. E., & Chamberlain, S. R. (2020). Neurocognitive findings in young adults with binge eating disorder. *International Journal of Psychiatry in Clinical Practice*, 24(1), 71–76. <https://doi.org/10.1080/13651501.2019.1687724>
- Gutiérrez-Zotes, J. A., Bayón, C., Montserrat, C., Valero, J., Labad, A., Cloninger, C. R., & Fernández-Aranda, F. (2004). Temperament and Character Inventory Revised (TCI-R). Standardization and normative data in a general population sample. *Actas Espanolas de Psiquiatria*, 32(1), 8–15
- Hardy, R., Fani, N., Jovanovic, T., & Michopoulos, V. (2018). Food addiction and substance addiction in women: Common clinical characteristics. *Appetite*, 120, 367–373. <https://doi.org/10.1016/j.appet.2017.09.026>
- Hebebrand, J., Albayrak, A., Adanb, R., Antel, J., Dieguez, C., De Jongh, J., Leng, G., Menzies, J., Mercer, J.G., Murphy, M., Van Der Plasse, G., & Dickson, S. L. (2014). “Eating addiction”, rather than “food addiction”, better captures addictive-like eating behavior. *Neuroscience and Biobehavioral Reviews* 47, 295–306. <https://doi.org/10.1016/j.neubiorev.2014.08.016>
- Hollingshead, A. B. (2011). Four factor index of social status. *Yale Journal of Sociology*, 8, 21–51
- Husky, M. M., Michel, G., Richard, J. B., Guignard, R., & Beck, F. (2015). Gender differences in the associations of gambling activities and suicidal behaviors with problem gambling in a nationally representative French sample. *Addictive Behaviors*, 45, 45–50. <https://doi.org/10.1016/j.addbeh.2015.01.011>
- Imperatori, C., Innamorati, M., Contardi, A., Continisio, M., Tamburello, S., Lamis, D.A., Tamburello, A. & Fabbriatore, M. (2014). The association among food addiction, binge eating severity and psychopathology in obese and overweight patients attending low-energy-diet therapy. *Comprehensive Psychiatry*, 55(6), 1358–1362. <https://doi.org/10.1016/j.comppsych.2014.04.023>
- Innamorati, M., Imperatori, C., Harnic, D., Erbuto, D., Patitucci, E., Janiri, L., Lamis, D.A., Pompili, M., Tamburello, S. & Fabbriatore, M. (2017). Emotion regulation and mentalization in people at risk for food addiction. *Behavioral Medicine*, 43(1), 21–30. <https://doi.org/10.1080/08964289.2015.1036831>
- Jiménez-Murcia, S., Stinchfield, R., Álvarez-Moya, E., Jaurrieta, N., Bueno, B., Granero, R., Aymamí, M.N., Gómez-Peña, M., Martínez-Giménez, R., Fernández-Aranda, F. & Vallejo, J. (2009). Reliability, validity, and classification accuracy of a Spanish translation of a measure of DSM-IV diagnostic criteria for pathological gambling. *Journal of Gambling Studies*, 25(1), 93–104. <https://doi.org/10.1007/s10899-008-9104-x>
- Jiménez-Murcia, S., Agüera, Z., Paslakis, G., Munguia, L., Granero, R., Sánchez-González, J., Sánchez, I., Riesco, N., Gearhardt, A.N., Dieguez, C. & Fazio, G., & Fernández-Aranda, F. (2019). Food


- Addiction in Eating Disorders and Obesity: Analysis of Clusters and Implications for Treatment. *Nutrients*, 11(11), 2633. <https://doi.org/10.3390/nu11112633>
- Jiménez-Murcia, S., Pino-Gutiérrez, D., Fernández-Aranda, A., Granero, F., Hakansson, R., Tárrega, A., Valdepérez, A., Aymamí, N., Gómez-Peña, M., Moragas, L., Baño, M., & Menchón, J. M., T. (2016). Treatment outcome in male Gambling Disorder patients associated with alcohol use. *Frontiers in Psychology*, 7, 1–11. <https://doi.org/10.3389/fpsyg.2016.00465>
- Jiménez-Murcia, S., Granero, R., Giménez, M., del Pino-Gutiérrez, A., Mestre-Bach, G., Mena-Moreno, T., Moragas, L., Baño, M., Sánchez-González, J., de Gracia, M., Baenas-Soto, I., & Fernández-Aranda, F. (2020). Contribution of sex on the underlying mechanism of the gambling disorder severity. *Scientific Reports*, 10(1), 1–13. <https://doi.org/10.1038/s41598-020-73806-6>
- Jiménez-Murcia, S., Granero, R., Wolz, I., Baño, M., Mestre-Bach, G., Steward, T., Agüera, Z., Hinney, A., Diéguez, C., Casanueva, F.F., Gearhardt, A.N., & Fernández-Aranda, F. (2017). Food addiction in gambling disorder: Frequency and clinical outcomes. *Frontiers in Psychology*, 8, 1–12. <https://doi.org/10.3389/fpsyg.2017.00473>
- Kandeger, A., Selvi, Y., & Tanyer, D. K. (2019). The effects of individual circadian rhythm differences on insomnia, impulsivity, and food addiction. *Eating and Weight Disorders*, 24(1), 47–55. <https://doi.org/10.1007/s40519-018-0518-x>
- Karlsson, J., Broman, N., & Håkansson, A. (2019). Associations between problematic gambling, gaming, and internet use: A cross-sectional population survey. *Journal of Addiction*. <https://doi.org/10.1155/2019/1464858>.
- Kelley, K., & Preacher, K. J. (2012). On effect size. *Psychological Methods*, 17(2), 137–152. <https://doi.org/10.1037/a0028086>
- Kim, H. S., von Ranson, K. M., Hodgins, D. C., & McGrath, D. S., & Tavares, H. (2018). Demographic, psychiatric, and personality correlates of adults seeking treatment for disordered gambling with a comorbid binge/purge type eating disorder. *European Eating Disorders Review*, 26(5), 508–518. <https://doi.org/10.1002/erv.2606>
- Kline, R. B. (2005). *Principles and practice of structural equation modeling* (2nd Edition). The Guilford Press.
- Lee, R. S. C., Hoppenbrouwers, S., & Franken, I. (2019). A systematic meta-review of impulsivity and compulsivity in addictive behaviors. *Neuropsychology Review*, 14–26. <https://doi.org/10.1007/s11065-019-09402-x>
- Lesieur, H. R., & Blume, S. B. (1987). The South Oaks Gambling Screen (SOGS): A new instrument for the identification of Pathological gamblers. *American Journal of Psychiatry*, 144(9), 1184–1188. <https://doi.org/10.1176/ajp.144.9.1184>
- MacCallum, R. C., & Austin, J. T. (2000). Applications of structural equation modeling in psychological research. *Annual Review of Psychology*, 51, 201–226
- Mestre-Bach, G., Fernández-Aranda, F., Jiménez-Murcia, S., & Potenza, M. N. (2020a). Emotional regulation in gambling disorder. *Current Opinion in Behavioral Sciences*, 31, 102–108. <https://doi.org/10.1016/j.cobeha.2020.03.004>
- Mestre-Bach, G., Fernández-Aranda, F., Jiménez-Murcia, S., & Potenza, M. N. (2020b). Decision-making in gambling disorder, problematic pornography use, and binge-eating disorder: Similarities and differences. In *Current Behavioral Neuroscience Reports*, 7(3), 97–108. <https://doi.org/10.1007/s40473-020-00212-7>
- Mestre-Bach, G., Steward, T., Granero, R., Fernández-Aranda, F., Mena-Moreno, T., Vintro-Alcaraz, C., Lozano-Madrid, M., Menchón, J.M., Potenza, M.N., & Jiménez-Murcia, S. (2020). Dimensions of Impulsivity in Gambling Disorder. *Scientific Reports*. <https://doi.org/10.1038/s41598-019-57117-z>
- Moragas, L., Granero, R., Stinchfield, R., Fernández-Aranda, F., Fröberg, F., Aymamí, N., Gómez-Peña, M., Fagundo, A.B., Islam, M.A., del Pino-Gutiérrez, A., Agüera, Z & Jiménez-Murcia, S. (2015). Comparative analysis of distinct phenotypes in gambling disorder based on gambling preferences. *BMC Psychiatry*. <https://doi.org/10.1186/s12888-015-0459-0>
- Murphy, C. M., Stojek, M. K., & MacKillop, J. (2014). Interrelationships among impulsive personality traits, food addiction, and Body Mass Index. *Appetite*. <https://doi.org/10.1016/j.appet.2013.10.008>
- Odaug, B. L., Marsh, P. J., Kim, S. W., & Grant, J. E. (2011). Strategic vs. nonstrategic gambling: Characteristics of pathological gamblers based on gambling preference. *Annals of Clinical Psychiatry*.
- Penzenstadler, L., Soares, C., Karila, L., & Khazaa, Y. (2018). Systematic review of food addiction as measured with the Yale food addiction scale: Implications for the food addiction construct. *Current Neuropharmacology*, 17(6), 526–538. <https://doi.org/10.2174/1570159X16666181108093520>.
- Petry, N. M., Stinson, F. S., & Grant, B. F. (2005). Comorbidity of DSM-IV pathological gambling and other psychiatric disorders: Results from the national epidemiologic survey on alcohol and related conditions. *Journal of Clinical Psychiatry*. <https://doi.org/10.4088/JCP.v66n0504>

- Potenza, Balodis, I. M., Derevensky, J., Grant, J. E., Petry, N. M., Verdejo-Garcia, A., & Yip, S. W. (2019). Gambling disorder. *Nature Reviews Disease Primers*, 5(1), 51. <https://doi.org/10.1038/s41572-019-0099-7>
- Potenza, M. N., Maciejewski, P. K., & Mazure, C. M. (2006). A gender-based examination of past-year recreational gamblers. *Journal of Gambling Studies*. <https://doi.org/10.1007/s10899-005-9002-4>
- Potenza, M. N., Steinberg, M. A., McLaughlin, S. D., Wu, R., Rounsaville, B. J., & O'Malley, S. S. (2001). Gender-related differences in the characteristics of problem gamblers using a gambling helpline. *American Journal of Psychiatry*. <https://doi.org/10.1176/appi.ajp.158.9.1500>
- Pursey, K. M., Contreras-Rodriguez, O., Collins, C. E., Stanwell, P., & Burrows, T. L. (2019). Food addiction symptoms and amygdala response in fasted and fed states. *Nutrients*. <https://doi.org/10.3390/nu11061285>
- Pursey, K. M., Stanwell, P., Gearhardt, A. N., Collins, C. E., & Burrows, T. L. (2014). The prevalence of food addiction as assessed by the yale food addiction scale: A systematic review. In *Nutrients*. <https://doi.org/10.3390/nu6104552>
- Meseri, R., & Akanalci, C. (2020). Food addiction: A key factor contributing to obesity?. *Journal of Research in Medical Sciences: The Official Journal of Isfahan University of Medical Sciences*, 25, 71. https://doi.org/10.4103/jrms.JRMS_971_19
- Romero, X., Agüera, Z., Granero, R., Sánchez, I., Riesco, N., Jiménez-Murcia, S., Gisbert-Rodríguez, M., Sánchez-González, J., Casalé, G., Baenas, I. and Valenciano-Mendoza, E., & Dieguez, C. (2019). Is food addiction a predictor of treatment outcome among patients with eating disorder? *European Eating Disorders Review*, 27(6), 700–711. <https://doi.org/10.1002/erv.2705>
- Saunders, J. B., Aasland, O. G., Babor, T. F., De la Fuente, J. R., & Grant, M. (1993). Development of the alcohol use disorders identification test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption-II. *Addiction*, 88(6), 791-804. <https://doi.org/10.1111/j.1360-0443.1993.tb02093.x>
- Saunders, J. B., Hao, W., Long, J., King, D. L., Mann, K., Fauth-Bühler, M., Rumpf, H.J., Bowden-Jones, H., Rahimi-Movaghar, A., Chung, T., Chan, E., & Poznyak, V. (2017). Gaming disorder: Its delineation as an important condition for diagnosis, management, and prevention. *Journal of Behavioral Addictions*, 6(3), 271–279. <https://doi.org/10.1556/2006.6.2017.039>
- Schulte, E. M., Avena, N. M., & Gearhardt, A. N. (2015). Which foods may be addictive? The roles of processing, fat content, and glycemic load. *PLoS ONE*. <https://doi.org/10.1371/journal.pone.01117959>
- Schulte, E. M., & Gearhardt, A. N. (2018). Associations of food addiction in a sample recruited to be nationally representative of the United States. *European Eating Disorders Review*. <https://doi.org/10.1002/erv.2575>
- Schulte, E. M., Potenza, M. N., & Gearhardt, A. N. (2017). A commentary on the “eating addiction” versus “food addiction” perspectives on addictive-like food consumption. *Appetite*, 115, 9–15. <https://doi.org/10.1016/j.appet.2016.10.033>
- Stata-Corp. (2019). *Stata Statistical Software: Release 16*. College Station, TX: StataCorp LLC
- Stevens, M., & Young, M. (2010). Who plays what? participation profiles in chance versus skill-based gambling. *Journal of Gambling Studies*. <https://doi.org/10.1007/s10899-009-9143-y>
- Steward, T., Mestre-Bach, G., Vintró-Alcaraz, C., Lozano-Madrid, M., Agüera, Z., Fernández-Formoso, J. A., Granero, R., Jiménez-Murcia, S., Vilarrasa, N., García-Ruiz-de-Gordejuela, A., Fernández-Aranda, F., & Veciana de las Heras, M. (2018). Food addiction and impaired executive functions in women with obesity. *European Eating Disorders Review*, 26(6), 574–584. <https://doi.org/10.1002/erv.2636>
- Stinchfield, R. (2003). Reliability, validity, and classification accuracy of a measure of DSM-IV diagnostic criteria for pathological gambling. *American Journal of Psychiatry*, 160(1), 180–182. <https://doi.org/10.1176/appi.ajp.160.1.180>
- Subramaniam, M., Wang, P., Soh, P., Vaingankar, J. A., Chong, S. A., Browning, C. J., & Thomas, S. A. (2015). Prevalence and determinants of gambling disorder among older adults: A systematic review. In *Addictive Behaviors*. <https://doi.org/10.1016/j.addbeh.2014.10.007>
- Tackett, J. L., Krieger, H., Neighbors, C., Rinker, D., Rodriguez, L., & Edward, G. (2017). Comorbidity of Alcohol and Gambling Problems in Emerging Adults: A Bifactor Model Conceptualization. *Journal of Gambling Studies*. <https://doi.org/10.1007/s10899-016-9618-6>
- The World Health Organisation. (2019). WHO | International Classification of Diseases, 11th Revision (ICD-11). *International Classification of Diseases 11th Revision (ICD-11)*.
- Treasure, J., Duarte, T. A., & Schmidt, U. (2020). Eating disorders. *The Lancet*, 395(10227), 899–911. [https://doi.org/10.1016/S0140-6736\(20\)30059-3](https://doi.org/10.1016/S0140-6736(20)30059-3)
- Valero-Solís, S., Granero, R., Fernández-Aranda, F., Steward, T., Mestre-Bach, G., Mallorquí-Bagué, N., Martín-Romera, V., Aymamí, N., Gómez-Peña, M., del Pino-Gutiérrez, A., Baño, M.,

- & Jiménez-Murcia, S. (2018). The contribution of sex, personality traits, age of onset and disorder duration to behavioral addictions. *Frontiers in Psychiatry*. <https://doi.org/10.3389/fpsy.2018.00497>
- Volkow, N. D., Wang, G. J., Fowler, J. S., Tomasi, D., & Baler, R. (2011). Food and drug reward: Overlapping circuits in human obesity and addiction. *Current Topics in Behavioral Neurosciences*. https://doi.org/10.1007/7854_2011_169
- Wolz, I., Hilker, I., Granero, R., Jiménez-Murcia, S., Gearhardt, A. N., Dieguez, C., Casanueva, F. F., Crujeiras, A. B., Menchón, J. M., Fernández-Aranda, F. (2016). “Food Addiction” in patients with eating disorders is associated with negative urgency and difficulties to focus on long-term goals. *Frontiers in Psychology*. <https://doi.org/10.3389/fpsyg.2016.00061>
- Zakiniacz, Y., Cosgrove, K. P., Mazure, C. M., & Potenza, M. N. (2017). Does telescoping exist in male and female gamblers? Does it matter? *Frontiers in Psychology*. <https://doi.org/10.3389/fpsyg.2017.01510>
- Zakiniacz, Y., & Potenza, M. N. (2018). Gender-related differences in addiction: A review of human studies. *Current Opinion in Behavioral Sciences*. <https://doi.org/10.1016/j.cobeha.2018.08.004>

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Authors and Affiliations

Mikel Etxandi¹ · Isabel Baenas^{1,2} · Lucero Munguía^{1,3} · Gemma Mestre-Bach^{1,4} · Roser Granero^{2,5} · Mónica Gómez-Peña¹ · Laura Moragas¹ · Amparo del Pino-Gutiérrez^{2,6} · Ester Codina¹ · Bernat Mora-Maltas¹ · Eduardo Valenciano-Mendoza¹ · Marc N. Potenza^{7,8,9,10} · Ashley N. Gearhardt¹¹ · Fernando Fernández-Aranda^{1,2,3} · Susana Jiménez-Murcia^{1,2,3} 

Mikel Etxandi
mikeletxandi@gmail.com

Isabel Baenas
ibaenas@bellvitgehospital.cat

Lucero Munguía
laarcreed_lm@hotmail.com

Gemma Mestre-Bach
gemma.mestre.bach@gmail.com

Roser Granero
Roser.granero@uab.cat

Mónica Gómez-Peña
monicagomez@bellvitgehospital.cat

Laura Moragas
lmoragas@bellvitgehospital.cat

Amparo del Pino-Gutiérrez
adelpino@ub.edu

Ester Codina
ecodina@bellvitgehospital.cat

Bernat Mora-Maltas
bmora@idibell.cat

Eduardo Valenciano-Mendoza
edu.valenciano@gmail.com

Marc N. Potenza
marc.potenza@yale.edu

Ashley N. Gearhardt
agearhar@umich.edu

Fernando Fernández-Aranda
ffernandez@bellvitgehospital.cat

- ¹ Department of Psychiatry, Bellvitge University Hospital-IDIBELL, Feixa Llarga s/n 08907. Hospitalet del Llobregat, Barcelona, Spain
- ² Ciber Fisiopatología Obesidad y Nutrición (CIBEROBn), Instituto de Salud Carlos III, Madrid, Spain
- ³ Department of Clinical Sciences, School of Medicine, University of Barcelona, Barcelona, Spain
- ⁴ Universidad Internacional de la Rioja, La Rioja, Spain
- ⁵ Department of Psychobiology and Methodology, Autonomous University of Barcelona, Barcelona, Spain
- ⁶ Department of Public Health, Mental Health and Perinatal Nursing, School of Nursing, University of Barcelona, Barcelona, Spain
- ⁷ Department of Psychiatry and Child Study Center, Yale University School of Medicine, New Haven, Connecticut, USA
- ⁸ Connecticut Council on Problem Gambling, Wethersfield, Connecticut, USA
- ⁹ Connecticut Mental Health Center, New Haven, Connecticut, USA
- ¹⁰ Department of Neuroscience, Yale University, New Haven, Connecticut, USA
- ¹¹ Department of Psychology, University of Michigan, Ann Arbor, USA