


## Meditation, Health and Scientific Investigations: Review of the Literature

Cynthia Vieira Sanches Sampaio<sup>1,2</sup>  · Manuela Garcia Lima<sup>1</sup> · Ana Marice Ladeia<sup>1</sup>

Published online: 25 February 2016  
© Springer Science+Business Media New York 2016

**Abstract** A growing number of people are seeking health recovery treatments with a holistic approach to the human being. Meditation is a mental training capable of producing connection between the mind, body and spirit. Its practice helps people to achieve balance, relaxation and self-control, in addition to the development of consciousness. At present, meditation is classified as a complementary and integrative technique in the area of health. The purpose of this review of the literature was to describe what meditation is, its practices and effects on health, demonstrated by consistent scientific investigations. Recently, the advances in researches with meditation, the discovery of its potential as an instrument of self-regulation of the human body and its benefits to health have shown that it is a consistent alternative therapy when associated with conventional medical treatments.

**Keywords** Health · Integrative treatment · Meditation · Spirituality · Well-being

In the East, meditation is an ancient practice, developed by various traditions to broaden consciousness and seek health. Ancient texts, with reference to the Indian Ayurvedic system of medicine, demonstrate that this practice formed part of medical procedures used for the recovery and maintenance of health that go back further than 3000 years ago (Carneiro 2009), whereas, in the West, the tradition of meditation is usually linked to religious experiences, especially Christian with a Catholic or Protestant connotation. However, as from the 1970s, Western groups and movement in contact with Oriental experiences opened themselves to understanding and make use of meditation—without a

---

✉ Cynthia Vieira Sanches Sampaio  
cysampaio@gmail.com

<sup>1</sup> Bahiana School of Medicine and Public Health, Bahia Foundation for the Development of Sciences, FBDC, Salvador, Bahia, Brazil

<sup>2</sup> Centro Médico Iguatemi, Av. Tancredo Neves, 805-A, Sala 301, Caminho das Árvores, 41820-021, Salvador, Bahia, Brazil

specifically religious nature—as a resource for the practice of a healthy lifestyle and even for the cure of psychological behavior patterns, which of themselves act on the general health of each individual.

The aim of the present study is to evaluate meditation as a useful and necessary resource for seeking and practicing a healthy lifestyle, based on quieting the mind, reducing stress and achieving inner harmony. This implies the better relationship of an individual with the world outside.

In the discourse that follows in this article, a description is given of what meditation is, its practices and effects on health, revealed by consistent scientific investigations.

## Meditation

The word meditation, which expresses the practice of meditating, is derived from Sanskrit, traditional language of India, from the word *dhyana*, which means attention, contemplation. The exercise of meditation is processed by a large variety of activities that range from techniques to promote relaxation through to exercises performed with objectives that are broader in scope, such as intensification of the feeling of wellbeing (Lutz et al. 2008). It also includes breathing techniques, repetition of sounds and/or observation of the process of thought to focus attention and promote a state of self-awareness and inner calm (Canter 2003).

There are different practices of meditation—those that specifically go through religious traditions, those that seek connection with spirituality without religious connotation and those that propose to be a purely mental training, unconnected to a spiritual proposal (Menezes et al. 2011).

The central point common to the numerous practices consists of temporarily withdrawing attention from the outside world and thoughts related to it, in order to focus on the chosen theme of meditation (Servan-Shreiber 2008). The common key to all the practices is silencing and harmonization of thoughts and judgments bubbling within each person. Meditation is based on control of attention.

Studies and researches about meditation in the area of health have led to the need for an operational definition for this term. A group of researchers established some elements as a parameter for a procedure to be characterized as meditation, composing a definition accepted in the scientific medium. These elements are as follows: the use of a clearly defined technique; production of muscle and psychic relaxation with reduction in logical thought; it must necessarily be a self-induced state; and developing the capacity to maintain the focus of attention on a certain point that functions as anchor (Cardoso et al. 2004).

For the majority of researchers, there are two general types of meditation: concentration meditation and mindfulness meditation (Lutz et al. 2008). The former emphasizes the need for attention focused on an object, and to sustain this process until the mind attains quieting of thoughts. Its continuous practice produces relaxation and mental clarity (Krisanaprakornkit et al. 2006). Mindfulness meditation is directed toward the opening of perception of contents that emerge in the mind, without the individual judging or reacting to his/her own thoughts and emotions. This technique favors the undoing of previous conditioned behavioral patterns, making it possible for the individual to create new strategies for coping with the events of life (Krisanaprakornkit et al. 2006). There is also a third type of meditation, denominated by some authors as the contemplative type, which

integrates the two previously mentioned types, stimulating both the capacity to focus and to open oneself (Menezes et al. 2009).

## **Meditation Recognized by Scientific Investigation as a Resource for a Healthy Life and Cure**

The scientific researches listed revealed how much meditation has to offer for attaining a healthy lifestyle and as a resource to seek cure in an integrative practice of medicine, in which all the dimensions of the human being are recognized and taken care of.

Despite its practice, each type of meditation identified above, in its own way, offers a possibility of entering into a state of inner coherence, which favors the integration of all the biological rhythms and harmonizing functions of the body (Servan-Shreiber 2008). Apart from this, the aim of the different types of meditative practices is to change the flow of thoughts, generating new patterns of behavior and awareness (Danucalov and Simões 2006).

When producing an altered state of consciousness, meditation facilitates the metacognitive mode of thought, making it possible to attain the perspective of behavioral cognitive benefits and healthy psychological functioning (Krisanaprakornkit et al. 2006; Menezes et al. 2011; Willis 1979).

The initial researches with meditation investigated its effects on human physiology. Robert Keith Wallace, of the University of California, one of the pioneers of this type of investigation, in 1970 conducted a classical study that was published in the prestigious journal *Science* (Wallace 1970). This study found that during meditation, a reduction occurs in oxygen consumption and cardiac frequency, and an increase in galvanic resistance of the skin. Furthermore, the electroencephalogram showed a predominance of alpha waves, thus it was concluded that these physiological alterations were compatible with changes in autonomic activity, indicative of reduction in sympathetic activities, and therefore, meditation could have applications in clinical medicine (Wallace 1970).

In the following year, Wallace et al. (1971) conducted a new clinical trial with a larger number of subjects and with a similar objective. In addition to confirming the previous results, they identified a reduction in respiratory frequency and elimination of CO<sub>2</sub>, as well as a reduction in pH and arterial lactate (Wallace et al. 1971).

These findings encouraged other researchers and showed that during the practice of meditation, the following occurred: diminishment of cardiac frequency, slower respiration, reduced electrical conductivity of the skin, reduction in blood lactate, variations in EEG frequency. In addition, it was verified hormonal variations, modifications in the concentrations of innumerable neurotransmitter substances, reduction in body temperature, alteration in the senses and perceptions, among others, indicating an increase in parasympathetic activity (Danucalov and Simões 2006).

These physiological changes characterized a state of relaxation and enabled some understanding to be gained about how meditation operates in the body. In addition, the relaxation and reduction in stress and anxiety, which are affirmed as results of meditation, may bring about prophylactic and therapeutic benefits to health (Canter 2003).

Later researchers, with the use of sophisticated equipment, in studies that used a variety of styles of meditation, showed that in a general manner, this practice could activate areas of the brain associated with well-being (Davidson 2003), regulation of the emotions (Tang et al. 2009) and the capacity to sustain attention (Lutz et al. 2009).

In addition, the scientific investigations signaled that the brain dynamics in those who meditate for many years is different from the brain dynamics of those who do not meditate (Lutz et al. 2004). Furthermore, they indicated that irrespective of the technique used, it is a mental training that acts on cerebral functions affecting the manner in which the stimuli are processed and perceived, thereby helping to undo conditioned behaviors (Allen et al. 2012; Kozasa et al. 2012; Newberg et al. 2001; Singh et al. 2012; Slagter et al. 2007).

At the same time, other researches revealed that meditation, in addition to producing neuroendocrine and neurochemical effects, which modified the brain activity and metabolism of the individual, may be associated with structural alterations in areas of the brain. A study with magnetic resonance (Lazar et al. 2005) compared the brain of experienced meditators with the brain of non-meditators. She found significant difference in the thickness of the cerebral cortex of the meditators, which was larger in the insula and prefrontal cortex, brain regions in which attention and the emotions are concentrated.

This study suggested that meditation might generate changes in the brain favoring improvement in the person's cognitive and emotional functions, in addition to retarding brain aging. However, because it was a cross-sectional study, it was not possible to demonstrate causality.

More recently, a controlled longitudinal study (Hölzel et al. 2011), aimed to investigate changes in the brain through the practice of meditation, proved that this activity increased the gray matter in certain regions of the brain, altering its structure.

During the study, sessions of magnetic resonance were performed of the brain of all the participants before and after the period of practices. The initial exams showed no difference between the intervention and control groups. However, the resonance exams performed after 30 min of meditative practice daily, for 8 weeks, showed increase in the concentration of gray matter of the left hippocampus, posterior cingulate cortex, temporoparietal junction and cerebellum in those who practiced meditation, in comparison with those who did not meditate. These regions of the brain are associated with processes of learning, memory, regulation of emotions and empathic capacity.

In spite of the innumerable researches conducted over the last few decades about the effects of meditation on the human body, yet little is known about the neural and biologic mechanisms associated with this practice. Therefore, the complex mental process that involves meditation is a fascinating field of research in progress.

However, as previously mentioned, the different studies have demonstrated that meditation is an active mental training, capable of modifying the functioning of the brain and mind, favoring the attentional skills, cognitive capacity and emotional regulation. This enables the person to respond better to day-to-day stressor stimuli. In addition, they have confirmed the understanding that the brain is modified according to the experiences of the individual, and support the notion that the practice of meditation may have a long-term effect, producing lasting changes by acting on brain plasticity.

Nowadays, it is known that mental attitudes may influence the body, generating health or diseases, to the extent that they balance or unbalance the release of innumerable hormones (Danucalov and Simões 2006). Studies have confirmed the link between mental processes and autonomic aspects relative to the functioning of the nervous system, which cause the creation of an entirely new discipline, known as psychoneuroimmunology (Lutgendorf and Costanzo 2003; Marques-Deak and Sternberg 2004; McnCain and Gray 2005). According to this science, a chronically altered mind may produce negative effects on the homeostatic mechanisms of the body, facilitating the appearance of somatic diseases (Humphreys and Lee 2009; Juster et al. 2010; Oppermann 2002). Thus, it emphasizes the

importance of the psychological aspects to treatment or cure (Basinski et al. 2013; Bertini et al. 2012; Pace and Heim 2011).

Therefore, meditation becomes a practice that is self-regulatory of the body and mind, with the potential of helping a person develop the capacity to obtain some degree of control over the psychophysiological autonomic processes. In addition, it is a means to re-establish the organic homeostatic mechanisms, and favors cognitive and behavioral changes that generate health and psychological well-being (Bogart 1991; Danucalov and Simões 2006; Davidson and Goleman 1977).

According to Doctor Herbert Benson (1998), of Harvard University, who studies the body–mind connection, emotional factors have great importance in the origin and evolution of innumerable diseases. A researcher since the 1970s about the effects of meditation on human physiology, he affirms that persons may learn to use the mind to combat stresses that generate physical diseases and promote health with better response to treatments (Benson and Stark 1998).

However, in spite of the innumerable benefits pointed out, it is important to emphasize that meditation may cause adverse effects, such depersonalization and derealization and is therefore not indicated for individuals with borderline or psychotic conditions (Canter 2003; McGee 2008).

Over the last 30 years, meditation has been the target of innumerable researches in the area of physical and mental health. In general, the studies suggest that the practice of meditation, in addition to promoting self-knowledge and spiritual growth, may be an important complementary tool to conventional treatment of different clinical conditions, potentiating the process of cure and well-being of patients.

Researches have indicated that meditation may help to strengthen the immune system. A study conducted in a work environment with healthy employees, vaccinated all the individuals against influenza after 8 weeks of training in meditation. The researchers found significant increases in the anterior activation on the left side of the frontal cortex, a pattern previously associated with positive affections, in addition to significant increases in the markers of antibodies for the influenza vaccine among mediators when compared with non-meditators. This suggested that the magnitude of the increase in the activation of the left side predicted the magnitude of the increase in the immunological response to the influenza vaccine (Davidson 2003).

Another study verified the effect of meditation on the T CD4+ lymphocytes of adults infected with HIV-1. The T CD4+ lymphocytes are the main type responsible for cell immunity, and in patients with HIV-1 may reach a very low level, making infections more frequent and difficult to treat. The measurements made after a period of 8 weeks training in meditation showed that the participants of the control group presented declines in the T CD4+ lymphocyte counts, while the counts among the participants in the meditation program remained unaltered from the beginning until the post-intervention measurement. This effect was irrespective of the use of the antiretroviral medication. Additional analyses demonstrated that with adherence to the meditation program, the results indicated that a damping effect on the reduction of T CD4+ lymphocytes occurred, suggesting that the practice may help the decline in these lymphocytes in adults infected with HIV-1 (Creswell et al. 2009).

In this same direction, a study conducted in a school health center with a sample of patients without any known autoimmune disturbance, investigated whether the increase in psychosocial well-being after 8 weeks of training in meditation would be associated with the corresponding alterations in the immunological activity markers. Analysis of the data demonstrated that the positive improvement in psychological well-being generated by

meditation was associated with an increase in NK cytolytic activity, and with reduction in the levels of C-Reactive Protein (Fang and Reibel 2010).

The practice of meditation also helps to regulate arterial pressure and increase cardiovascular efficiency. One study followed up individuals with coronary disease for a mean period of 5.4 years and randomized the patients for inclusion in a meditation or health education program. The study concluded that meditation significantly reduced the risk for mortality, infarction of the myocardium and cerebral vascular accident in patients with coronary disease. These alterations were associated with lower arterial pressure and psychosocial stress factors, and therefore, the practice of meditation could be clinically useful in the secondary prevention of cardiovascular disease (Schneider and Grim 2012).

In another study conducted with patients hospitalized in two health centers due to heterogeneous diseases, one group received treatment that included daily meditations, and the group that served as control, received only traditional treatment. Both groups were evaluated before and after a mean period of 8 days of treatment. The results of the measurements evaluated between the two groups suggested that the meditation treatment was associated with significant improvements in the quality of life, reduction of anxiety and control of arterial pressure (Chung et al. 2012).

Meditation has also shown an association with the quality of sleep and cognitive function in elderly persons. A randomized controlled clinical trial evaluated elderly persons who presented a reduction in sleep quality. The sleep quality and cognitive function of the two groups were measured before the training and at the end of time intervals of 3, 6 and 12 months using four validated questionnaires. The results indicated a significant improvement in sleep quality and cognitive functions of those who meditated when compared with those who did not meditate (Sun et al. 2013).

Furthermore, meditation improved the emotional condition of patients with cancer. Women recently diagnosed with breast cancer in the initial stage, who did not receive chemotherapy, participated in a non-randomized controlled study, which used a meditation program of 8 weeks. The results showed that the women who participated in the meditation program had reduced the cortisol levels, improved quality of life and increased efficacy of coping with the disease in comparison with the control group (Witek-Janusek et al. 2008).

Another study corroborated these results, evaluating 229 women after surgery, chemotherapy and radiotherapy for stage 0 to III breast cancer. The patients were randomly assigned to a program of 8 weeks meditation, or only to standard treatment. Evaluations as regards mood, well-being and quality of life related to the endocrine system and breast were made at time intervals of 0, 8 and 12 weeks. The results showed evidence that the meditation program associated with the standard treatment helped to alleviate the physical and emotional adverse effect of the medical treatments, including the endocrine treatments, and these effects were maintained in the long term (Hoffman et al. 2012).

Yet another study investigated the effects of three short-term interventions on sleep and psychological symptoms of comorbidity in cancer survivors. All the interventions, two body–mind interventions and one educational interventions for the control group, consisted of three sessions lasting 2 h, once a week for three consecutive weeks. The patients were randomized and evaluated as regards sleep, quality of life, stress, depression, self-pity and well-being. The study suggested that the two body–mind interventions were effective for post-treatment care of sleep disturbances in these patients, and could be an ideal vehicle for the management of multiple coexistent symptoms in cancer survivors (Nakamura et al. 2013).

Similarly, meditation has shown benefits in patients who had undergone organ transplants. A randomized controlled clinical trial evaluated the efficacy of an 8-week

meditation program in the reduction of symptoms of anxiety, depression and sleep disturbances in transplant recipients. Measurements were taken at the beginning of the study, and at time intervals of 8 weeks, 6 months and 1 year. The results showed that in the group that meditated there was a reduction in the symptoms evaluated and improvement in quality of life in comparison with the control group and that the benefits were maintained throughout the course of 1 year (Gross and Kreitzer 2010).

A systematic review evaluated the efficacy of meditation in the treatment of diseases. Studies with a healthy population were not included. Twenty randomized clinical trials with a total of 958 individuals met the criteria. The results supported the safety and potential efficacy of meditative practices in the treatment of epilepsy, symptoms of the premenstrual syndrome and menopause. Benefit was also demonstrated for mood, non-psychotic anxiety disturbances, autoimmune disease and emotional disturbances in neoplastic diseases (Arias and Steinberg 2006).

A meta-analysis evaluated the effects of a meditation program on stress in healthy persons. From these studies included, with a total of 671 individuals, it was shown that the meditation program reduced ruminative thought and trace of anxiety and increased empathy and self-pity, and was capable of reducing stress levels in healthy persons (Chiesa and Serretti 2009).

Another meta-analysis analyzed the effects of a meditation program on depression, anxiety and psychological anguish in adults with different chronic somatic diseases. Eight randomized controlled studies were included, with a total of 667 individuals, and found a positive effect of meditation on these symptoms (Bohlmeijer et al. 2010).

Furthermore, a systematic review with meta-analysis focused specifically on the efficacy of meditation for anxiety. A total of 36 randomized clinical trials were included in the meta-analysis, with a total of 2466 observations. In 25 studies, statistically superior results were reported in the meditation group in comparison with the control. The evaluation demonstrated the efficacy of meditation therapies in the reduction of symptoms of anxiety. However, it pointed out that the majority of studies measured only the improvement in the symptoms of anxiety, but not disturbances of anxiety as clinically diagnosed (Chen and Berger 2012).

A recent meta-analysis analyzed the efficacy of meditation programs for psychological stress and well-being. The study included 47 clinical trials with 3515 participants suffering from anxiety, depression, chronic pain, cancer and cardiovascular diseases, among others. The results showed that meditation might reduce the multiple negative dimensions of psychological stress, mainly having an effect on anxiety, depression and pain. In conclusion, the authors suggested that clinical physicians must be prepared to speak to their patients about the role of meditation in mental health and stress-related behaviors (Goyal et al. 2014).

Based on this body of scientific evidence, various hospitals and psychotherapy consulting rooms in various parts of the world have associated the practice of meditation with conventional treatments, in a complete and integral approach to the patient's process of cure (Davidson 2003; Hölzel et al. 2011).

In Brazil, following this world trend, since 2006 meditation has become part of the program of National Policy of Integrative and Complementary Practices of the national health service—SUS and has been defined as a procedure that focuses attention in a non-analytical or discriminative manner, promoting favorable alterations in mood and cognitive performance (Ministério da Saúde 2008).

**Table 1** Scientific research showing evidence of association between meditation and benefit in human body

Author/ year	Type of study	Population	N	Type of control	Type of meditation	Duration	Outcomes analyzed	Instruments for evaluating outcome
Wallace (1970)	Open-label uncontrolled clinical trial	Healthy university students with previous experience of meditation	15	—	Transcendental	30 min	O <sup>2</sup> consumption (↓) Cardiac frequency (↓) Skin resistance (↑) Electrical brain activity (↑ $\alpha$ waves)	Spirometry Blood gasometry Electrocardiography Galvanometry Electroencephalography
Wallace et al. (1971)	Open-label controlled non- randomized clinical trial	Healthy subjects with previous experience of meditation	36	Single- subject (subject as his/her own control)	Transcendental	20–30 min	O <sup>2</sup> consumption (↓) Cardiac frequency (↓) Skin resistance (↑) Electrical brain activity (↑ $\alpha$ waves) O <sup>2</sup> Elimination (↓) Respiratory quotient (N) Respiratory frequency (N) Ventilation minute (N) Arterial pressure (N) pH arterial (↓) PCO <sup>2</sup> —arterial (N) PO <sup>2</sup> —arterial (N) Arterial lactate (↓) Rectal temperature (N)	Spirometry Blood gasometry Electrocardiography Galvanometry Electroencephalography
Davidson (2003)	Open-label controlled randomized clinical trial	Healthy, right- handed subjects, without previous experience	41	Passive	<i>Mindfulness</i>	1 h/day 6 day/week 8 weeks	Anxiety (↓) Positive affection (N) Negative affection (↓) Electrical brain activity (↑ left anterior activation) Antibodies for influenza (↑)	<i>Spielberger State-Trait Anxiety Inventory Positive and Negative Affect Scale</i> Electroencephalogram



**Table 1** continued

Author/ year	Type of study	Population	N	Type of control	Type of meditation	Duration	Outcomes analyzed	Instruments for evaluating outcome
Tang et al. (2009)	Open-label controlled randomized clinical trial	Healthy university students, without previous experience	86	Active (relaxation)	Integrative body–mind training	20 min/day 5 days	Cardiac frequency (↓) Variability of CF (↑) Respiratory frequency (↓) Respiratory amplitude (↑) Skin conductance (↓) Electrical brain activity (↑ $\theta$ waves in ACC) Activ. metabolic brain activity (N- overall) (↑ in ACC)	Electroencephalography Single-photon emission computed tomography (SPECT)
Lutz et al. (2009)	Open-label controlled non-randomized clinical trial	Healthy subjects, meditators and non-meditators	40	Active (lesson and meditation for 20 min)	Vipassana	10–12 h/day 3 months	Electrical brain activity (↑ $\theta$ waves in anterior) Time of reaction (↓) Variability in reaction time (↓) Target detection accuracy rate (↑) Attention (↑) Electrical brain activity (↑ $\gamma$ )	Electroencephalography <i>Attention Blink Task</i> <i>Dichotic Listening Task</i>
Lutz et al. (2004)	Cross-sectional observational study	Experienced Buddhist monks and health students without previous experience	18	–	Buddhist ( <i>unconditional loving-kindness and compassion</i> )	–	Electrical brain activity (↑ $\gamma$ )	Electroencephalography
Newberg et al. (2001)	Open-label uncontrolled clinical trial	Healthy subjects with previous experience of meditation	8	–	Tibetan Buddhist	1 h	Activ. metabolic brain activity (↑ cingulate gyrus, prefrontal cortex, inferior frontal and orbital and dorsolateral cortex and thalamus) (verifier terminology por favor)	Single-photon emission computed tomography (SPECT)

Table 1 continued

Author/ year	Type of study	Population	N	Type of control	Type of meditation	Duration	Outcomes analyzed	Instruments for evaluating outcome
Slagter et al. (2007)	Open-label controlled non- randomized clinical trial	Healthy subjects, mediators and non- mediators	40	Active (lesson and meditation for 20 min)	Vipassana	10–12 h/day 3 months	Target detection accuracy (↑) Attention (↑) Allocation of brain resources for primary target detection (↓)	Electroencephalography <i>Attention Blink Task</i>
Kozasa et al. (2012)	Cross-sectional observational study	Healthy subjects, mediators and non- mediators	39	–	<i>Focus Attention</i> and/or <i>Open</i> <i>Monitoring</i>	–	Attention (N) Impulse control (N) Brain activity (↓ in right frontal medial, mid temporal, precentral and post-central gyri and in lentiform nucleus)	Functional nuclear magnetic resonance
Allen et al. (2012)	Open-label controlled randomized clinical trial	Healthy, right- handed subjects, without previous experience	61	Active (Group Lecture)	<i>Mindfulness</i>	2 h/week 6 weeks	Consciousness of errors (N) Signal dependent on blood oxygen level (↑ in left prefrontal dorsolateral cortex)	<i>Error-Awareness Task</i> (EAT) Functional nuclear magnetic resonance

**Table 1** continued

Author/ year	Type of study	Population	N	Type of control	Type of meditation	Duration	Outcomes analyzed	Instruments for evaluating outcome
Singh et al. (2012)	Open-label controlled non-randomized clinical trial	Healthy university students without previous experience	34	Single-subject (subject as his/her own control)	Not related	Phase 1: Daily for 1 month phase 2: 15 min	Cardiac frequency (N) Skin resistance (↓) salivary cortisol (N) Acute stress (↓) Memory (N) Time of reaction (↓) Intelligence quotient (↑)	Electrocardiography Galvanometry Salivary cortisol <i>Stanford Acute Stress Reaction</i> <i>Questionnaire (SASRQ)</i> <i>Cohen Perceived Stress Scale</i> <i>Sternberg memory test (MEMSCAN)</i> <i>Stroop color interference test</i> Intelligence quotient ( <i>Wechsler Adult Intelligence Scale-Performance Scale</i> ) <i>N. S. Schutte Emotional Intelligence Scale</i>
Lazar et al. (2005)	Cross-sectional observational study	Healthy subjects, meditators and non-meditators	35	–	<i>Insight</i>	–	Cortical thickness (↑ in prefrontal cortex and right anterior insula)	Nuclear magnetic resonance
Hölzel et al. (2011)	Open-label controlled non-randomized clinical trial	Healthy right-handed subjects, meditators and non-meditators	35	Passive	<i>Mindfulness</i>	2.5 h/week 8 weeks	Concentration of gray matter (↑ left hippocampus, posterior cingulate cortex, temporoparietal junction and cerebellum)	Nuclear magnetic resonance

Table 1 continued

Author/ year	Type of study	Population	N	Type of control	Type of meditation	Duration	Outcomes analyzed	Instruments for evaluating outcome
Creswell et al. (2009)	Uni-blind controlled randomized trial	HIV+ Subjects	48	Active (1- day seminar)	<i>Mindfulness</i>	2 h/week 8 weeks	Lymphocyte counts TCD4+ (↑)	–
Fang and Reibel (2010)	Open-label controlled non- randomized clinical trial	Healthy subjects	24	Single- subject (subject as his/her own control)	<i>Mindfulness</i>	2.5 h/week 8 weeks	Anxiety (↓) Quality of life (↑) Cytolytic cell activity <i>Natural-Killer</i> (↑) C-Reactive protein (↓)	<i>Brief Symptom Inventory-18 Medical Outcomes Survey Short-Form Health Survey</i>
Schneider and Grimm (2012)	Uni-blind controlled randomized trial	Negroes with coronary artery disease	201	Active (Health Education)	Transcendental	20 min 2 times/day 5.4 years	Composed of mortality by any cause whatever (AIM or CVA) (↓) Composed of cardiovascular death, revascularization or cardiovascular hospitalization (↓) Systolic blood pressure (↓) Psychosocial stress factors (↓)	<i>CESD Scale for depression Cook-Medley Hostility Inventory Anger Expression scale</i>
Chung et al. (2012)	Prospective observational study	Hospitalized patients	129	Passive	Sahaja Yoga	1 h 2 times/day 8.1 days	Quality of life (↑) Anxiety (↓) Arterial pressure (↓)	Abbreviated quality of life evaluation instrument (WHOQOL-BREF) WHOQOL-SRPB (Spirituality, Religion and Personal Beliefs) <i>Clinical Anxiety Scale (CAS)</i>

**Table 1** continued

Author/ year	Type of study	Population	N	Type of control	Type of meditation	Duration	Outcomes analyzed	Instruments for evaluating outcome
Sun et al. (2013)	Open-label controlled randomized clinical trial	The elderly (>60 years)	80	Active (sleep hygiene only)	Self-relaxation and sleep hygiene	30 min 3 times/day 1 year	Quality of sleep (↑) Cognitive functions (↑)	<i>Pittsburgh Sleep Quality Index</i> <i>Epworth Sleepiness Scale</i> Mini Mental State Exam <i>Wechsler Memory Scale</i>
Witek-Janusek et al. (2008)	Open-label controlled non-randomized clinical trial	Women with early breast cancer	75	Passive	<i>Mindfulness</i>	2.5 h/week 8 weeks	Quality of life (↑) Coping (↑) Cortisol (↓) IFN-gamma (↑) IL-4, IL-6, IL-10 (↓) Cytolytic cell activity <i>Natural-Killer</i> (↑)	<i>Quality of Life Index Cancer Vers. III</i> <i>Jalowiec Coping Scale</i> (JCS) <i>Mindful Attention Awareness Scale</i> (MAAS)
Hoffman et al. (2012)	Uni-blind controlled randomized trial	Women with Stage 0–III breast cancer	229	Passive	<i>Mindfulness</i>	2.5 h/week 8 weeks	Mood (↑) Wellbeing (↑)	<i>Profile of Mood States</i> (POMS) <i>Functional Assessment of Cancer Therapy—Breast</i> (FACT-B) <i>Functional Assessment of Cancer Therapy—Endocrine Symptoms</i> (FACT-ES) <i>WHO five-item well-being questionnaire</i> (WHO-5)
Nakamura et al. (2013)	Open-label controlled randomized clinical trial	Survivors of any type of cancer with sleep disturbance	57	Active (sleep hygiene)	<i>Mindfulness e Mind–Body Bridging</i> (MBB)	2 h/week 3 weeks	Sleep disturbance (↓)	<i>Medical Outcomes Study Sleep Scale</i>

Table 1 continued

Author/ year	Type of study	Population	N	Type of control	Type of meditation	Duration	Outcomes analyzed	Instruments for evaluating outcome
Gross and Kretzler (2010)	Open-label controlled randomized clinical trial	Solid organ transplant recipients	138	Active (Health Education)	<i>Mindfulness- Based Stress Reduction (MBSR)</i>	2.5 h/week 8 weeks	Anxiety (↓) Depression (↓) Quality of sleep (↑)	<i>State-Trait Anxiety Inventory—State Version (STAI) Center for Epidemiological Studies -Depression Scale (CES-D) Pittsburgh Sleep Quality Index (PSQI)</i>

\*(N) without alteration, (↑) increase, (↓) reduction

Table 1 presents a summary of the characteristics of the studies that were included in order to demonstrate the associations between the practice of meditation and its effects on the human body.

## Conclusion

The growing scientific interest in meditation showed that this practice is a mental training associated with lasting changes in cognition and emotion. Different methodologies have shown that this association and its impact on self-regulation are well established. However, the changes in physical and mental health reported in response to meditation need to be better explored, and their impact on the brain requires further studies.

Nevertheless, the body of scientific evidence that has already been obtained with respect to the benefits arising from meditation for the recovery of health encourage a change to more integrated forms of treatment, and therapeutic interventions that incorporate the practice of meditation are becoming increasingly popular.

Health begins to be understood as ample well-being that involves interaction between the physical, emotional, mental and spiritual aspects of the individual. This new paradigm has a concept of the human being as an indivisible whole, in which all its dimensions need to be cared for. It emphasizes the need for a multidisciplinary approach to the patient's cure.

Meditation, by supporting the individual to develop his/her own resources for self-regulation, is at present accepted within a context of integrative treatment in the area of health.

**Acknowledgments** Sampaio CVS was supported by a master's degree scholarship from Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES).

## References

- Allen, M., Dietz, M., Blair, K. S., van Beek, M., Rees, G., Vestergaard-Poulsen, P., et al. (2012). Cognitive-affective neural plasticity following active-controlled mindfulness intervention. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 32(44), 15601–15610.
- Arias, A., & Steinberg, K. (2006). Systematic review of the efficacy of meditation techniques as treatments for medical illness. *Journal of Alternative and Complementary Medicine (New York, N.Y.)*, 12(8), 817–832.
- Basiński, A., Stefaniak, T., & Stadnyk, M. (2013). Influence of religiosity on the quality of life and on pain intensity in chronic pancreatitis patients after neurolytic celiac plexus block: Case-controlled study. *Journal of Religion and Health*, 52(1), 276–284.
- Benson, H., & Stark, M. (1998). *Medicina espiritual: o poder essencial da cura*. Rio de Janeiro: Campus.
- Bertini, M., Conti, C., & Fulcheri, M. (2012). Psychoneuroimmunology and health psychology: Inflammation and protective factors. *Journal of Biological Regulators and Homeostatic Agents*, 27, 637–645.
- Bogart, G. (1991). The use of meditation in psychotherapy: A review of the literature. *American Journal of Psychotherapy*, 45(3), 383–412.
- Bohlmeijer, E., Prenger, R., Taal, E., & Cuijpers, P. (2010). The effects of mindfulness-based stress reduction therapy on mental health of adults with a chronic medical disease: A meta-analysis. *Journal of Psychosomatic Research*, 68(6), 539–544.
- Canter, P. (2003). The therapeutic effects of meditation. *BMJ*, 326, 1049–1050.
- Cardoso, R., de Souza, E., Camano, L., & Leite, J. R. (2004). Meditation in health: An operational definition. *Brain Research Protocols*, 14(1), 58–60.
- Carneiro, D. (2009). *Ayurveda: saúde e longevidade na tradição milenar da Índia*. São Paulo: Pensamento.

- Chen, K., & Berger, C. (2012). Meditative therapies for reducing anxiety: A systematic review and meta-analysis of randomized controlled trials. *Depression and Anxiety*, 29(7), 545–562.
- Chiesa, A., & Serretti, A. (2009). Mindfulness-based stress reduction for stress management in healthy people: A review and meta-analysis. *The Journal of Alternative and Complementary Medicine*, 15(5), 593–600.
- Chung, S.-C., Brooks, M. M., Rai, M., Balk, J. L., & Rai, S. (2012). Effect of Sahaja yoga meditation on quality of life, anxiety, and blood pressure control. *The Journal of Alternative and Complementary Medicine*, 18(6), 589–596.
- Creswell, J., Myers, H., Cole, S., & Irwin, M. (2009). Mindfulness meditation training effects on CD4+ T lymphocytes in HIV-1 infected adults: A small randomized controlled trial. *Brain, Behavior, and Immunity*, 23(2), 184–188.
- Danucalov, M., & Simões, R. (2006). *Neurofisiologia da meditação*. São Paulo: Phorte.
- Davidson, R. (2003). Alterations in brain and immune function produced by mindfulness meditation. *Psychosomatic Medicine*, 66(1), 148–149.
- Davidson, R., & Goleman, D. (1977). The role of attention in meditation and hypnosis: A psychobiological perspective on transformations of consciousness. *International Journal of Clinical and Experimental Hypnosis*, 25(4), 291–308.
- Fang, C., & Reibel, D. (2010). Enhanced psychosocial well-being following participation in a mindfulness-based stress reduction program is associated with increased natural killer cell activity. *The Journal of Alternative and Complementary Medicine*, 16(5), 531–538.
- Goyal, M., Singh, S., Sibinga, E. M. S., Gould, N. F., Rowland-Seymour, A., Sharma, R., et al. (2014). Meditation programs for psychological stress and well-being: A systematic review and meta-analysis. *JAMA Internal Medicine*, 174(3), 357–368.
- Gross, C., & Kreitzer, M. (2010). Mindfulness-based stress reduction for solid organ transplant recipients: A randomized controlled trial. *Alternative Therapies Health Medicine*, 16(5), 30–38.
- Hoffman, C. J., Ersser, S. J., Hopkinson, J. B., Nicholls, P. G., Harrington, J. E., & Thomas, P. W. (2012). Effectiveness of mindfulness-based stress reduction in mood, breast- and endocrine-related quality of life, and well-being in stage 0 to III breast cancer: A randomized, controlled trial. *Journal of Clinical Oncology: Official Journal of the American Society of Clinical Oncology*, 30(12), 1335–1342.
- Hölzel, B. K., Carmody, J., Vangel, M., Congleton, C., Yerramsetti, S. M., Gard, T., & Lazar, S. W. (2011). Mindfulness practice leads to increases in regional brain gray matter density. *Psychiatry Research*, 191(1), 36–43.
- Humphreys, J., & Lee, K. (2009). Interpersonal violence is associated with depression and chronic physical health problems in midlife women. *Issues in Mental Health Nursing*, 30(4), 206–213.
- Juster, R.-P., McEwen, B. S., & Lupien, S. J. (2010). Allostatic load biomarkers of chronic stress and impact on health and cognition. *Neuroscience and Biobehavioral Reviews*, 35(1), 2–16.
- Kozasa, E. H., Sato, J. R., Lacerda, S. S., Barreiros, M. A., Radvany, J., Russell, T. A., et al. (2012). Meditation training increases brain efficiency in an attention task. *NeuroImage*, 59(1), 745–749.
- Krisanaprakornkit, T., Krisanaprakornkit, W., Piyavhatkul, N., & Laopaiboon, M. (2006). Meditation therapy for anxiety disorders. *Cochrane Database System Reviews*, (1), CD004998.
- Lazar, S., Kerr, C., & Wasserman, R. (2005). Meditation experience is associated with increased cortical thickness. *NeuroReport*, 16(17), 1893–1897.
- Lutgendorf, S. K., & Costanzo, E. S. (2003). Psychoneuroimmunology and health psychology: An integrative model. *Brain, Behavior, and Immunity*, 17(4), 225–232.
- Lutz, A., Greischar, L. L., Rawlings, N. B., Ricard, M., & Davidson, R. J. (2004). gamma synchrony during mental practice. *Proceedings of the National Academy of Sciences of the United States of America*, 101(46), 16369–16373.
- Lutz, A., Slagter, H. A., Dunne, J. D., & Davidson, R. J. (2008). Attention regulation and monitoring in meditation. *Trends in Cognitive Sciences*, 12(4), 163–169.
- Lutz, A., Slagter, H. A., Rawlings, N. B., Francis, A. D., Greischar, L. L., & Davidson, R. J. (2009). Mental training enhances attentional stability: Neural and behavioral evidence. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 29(42), 13418–13427.
- Marques-Deak, A., & Sternberg, E. (2004). Psychoneuroimmunology: The relation between the central nervous system and the immune system. *Revista Brasileira de Psiquiatria*, 26(3), 143–144.
- McGee, M. (2008). Meditation and psychiatry. *Psychiatry (Edgmont)*, 5(1), 28–41.
- McN Cain, N., & Gray, D. (2005). Implementing a comprehensive approach to the study of health dynamics using the psychoneuroimmunology paradigm. *ANS. Advances in Nurse's Science*, 28(4), 320–332.
- Menezes, C., Dell'Aglio, & Dalbosco, D. (2009). Os efeitos da meditação à luz da investigação científica em Psicologia: Revisão de literatura. *Psicologia: Ciência E Profissão*, 29(2), 276–89.



- Menezes, C., Dell’Aglío, D., & Bizarro, L. (2011). Meditação, bem-estar ea ciência psicológica: revisão de estudos empíricos. *Interação Em Psicologia*, 15(2), 239–248.
- Ministério da Saúde, S. de A. à S. (2008). *Política Nacional de Práticas Integrativas e Complementares no SUS*. Brasília.
- Nakamura, Y., Lipschitz, D. L., Kuhn, R., Kinney, A. Y., & Donaldson, G. W. (2013). Investigating efficacy of two brief mind-body intervention programs for managing sleep disturbance in cancer survivors: A pilot randomized controlled trial. *Journal of Cancer Survivorship: Research and Practice*, 7(2), 165–182.
- Newberg, A., Alavi, A., & Baime, M. (2001). The measurement of regional cerebral blood flow during the complex cognitive task of meditation: A preliminary SPECT study. *Psychiatry Research: Neuroimaging Section*, 106(2), 113–122.
- Oppermann, R. (2002). Efeitos do estresse sobre a imunidade ea doença periodontal. *Revista Faculdade de Odontologia Porto Alegre*, 43(2), 52–59.
- Pace, T. W. W., & Heim, C. M. (2011). A short review on the psychoneuroimmunology of posttraumatic stress disorder: From risk factors to medical comorbidities. *Brain, Behavior, and Immunity*, 25(1), 6–13.
- Schneider, R., & Grim, C. (2012). Stress reduction in the secondary prevention of cardiovascular disease randomized, controlled trial of transcendental meditation and health education in Blacks. *Circulation Cardiovascular Quality Outcomes*, 5(6), 750–758.
- Servan-Shreiber, D. (2008). *Anticâncer: prevenir e vencer usando nossas defesas naturais*. Rio de Janeiro: Objetiva.
- Singh, Y., Sharma, R., & Talwar, A. (2012). Immediate and long-term effects of meditation on acute stress reactivity, cognitive functions, and intelligence. *Alternative Therapies Health Medicine*, 18, 46–53.
- Slagter, H. A., Lutz, A., Greischar, L. L., Francis, A. D., Nieuwenhuis, S., Davis, J. M., & Davidson, R. J. (2007). Mental training affects distribution of limited brain resources. *PLoS Biology*, 5(6), e138.
- Sun, J., Kang, J., Wang, P., & Zeng, H. (2013). Self-relaxation training can improve sleep quality and cognitive functions in the older: A one-year randomised controlled trial. *Journal of Clinical Nursing*, 22(9–10), 1270–1280.
- Tang, Y.-Y., Ma, Y., Fan, Y., Feng, H., Wang, J., Feng, S., et al. (2009). Central and autonomic nervous system interaction is altered by short-term meditation. *Proceedings of the National Academy of Sciences of the United States of America*, 106(22), 8865–8870.
- Wallace, R. (1970). Physiological effects of transcendental meditation. *Science*, 167(March), 1751–1754.
- Wallace, R., Benson, H., & Wilson, A. (1971). A wakeful hypometabolic physiologic state. *American Journal of Psychotherapy*, 221, 795–799.
- Willis, R. (1979). Meditation to fit the person: Psychology and the meditative way. *Journal of Religion and Health*, 18(2), 93–119.
- Witek-Janusek, L., Albuquerque, K., Chroniak, K. R., Chroniak, C., Durazo-Arvizu, R., & Mathews, H. L. (2008). Effect of mindfulness based stress reduction on immune function, quality of life and coping in women newly diagnosed with early stage breast cancer. *Brain, Behavior, and Immunity*, 22(6), 969–981.