

Numerical Convergence of the Block-Maxima Approach to the Generalized Extreme Value Distribution

Davide Faranda · Valerio Lucarini · Giorgio Turchetti · Sandro Vaienti

Received: 4 March 2011 / Accepted: 23 May 2011 / Published online: 27 September 2011
© Springer Science+Business Media, LLC 2011

Abstract In this paper we perform an analytical and numerical study of Extreme Value distributions in discrete dynamical systems. In this setting, recent works have shown how to get a statistics of extremes in agreement with the classical Extreme Value Theory. We pursue these investigations by giving analytical expressions of Extreme Value distribution parameters for maps that have an absolutely continuous invariant measure. We compare these analytical results with numerical experiments in which we study the convergence to limiting distributions using the so called block-maxima approach, pointing out in which cases we obtain robust estimation of parameters. In regular maps for which mixing properties do not hold, we show that the fitting procedure to the classical Extreme Value Distribution fails, as expected. However, we obtain an empirical distribution that can be explained starting from a different observable function for which Nicolis et al. (Phys. Rev. Lett. 97(21): 210602, 2006) have found analytical results.

Keywords Extreme values · Dynamical systems · EVT · GEV · Mixing · Logistic map · Chaos

D. Faranda (✉) · V. Lucarini
Department of Mathematics and Statistics, University of Reading, Whiteknights, PO Box 220,
Reading RG6 6AX, UK
e-mail: d.faranda@pgr.reading.ac.uk

V. Lucarini
Department of Meteorology, Department of Mathematics, University of Reading, Whiteknights, PO
Box 220, Reading RG6 6AX, UK
e-mail: v.lucarini@reading.ac.uk

G. Turchetti
Department of Physics, INFN-Bologna, University of Bologna, Via Irnerio 46, Bologna 40126, Italy
e-mail: turchett@bo.infn.it

S. Vaienti
UMR-6207, Centre de Physique Théorique, CNRS, Universités d'Aix-Marseille I, II, Université du Sud
Toulon-Var and FRUMAM (Fédération de Recherche des Unités de Mathématiques de Marseille), CPT,
Luminy, Case 907, 13288 Marseille Cedex 09, France
e-mail: vaienti@cpt.univ-mrs.fr

1 Introduction

Extreme Value Theory (EVT) was first developed by Fisher and Tippett [21] and formalized by Gnedenko [29] which showed that the distribution of the block-maxima of a sample of independent identically distributed (i.i.d.) variables converges to a member of the so-called Extreme Value (EV) distribution. It arises from the study of stochastic series that is of great interest in different disciplines: it has been applied to extreme floods [30, 50, 56], amounts of large insurance losses [6, 14]; extreme earthquakes [8, 13, 55]; meteorological and climate events [1, 19, 20, 48, 54, 60]. All these events have a relevant impact on socioeconomic activities and it is crucial to find a way to understand and, if possible, forecast them [34, 39].

The attention of the scientific community to the problem of modeling extreme values is growing. An extensive account of recent results and relevant applications is given in Ghil et al. [27]. Such an interest is mainly due to the fact that this theory is also important in defining risk factor in a wide class of applications such as the modeling of financial risk after the significant instabilities in financial markets worldwide [18, 28, 45], the analysis of seismic and hydrological risk [8, 47]. Even if the probability of extreme events decreases with their magnitude, the damage that they may bring increases rapidly with the magnitude as does the cost of protection against them Nicolis et al. [49].

From a theoretical point of view, extreme values represent extreme fluctuations of a system. Very recently, many authors have shown clearly how the statistics of global observables in correlated systems can be related to EV statistics [5, 15]. Clusel and Bertin [9] have shown how to connect fluctuations of global additive quantities, like total energy or magnetization, by statistics of sums of random variables in such a way that it is possible to identify a class of random variables whose sum follows an extreme value distributions.

The so called block-maxima approach is widely used in EVT since it represents a very natural way to look at extremes. It consists of dividing the data series of some observable into bins of equal length and selecting the maximum (or the minimum) value in each of them [11]. When dealing with climatological or financial data, since we usually have limited dataset, the main problem in applying EVT is related to the choice of a sufficiently large statistics of extremes provided that each bin contains a suitable number of observations. Therefore a smart balance between number of maxima and observations per bin is needed [19, 40–42].

Recently a number of alternative approaches have been studied. One consists in looking at exceedance over high thresholds rather than maxima over fixed time periods. While the idea of looking at extreme value problems from this point of view is very old, the development of a modern theory has started with Todorovic and Zelenhasic [57] that have proposed the so called Peaks Over Threshold approach. At the same time there was a mathematical development of procedures based on a certain number of extreme order statistics [36, 52] and the Generalized Pareto distribution for excesses over thresholds [16, 17, 53].

Since dynamical systems theory can be used to understand features of physical systems like climate and forecast financial behaviors, many authors have studied how to extend EVT to these field. When dealing with dynamical systems we have to know what kind of properties (i.e. stability, degree of mixing, correlations decay) are related to Gnedenko's hypotheses and also which observables we must consider in order to obtain an EV distribution. Furthermore, even if the convergence is achieved, we should evaluate how fast it is depending on all parameters and properties used. Empirical studies show that in some cases a dynamical observable obeys to the extreme value statistics even if the convergence is highly dependent on the kind of observable we choose [58–60]. For example, Balakrishnan et al. [3] and more recently Nicolis et al. [49] and Haiman [33] have shown that for regular orbits of dynamical systems we don't expect to find convergence to EV distribution.

The first rigorous mathematical approach to extreme value theory in dynamical systems goes back to the pioneer paper by P. Collet in 2001 [12]. Collet got the Gumbel Extreme Value Law (see below) for certain one-dimensional non-uniformly hyperbolic maps which admit an absolutely continuous invariant measure and exhibit exponential decay of correlations. Collet's approach used Young towers [61, 62] and his suggestion was successively applied to other systems. Before quoting them, we would like to point out that Collet was able to establish a few conditions (usually called D and D') and which have been introduced by Leadbetter [43] with the aim to associate to the stationary stochastic process given by the dynamical system, a new stationary independent sequence which enjoyed one of the classical three extreme value laws, and this law could be pulled back to the original dynamical sequence. Conditions D and D' require a sort of independence of the stochastic dynamical sequence in terms of uniform mixing condition on the distribution functions. Condition D was successively improved by Freitas and Freitas [22], in the sense that they introduced a new condition, called D_2 , which is weaker than D and that could be checked directly by estimating the rate of decay of correlations for Hölder observables.¹ We notice that conditions D_2 and D' allow immediately to get Extreme Value Laws for absolutely continuous invariant measures for uniformly one-dimensional expanding dynamical systems: this is the case for instance of the 1-D maps with constant density studied in Sect. 3 below. Another interesting issue of Collet's paper was the choice of the observables g 's whose values along the orbit of the dynamical systems constitute the sequence of events upon which we successively search for the partial maximum. Collet considered a function $g(\text{dist}(x, \zeta))$ of the distance with respect to a given point ζ , with the aim that g achieves a global maximum at almost all points ζ in the phase space; for example $g(x) = -\log x$. Using a different g , Freitas and Freitas [23] were able to get the Weibull law for the family of quadratic maps with the Benedicks-Carleson parameters and for ζ taken as the critical point or the critical value, so improving the previous results by Collet who did not keep such values in his set of full measure.

The latter paper [23] strongly relies on condition D_2 ; this condition has also been invoked to establish the extreme value laws on towers which model dynamical systems with stable foliations (hyperbolic billiards, Lozi maps, Hénon diffeomorphisms, Lorenz maps and flows). This is the content of the paper by Gupta, Holland and Nicol [32]. We point out that the observable g was taken in one of three different classes g_1, g_2, g_3 , see Sect. 2 below, each one being again a function of the distance with respect to a given point ζ . The choice of these particular forms for the g 's is just to fit with the necessary and sufficient condition on the tail of the distribution function $F(u)$, see next section, in order to exist a non-degenerate limit distribution for the partial maxima [24, 37]. The paper Gupta et al. [32] also covers the easier case of uniformly hyperbolic diffeomorphisms, for instance the Arnold Cat map which we studied in Sect. 3.2.

Another major step in this field was achieved by establishing a connection between the extreme value laws and the statistics of first return and hitting times, see the papers by

¹We briefly state here the two conditions, we defer to the next section for more details about the quantities introduced. If $X_n, n \geq 0$ is a stochastic process, we define $M_{j,l} \equiv \{X_j, X_{j+1}, \dots, X_{j+l}\}$ and we put $M_{0,m} = M_m$. Moreover we set a_m and b_m two normalizing sequences and $u_m = x/a_m + b_m$, where x is a real number, cf. next section for the meaning of these variables. The condition $D_2(u_m)$ holds for the sequence X_m if for any integer l, t, m we have $|\nu(X_0 > u_m, M_{t,l} \leq u_m) - \nu(X_0 > u_m)\nu(M_{t,l} \leq u_m)| \leq \gamma(m, t)$, where $\gamma(m, t)$ is non-increasing in t for each m and $m\gamma(m, t_m) \rightarrow 0$ as $m \rightarrow \infty$ for some sequence $t_m = o(m), t_m \rightarrow \infty$.

We say condition $D'(u_m)$ holds for the sequence X_m if $\lim_{k \rightarrow \infty} \limsup_m m \sum_{j=1}^{\lfloor m/k \rfloor} \nu(X_0 > u_m, X_j > u_m) = 0$. Whenever the process is given by the iteration of a dynamical system, the previous two conditions could also be formulated in terms of decay of correlation integrals, see Freitas and Freitas [22], Gupta [31].

Freitas, Freitas and Todd [24, 26]. They showed in particular that for dynamical systems preserving an absolutely continuous invariant measure or a singular continuous invariant measure ν , the existence of an exponential hitting time statistics on balls around ν almost any point ζ implies the existence of extreme value laws for one of the observables of type g_i , $i = 1, 2, 3$ described above. The converse is also true, namely if we have an extreme value law which applies to the observables of type g_i , $i = 1, 2, 3$ achieving a maximum at ζ , then we have exponential hitting time statistics to balls with center ζ . Recently these results have been generalized to local returns around balls centered at periodic points [25]. We would like to point out that the equivalence between extreme values laws and the hitting time statistics allowed to prove the former for broad classes of systems for which the statistics of recurrence were known, for instance for expanding maps in higher dimension.

In this work we consider a few aspects of the extreme value theory applied to dynamical systems throughout both analytical results and numerical experiments. In particular we analyze the convergence to EV limiting distributions pointing out how robust are parameters estimations. Furthermore, we check the consistency of block-maxima approach highlighting deviations from theoretical expected behavior depending on the number of maxima and number of block-observation. To perform our analysis we use low dimensional maps with different properties: mixing maps in which we expect to find convergence to EV distributions and regular maps where the convergence is not ensured.

The work is organized as follows: in Sect. 2 we briefly recall methods and results of EVT for independent and identical distributed (i.i.d.) variables and dynamical systems. In Sect. 3 we explicitly compute theoretical expected distributions parameter in respect to the observable functions of type g_i , $i = 1, 2, 3$ for map that have constant density measure. Numerical experiments on low dimensional maps are presented. In Sect. 4 we show that it is possible to derive an asymptotic expression of normalizing sequences when the density measure is not constant. As an example we derive the explicit expressions for the Logistic map. Eventually, in Sect. 5 we repeat the experiment for regular maps showing that extreme values laws do not follow from numerical experiments.

2 Background on EVT

Gnedenko [29] studied the convergence of maxima of i.i.d. variables

$$X_0, X_1, \dots, X_{m-1}$$

with cumulative distribution (cdf) $F(x)$ of the form:

$$F(x) = P\{a_m(M_m - b_m) \leq x\}$$

Where a_m and b_m are normalizing sequences and $M_m = \max\{X_0, X_1, \dots, X_{m-1}\}$. It may be rewritten as $F(u_m) = P\{M_m \leq u_m\}$ where $u_m = x/a_m + b_m$. Such types of normalizing sequences converge to one of the three type of Extreme Value (EV) distribution if necessary and sufficient conditions on parent distribution of X_i variables are satisfied [43]. EV distributions include the following three families:

- Gumbel distribution (type 1):

$$F(x) = \exp\{-e^{-x}\} \quad x \in \mathbb{R} \tag{1}$$

- Fréchet distribution (type 2):

$$\begin{cases} F(x) = 0 & x \leq 0 \\ F(x) = \exp\{-x^{1/\xi}\} & x > 0 \end{cases} \tag{2}$$

- Weibull distribution (type 3):

$$\begin{cases} F(x) = \exp\{-(-x)^{1/\xi}\} & x < 0 \\ F(x) = 0 & x \geq 0 \end{cases} \tag{3}$$

Let us define the right endpoint x_F of a distribution function $F(x)$ as:

$$x_F = \sup\{x : F(x) < 1\} \tag{4}$$

then, it is possible to compute normalizing sequences a_m and b_m using the following corollary of Gnedenko’s theorem:

Corollary (Gnedenko) *The normalizing sequences a_m and b_m in the convergence of normalized maxima $P\{a_m(M_m - b_m) \leq x\} \rightarrow F(x)$ may be taken (in order of increasing complexity) as:*

- Type 1: $a_m = [G(\gamma_m)]^{-1}$, $b_m = \gamma_m$;
- Type 2: $a_m = \gamma_m^{-1}$, $b_m = 0$ or $b_m = c \cdot m^{-\xi}$;
- Type 3: $a_m = (x_F - \gamma_m)^{-1}$, $b_m = x_F$;

where

$$\gamma_m = F^{-1}(1 - 1/m) = \inf\{x; F(x) \geq 1 - 1/m\} \tag{5}$$

$$G(t) = \int_t^{x_F} \frac{1 - F(u)}{1 - F(t)} du \quad t < x_F \tag{6}$$

and $c \in \mathbb{R}$ is a constant. It is important to remark that the choice of normalizing sequences is not unique [43]. For example for b_m of type 2 distribution it is possible to choose either $b_m = 0$ or $b_m = c \cdot m^{-\xi}$. In particular, we will use the last one since it is a more general choice that ensure the convergence for a much broader class of initial distributions [4].

Instead of Gnedenko’s approach it is possible to fit unnormalized data directly to a single family of generalized distribution called GEV distribution with cdf:

$$F_G(x; \mu, \sigma, \xi) = \exp\left\{-\left[1 + \xi\left(\frac{x - \mu}{\sigma}\right)\right]^{-1/\xi}\right\} \tag{7}$$

which holds for $1 + \xi(x - \mu)/\sigma > 0$, using $\mu \in \mathbb{R}$ (location parameter) and $\sigma > 0$ (scale parameter) as scaling constants in place of b_m , and a_m [51]. $\xi \in \mathbb{R}$ is the shape parameter also called the tail index: when $\xi \rightarrow 0$, the distribution corresponds to a Gumbel type. When the index is negative, it corresponds to a Weibull; when the index is positive, it corresponds to a Fréchet.

In order to adapt the extreme value theory to dynamical systems, we will consider the stationary stochastic process X_0, X_1, \dots given by:

$$X_m(x) = g(\text{dist}(f^m(x), \zeta)) \quad \forall m \in \mathbb{N} \tag{8}$$

where ‘dist’ is a Riemannian metric on Ω , ζ is a given point and g is an observable function, and whose partial maximum is defined as:

$$M_m = \max\{X_0, \dots, X_{m-1}\} \tag{9}$$

The probability measure will be here an invariant measure ν for the dynamical system. As we anticipated in the Introduction, we will use three types of observables $g_i, i = 1, 2, 3$, suitable to obtain one of the three types of EV distribution for normalized maxima:

$$g_1(x) = -\log(\text{dist}(x, \zeta)) \tag{10}$$

$$g_2(x) = \text{dist}(x, \zeta)^{-1/\alpha} \tag{11}$$

$$g_3(x) = C - \text{dist}(x, \zeta)^{1/\alpha} \tag{12}$$

where C is a constant and $\alpha > 0 \in \mathbb{R}$.

These three type of functions are representative of broader classes which are defined, for instance, throughout (1.11) to (1.13) in Freitas et al. [24]; we now explain the reasons and the meaning of these choices. First of all these functions have in common the following properties: (i) they are defined on the positive semi-axis $[0, \infty]$ with values into $\mathbb{R} \cup \{+\infty\}$; (ii) 0 is a global maximum, possibly equal to $+\infty$; (iii) they are a strictly decreasing bijection in a neighborhood V of 0 with image W . Then we consider three types of behavior which generalize the previous specific choices:

Type 1: there is a strictly positive function $p : W \rightarrow \mathbb{R}$ such that $\forall y \in \mathbb{R}$ we have

$$\lim_{s \rightarrow g_1(0)} \frac{g_1^{-1}(s + yp(s))}{g_1^{-1}(s)} = e^{-y}$$

Type 2: $g_2(0) = +\infty$ and there exists $\beta > 0$ such that $\forall y > 0$ we have

$$\lim_{s \rightarrow \infty} \frac{g_2^{-1}(sy)}{g_2^{-1}(s)} = y^{-\beta}$$

Type 3: $g_3(0) = D < +\infty$ and there exists $\gamma > 0$ such that $\forall y > 0$ we have

$$\lim_{s \rightarrow 0} \frac{g_3^{-1}(D - sy)}{g_3^{-1}(D - s)} = y^\gamma$$

The Gnedenko corollary says that the different kinds of extreme value laws are determined by the distribution of $F(u) = \nu(X_0 \leq u)$ and by the right endpoint of F, x_F . We will see in the next section that the local invertibility of $g_i, i = 1, 2, 3$ in the neighborhood of 0 together with the Lebesgue’s differentiation theorem (which basically says that whenever the measure ν is absolutely continuous with respect to Lebesgue with density ρ , the measure of a ball $B_\delta(x_0)$ of radius δ centered around almost any point x_0 scales like $\delta\rho(x_0)$), allow us to compute the tail of F , in fact we have

$$1 - F(u) \sim \rho(\zeta)|B_{g^{-1}(u)}(\zeta)|,$$

where g is any of the three types of functions introduced in (10) to (12) and $|A|$ denotes the diameter of the set A . As we said above the tail of F determines the three limit laws for partial maximum of i.i.d. sequences. In particular Th. 1.6.2 in Leadbetter et al. [43] specifies what kind of conditions the distribution function F must verify to get one specific law: the above type 1, 2, 3 assumptions are just the translation in terms of the shape of g_i of the conditions on the tail of F .

3 Distribution of Extremes in Mixing Maps with Constant Density Measure

Our goal is to use a block-maxima approach and fit our unnormalized data to a GEV distribution; for that it will be necessary to find a linkage among a_m, b_m, μ and σ . At this regard we will use Gnedenko’s corollary to compute normalizing sequences showing that they correspond to the parameter we obtain fitting directly data to GEV distribution.

We derive the correct expression for mixing maps with constant density measure and the asymptotic behavior for logistic map that is a case of non-constant density measure.

3.1 Asymptotic Sequences

In this section we will consider the case of uniformly hyperbolic maps which preserve the Lebesgue measure (the density $\rho = 1$) and satisfy the conditions D_2 and D' , sufficient to get extreme values distributions. For the second map, the algebraic automorphisms of the torus better known as the Arnold cat map, the existence of extreme value laws follows from the theory developed in Gupta et al. [32]. Starting from the definitions provided by Gnedenko we derive as a novel result the exact expression for the normalizing sequences a_m and b_m .

Case 1 ($g_1(x) = -\log(\text{dist}(x, \zeta))$) By (8) and (9) we know that:

$$\begin{aligned} 1 - F(u) &= 1 - \nu(g(\text{dist}(x, \zeta)) \leq u) \\ &= 1 - \nu(-\log(\text{dist}(x, \zeta)) \leq u) \\ &= 1 - \nu(\text{dist}(x, \zeta) \geq e^{-u}) \end{aligned} \tag{13}$$

and the last line is justified by using Lebesgue’s Differentiation Theorem. Then, for maps with constant density measure, we can write:

$$1 - F(u) \simeq \nu(B_{e^{-u}}(\zeta)) = \Omega_d e^{-ud} \tag{14}$$

where d is the dimension of the space and Ω_d is a constant. To use Gnedenko corollary it is necessary to calculate u_F

$$u_F = \sup\{u; F(u) < 1\}$$

in this case $u_F = +\infty$.

Using Gnedenko (6) we can calculate $G(t)$ as follows:

$$G(t) = \int_t^\infty \frac{1 - F(u)}{1 - F(t)} du = \int_t^\infty \frac{e^{-ud}}{e^{-td}} du = \frac{1}{d} \int_{td}^\infty \frac{e^{-v}}{e^{-td}} dv = \frac{1}{d} \tag{15}$$

According to the Leadbetter et al. [43] proof of Gnedenko theorem we can study both a_m and b_m or γ_m convergence as:

$$\begin{aligned} \lim_{m \rightarrow \infty} m(1 - F\{\gamma_m + xG(\gamma_m)\}) &= e^{-x} \\ \lim_{m \rightarrow \infty} m\Omega_d e^{-d(\gamma_m + xG(\gamma_m))} &= e^{-x} \end{aligned} \tag{16}$$

then we can use the connection between γ_m and normalizing sequences to find a_m and b_m . By (5) or using relation (16):

$$\gamma_m \simeq \frac{\ln(m\Omega_d)}{d}$$

so that:

$$a_m = d \quad b_m = \frac{1}{d} \ln(m) + \frac{\ln(\Omega_d)}{d}$$

Case 2 ($g_2(x) = \text{dist}(x, \zeta)^{-1/\alpha}$) We can proceed as for g_1 :

$$\begin{aligned} 1 - F(u) &= 1 - \nu(\text{dist}(x, \zeta)^{-1/\alpha} \leq u) \\ &= 1 - \nu(\text{dist}(x, \zeta) \geq u^{-\alpha}) \\ &= \nu(B_{u^{-\alpha}}(\zeta)) = \Omega_d u^{-\alpha d} \end{aligned} \tag{17}$$

in this case $u_F = +\infty$.

$$\gamma_m = F^{-1}(1 - 1/m) = (m\Omega_d)^{1/(\alpha d)} \tag{18}$$

and, as discussed in Sect. 2, using Beirlant [4] choice of normalizing sequences we expect:

$$b_m = c \cdot m^{-\xi}$$

where $c \in \mathbb{R}$ is a constant.

Case 3 ($g_3(x) = C - \text{dist}(x, \zeta)^{1/\alpha}$) Eventually we compute a_m and b_m for the g_3 observable class:

$$\begin{aligned} 1 - F(u) &= 1 - \nu(C - \text{dist}(x, \zeta)^{1/\alpha} \leq u) \\ &= 1 - \nu(\text{dist}(x, \zeta) \geq (C - u)^\alpha) \\ &= \nu(B_{(C-u)^\alpha}(\zeta)) = \Omega_d (C - u)^{\alpha d} \end{aligned} \tag{19}$$

in this case $u_F = C$.

$$\gamma_m = F^{-1}(1 - 1/m) = C - (m\Omega_d)^{-1/(\alpha d)} \tag{20}$$

For type 3 distribution:

$$a_m = (u_F - \gamma_m)^{-1} \quad b_m = u_F \tag{21}$$

3.2 Numerical Experiments

Since we want to show that unnormalized data may be fitted by using the GEV distribution $F_G(x; \mu, \sigma, \xi)$ we expect to find the following equivalence:

$$a_m = 1/\sigma \quad b_m = \mu$$

where, clearly, $\mu = \mu(m)$ and $\sigma = \sigma(m)$. This fact can be seen as a linear change of variable: the variable $y = a_m(x - b_m)$ has a GEV distribution $F_G(y; \mu = 0, \sigma = 1, \xi)$ (that is an EV one parameter distribution with a_m and b_m normalizing sequences) while x is GEV distributed $F_G(x; \mu = b_m, \sigma = 1/a_m, \xi)$.

As we said above we now apply the previous considerations to two maps which enjoy extreme values laws and have constant density: we summarize below the theoretical results we obtained for all three type of observables. We have obtained the results in terms of m but, since we fix $k = n \cdot m$, the previous results can be translated in terms of n as follows:

For g_1 type observable:

$$\sigma = \frac{1}{d} \quad \mu \propto \frac{1}{d} \ln\left(\frac{k}{n}\right) \tag{22}$$

For g_2 type observable:

$$\sigma \propto n^{-1/(\alpha d)} \quad \mu \propto n^{-1/(\alpha d)} \tag{23}$$

For g_3 type observable:

$$\sigma \propto n^{1/(\alpha d)} \quad \mu = C \tag{24}$$

Following Freitas et al. [24] we obtain the expression for the shape parameters: $\xi = 0$ for g_1 type, $\xi = 1/(\alpha d)$ for g_2 type and $\xi = -1/(\alpha d)$ for g_3 type.

In order to provide a numerical test of our results we consider a one-dimensional and a two dimensional map. The one dimensional map used is a Bernoulli Shift map:

$$x_{t+1} = qx_t \pmod 1 \quad q > 1 \in \mathbb{N} \tag{25}$$

with $q = 3$.

The considered two dimensional map is the famous Arnold’s cat map defined on the 2-torus by:

$$\begin{bmatrix} x_{t+1} \\ y_{t+1} \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x_t \\ y_t \end{bmatrix} \pmod 1 \tag{26}$$

A wide description of properties of these maps can be found in Arnold and Avez [2] and Hasselblatt and Katok [35].

We proceed as follows. For each map we run a long simulation up to k iterations starting from a given initial condition ζ . Note that the results—as we tested—do not depend on the choice of ζ . From the trajectory we compute the sequence of observables g_1, g_2, g_3




Fig. 1 *Left:* g_1 observable empirical histogram and fitted GEV pdf. *Right:* g_1 observable empirical cdf and fitted GEV cdf. Logistic map, $n = 10^4$, $m = 10^4$

as follows dividing it into n bins each containing $m = k/n$ observations. Then, we test the degree of agreement between the empirical distribution of the maxima and the GEV distribution according to the theoretical values presented above. A priori, it is reasonable to assume GEV as a suitable family of statistical models. For some selected values of n , the maxima are normalized and fitted to GEV distributions $F_G(x; \mu, \sigma, \xi)$ using a maximum likelihood method which selects values of the model parameters that produce the distribution most likely to have resulted in the observed data.

All the numerical analysis contained in this work has been performed using MATLAB Statistics Toolbox functions such as *gevfit* and *gevcdf*. These functions return maximum likelihood estimates of the parameters for the generalized extreme value (GEV) distribution giving 95% confidence intervals for estimates [46].

As in every fitting procedure, it is necessary to test the a posteriori goodness of fit. We anticipate that in every case considered, fitted distributions passed, with maximum confidence interval, the Kolmogorov-Smirnov test described in Lilliefors [44]. For illustration purposes, we present in Fig. 1 an empirical pdf and cdf with the corresponding fits.

Once k is set to a given value (in our case $k = 10^7$), the numerical simulations allow us to explore two limiting cases of great interest in applications where the statistical inference is intrinsically problematic:

1. n is small (m is large), so that we extract only few maxima, each corresponding to a *very* extreme event.
2. m is small (n is large), so that we extract many maxima but most of those will not be as extreme as in case (1).

In case (1), we have only few data—of high quality—to fit our statistical models whereas in case (2) we have many data but the sampling may be spoiled by the inclusion of data not giving a good representation of extreme events. We have in general that in order to obtain a reliable fit for a distribution with p parameters we need 10^p independent data [19] so that we expect that fit procedure gives reliable results for $n > 10^3$. As the value of m determines to which extent the extracted bin maximum is representative of an extreme, below a certain value m_{\min} our selection procedure will be unavoidably misleading. We have no obvious theoretical argument to define the value of m_{\min} . We expect to obtain good fits throughout




Fig. 2 (Color online) g_1 observable, $\zeta \simeq 0.51$. (a) ξ VS $\log_{10}(n)$; (b) σ VS $\log_{10}(n)$; (c) μ VS $\log_{10}(n)$. *Left*: Bernoulli Shift map. *Right*: Arnold Cat Map. *Dotted lines* represent computed confidence interval, *red lines* represent a linear fit, *blue lines* are theoretical values

the parametric region where the constraints on n, m are satisfied. Therefore, our flexibility in choosing satisfying pairs (n, m) increases with larger values of k .

For a g_1 type observable function the behavior against n of the three parameters is presented in Fig. 2. According to (22) we expect to find $\xi = 0$. For relatively small values of n the sample is too small to ensure a good convergence to analytical ξ and confidence intervals are wide. On the other hand we see deviations from expected value as $m < 10^3$ that is when $n > 10^4$. For the scale parameter a similar behavior is achieved and deviations from expected theoretical values $\sigma = 1/2$ for Arnold Cat Map and $\sigma = 1$ for Bernoulli Shift are




Fig. 3 (Color online) g_2 observable, $\zeta \simeq 0.51$. (a) ξ VS $\log_{10}(n)$; (b) $\log_{10}(\sigma)$ VS $\log_{10}(n)$; (c) $\log_{10}(\mu)$ VS $\log_{10}(n)$. *Left*: Bernoulli Shift map. *Right*: Arnold Cat Map. *Dotted lines* represent computed confidence interval, *red lines* represent a linear fit, *blue lines* are theoretical values

found when $n < 10^3$ or $m < 10^3$. Location parameter μ shows a logarithm decay with n as expected from (22). A linear fit of μ in respect to $\log(n)$ is shown with a red line in Fig. 2. The linear fit computed angular coefficients K^* of (22) well approximate $1/d$: for Bernoulli Shift map we obtain $|K^*| = 1.001 \pm 0.001$ while for Arnold Cat map $|K^*| = 0.489 \pm 0.001$. We find that ξ values have best matching with theoretical ones with reliable confidence interval when both $n > 10^3$ and $m > 10^3$. These results are confirmed even for g_2 type and g_3 type observable functions as shown in Figs. 3(a) and 4(a) respectively. We present the fit results for $\alpha = 3$ but we have done tests for different α and for fixed n and different α .




Fig. 4 (Color online) g_3 observable, $\zeta \simeq 0.51$. **(a)** ξ VS $\log_{10}(n)$; **(b)** $\log_{10}(\sigma)$ VS $\log_{10}(n)$; **(c)** $\log_{10}(\mu)$ VS $\log_{10}(n)$. *Left:* Bernoulli Shift map. *Right:* Arnold Cat Map. *Dotted lines* represent computed confidence interval, *red lines* represent a linear fit, *blue lines* are theoretical values

For g_2 observable function we can also check that μ and σ parameters follow a power law as described in (23). In the log-log plot in Fig. 3(b), 3(c), we can see a very clear linear behavior. For the Bernoulli Shift map we obtain $|K^*| = 0.330 \pm 0.001$ for μ series, $|K^*| = 0.341 \pm 0.001$ for σ in good agreement with theoretical value of $1/3$. For Arnold Cat map we expect to find $K^* = 1/6$, from the experimental data we obtain $|K^*| = 0.163 \pm 0.001$ for μ and $|K^*| = 0.164 \pm 0.001$ for σ .

Eventually, computing g_3 as observable function we expect to find a constant value for μ while σ has to grow with a power law in respect to n as expected from (24). As in g_2 case we expect $|K^*| = 1/(\alpha d)$ and numerical results shown in Fig. 4(b), 4(c) are consistent with the

theoretical one since $|K^*| = 0.323 \pm 0.006$ for Bernoulli shift map and $|K^*| = 0.162 \pm 0.006$ for Arnold Cat map.

In all cases considered the analytical behavior described in (23) and (24) is achieved and the fit quality improves if $n > 10^3$ and $m > 10^3$. The g_3 type observable constant has been chosen $C = 10$. The nature of these lower bound is quite different.

4 Distributions of Extremes in Mixing Map with Non-constant Density Measure

4.1 Asymptotic Sequences

The main problem when dealing with maps that have absolutely continuous but non-constant density measure $\rho(\zeta)$ is in the computation of the integral:

$$v(B_\delta(\zeta)) = \int_{B_\delta(\zeta)} \rho(x) dx \tag{27}$$

where $B_\delta(\zeta)$ is the d -dimensional ball of radius δ centered in ζ .

We have to know the value of this integral in order to evaluate $F(u)$ and, therefore, the sequences a_m and b_m .

As shown in the previous section δ is linked to the observable type: in all cases, since we substitute $u = 1 - 1/m$, $\delta \rightarrow 0$ means that we are interested in $m \rightarrow \infty$.

In this limit, a first order approximation of the previous integral is:

$$v(B_\delta(\zeta)) \simeq \rho(\zeta)\delta^d + \mathcal{O}(\delta^{d+1}) \tag{28}$$

that is valid if we are not in a neighborhood of a singular point of $\rho(\zeta)$.

As an example we compute the asymptotic sequences for a logistic map:

$$x_{t+1} = rx_t(1 - x_t) \tag{29}$$

with $r = 4$. This map satisfies hypothesis described in the analysis performed for Benedicks-Carleson maps in Moreira Freitas and Freitas [23].

For this map the density of the absolutely continuous invariant measure is explicit and reads:

$$\rho(\zeta) = \frac{1}{\pi\sqrt{\zeta(1-\zeta)}} \quad \zeta \in (0, 1) \tag{30}$$

So that:

$$\int_{B_\delta(\zeta)} \rho(\zeta) d\zeta = \frac{2}{\pi} [\arcsin(\sqrt{\zeta + \delta}) - \arcsin(\sqrt{\zeta - \delta})] \tag{31}$$

where $\zeta + \delta < 1$ and $\zeta - \delta > 0$. Since Extreme Value Theory effectively works only if n, m are large enough, the results in (31) can be replaced by a series expansion for $\delta \rightarrow 0$:

$$\frac{2}{\pi} [\arcsin(\sqrt{\zeta + \delta}) - \arcsin(\sqrt{\zeta - \delta})] = \frac{1}{\pi} \frac{2\delta}{\sqrt{\zeta(1-\zeta)}} [1 + \delta^2 P(\zeta) + \dots] \tag{32}$$

up to order δ^3 , where:

$$P(\zeta) = \frac{1}{8\zeta^2} - \frac{2}{\zeta(1-\zeta)} + \frac{2}{\zeta^2(1-\zeta)} + \frac{6}{\zeta^2(1-\zeta)^2} \tag{33}$$

Using the last two equations we are able to compute asymptotic normalizing sequences a_m and b_m for all g_i observables.

Case 1 ($g_1(x) = -\log(\text{dist}(x, \zeta))$) For g_1 observable functions we set $\delta = e^{-ud}$. In case of logistic map $d = 1$. First we have to compute $G(t)$ using (15) and the expansion in equation:

$$G(t) = \frac{\int_t^\infty du(e^{-u} + e^{-3u}P(\zeta))}{e^{-t} + e^{-3t}P(\zeta)} \simeq 1 - \frac{2}{3}e^{-2t}P(\zeta) \tag{34}$$

We can compute γ_m , if $m \gg 1$, as follows:

$$F(\gamma_m) \simeq 1 - \frac{1}{m} \tag{35}$$

At the first order in (32) we get

$$\frac{1}{m} \simeq \frac{1}{\pi} \frac{2e^{-\gamma_m}}{\sqrt{\zeta(1-\zeta)}} \tag{36}$$

so that:

$$\gamma_m \simeq \ln(m) + \ln\left(\frac{2}{\pi\sqrt{\zeta(1-\zeta)}}\right) \tag{37}$$

Therefore, the sequences a_m and b_m if $m \gg 1$ are:

$$a_m \simeq [G(\gamma_m)]^{-1} \simeq 1 + \frac{2}{3} \frac{\pi^2}{4m^2} \zeta(\zeta - 1)P(\zeta) \tag{38}$$

$$b_m \simeq \gamma_m \simeq \ln(m) + \ln(2\rho(\zeta)) \tag{39}$$

Case 2 ($g_2(x) = \text{dist}(x, \zeta)^{-1/\alpha}$) We can proceed as for g_1 setting $\delta = (\alpha u)^{-\alpha}$, computing γ_m we get at the first order in (32):

$$\frac{1}{m} \simeq \frac{1}{\pi} \frac{2\gamma_m^{-\alpha}}{\sqrt{\zeta(1-\zeta)}} = 2\rho(\zeta)(\alpha\gamma_m)^{-\alpha} \tag{40}$$

$$\gamma_m = \frac{1}{\alpha} \left(\frac{1}{2m\rho(\zeta)}\right)^{-1/\alpha} \tag{41}$$

We can respectively compute a_m and b_m as:

$$a_m = \gamma_m^{-1} \quad b_m = (2m\rho(\zeta))^{-\xi} \tag{42}$$

Case 3 ($g_3(x) = C - \text{dist}(x, \zeta)^{1/\alpha}$) As in the previous cases, we compute γ_m up to the first order setting $\delta = [\alpha(C - \gamma_m)]^\alpha$:

$$\frac{1}{m} \simeq \frac{1}{\pi} \frac{2[\alpha(C - \gamma_m)]^\alpha}{\sqrt{\zeta(1 - \zeta)}} = 2\rho(\zeta)[\alpha(C - \gamma_m)]^\alpha \tag{43}$$

$$\gamma_m = C - \frac{1}{\alpha} \left(\frac{1}{2m\rho(\zeta)} \right)^{1/\alpha} \tag{44}$$

For type 3 distribution:

$$a_m = (u_F - \gamma_m)^{-1} \quad b_m = u_F; \tag{45}$$

where $u_f = C$.

4.2 Numerical Experiment on the Logistic Map

Following the same procedure detailed in Sect. 3.2, we want to show the equivalence between EV computed normalizing sequences a_m and b_m and the parameters of a GEV distribution obtained directly fitting the data even in case of logistic map that has not constant density measure. Using (38)–(39) for g_1 , we obtain the following theoretical expression:

$$\sigma(m, \zeta) \simeq 1 + \frac{2}{3} \frac{\pi^2}{4m^2} \zeta(\zeta - 1)P(\zeta) \quad \mu(m, \zeta) \simeq \ln(m) + \ln(2\rho(\zeta)) \tag{46}$$

From (39), for g_2 observable type, we write:

$$\sigma(m, \zeta) \simeq \frac{1}{\alpha} (2m\rho(\zeta))^{1/\alpha} \quad \mu(m, \zeta) \simeq (2m\rho(\zeta))^{1/\alpha} \tag{47}$$

and in g_3 case using (45), we expect to find:

$$\sigma(m, \zeta) \simeq \frac{1}{\alpha} (2m\rho(\zeta))^{-1/\alpha} \quad \mu(m, \zeta) \simeq C = u_F \tag{48}$$

Values of ξ are independent on density and, as stated in Freitas' $\xi = 0$ for g_1 type, $\xi = 1/(\alpha d)$ for g_2 type and $\xi = -1/(\alpha d)$ for g_3 type.

In Figs. 5, 6, 7 we presents a numerical test of the asymptotic behavior described in (46)–(48) on logistic map for $d = 1, a = 3, C = u_F = 10, \zeta = 0.3$ against the variable n . As shown in previous section, block maxima approach works well with maps with constant density measure when n and m are at least 10^3 : In fact, regarding ξ parameter. Significant deviations from the theoretical value are achieved when $n < 1000$ or $m < 1000$ even in the case of the Logistic Map.

Regarding μ and σ , for g_1 observable a linear fit of μ in respect to $\log(n)$ give us $|K^*| = 0.999 \pm 0.002$, while σ shows the same behavior of ξ since the best agreement with theoretical value $\sigma = 1$ is achieved when $n, m > 10^3$. In the log-log plots of Fig. 6(b), 6(c) for g_2 observable, we can observe again the expected linear behavior for μ and σ with $|K^*|$ corresponding to $1/(\alpha d)$. From numerical fit we obtain $|K^*| = 0.3334 \pm 0.0007$ for μ series and $|K^*| = 0.337 \pm 0.002$ for σ in good agreement with theoretical value of $1/3$. By applying a linear fit to the log-log plot in Fig. 7(b), the angular coefficient corresponding to σ series is $|K| = 0.323 \pm 0.003$ again consistent with the theory.




Fig. 5 (Color online) g_1 observable, $\zeta = 0.31$. (a) ξ VS $\log_{10}(n)$; (b) σ VS $\log_{10}(n)$; (c) μ VS $\log_{10}(n)$. Logistic map. Dotted lines represent computed confidence interval, red lines represent a linear fit, blue lines are theoretical values

For a logistic map we can also check the GEV behavior in respect to initial conditions. If we fix $n^* = m^* = 10^3$ and fit our data to GEV distribution for 10^3 different $\zeta \in (0, 1)$ an asymptotic behavior is reached as shown from the previous analysis. For g_1 observable function we have observed that the first order approximation works well for all three parameters. Deviation from this behavior are achieved for $\zeta \rightarrow 1$ and $\zeta \rightarrow 0$ as the measure become singular when we move to these points and we should take in account other terms of the series expansion. Numerically, we found that deviations from first order approximation are meaningful only if $\zeta < 10^{-3}$ and $\zeta > 1 - 10^{-3}$. Averaging over ζ both ξ and σ we




Fig. 6 (Color online) g_2 observable, $\zeta = 0.3$. **(a)** ξ VS $\log_{10}(n)$; **(b)** $\log_{10}(\sigma)$ VS $\log_{10}(n)$; **(c)** $\log_{10}(\mu)$ VS $\log_{10}(n)$. Logistic map. Dotted lines represent computed confidence interval, red lines represent a linear fit, blue lines are theoretical values

obtain $\langle \xi \rangle = 1.000 \pm 0.009$ and $\langle \sigma \rangle = 1.00 \pm 0.03$ where the uncertainties are computed with respect to the estimator. Since we expect $\xi = 0$ and $\sigma = 1$ at zero order approximation, numerical results are consistent with the theoretical ones; furthermore, experimental data are normally distributed around theoretical values.

Asymptotic expansion also works well for g_2 observables: we obtain $\langle \xi \rangle = 0.334 \pm 0.001$ in excellent agreement with theoretical value $\xi = 1/3$. Eventually, in g_3 , averaging ξ over different initial conditions we get $\langle \xi \rangle = -0.334 \pm 0.002$ that is again consistent to theoretical value $-1/3$.




Fig. 7 (Color online) g_3 observable, $\zeta = 0.3$. **(a)** ξ VS $\log_{10}(n)$; **(b)** $\log_{10}(\sigma)$ VS $\log_{10}(n)$; **(c)** $\log_{10}(\mu)$ VS $\log_{10}(n)$. Logistic map. *Dotted lines* represent computed confidence interval, *red lines* represent a linear fit, *blue lines* are theoretical values

5 The Case of Regular Maps

Freitas and Freitas [22] have posed the problem of dependent extreme values in dynamical systems that show uniform quasi periodic motion. Here we try to investigate this problem numerically. We have used a one-dimensional and a bi-dimensional discrete map. The first one is the irrational translation on the torus defined by:

$$x_{t+1} = x_t + \beta \pmod 1 \quad \beta \in [0, 1] \setminus \mathbb{Q}. \tag{49}$$

And for the bidimensional case, we use the so called standard map:

$$y_{t+1} = y_t + \frac{\lambda}{2\pi} \sin(2\pi x_t) \pmod 1 \quad x_{t+1} = x_t + y_{t+1} \pmod 1 \tag{50}$$

with $\lambda = 10^{-4}$. For this value of λ , the standard map exhibits a regular behavior and it is not mixing, as well as torus translations. This means that these maps fail in satisfying hypothesis D_2 and D' and moreover they do not enjoy as well an exponential hitting time statistics. About this latter statistics, it is however known that it exists for torus translation and it is given by a particular piecewise linear function or a uniform distribution depending on which sequence of sets A_k is considered [10]. In a similar way, a non-exponential Hitting Time Statistics (HTS) is achieved for standard map when $\lambda \ll 1$ as well as for a skew map, that is a standard map with $\lambda = 0$ [7]. Therefore we expect not to obtain a GEV distribution of any type using g_i observables.

We have pointed out that the observable functions choice is crucial in order to observe some kind of distribution of extreme values when we are dealing with dynamical systems instead of stochastic series. Nicolis et al. [49] have shown how it is possible to obtain an analytical EV distribution which does not belong to GEV family choosing a simple observable: they considered the series of distances between the iterated trajectory and the initial condition. Using the same notation of Sect. 2 we can write:

$$Y_m(x = f^t \zeta) = \text{dist}(f^t \zeta, \zeta) \quad \hat{M}_m = \min\{Y_0, \dots, Y_{m-1}\}$$

For this observable they have shown that the cumulative distribution $F(x) = P\{a_m(\hat{M}_m - b_m) \leq x\}$ of a uniform quasi periodic motions is not smooth but piecewise linear (Nicolis et al. [49], Fig. 3). Furthermore slop changes of $F(x)$ can be explained by constructing the intersections between different iterates of (49). $F(x)$ must correspond to a density distribution continuous obtained as a composition of box functions: each box must be related to a change in the slope of $F(x)$.

The numerical results we report below confirm that for the maps (49) and (50) the distributions of maxima for various observables cannot be fitted with a GEV since they are multi modal. We recall that the return times into a sphere of vanishing radius do not have a spectrum, if the orbits have the same frequency, whereas a spectrum appears if the frequency varies continuously with the action, as in the standard map for λ close to zero [38]. Since the EV statistics refers to a single orbit, no change due to the local mixing, which insures the existence of a return times spectrum [38], can be observed. Considering that the GEV exists when the system is mixing and does not when it is integrable, one might use the quality of fit to GEV as a dynamical indicator, for systems which exhibit regions with different dynamical properties, ranging from integrable to mixing as it occurs for the standard map when λ is order 1. Indeed we expect that in the neighborhood of a low order resonance, where the homoclinic tangle of intersecting separatrices appears, a GEV fit is possible. Preliminary computations carried out for the standard map and for a model with parametric resonance confirm this claim, that will be carefully tested in the near future.

Using the theoretical framework provided in Nicolis et al. [49] we check numerically the behavior of maps described in (49)–(50) analyzing EV distributions for g_i observable functions. Proceeding as in Sect. 3 for mixing maps, we try to perform a fit to GEV distribution starting with different initial conditions ζ , a set of different α values and (n, m) combinations. In all cases analyzed the Kolmogorov-Smirnov test fails and this means that GEV distribution is not useful to describe the behavior of this kind of statistics. This result is in

agreement with Freitas et al. [26] but we may find out which kind of empirical distribution is obtained.

Looking in details at M_m histograms that correspond to empirical density distributions, they appear always to be multi modal and each mode have a well defined shape: for g_1 type observable function modes are exponential while, for g_2 and g_3 , their shape depends on α value of observable function. Furthermore, the number of modes and their positions are highly dependent on both n, m and initial conditions.

Using Nicolis et al. [49] results it is possible to understand why we obtain this kind of histograms: since density distribution of \hat{M}_m is a composition of box functions, when we apply g_i observables we modulate it changing the shape of the boxes. Therefore, we obtain a multi modal distribution modified according to the observable functions g_i .

An example is shown in Fig. 8 for standard map: the left figures correspond to the histogram of the minimum distance obtained without computing g_i observable and reproduce a composition of box functions. The figures in the right show how this distribution is modified by applying g_1 observable to the series of minimum distances. We can see two exponential modes, while the third is hidden in the linear scale but can be highlighted using a log-scale. The upper figures are drawn using $n = 3300, m = 3300$, the lower with $n = 10000, m = 1000$.

6 Concluding Remarks

EVT was developed to study a wide class of problems of great interest in different disciplines: the need of modeling events that occur with very small probability comes from the fact that they can affect in a strong way several socioeconomic activities: floods, insurance losses, earthquakes, catastrophes. A very extensive account of EVT applications has been recently given in Ghil et al. [27]. EVT was applied on limited data series using the block-maxima approach facing the problem of having a good statistics of extreme values retaining a sufficient number of observation in each bin. Often, since no theoretical a priori values of GEV parameters are available for this kind of applications, we may obtain a biased fit to GEV distribution even if tests of statistical significance succeed. The recent development of an extreme value theory in dynamical systems give us the theoretical framework to test the consistency of block-maxima approach when analytical results for distribution parameters are available. This theory relies on the global properties of the dynamical systems considered (such as the degree of mixing or the decay rate of the Hitting Time Statistics) but also on the observable functions we chose.

Our main finding is that a block-maxima approach for GEV distribution is totally equivalent to fit an EV distribution after normalizing sequences are computed. To prove this we have derived analytical expressions for a_m and b_m normalizing sequences, showing that μ and σ of fitted GEV distribution can replace them. This approach works for maps that have an absolutely continuous invariant measure and retain some mixing properties that can be directly related to the exponential decay of HTS. Since GEV approach does not require the a priori knowledge of the measure density that is instead require by the EV approach, it is possible to use it in many numerical applications.

Furthermore, if we compare analytical and numerical results we can study what is the minimum number of maxima and how big the set of observations in which the maximum is taken has to be. To accomplish this goal we have analyzed maps with constant density measure finding that a good agreement between numerical and analytical value is achieved when both the number of maxima n and the observations per bin m are at least 10^3 . We remark that the fits have passed Kolmogorov-Smirnov test with maximum confidence interval




Fig. 8 Histogram of maxima for g_1 type observable function, standard map, $x_0 = y_0 = \sqrt{2} - 1$. *Left:* series of $\min(\text{dist}(f^t \zeta, \zeta))$. *Right:* series of $g_1 = -\log(\min(\text{dist}(f^t \zeta, \zeta)))$. **(a)** $n = 3300, m = 3300$. **(b)** $n = 10000, m = 1000$

even if $n < 10^3$ or $m < 10^3$ so that parametric or non-parametric tests are not the only thing to take in account when dealing with extreme value distributions: if maxima are not proper extreme values (which means m is not large enough) the fit is good but parameters are different from expected values. The lower bound of n can be explained using the argument that a fit to a 3-parameters distribution needs at least 10^3 independent data to give reliable informations.

Therefore, we checked that in case of non-constant absolutely continuous density measure the asymptotic expressions used to compute μ and σ works when we consider n and m of order 10^3 . For logistic map the numerical values of parameters we obtain averaging over different initial conditions are totally in agreement with the theoretical ones. In regu-

lar maps, as expected, the fit to a GEV distribution is unreliable. We obtain a multi modal distribution, that, for the analyzed maps, is the result of a composition of modes in which the shape depends on observable types. This behavior can be explained pointing out that this kind of systems have not an exponential HTS decay and therefore have no EV law for observables of type g_i .

To conclude, we claim that we have provided a reliable way to investigate properties of extreme values in mixing dynamical systems which may satisfy mixing conditions (like D_2 and D'), finding an equivalence among a_m , b_m , μ and σ behavior for absolutely continuous measures. In our future work we intend to address the case of singular measure. Recently the theorem was generalized to the case of non-smooth observations and therefore it holds also with non-absolutely continuous invariant probability measure [26]. In this case we expect the same for all the procedure described here. Understanding the extreme values behavior for singular measures will be crucial to apply proficiently this analysis to operative geophysical models since in these case we are always dealing with singular measures. In this way we will provide a complete tool to study extreme events in complex dynamical systems used in geophysical or financial applications.

Acknowledgements S.V. was supported by the CNRS-PEPS Project *Mathematical Methods of Climate Models*, and he thanks the GDRE Grefi-Mefi for having supported exchanges with Italy. V.L. and D.F. acknowledge the financial support of the EU FP7-ERC project NAMASTE: Thermodynamics of the Climate System.

References

1. Altmann, E.G., Hallerberg, S., Kantz, H.: Reactions to extreme events: moving threshold model. *Phys. A, Stat. Mech. Appl.* **364**, 435–444 (2006)
2. Arnold, V.I., Avez, A.: *Ergodic Problems of Classical Mechanics*. Benjamin, New York (1968)
3. Balakrishnan, V., Nicolis, C., Nicolis, G.: Extreme value distributions in chaotic dynamics. *J. Stat. Phys.* **80**(1), 307–336 (1995). ISSN 0022-4715
4. Beirlant, J.: *Statistics of Extremes: Theory and Applications*. Wiley, New York (2004). ISBN 0471976474
5. Bertin, E.: Global fluctuations and Gumbel statistics. *Phys. Rev. Lett.* **95**(17), 170601 (2005). ISSN 1079-7114
6. Brodin, E., Kluppelberg, C.: Extreme value theory in finance. In: *Encyclopedia of Quantitative Risk Analysis and Assessment* (2008). doi:[10.1002/9780470061596.risk0431](https://doi.org/10.1002/9780470061596.risk0431). ISBN:0-470-03549-8, 978-0-470-03549-8
7. Buric, N., Rampioni, A., Turchetti, G.: Statistics of Poincaré recurrences for a class of smooth circle maps. *Chaos Solitons Fractals* **23**(5), 1829–1840 (2005)
8. Burton, P.W.: Seismic risk in southern Europe through to India examined using Gumbel's third distribution of extreme values. *Geophys. J. R. Astron. Soc.* **59**(2), 249–280 (1979). ISSN 1365-246X
9. Clusel, M., Bertin, E.: Global fluctuations in physical systems: a subtle interplay between sum and extreme value statistics. *Int. J. Mod. Phys. B* **22**(20), 3311–3368 (2008). ISSN 0217-9792
10. Coelho, Z., De Faria, E.: Limit laws of entrance times for homeomorphisms of the circle. *Isr. J. Math.* **93**(1), 93–112 (1996). ISSN 0021-2172
11. Coles, S., Heffernan, J., Tawn, J.: Dependence measures for extreme value analyses. *Extremes* **2**(4), 339–365 (1999). ISSN 1386-1999
12. Collet, P.: Statistics of closest return for some non-uniformly hyperbolic systems. *Ergod. Theory Dyn. Syst.* **21**(02), 401–420 (2001)
13. Cornell, C.A.: Engineering seismic risk analysis. *Bull. Seismol. Soc. Am.* **58**(5), 1583 (1968). ISSN 0037-1106
14. Cruz, M.G.: *Modeling, Measuring and Hedging Operational Risk*. Wiley, New York (2002). ISBN 0471515604
15. Dahlstedt, K., Jensen, H.J.: Universal fluctuations and extreme-value statistics. *J. Phys. A, Math. Gen.* **34**, 11193 (2001)

16. Davison, A.C.: Modelling excesses over high thresholds, with an application. In: *Statistical Extremes and Applications*, pp. 461–482. Reidel, Dordrecht (1984)
17. Davison, A.C., Smith, R.L.: Models for exceedances over high thresholds. *J. R. Stat. Soc., Ser. B, Methodol.* **52**(3), 393–442 (1990). ISSN 0035-9246
18. Embrechts, P., Resnick, S.I., Samorodnitsky, G.: Extreme value theory as a risk management tool. *N. Am. Actuar. J.* **3**, 30–41 (1999). ISSN 1092-0277
19. Felici, M., Lucarini, V., Speranza, A., Vitolo, R.: Extreme value statistics of the total energy in an intermediate complexity model of the mid-latitude atmospheric jet. Part I: Stationary case. *J. Atmos. Sci.* **64**, 2137–2158 (2007)
20. Felici, M., Lucarini, V., Speranza, A., Vitolo, R.: Extreme value statistics of the total energy in an intermediate complexity model of the mid-latitude atmospheric jet. Part II: Trend detection and assessment. *J. Atmos. Sci.* **64**, 2159–2175 (2007)
21. Fisher, R.A., Tippett, L.H.C.: Limiting forms of the frequency distribution of the largest or smallest member of a sample. *Proc. Camb. Philos. Soc.* **24**, 180 (1928)
22. Freitas, A.C.M., Freitas, J.M.: On the link between dependence and independence in extreme value theory for dynamical systems. *Stat. Probab. Lett.* **78**(9), 1088–1093 (2008). ISSN 0167-7152
23. Freitas, A.C.M., Freitas, J.M.: Extreme values for Benedicks-Carleson quadratic maps. *Ergod. Theory Dyn. Syst.* **28**(04), 1117–1133 (2008). ISSN 0143-3857
24. Freitas, A.C.M., Freitas, J.M., Todd, M.: Hitting time statistics and extreme value theory. *Probab. Theory Relat. Fields* **147**(3–4), 675–710 (2010)
25. Freitas, A.C.M., Freitas, J.M., Todd, M.: Extremal index, hitting time statistics and periodicity. *Arxiv preprint, arXiv:1008.1350* (2010)
26. Freitas, A.C.M., Freitas, J.M., Todd, M., Gardas, B., Drichel, D., Flohr, M., Thompson, R.T., Cummer, S.A., Frauendiener, J., Doliwa, A., et al.: Extreme value laws in dynamical systems for non-smooth observations. *Arxiv preprint, arXiv:1006.3276* (2010)
27. Ghil, M., et al.: Extreme events: dynamics, statistics and prediction. *Nonlinear Process. Geophys.* **18**, 295–350 (2011)
28. Gilli, M., Kőllezi, E.: An application of extreme value theory for measuring financial risk. *Comput. Econ.* **27**(2), 207–228 (2006). ISSN 0927-7099
29. Gnedenko, B.: Sur la distribution limite du terme maximum d’une série aléatoire. *Ann. Math.* **44**(3), 423–453 (1943)
30. Gumbel, E.J.: The return period of flood flows. *Ann. Math. Stat.* **12**(2), 163–190 (1941). ISSN 0003-4851
31. Gupta, C.: Extreme-value distributions for some classes of non-uniformly partially hyperbolic dynamical systems. *Ergod. Theory Dyn. Syst.* **30**(03), 757–771 (2010)
32. Gupta, C., Holland, M., Nicol, M.: Extreme value theory for dispersing billiards and a class of hyperbolic maps with singularities. *Preprint* (2009)
33. Haiman, G.: Extreme values of the tent map process. *Stat. Probab. Lett.* **65**(4), 451–456 (2003). ISSN 0167-7152
34. Hallerberg, S., Kantz, H.: Influence of the event magnitude on the predictability of an extreme event. *Phys. Rev. E* **77**(1), 11108 (2008). ISSN 1550-2376
35. Hasselblatt, B., Katok, A.B.: *A First Course in Dynamics: With a Panorama of Recent Developments*. Cambridge University Press, Cambridge (2003)
36. Hill, B.M.: A simple general approach to inference about the tail of a distribution. *Ann. Stat.* **3**(5), 1163–1174 (1975). ISSN 0090-5364
37. Holland, M., Nicol, M., Török, A.: Extreme value distributions for non-uniformly hyperbolic dynamical systems. *Preprint* (2008)
38. Hu, H., Rampioni, A., Rossi, L., Turchetti, G., Vaienti, S.: Statistics of Poincaré recurrences for maps with integrable and ergodic components. *Chaos, Interdiscip. J. Nonlinear Sci.* **14**, 160 (2004)
39. Kantz, H., Altmann, E., Hallerberg, S., Holstein, D., Riegert, A.: Dynamical interpretation of extreme events: predictability and predictions. In: *Extreme Events in Nature and Society*, pp. 69–93. Springer, Berlin (2006)
40. Katz, R.W.: Extreme value theory for precipitation: sensitivity analysis for climate change. *Adv. Water Resour.* **23**(2), 133–139 (1999). ISSN 0309-1708
41. Katz, R.W., Brown, B.G.: Extreme events in a changing climate: variability is more important than averages. *Clim. Change* **21**(3), 289–302 (1992). ISSN 0165-0009
42. Katz, R.W., Brush, G.S., Parlange, M.B.: Statistics of extremes: modeling ecological disturbances. *Ecology* **86**(5), 1124–1134 (2005). ISSN 0012-9658
43. Leadbetter, M.R., Lindgren, G., Rootzen, H.: *Extremes and Related Properties of Random Sequences and Processes*. Springer, New York (1983)

44. Lilliefors, H.W.: On the Kolmogorov-Smirnov test for normality with mean and variance unknown. *J. Am. Stat. Assoc.* **62**(318), 399–402 (1967). ISSN 0162-1459
45. Longin, F.M.: From value at risk to stress testing: the extreme value approach. *J. Bank. Finance* **24**(7), 1097–1130 (2000). ISSN 0378-4266
46. Martinez, W.L., Martinez, A.R.: *Computational Statistics Handbook with MATLAB*. CRC Press, Boca Raton (2002)
47. Martins, E.S., Stedinger, J.R.: Generalized maximum-likelihood generalized extreme-value quantile estimators for hydrologic data. *Water Resour. Res.* **36**(3), 737–744 (2000). ISSN 0043-1397
48. Nicholls, N.: CLIVAR and IPCC interests in extreme events. In: *Workshop Proceedings on Indices and Indicators for Climate Extremes*, Asheville, NC. Sponsors, CLIVAR, GCOS and WMO (1997)
49. Nicolis, C., Balakrishnan, V., Nicolis, G.: Extreme events in deterministic dynamical systems. *Phys. Rev. Lett.* **97**(21), 210602 (2006). ISSN 1079-7114
50. Friederichs, P., Hense, A.: Statistical downscaling of extreme precipitation events using censored quantile regression. *Mon. Weather Rev.* **135**(6), 2365–2378 (2007). ISSN 0027-0644
51. Pickands III, J.: Moment convergence of sample extremes. *Ann. Math. Stat.* **39**(3), 881–889 (1968)
52. Pickands III, J.: Statistical inference using extreme order statistics. *Ann. Stat.* **3**(1), 119–131 (1975). ISSN 0090-5364
53. Smith, R.L.: Threshold methods for sample extremes. *Stat. Extremes Appl.* **621**, 638 (1984)
54. Smith, R.L.: Extreme value analysis of environmental time series: an application to trend detection in ground-level ozone. *Stat. Sci.* **4**(4), 367–377 (1989). ISSN 0883-4237
55. Sornette, D., Knopoff, L., Kagan, Y.Y., Vanneste, C.: Rank-ordering statistics of extreme events: application to the distribution of large earthquakes. *J. Geophys. Res.* **101**(B6), 13883 (1996). ISSN 0148-0227
56. Sveinsson, O.G.B., Boes, D.C.: Regional frequency analysis of extreme precipitation in Northeastern Colorado and Fort Collins flood of 1997. *J. Hydrol. Eng.* **7**, 49 (2002)
57. Todorovic, P., Zelenhasic, E.: A stochastic model for flood analysis. *Water Resour. Res.* **6**(6), 1641–1648 (1970). ISSN 0043-1397
58. Vannitsem, S.: Statistical properties of the temperature maxima in an intermediate order Quasi-Geostrophic model. *Tellus A* **59**(1), 80–95 (2007). ISSN 1600-0870
59. Vitolo, R., Holland, M.P., Ferro, C.A.T.: Robust extremes in chaotic deterministic systems. *Chaos, Interdiscip. J. Nonlinear Sci.* **19**, 043127 (2009)
60. Vitolo, R., Ruti, P.M., Dell’Aquila, A., Felici, M., Lucarini, V., Speranza, A.: Accessing extremes of mid-latitude wave activity: methodology and application. *Tellus A* **61**(1), 35–49 (2009). ISSN 1600-0870
61. Young, L.S.: Statistical properties of dynamical systems with some hyperbolicity. *Ann. Math.* **147**(3), 585–650 (1998)
62. Young, L.S.: Recurrence times and rates of mixing. *Isr. J. Math.* **110**(1), 153–188 (1999)