



Complex hyperkähler structures defined by Donaldson–Thomas invariants

Tom Bridgeland¹ · Ian A. B. Strachan²

Received: 16 November 2020 / Revised: 11 March 2021 / Accepted: 18 March 2021 /

Published online: 22 April 2021

© The Author(s) 2021

Abstract

The notion of a Joyce structure was introduced in Bridgeland (Geometry from Donaldson–Thomas invariants, preprint [arXiv:1912.06504](https://arxiv.org/abs/1912.06504)) to describe the geometric structure on the space of stability conditions of a CY_3 category naturally encoded by the Donaldson–Thomas invariants. In this paper we show that a Joyce structure on a complex manifold defines a complex hyperkähler structure on the total space of its tangent bundle, and give a characterisation of the resulting hyperkähler metrics in geometric terms.

Keywords Hyperkähler geometry · Donaldson–Thomas invariants · Stability conditions · Integrable systems

Mathematics Subject Classification Primary 14N35; Secondary 53C26

1 Introduction

In the recent paper [6] it was argued that the Donaldson–Thomas (DT) invariants of a CY_3 triangulated category \mathcal{D} should encode a certain geometric structure on its space of stability conditions $M = \text{Stab}(\mathcal{D})$. These structures were called Joyce structures, since the most important ingredients already appear in the paper [18]. The definition

¹ This was denoted by the letter J in [6], but to avoid confusion with the standard notation I, J, K in hyperkähler geometry, we switch to W here.

Boris Dubrovin Memorial Issue Guest Editors: G. Dito, P. Lorenzoni, D. Masoero, P. Rossi, Y. Zhang.

✉ Ian A. B. Strachan
Ian.Strachan@glasgow.ac.uk
Tom Bridgeland
t.bridgeland@sheffield.ac.uk

¹ Royal Society Research Professor, University of Sheffield, Sheffield, UK

² University of Glasgow, Glasgow, UK

of a Joyce structure on a complex manifold M given in [6] is rather ungeometric in nature. In this paper we explain that it can be re-interpreted as the existence of a complex hyperkähler metric of a particular kind on the total space $X = \mathcal{T}_M$ of the holomorphic tangent bundle of M . We include a brief summary of the most important ideas of [6] in the ‘‘Appendix’’ section.

In terms of a local system of co-ordinates (z_1, \dots, z_n) on M , the main ingredient of a Joyce structure is a function $W : X \rightarrow \mathbb{C}$ satisfying the partial differential equations

$$\frac{\partial^2 W}{\partial \theta_i \partial z_j} - \frac{\partial^2 W}{\partial \theta_j \partial z_i} = \sum_{p,q} \eta^{pq} \cdot \frac{\partial^2 W}{\partial \theta_i \partial \theta_p} \cdot \frac{\partial^2 W}{\partial \theta_j \partial \theta_q}, \tag{1}$$

where η^{pq} is a constant non-degenerate skew-symmetric matrix, and $(\theta_1, \dots, \theta_n)$ are the natural linear co-ordinates on the tangent spaces $\mathcal{T}_{M,p}$ obtained by writing a tangent vector in the form $v = \sum_{i=1}^n \theta_i \cdot \frac{\partial}{\partial z_i}$. We shall explain below that (1) is the condition for the expression

$$g = \sum_{i,j} \omega_{ij} \cdot (d\theta^i \otimes dz^j + dz^j \otimes d\theta^i) - \sum_{i,j} \frac{\partial^2 W}{\partial \theta_i \partial \theta_j} \cdot (dz^i \otimes dz^j + dz^j \otimes dz^i). \tag{2}$$

to define a complex hyperkähler metric on X . Here ω_{ij} is the inverse matrix to η^{ij} . In terms of the basis of vector fields

$$v_i = \frac{\partial}{\partial \theta_i}, \quad h_i = \frac{\partial}{\partial z_i} + \sum_{p,q} \eta^{pq} \cdot \frac{\partial^2 W}{\partial \theta_i \partial \theta_p} \cdot \frac{\partial}{\partial \theta_q},$$

the complex structures I, J, K are defined by the block matrices

$$I = \begin{pmatrix} i \cdot \mathbb{1} & 0 \\ 0 & -i \cdot \mathbb{1} \end{pmatrix}, \quad J = \begin{pmatrix} 0 & -\mathbb{1} \\ \mathbb{1} & 0 \end{pmatrix}, \quad K = \begin{pmatrix} 0 & -i \cdot \mathbb{1} \\ -i \cdot \mathbb{1} & 0 \end{pmatrix},$$

and the metric is given by $g = \sum_{i,j} \omega_{ij} \cdot (v^i \otimes h^j + h^j \otimes v^i)$.

The main characteristic of the complex hyperkähler structures defined by this construction is that the holomorphic 2-form

$$\Omega_-(v, w) = g(v, (J - iK)(w)),$$

on X is the pull-back via the natural projection $\pi : X \rightarrow M$ of a holomorphic symplectic form on M , namely $\omega = \sum_{i,j} \omega_{ij} dz_i \wedge dz_j$. Conversely, as we show in Sect. 2, all complex hyperkähler structures with this property are locally given by the above construction for some function W satisfying (1). We leave to future work the problem of describing interesting classes of examples of Joyce structures, and hence of functions W satisfying the partial differential Eq. (1). One non-trivial example is discussed in detail in [8], and a large class of examples will be described in [9].

In a recent preprint [10], Dunajski showed that the form (2) of the metric follows from a weaker set of conditions than complex hyperkähler, namely null Kähler. Such

a structure consists of a metric g and an endomorphism N with the property that its kernel is half-dimensional, together with the conditions

$$g(X, NY) + g(NX, Y) = 0, \quad N^2 = 0$$

and $\nabla N = 0$ (with ∇ being the Levi-Civita connection of g). Since $(J - iK)^2 = 0$ the constructions in this paper provide examples of such null-Kähler structures. This work also provides an explanation of the appearance of isomonodromy problems.

1.1 Relations with previous work

Since hyperkähler metrics may be written in terms of a Kähler potential, the geometric conditions of being hyperkähler result in differential equations for such a potential, and these take the form of Monge–Ampère-type equations. In 4-dimensions there is a single equation which is known as Plebański’s first heavenly Eq. [25] (the original motivation coming from the equivalent description of Ricci-flat metrics with anti-self-dual Weyl tensor). Many other forms of the equations exist, and Eq. (1), in 4-dimensions, is known as Plebański’s second heavenly equation. From the work of Penrose [24]—the original nonlinear graviton construction—it immediately follows that there is an associated twistor space, and a crucial property of this space is the family of curves with normal bundle $\mathcal{O}(1) \oplus \mathcal{O}(1)$.

This construction can be generalized in many ways, and the simplest is to generalize the normal bundle structure to $\bigoplus_i \mathcal{O}(n_i)$. Equation (1) first appear *explicitly* in the literature in the work of Takasaki [29], who used the bundle of 2-forms construction of Gindikin [15] to write down the associated hierarchies of integrable equations. Equation (1) corresponds to the case where the manifold is hyperkähler, with the associated family of curves in the twistor space having normal bundle structure $\mathcal{O}(1) \oplus \dots \oplus \mathcal{O}(1)$ with $2r$ -terms.

With the development of the links between twistor theory and the theory of integrable systems initiated by Ward [31], these curved twistor space constructions and their pencils of commuting vector fields were reinterpreted in terms of Lax equations with ‘gauge fields’ taking values in, for example, the Lie algebra $\mathfrak{sdiff}(\Sigma)$ of volume preserving diffeomorphisms of some associated manifold Σ (see, for example, [22]). Thus, for example, the original 4D self-duality equation may be interpreted as a two-dimensional σ -model [23] with fields taking values in the Lie algebra $\mathfrak{sdiff}(\Sigma^2)$. This splitting into two distinct sets of coordinates is mirrored here with the coordinates z_i on the base space M and the fibre coordinates θ_i . Indeed, here the Lie algebra $\mathfrak{sdiff}(\Sigma^2)$ is replaced by the algebra of Poisson-preserving symplectic vector fields on the algebraic torus \mathbb{T} , and the Lie group $\text{SDiff}(\Sigma)$ by the group of automorphisms of this torus.

Connections between hyperkähler geometry and Donaldson–Thomas theory are also not new. Gaiotto, Moore and Neitzke explained a beautiful connection between these subjects [13], and went on to describe explicit examples leading to the Hitchin metric on the moduli space of Higgs bundles [14]. The approach of [6] is inspired by, and closely-related to, their work, although it is strictly different, since it deals with

the ‘conformal limit’, and hence leads to complex hyperkähler structures rather than real ones.

As will be explained below, and as explained in [6], the function W has also to satisfy certain homogeneity conditions in order for it to define a Joyce structure, and geometrically these imply the existence of a conformal (actually a homothetic) Killing vector on the hyperkähler manifold $X = \mathcal{T}_M$. In 4-dimensions, hyperkähler manifolds with such a conformal Killing vector were studied by Dunajski and Tod [11], with the corresponding Einstein-Weyl structures on the orbit space being constructed, following Hitchin, and the associated mini-twistor space defined. Such constructions will obviously extend to arbitrary dimensional hyperkähler manifolds with a conformal Killing vector, but the details do not appear to have been explicitly written down.

As well as bringing together some of the disparate sources, which span 50 years of research in several different areas, the reason for writing the paper is to give a precise geometric description of the structure on stability space one expects to be encoded by DT theory. This is a complex hyperkähler structure on the tangent bundle, but with certain extra features which it seems worthwhile making explicit. We also took the opportunity to give a geometric characterisation of the hyperkähler structures arising from the above construction.

Notation and terminology

We use the notation \mathcal{T}_M to denote the holomorphic tangent bundle of a complex manifold M . Given a holomorphic map of complex manifolds $\pi: X \rightarrow M$, a tangent vector $v \in \mathcal{T}_{X,x}$ at a point $x \in X$ will be called vertical if $\pi_*(v) = 0 \in \mathcal{T}_{M,\pi(x)}$, and a holomorphic vector field on X will be called vertical if its value at each point is vertical.

A system of co-ordinates (z_1, \dots, z_n) on a complex manifold M gives natural linear co-ordinates $(\theta_1, \dots, \theta_n)$ on the tangent spaces $\mathcal{T}_{M,p}$ by writing a tangent vector in the form $v = \sum_i \theta_i \cdot \frac{\partial}{\partial z_i}$. We thus obtain a system of co-ordinates $(z_1, \dots, z_n, \theta_1, \dots, \theta_n)$ on the total space of the tangent bundle \mathcal{T}_M , which we refer to as being induced by the co-ordinate system (z_1, \dots, z_n) on M .

2 Complex hyperkähler manifolds and affine symplectic fibrations

In this section we introduce the notion of an affine symplectic fibration of a complex hyperkähler manifolds, and give a local description of an arbitrary complex hyperkähler manifold admitting such a map.

2.1 Affine symplectic fibrations

A *complex hyperkähler structure* on a complex manifold X consists of a non-degenerate symmetric bilinear form $g: \mathcal{T}_X \otimes \mathcal{T}_X \rightarrow \mathcal{O}_X$, together with three

holomorphic endomorphisms $I, J, K \in \text{End}(\mathcal{T}_X)$ satisfying the quaternion relations

$$I^2 = J^2 = K^2 = IJK = -1,$$

which preserve the form g , and which are covariantly constant with respect to the associated holomorphic Levi-Civita connection. By a *complex hyperkähler manifold* we mean a complex manifold equipped with a complex hyperkähler structure.

Remark 2.1 Viewing the complex manifold X as consisting of the underlying smooth manifold $X_{\mathbb{R}}$ equipped with a complex structure $S \in \text{End}(T_{X_{\mathbb{R}}})$, we can use the identification $T_{X_{\mathbb{R}}} \otimes_{\mathbb{R}} \mathbb{C} \cong \mathcal{T}_X \oplus \overline{\mathcal{T}}_X$ to define endomorphisms $I_{\mathbb{R}}, J_{\mathbb{R}}, K_{\mathbb{R}} \in \text{End}(T_{X_{\mathbb{R}}})$ which commute with S and satisfy the quaternion relations, together with a non-degenerate symmetric bilinear form $g_{\mathbb{R}}: T_{X_{\mathbb{R}}} \otimes T_{X_{\mathbb{R}}} \rightarrow \mathbb{R}$ satisfying $g_{\mathbb{R}}(SX, SY) = -g_{\mathbb{R}}(X, Y)$. This defines a hyperkähler structure on the smooth manifold $X_{\mathbb{R}}$ in the usual sense, except that the metric $g_{\mathbb{R}}$ is indefinite.

Given a complex hyperkähler structure on a complex manifold X we define a holomorphic symplectic form Ω_I on X by the rule $\Omega_I(X, Y) = g(X, I(Y))$. The holomorphic symplectic forms Ω_J and Ω_K are defined analogously. We also introduce the combinations $\Omega_{\pm} = \Omega_J \pm i\Omega_K$. These are closed, holomorphic 2-forms, but are not symplectic. Indeed, since

$$g(v_1, (I \pm i)v_2) = g(K(v_1), (J \pm iK)(v_2)) = \Omega_{\pm}(K(v_1), v_2), \tag{3}$$

the kernels of the forms Ω_{\pm} are precisely the eigenspaces of $\mp i$ with eigenvalues $\mp pi$. These two subspaces are half-dimensional, and are exchanged by the action of J .

Definition 2.2 Let X be a complex hyperkähler manifold. A holomorphic map $\pi: X \rightarrow M$ is an *affine symplectic fibration* if there exists a holomorphic symplectic form ω on M such that $\Omega_{-} = \pi^*(\omega)$.

We call the symplectic form ω the *base symplectic form* of the affine symplectic fibration. For an explanation of the name affine symplectic see Remark 2.4 below.

Remark 2.3 The vertical tangent vectors for π are clearly contained in the kernel of the form $\Omega_{-} = \pi^*(\omega)$. The assumption that ω is non-degenerate ensures that on the open dense subset where π is a submersion this inclusion is an equality. But by (3), the kernel of Ω_{-} also coincides with the $+i$ eigenspace of the operator I , and is therefore everywhere half-dimensional. It follows that π is a submersion with half-dimensional fibres.

Let $\mathcal{V}_{\pi} \subset \mathcal{T}_X$ denote the bundle of vertical tangent vectors for the map $\pi: X \rightarrow M$. Consider a nonzero vertical tangent vector $v \in \mathcal{T}_{X,x}$ at some point $x \in X$. Since J switches the two eigenspaces of I , the tangent vector $J(v)$ is not vertical. Thus the map $v \mapsto \pi_*(J(v))$ induces an isomorphism $\mathcal{V}_{\pi,x} \rightarrow \mathcal{T}_{M,\pi(x)}$. Putting these maps together gives a bundle isomorphism

$$b: \mathcal{V}_{\pi} \rightarrow \pi^*(\mathcal{T}_M)$$

which we will call the *basing map* of the affine symplectic fibration.

Remark 2.4 Note that by (3), the kernel of the form Ω_+ is precisely the $-i$ eigenspace of I . Thus Ω_+ restricts to a holomorphic symplectic form on each fibre of the map $\pi: X \rightarrow M$. The basing map relates this symplectic form to the symplectic form ω on the base M . Indeed, if $v_1, v_2 \in \mathcal{T}_{X,x}$ are vertical tangent vectors, then since $(J - iK)J = J(J + iK)$, we have

$$\Omega_+(v_1, v_2) = \Omega_-(J(v_1), J(v_2)) = (\pi^*\omega)(J(v_1), J(v_2)) = \omega(b(v_1), b(v_2)).$$

This now gives the reason for the name affine symplectic fibration: any given fibre $\pi^{-1}(m)$ of the map $\pi: X \rightarrow M$ has the property that each of its tangent spaces is identified via the basing map with the fixed symplectic vector space $\mathcal{T}_{M,m}$.

2.2 Standard example

Let $n \geq 2$ be an even integer, and take $M = \mathbb{C}^n$. Let X denote the total space of the holomorphic tangent bundle \mathcal{T}_M , with its canonical projection $\pi: X \rightarrow M$. Let (z_1, \dots, z_n) be standard linear co-ordinates on M , and let $(z_1, \dots, z_n, \theta_1, \dots, \theta_n)$ be the induced co-ordinate system on X , as explained in the notation and terminology section.

Choose an $n \times n$ non-degenerate, skew-symmetric matrix ω_{pq} and introduce the holomorphic symplectic form

$$\omega = \sum_{p,q} \omega_{pq} \cdot dz^p \wedge dz^q,$$

on M . Let $X^0 \subset X$ be an open subset, and let $W: X^0 \rightarrow \mathbb{C}$ be a holomorphic function satisfying the partial differential equations

$$\frac{\partial}{\partial \theta_k} \left(\frac{\partial^2 W}{\partial \theta_i \partial z_j} - \frac{\partial^2 W}{\partial \theta_j \partial z_i} - \sum_{p,q} \eta^{pq} \cdot \frac{\partial^2 W}{\partial \theta_i \partial \theta_p} \cdot \frac{\partial^2 W}{\partial \theta_j \partial \theta_q} \right) = 0, \tag{4}$$

where η^{ij} is the inverse matrix to ω_{ij} . Then define vector fields

$$v_i = \frac{\partial}{\partial \theta_i}, \quad h_i = \frac{\partial}{\partial z_i} + \sum_{p,q} \eta^{pq} \cdot \frac{\partial^2 W}{\partial \theta_i \partial \theta_p} \cdot \frac{\partial}{\partial \theta_q}. \tag{5}$$

We define a complex hyperkähler structure on X^0 by setting

$$I(v_j) = i \cdot v_j, \quad J(v_j) = h_j, \quad K(v_j) = -i h_j, \tag{6}$$

$$I(h_j) = -i \cdot h_j, \quad J(h_j) = -v_j, \quad K(h_j) = -i v_j. \tag{7}$$

$$g(v_i, v_j) = 0, \quad g(v_i, h_j) = \omega_{ij}, \quad g(h_i, h_j) = 0. \tag{8}$$

The quaternion relations are immediate, as is the fact that I, J and K preserve the metric. It is clear that $[v_i, v_j] = 0$, and the Eq. (4) implies that $[h_i, h_j] = 0$. Note that

$$g([v_i, h_j], h_k) = \frac{\partial^3 W}{\partial \theta_i \partial \theta_j \partial \theta_k}.$$

Using the Koszul formula to compute the Levi-Civita connection, most terms vanish, and we find that

$$\nabla_{h_i}(h_j) = - \sum_{p,q} \eta^{pq} \cdot \frac{\partial^3 W}{\partial \theta_i \partial \theta_j \partial \theta_p} \cdot h_q, \quad \nabla_{h_i}(v_j) = - \sum_{p,q} \eta^{pq} \cdot \frac{\partial^3 W}{\partial \theta_i \partial \theta_j \partial \theta_p} \cdot v_q, \tag{9}$$

and $\nabla_{v_i}(h_j) = 0 = \nabla_{v_i}(v_j)$. It is then easy to check that ∇ preserves I, J and K .

The associated holomorphic symplectic forms are

$$\begin{aligned} \Omega_I &= -i \cdot \sum_{p,q} \omega_{pq} \cdot v^p \wedge h^q, \\ \Omega_J &= \frac{1}{2} \cdot \sum_{p,q} \omega_{pq} (v^p \wedge v^q + h^p \wedge h^q), \quad \Omega_K = -\frac{i}{2} \cdot \sum_{p,q} \omega_{pq} (v^p \wedge v^q - h^p \wedge h^q) \end{aligned}$$

where we used the dual bases of covectors

$$h^j = dz^j, \quad v^j = d\theta^j + \sum_{r,s} \eta^{jr} \cdot \frac{\partial^2 W}{\partial \theta_r \partial \theta_s} \cdot dz^s$$

so that $(h^j, v_i) = 0 = (v^j, h_i)$ and $(h^j, h_i) = \delta_{ij} = (v^j, v_i)$. In particular

$$\Omega_- = \Omega_J - i\Omega_K = \sum_{p,q} \omega_{pq} \cdot h^p \wedge h^q = \pi^*(\omega),$$

which shows that the restriction $\pi : X^0 \rightarrow M$ is an affine symplectic fibration.

Remark 2.5 Note that since only the second derivatives of the function W with respect to the fibre variables θ_i play any role in the definition of the hyperkähler structure, this function is only well-defined up to the addition of functions which are at most linear in the fibre directions. The transformation of W under symplectic co-ordinate changes $(z_1, \dots, z_n) \mapsto (w_1, \dots, w_n)$ on the base M is written out explicitly in [8, Section 4.2]. In addition to the obvious substitutions, the function W picks up an extra term which is cubic in the fibre directions.

Remark 2.6 A calculation with the Eq. (4) shows that the curvature component

$$R(h_i, h_j) = \nabla_{h_i} \circ \nabla_{h_j} - \nabla_{h_j} \circ \nabla_{h_i} = 0,$$

and it is immediate that $R(v_i, v_j) = 0$. On the other hand

$$R(h_j, v_i)(h_k) = \sum_{p,q} \eta^{pq} \cdot \frac{\partial^4 W}{\partial \theta_i \partial \theta_j \partial \theta_k \partial \theta_p} \cdot h_q,$$

$$R(h_j, v_i)(v_k) = \sum_{p,q} \eta^{pq} \cdot \frac{\partial^4 W}{\partial \theta_i \partial \theta_j \partial \theta_k \partial \theta_p} \cdot v_q.$$

Thus we conclude that the metric is flat precisely if the restriction of the function $W : X \rightarrow \mathbb{C}$ to each fibre $\pi^{-1}(m) = \mathcal{T}_{M,p}$ is a polynomial function of degree ≤ 3 .

2.3 Normalised affine symplectic fibrations

Consider the total space $X = \mathcal{T}_M$ of the holomorphic tangent bundle of M , with its natural projection $\pi : X \rightarrow M$. Denote by $\mathcal{V}_\pi \subset \mathcal{T}_X$ the bundle of vertical tangent vectors. There is a canonical bundle isomorphism

$$\nu : \mathcal{V}_\pi \rightarrow \pi^*(\mathcal{T}_M)$$

sending a vertical tangent vector $v \in \mathcal{T}_{X,x}$ to the corresponding tangent vector $\nu(v) \in \mathcal{T}_{M,\pi(x)}$.

Definition 2.7 Let M be a complex manifold. Suppose given a complex hyperkähler structure on an open subset X^0 of the total space $X = \mathcal{T}_M$, for which the restriction of the projection map $\pi : X \rightarrow M$ is an affine symplectic fibration. Then we call this affine symplectic fibration *normalised* if the basing map b coincides with the natural map ν .

Note that the examples of Example 2.2 are normalised in this sense, since in the co-ordinate system considered there, the map ν is defined by $\nu(\frac{\partial}{\partial \theta_i}) = \frac{\partial}{\partial z_i}$, and it is immediate from the definition that $\pi_*(J(\frac{\partial}{\partial \theta_i})) = \frac{\partial}{\partial z_i}$. Conversely, we have the following:

Proposition 2.8 *Let M be a complex manifold, and suppose given a complex hyperkähler structure on an open subset X^0 of the total space $X = \mathcal{T}_M$, for which the restriction of the projection map $\pi : X \rightarrow M$ is a normalised affine symplectic fibration. Then, shrinking X^0 if necessary, the hyperkähler structure arises via the construction of Sect. 2.2.*

Proof By the holomorphic Darboux theorem, we can find local co-ordinates z_i on the base M such that the base symplectic form can be written in the form $\sum_{i,j} \omega_{ij} \cdot dz^i \wedge dz^j$ for some constant skew-symmetric matrix ω_{ij} . Take the induced co-ordinate system (z_i, θ_j) on X and set $v_i = \frac{\partial}{\partial \theta_i}$ and $h_i = J(v_i)$. Then, as in Remark 2.3, since the v_i are vertical tangent vector fields they satisfy $I(v_i) = i v_i$, and it follows that in the basis of vector fields (v_i, h_j) the operators I, J, K are given by the formulae of Sect. 2.2.

The normalisation condition is the statement that $\pi_*(h_i) = \frac{\partial}{\partial z_i}$, and we can therefore write

$$h_j = \frac{\partial}{\partial z_j} + \sum_{p,q} \eta^{pq} \cdot c_{jp}(z, \theta) \cdot \frac{\partial}{\partial \theta_q}$$

for some locally-defined holomorphic functions $c_{jp}(z, \theta)$. Since I preserves the metric g , the eigenspaces of I are necessarily isotropic. Thus the metric is determined by

$$2g(v_i, h_j) = -g(h_j, (J - iK)(h_i)) = \Omega_-(h_i, h_j) = \omega(\pi_*(h_i), \pi_*(h_j)) = 2\omega_{ij},$$

and therefore coincides with that of Sect. 2.2.

Consider next the expression

$$g([v_i, h_j], h_k) = \frac{\partial}{\partial \theta_i} c_{jk}(z, \theta).$$

We claim that the expression on the right is completely symmetric in i, j, k . To prove this, note first that $[v_i, h_j]$ is a vertical vector field. Since ∇ preserves the eigenspace decomposition of I , the relation $\nabla_{v_i}(h_j) - \nabla_{h_j}(v_i) = [v_i, h_j]$ implies that $\nabla_{v_i}(h_j) = 0$, and so

$$g([v_i, h_j], h_k) = -g(\nabla_{h_j}(v_i), h_k).$$

Similarly, since $[h_i, h_j]$ is vertical, both sides of the expression $\nabla_{h_i}(h_j) - \nabla_{h_j}(h_i) = [h_i, h_j]$ must vanish. Since J is covariantly constant, and $g(h_j, v_k)$ is constant, we therefore have

$$g(\nabla_{h_j}(v_i), h_k) = g(\nabla_{h_j}(J(v_i)), J(h_k)) = -g(\nabla_{h_j}(h_i), v_k) = g(h_i, \nabla_{h_j}(v_k)),$$

which together with $\nabla_{h_i}(h_j) = \nabla_{h_j}(h_i)$ proves the required symmetry property.

It now follows that we can write h_i in the form (5) for some locally-defined function $W = W(z, \theta)$. The relations $[h_i, h_j] = 0$ obtained above then imply that W satisfies the partial differential Eq. (4), which completes the proof. \square

2.4 Developing maps

In this section we prove that an arbitrary complex hyperkähler manifold with an affine symplectic fibration is locally isomorphic to one arising from the construction of Sect. 2.2. For this purpose we introduce the following notion:

Definition 2.9 Let $\pi : X \rightarrow M$ be an affine symplectic fibration. A *developing map* defined on an open subset $U \subset X$ is defined to be an open embedding $f : U \rightarrow \mathcal{T}_M$, commuting with the projections to M , such that

$$v(f_*(v)) = \pi_*(J(v)), \tag{10}$$

for any vector field v on U which is vertical for the restriction of the map π .

Proposition 2.10 *Let $\pi : X \rightarrow M$ be an affine symplectic fibration. Then a developing map exists in a neighbourhood of any given point $x \in X$.*

Proof Let z_i be local Darboux co-ordinates on M for the base symplectic structure ω , as in the proof of the previous result. Since π is a submersion, we can complete these to local co-ordinates (z_i, ϕ_i) on X so that the vertical tangent spaces are spanned by the vector fields $v_i = \frac{\partial}{\partial \phi_i}$. Let us write

$$h_i = J(v_i) = \sum_j a_{ij}(z, \phi) \cdot \frac{\partial}{\partial z_j} + \sum_j b_{ij}(z, \phi) \cdot \frac{\partial}{\partial \phi_j}$$

for some locally-defined holomorphic functions $a_{ij}(z, \phi)$ and $b_{ij}(z, \phi)$.

Consider the induced co-ordinate system (z_i, θ_j) on the total space \mathcal{T}_M . Then $v(\frac{\partial}{\partial \theta_j}) = \frac{\partial}{\partial z_j}$, and the defining property of the developing map $\theta_j = \theta_j(z_i, \phi_i)$ becomes the condition that

$$\sum_j \frac{\partial \theta_j}{\partial \phi_i} \cdot \frac{\partial}{\partial z_j} = \sum_j a_{ij}(z, \phi) \cdot \frac{\partial}{\partial z_j},$$

for all i . To show the local existence of such a map we must check that $\frac{\partial a_{ij}}{\partial \phi_k} = \frac{\partial a_{kj}}{\partial \phi_i}$. In more intrinsic terms this is the statement that the expression

$$\pi_*([v_k, h_i]) = \sum_j \frac{\partial a_{ij}}{\partial \phi_k} \cdot \frac{\partial}{\partial z_j}$$

is symmetric under exchanging i and k . Since ∇ preserves the eigenspaces of I , and hence the sub-bundle of vertical vector fields, the relation $[v_k, h_i] = \nabla_{v_k}(h_i) - \nabla_{h_i}(v_k)$ implies that

$$\pi_*([v_k, h_i]) = \pi_*(\nabla_{v_k}(h_i)).$$

Using the fact that J is covariantly constant we get

$$\nabla_{v_k}(h_i) - \nabla_{v_i}(h_k) = J(\nabla_{v_k}(v_i) - \nabla_{v_i}(v_k)) = J([v_i, v_k]) = 0,$$

which gives the required symmetry. It is easy to see from the relation (10) that the derivative of the developing map is an isomorphism, so restricting its domain if necessary we can assume that it is a local embedding. \square

The next result gives the promised local description of complex hyperkähler manifolds with an affine symplectic fibration.

Proposition 2.11 *Let $\pi : X \rightarrow M$ be a complex hyperkähler manifold equipped with an affine symplectic fibration. Take a point $x \in X$ and let $f : U \rightarrow \mathcal{T}_M$ be a developing map defined on an open neighbourhood $x \in U \subset X$. Set $V = f(U) \subset \mathcal{T}_M$, and use the resulting isomorphism $f : U \rightarrow V$ to transfer the complex hyperkähler structure from U to V . Then, after possibly shrinking U , the hyperkähler structure on V arises from the construction of Sect. 2.2.*

Proof Since the developing map commutes with the projections to M it is clear that the restriction of the projection $\pi : \mathcal{T}_M \rightarrow M$ is an affine symplectic fibration for the transferred hyperkähler structure on V . By definition of the developing map it is moreover normalised. The result therefore follows from Proposition 2.8. \square

3 Joyce structures

In this section we give a more geometric definition of the notion of a Joyce structure from [6], at least in the case when the form η appearing there is non-degenerate. A Joyce structure is a rather complicated combination of geometric structures which one expects to find on the space of stability conditions on a CY_3 triangulated category. The simplest part of this structure, explained in Sect. 3.1, is obtained from the basic properties of spaces of stability conditions, in particular the existence of a local isomorphism to a vector space. The rest of the data of a Joyce structure, discussed in Sect. 3.2, is induced by the Donaldson–Thomas invariants of the category in a rather indirect and involved way which is explained in [6] and briefly summarised in the “Appendix” section.

3.1 Spaces with period maps

Let M be a complex manifold and \mathcal{H} a holomorphic vector bundle on M . By a *bundle of lattices* in \mathcal{H} we mean a holomorphically-varying collection of subgroups $\mathcal{L}_p \subset \mathcal{H}_p$ in the fibres of \mathcal{H} such that the induced maps $\mathcal{L}_p \otimes_{\mathbb{Z}} \mathbb{C} \rightarrow \mathcal{H}_p$ are all isomorphisms. Any such bundle of lattices determines a locally-constant subsheaf $\mathcal{L} \subset \mathcal{T}_M$ consisting of the holomorphic sections of \mathcal{T}_M whose values at each point $p \in M$ lie in the subgroup \mathcal{L}_p . This local system in turn determines a unique flat connection on the bundle \mathcal{H} , whose flat sections are precisely the \mathbb{C} -linear combinations of the sections of the subsheaf \mathcal{L} .

By a *period map* on a complex manifold M we mean a local isomorphism

$$\varpi : M \rightarrow \text{Hom}_{\mathbb{Z}}(\Gamma, \mathbb{C}), \tag{11}$$

where $\Gamma \cong \mathbb{Z}^{\oplus n}$ is a free abelian group of finite rank. Thus ϖ is a holomorphic map from M to the vector space $\text{Hom}_{\mathbb{Z}}(\Gamma, \mathbb{C})$ whose derivative

$$(D\varpi)_p : \mathcal{T}_{M,p} \rightarrow \text{Hom}_{\mathbb{Z}}(\Gamma, \mathbb{C}), \tag{12}$$

at each point $p \in M$ is an isomorphism. The inverse images of the subgroup $\Gamma^* = \text{Hom}_{\mathbb{Z}}(\Gamma, \mathbb{Z})$ under the maps (12) define a bundle of lattices in \mathcal{T}_M , and we therefore obtain a flat connection ∇ . The period map ϖ now gives a distinguished vector field $Z \in \Gamma(M, \mathcal{T}_M)$ via the assignment

$$p \mapsto Z_p = (D\varpi)_p^{-1}(\varpi(p)) \in \mathcal{T}_{M,p}.$$

Let us express all this in co-ordinates. Take a basis $(\gamma_1, \dots, \gamma_n)$ for the lattice Γ . We obtain holomorphic functions

$$z_i : M \rightarrow \mathbb{C}, \quad z_i(p) = \varpi(p)(\gamma_i),$$

and the assumption that ϖ is a local isomorphism ensures that (z_1, \dots, z_n) form a system of co-ordinates on M . The dual of the derivative of ϖ at a point $p \in M$ sends $\gamma_i \in \Gamma$ to the element $dz^i \in \mathcal{T}_{M,p}^*$. This implies that the connection ∇ is torsion-free and that the z_i are flat co-ordinates for this connection. The distinguished vector field is $Z = \sum_i z_i \cdot \frac{\partial}{\partial z_i}$. Note that it follows that $\nabla(Z) = \text{id}$.

In general we would like to consider complex manifolds M with a locally-defined period map ϖ . Such spaces arise for example as discrete quotients of manifolds with a globally-defined period map. We therefore make the following

Definition 3.1 A *period structure* on a complex manifold M consists of data

- (P1) a bundle of lattices $\mathcal{L} \subset \mathcal{T}_M$ whose associated flat connection we denote by ∇ ;
- (P2) a distinguished vector field $Z \in \Gamma(M, \mathcal{T}_M)$ satisfying $\nabla(Z) = \text{id}$.

Take a point $p \in M$ and set $\Gamma^* = \mathcal{L}_p$. A basis of the free abelian group Γ^* extends uniquely to a basis of ∇ -flat sections ϕ_1, \dots, ϕ_n of the tangent bundle \mathcal{T}_M over a contractible open neighbourhood $p \in M^0 \subset M$. Writing the vector field Z in the form $Z = \sum_i z_i \cdot \phi_i$ then defines holomorphic functions $z_i : M^0 \rightarrow \mathbb{C}$, and condition (P2) then implies that $\phi_i = \frac{\partial}{\partial z_i}$. It follows that on the open neighbourhood $p \in M^0 \subset M$ the structure arises from a period map as explained above. Note in particular that the connection ∇ is necessarily torsion-free.

In the situations of interest below we consider period maps (11) in which the lattice Γ has a natural skew-symmetric integral form $\eta : \Gamma \times \Gamma \rightarrow \mathbb{Z}$. The local version of this is

Definition 3.2 A *period structure with skew-form* consists of a period structure as above, together with a ∇ -flat skew-symmetric form

$$\eta : \mathcal{T}_M^* \times \mathcal{T}_M^* \rightarrow \mathcal{O}_M,$$

which takes integral values on the lattices $\mathcal{L}^* \subset \mathcal{T}_M^*$.

We will be particularly interested in the case when the form η is non-degenerate. The inverse then defines a complex symplectic form

$$\omega : \mathcal{T}_M \times \mathcal{T}_M \rightarrow \mathcal{O}_M,$$

taking rational values on the lattices $\mathcal{L} \subset \mathcal{T}_M$.

3.2 Joyce structures

We can now give a geometric definition which is a slight strengthening of that of a Joyce structure given in [6]. We discuss the differences in Remark 3.4 below.

Definition 3.3 Let M be a complex manifold, and let X denote the total space of the holomorphic tangent bundle. A *strong Joyce structure* on M consists of

- (a) a period structure with skew-form (\mathcal{L}, Z, η) on M ;
- (b) a complex hyperkähler structure (g, I, J, K) on X ;

satisfying the following conditions

- (J1) the canonical projection $\pi : X \rightarrow M$ is a normalised affine symplectic fibration;
- (J2) the base symplectic structure $\omega \in \Gamma(M, \wedge^2 \mathcal{T}_M)$ is the inverse to the form η ;
- (J3) the involution $\iota : X \rightarrow X$ acting by -1 on the fibres of π satisfies

$$\iota^*(g) = -g, \quad \iota^*(I) = I, \quad \iota^*(J \pm iK) = -(J \pm iK);$$

- (J4) the ∇ -horizontal lift $E \in \Gamma(X, \mathcal{T}_X)$ of the vector field Z satisfies

$$\mathcal{L}_E(g) = g, \quad \mathcal{L}_E(I) = 0, \quad \mathcal{L}_E(J \pm iK) = mpJ \pm iK);$$

- (J5) the hyperkähler structure is invariant under translations by the lattice $(2\pi i) \cdot \mathcal{L} \subset \mathcal{T}_M$.

As in Sect. 3.1, we can take ∇ -flat local co-ordinates z_i on M such that the lattice $\mathcal{L} \subset \mathcal{T}_{M,p}$ is spanned by the vector fields $\frac{\partial}{\partial z_i}$. Let (z_i, θ_j) be the induced co-ordinate system on X . Condition (J2) together with Definition 3.2 ensure that the symplectic form ω is covariantly constant and can therefore be written in the form $\omega = \sum_{i,j} \omega_{ij} \cdot dz_i \wedge dz_j$, with ω_{ij} a non-degenerate, skew-symmetric matrix. Moreover, the inverse matrix η^{ij} is integral. Condition (J1) and Proposition 2.8 then imply that the complex hyperkähler structure (g, I, J, K) arises from the formulae of Example 2.2 for some locally-defined holomorphic function $W : X \rightarrow \mathbb{C}$ satisfying the equations

$$\frac{\partial}{\partial \theta_k} \left(\frac{\partial^2 W}{\partial \theta_i \partial z_j} - \frac{\partial^2 W}{\partial \theta_j \partial z_i} - \sum_{p,q} \eta^{pq} \cdot \frac{\partial^2 W}{\partial \theta_i \partial \theta_p} \cdot \frac{\partial^2 W}{\partial \theta_j \partial \theta_q} \right) = 0. \tag{13}$$

As in Remark 2.5, the function $W = W(z, \theta)$ is only well-defined up to the addition of functions which are at most linear in the θ_i variables. Condition (J3) is equivalent to the statement that it may be taken to be an odd function of the θ_i co-ordinates, in the sense that it satisfies $i^*(W) = -W$. To see this note first that $i^*(z_j) = z_j$ and $i^*(\theta_j) = -\theta_j$. Consulting (5) we see that the condition $i^*(J) = -J$ is equivalent to the second derivatives appearing there being odd. If this holds, then modifying W by functions at most linear in the co-ordinates θ_i we can assume that W has the same property.

Definition 3.1 shows that $Z = \sum_i z_i \cdot \frac{\partial}{\partial z_i}$, and the ∇ -horizontal lift E is given by the same formula. Condition (J4) then becomes the statement that W can be taken

to be homogeneous of degree -1 under simultaneous rescaling of the variables z_i . Finally, condition (J5) is equivalent to the statement that the second derivatives of W with respect to the fibre variables θ_j are invariant under transformations of the form $\theta_j \mapsto \theta_j + 2\pi i$.

We can in fact assume that W satisfies the simplified equation appearing in [6]:

$$\frac{\partial^2 W}{\partial \theta_i \partial z_j} - \frac{\partial^2 W}{\partial \theta_j \partial z_i} = \sum_{p,q} \eta^{pq} \cdot \frac{\partial^2 W}{\partial \theta_i \partial \theta_p} \cdot \frac{\partial^2 W}{\partial \theta_j \partial \theta_q}. \tag{14}$$

Indeed, if we replace W by the expression

$$W(z_i, \theta_j) - \sum_k \theta_k \cdot \frac{\partial W}{\partial \theta_k}(z_i, 0),$$

then each of the first two terms on the left-hand side of (14) vanishes along the locus where all $\theta_i = 0$. But the third term also vanishes along this locus, because W is odd in the co-ordinates θ_j . Since the Eq. (13) state that the left-hand side of (14) is independent of the co-ordinates θ_j , after the above modification it must vanish identically.

Remark 3.4 There are three differences between the above definition of a strong Joyce structure and the definition of a Joyce structure in [6].

- (i) The above formulation assumes that the form η is non-degenerate. This is not required in the definition of [6]. To include degenerate forms in the framework of this paper we would have to consider generalisations of the notion of a hyperkähler structure which have the endomorphisms I, J, K , but in which the metric is dropped and replaced with a symmetric, bilinear form on the cotangent bundle.
- (ii) In [6] it was only assumed that the third, rather than the second, derivatives of the function W in the fibre directions were periodic. This was to allow the inclusion of certain examples obtained by doubling spaces of stability conditions for which the form η is degenerate, and even vanishing.
- (iii) The definition in [6] allows the function W to be meromorphic. Of course, we could also introduce meromorphic complex hyperkähler structures to deal with this generalisation.

Let us briefly consider the linearisation procedure of [6, Section 7]. Identifying M with the zero-section $M \subset X$, the tangent bundle \mathcal{T}_M becomes a sub-bundle of the restriction $\mathcal{T}_X|_M$. The fact that W is an odd function of the θ_i co-ordinates ensures that along the zero section $M \subset X$ we have $h_i = \frac{\partial}{\partial z_i}$. It follows that the Levi-Civita connection preserves the tangents to the zero section and hence induces a torsion-free connection ∇^J on \mathcal{T}_M . Moreover, Remark 2.6 together with the oddness of the function W ensure that this connection is flat. From Eq. (9) we see that it is given explicitly by

$$\nabla_{\frac{\partial}{\partial z_i}} \left(\frac{\partial}{\partial z_j} \right) = - \sum_{p,q} \eta^{pq} \cdot \frac{\partial^3 W}{\partial \theta_i \partial \theta_j \partial \theta_p} \Big|_{\theta=0} \cdot \frac{\partial}{\partial z_q}. \tag{15}$$

This is what was referred to as the linearised Joyce connection in [6].

3.3 Twistor space

We conclude by making some brief remarks on the twistor space of a complex hyperkähler manifold. Note that the existence of such a twistor space is not obvious since the quotienting out by a distribution could potentially result in an object that is not Hausdorff. Conditions guaranteeing that the space of complex null geodesics is Hausdorff are described in [21].

Consider the quadric in the complex projective plane

$$Q = \{[a : b : c] \in \mathbb{P}^2 : a^2 + b^2 + c^2 = 0\}.$$

For each point $q = [a : b : c] \in Q$ the kernel of the operator $aI + bJ + cK$ defines a half-dimensional sub-bundle $\mathcal{H}(q) \subset \mathcal{T}_X$. The fact that I, J, K are flat for the Levi-Civita connection ∇ ensures that this sub-bundle is involutive, since if v, w are sections of $\mathcal{H}(q)$ then so is $[v, w] = \nabla_v(w) - \nabla_w(v)$. Thus there is a corresponding foliation $\mathcal{F}(q)$ of the space X by complex submanifolds. We define the twistor space Z to be the set of pairs (q, \mathcal{L}) , where $q \in Q$, and \mathcal{L} is a leaf of the foliation $\mathcal{F}(q)$. There is an obvious projection $\pi : Z \rightarrow Q \cong \mathbb{P}^1$. Each point $x \in X$ then defines a section of this projection by sending a point $q \in Q$ to the unique leaf of the foliation \mathcal{F}_q containing the point $x \in X$.

Note that if $\pi : X \rightarrow M$ is an affine symplectic fibration, the sub-bundle $\mathcal{H}(q) \subset \mathcal{T}_X$ corresponding to the point $q = [0 : 1 : -i] \in Q$ is the vertical sub-bundle defined by the map π , and so the corresponding fibre of the twistor space $Z_q = \pi^{-1}(q)$ is naturally identified with M .

If we identify \mathbb{P}^1 with the quadric Q via the isomorphism

$$[s : t] \mapsto [2ist, s^2 + t^2, i(s^2 - t^2)],$$

then the sub-bundle $\mathcal{H}(q)$ becomes the kernel of the operator

$$2istI + s^2(J + iK) + t^2(J - iK).$$

Given a local basis of vector fields v_i and h_j on X in which the operators I, J, K take the simple form (6)–(7), this sub-bundle is spanned by the vector fields $sv_i + th_i$.

Remark 3.5 In the context of Sect. 2.2 the sub-bundle $\mathcal{H}(q)$ is spanned by

$$\epsilon^{-1} \cdot \frac{\partial}{\partial \theta_i} + \frac{\partial}{\partial z_i} + \sum_{p,q} \eta^{pq} \cdot \frac{\partial^2 W}{\partial \theta_i \partial \theta_p} \cdot \frac{\partial}{\partial \theta_q}, \tag{16}$$

where we set $\epsilon = t/s$. In [6] these sub-bundles $\mathcal{H}(\epsilon)$ played a central role, and were viewed as defining a pencil of Ehresmann connections on the map $\pi : X \rightarrow M$.

It seems interesting to try to relate the Riemann–Hilbert problems considered in [6] to the geometry of the twistor space Z . Note in particular, that the solutions to the Riemann–Hilbert problem at a particular value $\epsilon \in \mathbb{C}^*$ are annihilated by the flows (16), and are therefore constant on the leaves of the corresponding foliation $\mathcal{F}(q)$. Thus such

solutions naturally define functions on the fibre $Z_q = \pi^{-1}(q)$ of the twistor space. In the context of categories defined by quivers with potential this suggests a relation between these fibres and the associated cluster Poisson variety.

4 Quantum DT invariants and deformations of anti-self-duality

In [27] an integrable deformation of Plebański's first heavenly equation was constructed using ideas from the deformation quantization programme. The nonlinear terms in both the first and second Plebański equations comes from the Poisson bracket

$$\{f, g\} = \sum_{i,j=1}^2 \eta^{ij} \frac{\partial f}{\partial \theta_i} \frac{\partial g}{\partial \theta_j}$$

and deformations of this bracket - preserving the Jacobi identity - date back to the work of Moyal who introduced such a structure in his phase space approach to quantum mechanics. The Moyal bracket can be defined in terms of the operator

$$P = \exp \left[\frac{i\hbar}{2} \eta^{ij} \overleftarrow{\partial}_i \overrightarrow{\partial}_j \right]$$

(where the arrows show which direction the derivatives are to be taken) and with this one defines the non-commutative but associative product $f * g = fPg$ and the corresponding Moyal bracket

$$\{f, g\}_M = \frac{f * g - g * f}{i\hbar}.$$

Since

$$\lim_{\hbar \rightarrow 0} \{f, g\}_M = \{f, g\}$$

this is a deformation of the normal Poisson bracket with \hbar being the deformation parameter. The Jacobi identity for the Moyal bracket follows trivially from the associative property of the underlying $*$ -product.

One can also introduce differential operators \hat{X}_f which contain higher derivatives (formally, as these are to all orders in \hbar) that are deformations of Hamiltonian vector fields, and have the property

$$[\hat{X}_f, \hat{X}_g] = \hat{X}_{\{f, g\}_M}$$

where the left-hand side is commutator of operators. These operator no longer have a clear geometric interpretation but one can use them instead of Hamiltonian vector fields in Lax equations. Applying this idea results in the Moyal deformation of the first Plebański equation

$$\left\{ \frac{\partial W}{\partial z_1}, \frac{\partial W}{\partial z_2} \right\}_M = 1$$

and, in the same way, to the Moyal deformed version of Plebański’s second heavenly equation [26]

$$\frac{\partial^2 W}{\partial z_1 \partial \theta_2} - \frac{\partial^2 W}{\partial z_2 \partial \theta_1} = \left\{ \frac{\partial W}{\partial \theta_1}, \frac{\partial W}{\partial \theta_2} \right\}_M. \tag{17}$$

The integrability of such equations remains, though in a formal setting. In [27] solutions were constructed as a formal power series in \hbar and in [30] the equations were derived from a Riemann–Hilbert splitting problem in the associated Moyal loop group. The ideas are easily extended to give deformation of the system of equations (1).

The $*$ -product, and hence the Moyal bracket, has a very rich structure. In terms of a torus basis,

$$e^{\theta(\alpha)} * e^{\theta(\beta)} = (-q^{\frac{1}{2}})^{\langle \alpha, \beta \rangle} e^{\theta(\alpha+\beta)},$$

where $\langle \alpha, \beta \rangle = \eta^{ij} \alpha_i \beta_j$, and hence

$$\{e^{\theta(\alpha)}, e^{\theta(\beta)}\}_M = \frac{1}{i\hbar} \left[(-q^{\frac{1}{2}})^{\langle \alpha, \beta \rangle} - (-q^{-\frac{1}{2}})^{\langle \alpha, \beta \rangle} \right] e^{\theta(\alpha+\beta)}, \tag{18}$$

$$= \frac{2}{\hbar} \sin \left[\frac{\hbar \langle \alpha, \beta \rangle}{2} \right] e^{\theta(\alpha+\beta)}, \tag{19}$$

where $q = e^{i(\hbar+2\pi)}$ (the $e^{2\pi i}$ term is to ensure the correct sign for the square root). This algebra has a number of names: the sine-algebra [12] and the quantum torus algebra [20]. Since

$$\lim_{\hbar \rightarrow 0} \{e^{\theta(\alpha)}, e^{\theta(\beta)}\}_M = \langle \alpha, \beta \rangle e^{\theta(\alpha+\beta)}.$$

in the classical limit the quantum torus algebra becomes the twisted torus algebra.

With the same ansatz

$$W = \sum_{\alpha} F_{\alpha}(z) \frac{e^{\theta(\alpha)}}{z(\alpha)}$$

with F_{α} of degree zero, one obtains the isomonodromy equation

$$dF_{\gamma} = \sum_{\alpha+\beta=\gamma} \frac{1}{i\hbar} \left\{ \mathbb{L}^{\frac{1}{2}\langle \alpha, \beta \rangle} - \mathbb{L}^{-\frac{1}{2}\langle \alpha, \beta \rangle} \right\} F_{\alpha} F_{\beta} d \log z(\beta).$$

where $\mathbb{L} = e^{i\hbar}$. The construction of solutions to such equations is still at a very early stage, and even in the simplest case, it involves highly intricate calculations [7]. But it appears that the quantum DT invariants play the same role in the construction

of solutions to the deformed Eq. (17) as DT invariants play in the construction of solutions to Eq. (1).

With the replacement of Hamilton vector fields with infinite order differential operators the link with geometry becomes less clear. At the level of Lax pairs, the replacement of Hamiltonian vector fields with operators seems minor, but these are infinite-order operators, and so the resulting differential equations are of infinite order. Also, any notion of hyperkähler geometry is lost - though it is intriguing to speculate where there is a notion of q -deformed hyperkähler structures and q -deformed twistor spaces (see, for example, [19,28]).

Acknowledgements The first author is grateful to the Sydney Mathematical Research Institute for support and hospitality while this research was being carried out, and to the Royal Society for financial support. He is also grateful to Dominic Joyce for explaining the twistor construction of Section 3.3. Both authors would like to thank Maciej Dunajski for his comments on an earlier version of this paper.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Appendix A. Summary of [6]

For the convenience of the reader we give a brief summary of the main ideas of [6]. The key point is that the Donaldson-Thomas (DT) invariants of a CY_3 triangulated category can be interpreted as Stokes data for a family of connections over the space of stability conditions. These connections take values in an infinite-dimensional group G of automorphisms of the space $(\mathbb{C}^*)^n$ equipped with a constant Poisson structure.

A.1. Stokes data

We will start by considering Stokes data in the context of the finite-dimensional group $GL_n(\mathbb{C})$. Our treatment is based on that in [4], to which we refer the reader for references to the original literature. Set $G = GL_n(\mathbb{C})$ and $\mathfrak{g} = \mathfrak{gl}_n(\mathbb{C})$, and let $\mathfrak{h} \subset \mathfrak{g}$ denote the Cartan subalgebra of diagonal matrices. Introduce the standard root decomposition

$$\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{g}^{\text{od}}, \quad \mathfrak{g}^{\text{od}} = \bigoplus_{\alpha \in \Phi} \mathfrak{g}_{\alpha}, \quad \Phi = \{e_i^* - e_j^*\} \subset \mathfrak{h}^*.$$

Let us consider a meromorphic connection on the trivial G -bundle over \mathbb{P}^1 of the form

$$\nabla = d - \left(\frac{U}{\epsilon^2} + \frac{V}{\epsilon} \right), \quad (20)$$

where $U, V \in \mathfrak{g}$ are constant matrices such that

- (i) $U = \text{diag}(u_1, \dots, u_n) \in \mathfrak{h}^{\text{reg}}$ is diagonal with distinct eigenvalues,
- (ii) $V \in \mathfrak{g}^{\text{od}}$ has zeroes on the diagonal.

The connection ∇ has a regular singularity at $\epsilon = \infty$, but the singularity at $\epsilon = 0$ is irregular, and is essentially the simplest example of such a singularity. It is a classical fact that the gauge equivalence of a connection in a neighbourhood of an irregular singularity is not determined solely by the monodromy: we must also consider its Stokes data.

For any ray $r = \mathbb{R}_{>0} \cdot z \subset \mathbb{C}^*$ we denote by $\mathbb{H}_r \subset \mathbb{C}^*$ the half-plane centered on it. The Stokes rays of the connection (20) at $\epsilon = 0$ are defined to be the rays

$$\mathbb{R}_{>0} \cdot (u_i - u_j) = \mathbb{R}_{>0} \cdot U(\alpha) \subset \mathbb{C}^*, \quad \alpha = e_i^* - e_j^*.$$

We then have the following fundamental existence result [1]:

Theorem A.1 (Balsler, Jurkat, Lutz) *For any non-Stokes ray $r \subset \mathbb{C}^*$ there is a unique flat section $Y_r : \mathbb{H}_r \rightarrow G$ of the connection (20) such that*

$$Y_r(\epsilon) \cdot \exp(U/\epsilon) \rightarrow 1 \text{ as } \epsilon \rightarrow 0.$$

Using this result we can associate to each Stokes ray ℓ a Stokes factor

$$\mathbb{S}_\ell = Y_{r_+}(\epsilon) \cdot Y_{r_-}(\epsilon)^{-1} \in \exp \left(\bigoplus_{U(\alpha) \in \ell} \mathfrak{g}_\alpha \right) \subset G,$$

where r_+ and r_- are small clockwise and anti-clockwise perturbations of the ray ℓ .

Let us now consider varying the diagonal matrix $U \in \mathfrak{h}^{\text{reg}}$. It turns out that we can uniquely deform the matrix $V \in \mathfrak{g}^{\text{od}}$ so that the Stokes factors $\mathbb{S}(\ell)$ remain constant. Such deformations are called *isomonodromic* or *iso-Stokes*, and the variation of $V = V(U)$ is described by the partial differential equation

$$d \log V_\gamma = \sum_{\alpha+\beta=\gamma} [V_\alpha, V_\beta] \cdot d \log U(\beta), \quad V = \sum_{\gamma \in \Phi} V_\gamma \in \mathfrak{g}^{\text{od}}. \tag{21}$$

In fact the isomonodromy condition stated above is not sufficiently precise. As $U \in \mathfrak{h}^{\text{reg}}$ varies, the Stokes rays $\mathbb{R}_{>0} \cdot U(\alpha)$ may cross, at which point the statement that the Stokes factors are constant ceases to make sense. The correct condition to impose is that for any convex sector $\Delta \subset \mathbb{C}^*$, the clockwise product

$$\mathbb{S}_p(\Delta) = \prod_{\ell \in \Delta} \mathbb{S}_p(\ell) \in G,$$

of Stokes factor associated to rays in the sector should be constant as $U \in \mathfrak{h}^{\text{reg}}$ varies, providing that no Stokes ray crosses the boundary of Δ .

Let us suppose given the matrix $U \in \mathfrak{h}^{\text{reg}}$ and the Stokes factors $\mathbb{S}(\ell) \in G$, and consider the inverse problem of reconstructing the connection (20). To do this we can

first try to construct the canonical half-plane solutions $Y_r(t)$ and then differentiate to obtain the operator V . This leads to the following Riemann–Hilbert boundary-value problem:

Riemann–Hilbert problem. For each non-Stokes ray $r \subset \mathbb{C}^*$ find a holomorphic function $Y_r : \mathbb{H}_r \rightarrow G$ such that the following three properties hold:

- (RH1) $Y_r(\epsilon) \cdot \exp(U/\epsilon) \rightarrow 1$ as $t \rightarrow 0$ in \mathbb{H}_r ,
- (RH2) there exists $k > 0$ such that $|\epsilon|^{-k} < \|Y_r(\epsilon)\| < |\epsilon|^k$ as $\epsilon \rightarrow \infty$ in \mathbb{H}_r ,
- (RH3) if $\Delta \subset \mathbb{C}^*$ is a convex sector with $\partial\Delta = \{r_+\} \cup \{r_-\}$ then

$$Y_{r_+}(\epsilon) = Y_{r_-}(\epsilon) \cdot \mathbb{S}(\Delta) \text{ for } \epsilon \in \mathbb{H}_{r_+} \cap \mathbb{H}_{r_-}.$$

Note that the canonical half-plane solutions of Theorem A.1 satisfy (RH1) by definition, and (RH3) holds by the definition of the Stokes factors. The condition (RH2) follows from the fact that the Eq. (20) has a regular singularity at $\epsilon = \infty$, so that solutions have moderate growth at this point.

A.2. Stability conditions and DT invariants

Let \mathcal{D} be a \mathbb{C} -linear triangulated category of finite type. We assume for simplicity that the Grothendieck group

$$\Gamma := K_0(\mathcal{D}) \cong \mathbb{Z}^{\oplus n}$$

is free of finite rank. The expression

$$\langle [E], [F] \rangle = \sum_i (-1)^i \dim_{\mathbb{C}} \text{Hom}^i(E, F[i]),$$

defines a bilinear form $\langle -, - \rangle : \Gamma \times \Gamma \rightarrow \mathbb{Z}$ known as the *Euler form*.

The data of a stability condition on \mathcal{D} consists of a group homomorphism $Z : \Gamma \rightarrow \mathbb{C}$ called the *central charge*, and for each $\phi \in \mathbb{R}$ a full subcategory $\mathcal{P}(\phi) \subset \mathcal{D}$ whose objects are said to be *semistable* of phase ϕ . This data is required to satisfy a simple set of axioms [3]. The important fact is then that the space $\text{Stab}(\mathcal{D})$ of all such stability conditions on the category \mathcal{D} is a complex manifold, and the forgetful map

$$\varpi : \text{Stab}(\mathcal{D}) \rightarrow \text{Hom}_{\mathbb{Z}}(\Gamma, \mathbb{C}) \tag{22}$$

sending a stability condition to its central charge is a local isomorphism of complex manifolds.

To be able to define Donaldson-Thomas (DT) invariants we must assume that the category \mathcal{D} satisfies the CY_3 condition

$$\text{Hom}_{\mathcal{D}}^i(A, B) \cong \text{Hom}_{\mathcal{D}}^{3-i}(B, A)^*.$$

This implies that the Euler form is skew-symmetric. Thus in the terminology of Sect. 3.1 the manifold $\text{Stab}(\mathcal{D})$ is naturally equipped with a period structure with skew-form. Our aim is to use DT theory to enrich this to a Joyce structure.

Let us fix a stability condition $\sigma \in \text{Stab}(\mathcal{D})$. Given further assumptions on the pair (\mathcal{D}, σ) it is possible to define DT invariants $\text{DT}_\sigma(\gamma) \in \mathbb{Q}$ for each class $\gamma \in \Gamma$. In the simplest case, when there are no strictly-semistable objects of class $\gamma \in \Gamma$, and the moduli stack $\mathcal{M}^\sigma(\gamma)$ of semistable objects is smooth, the invariant $\text{DT}_\sigma(\gamma)$ coincides up to sign with the Euler characteristic of the coarse moduli space of $\mathcal{M}^\sigma(\gamma)$ viewed as a complex manifold. The definition in the general case is due to Joyce and Song [17], and Kontsevich and Soibelman [20]. There is an equivalent system of invariants $\Omega_\sigma(\gamma) \in \mathbb{Q}$ defined by

$$\text{DT}_\sigma(\alpha) = \sum_{\alpha=k\beta} \frac{1}{k^2} \cdot \Omega_\sigma(\beta),$$

These appear in physics as BPS invariants, and in many cases are known to be integers.

A.3. The wall-crossing formula

Continuing with the notation of the previous section, the next step is to consider the dependence of the DT invariants on the stability condition $\sigma \in \text{Stab}(\mathcal{D})$. It turns out that for a fixed class $\gamma \in \Gamma$, the invariant $\text{DT}_\sigma(\gamma)$ is constant in the complement of a collection of real codimension-one submanifolds in the space $\text{Stab}(\mathcal{D})$, but jumps discontinuously as the stability condition crosses one of these walls. Joyce [16,17], and Kontsevich and Soibelman [20], were able to describe this wall-crossing behaviour exactly, in such a way that knowledge of all invariants $\text{DT}_\sigma(\gamma)$ at one point $\sigma \in \text{Stab}(\mathcal{D})$ determines them at all other points. Even more remarkably, in the formulation of [20], the resulting wall-crossing formula is exactly the iso-Stokes condition for a family of differential equations of the form (20), but with the finite-dimensional group $\text{GL}_n(\mathbb{C})$ replaced by an infinite-dimensional group of Poisson automorphisms of the space $(\mathbb{C}^*)^n$.

To explain this in more detail, introduce the algebraic torus

$$\mathbb{T} = \text{Hom}_{\mathbb{Z}}(\Gamma, \mathbb{C}^*) \cong (\mathbb{C}^*)^n, \quad \mathbb{C}[\mathbb{T}] = \bigoplus_{\gamma \in \Gamma} \mathbb{C} \cdot x_\gamma,$$

whose character lattice is Γ . The Euler form defines an invariant Poisson structure on \mathbb{T} given on characters by

$$\{x_\alpha, x_\beta\} = \langle \alpha, \beta \rangle \cdot x_{\alpha+\beta}.$$

To make connection with the material of Section A.1. we would like to consider the group

$$G = \text{Aut}_{\{-,-\}}(\mathbb{T})$$

of algebraic Poisson automorphisms of the variety \mathbb{T} . Note that the group structure on the torus \mathbb{T} plays no role in this definition. The corresponding Lie algebra \mathfrak{g} consists of algebraic vector fields on \mathbb{T} whose flows preserve the Poisson structure. The fact that these objects are infinite-dimensional will make some aspects of the following discussion heuristic. Precise discussions can be found in [5].

For simplicity we assume that the form $\langle -, - \rangle$ is non-degenerate. There is then a root decomposition

$$\mathfrak{g} = \text{Vect}_{\langle -, - \rangle}(\mathbb{T}) = \mathfrak{h} \oplus \mathfrak{g}^{\text{od}}, \tag{23}$$

where the Cartan subalgebra $\mathfrak{h} \cong \text{Hom}_{\mathbb{Z}}(\Gamma, \mathbb{C})$ consists of translation-invariant vector fields on \mathbb{T} , and the subspace $\mathfrak{g}^{\text{od}} \subset \mathfrak{g}$ consists of Hamiltonian vector fields, and can be identified with the Poisson algebra of non-constant algebraic functions on \mathbb{T} :

$$\mathfrak{g}^{\text{od}} = \bigoplus_{\alpha \in \Gamma \setminus \{0\}} \mathfrak{g}_{\alpha} = \bigoplus_{\alpha \in \Gamma \setminus \{0\}} \mathbb{C} \cdot x^{\alpha}.$$

Fix a stability condition $\sigma \in \text{Stab}(\mathcal{D})$. For each ray $\ell \subset \mathbb{C}^*$ we can attempt to define an automorphism of \mathbb{T} by taking the time 1 Hamiltonian flow of the corresponding DT generating function viewed as a regular function on \mathbb{T} . The action of this automorphism on characters is

$$\mathbb{S}_{\sigma}(\ell)^*(x_{\beta}) = \exp \left\{ - \sum_{Z(\gamma) \in \ell} \text{DT}_{\sigma}(\gamma) \cdot x_{\gamma}, - \right\} (x_{\beta}) = x_{\beta} \cdot \prod_{Z(\gamma) \in \ell} (1 - x_{\gamma})^{\Omega_{\sigma}(\gamma) \cdot \langle \gamma, \beta \rangle}. \tag{24}$$

Since the sum and product here could be infinite, making rigorous sense of this requires further work [5,20]. Let us call a ray $\ell \subset \mathbb{C}^*$ active if $\mathbb{S}(\ell)$ is not the identity. The wall-crossing formula can now be stated as follows: for any convex sector $\Delta \subset \mathbb{C}^*$ the product

$$\mathbb{S}_{\sigma}(\Delta) = \prod_{\ell \in \Delta} \mathbb{S}_{\sigma}(\ell) \in G$$

is constant as $\sigma \in \text{Stab}(\mathcal{D})$ varies, providing the boundary of Δ remains non-active.

A.4. Joyce structures

Comparing the results of the last two subsections we come to the remarkable conclusion that the wall-crossing formula is the isomonodromy condition for a family of meromorphic connections on \mathbb{P}^1 of the form

$$\nabla = d - \left(\frac{Z}{\epsilon^2} + \frac{\text{Ham}_F}{\epsilon} \right) d\epsilon, \tag{25}$$

parameterised by the points of $\text{Stab}(\mathcal{D})$ and taking values in the group $G = \text{Aut}_{\{-,-\}}(\mathbb{T})$ of Section A.3. Here Z and Ham_F are constant elements of the Lie algebra (23) such that

- (i) $Z \in \mathfrak{h}$ is the central charge $Z: \Gamma \rightarrow \mathbb{C}$, viewed as an invariant vector field on \mathbb{T} ,
- (ii) $\text{Ham}_F \in \mathfrak{g}^{\text{od}}$ is the Hamiltonian vector field of a regular function $F = \sum_{\gamma \in \Gamma} F_\gamma \cdot x^\gamma$.

Let us choose a basis $(\gamma_1, \dots, \gamma_n) \subset \Gamma$ giving co-ordinates $z_i = Z(\gamma_i)$ on $\text{Stab}(\mathcal{D})$, and $\theta_i = \theta(\gamma_i)$ on its tangent spaces, which are all identified with the fixed vector space $\text{Hom}_{\mathbb{Z}}(\Gamma, \mathbb{C})$ by the derivative of the period map (22). Set $\eta^{ij} = \langle \gamma_i, \gamma_j \rangle$.

Let X be the total space of the tangent bundle of $\text{Stab}(\mathcal{D})$ and define a holomorphic function $W: X \rightarrow \mathbb{C}$ by the formula

$$W(z_i, \theta_j) = \sum_{\gamma \in \Gamma^\times} F_\gamma(z_1, \dots, z_n) \cdot \frac{e^{\theta(\gamma)}}{Z(\gamma)}.$$

The variation of $F = F(Z)$ is controlled by the isomonodromy equation (21), which in the Lie algebra (23) takes the form

$$dF_\gamma = \sum_{\alpha+\beta=\gamma} \langle \alpha, \beta \rangle \cdot F_\alpha F_\beta \cdot d \log Z(\beta).$$

when written in terms of the Joyce function W this becomes

$$\frac{\partial^2 W}{\partial \theta_i \partial z_j} - \frac{\partial^2 W}{\partial \theta_j \partial z_i} = \sum_{p,q} \eta^{pq} \cdot \frac{\partial^2 W}{\partial \theta_i \partial \theta_p} \cdot \frac{\partial^2 W}{\partial \theta_j \partial \theta_q}, \tag{26}$$

which obviously implies the Eq. (4).

Reconstructing the connection (25), and hence the Joyce function W , from the Stokes data (24) involves solving a Riemann–Hilbert boundary-value problem analogous to the one stated in Section A.1. In the context of the infinite-dimensional Lie algebra (23) these problems were considered in detail in [5,6].

References

1. Balsler, W., Jurkat, W.B., Lutz, D.A.: Birkhoff invariants and Stokes’ multipliers for meromorphic linear differential equations. *J. Math. Anal. Appl.* **71**(1), 48–94 (1979)
2. Boyer, C., Plebański, J.: An infinite hierarchy of conservation laws and nonlinear superposition principles for self-dual Einstein spaces. *J. Math. Phys.* **26**, 229 (1985)
3. Bridgeland, T.: Stability conditions on triangulated categories. *Ann. Math.* **166**, 31–345 (2007)
4. Bridgeland, T., Toledano Laredo, V.: Stability conditions and Stokes factors. *Invent. Math.* **187**(1), 61–98 (2012)
5. Bridgeland, T.: Riemann–Hilbert problems from Donaldson–Thomas theory. *Invent. Math.* **216**, 69–124 (2019)
6. Bridgeland, T.: Geometry from Donaldson–Thomas invariants, preprint [arXiv:1912.06504](https://arxiv.org/abs/1912.06504)
7. Bridgeland, T., Barbieri, A., Stoppa, J.: A quantized Riemann–Hilbert problem in Donaldson–Thomas theory, preprint [arXiv:1905.00748](https://arxiv.org/abs/1905.00748) (to appear IMRN)

8. Bridgeland, T.: On the monodromy of the deformed cubic oscillator, with an appendix by D. Masoero, preprint [arXiv:2006.10648](https://arxiv.org/abs/2006.10648)
9. Bridgeland, T.: A complexification of the Hitchin system, in preparation
10. Dunajski, M.: Null Kähler geometry and isomonodromic deformations, preprint [arXiv:2010.11216](https://arxiv.org/abs/2010.11216)
11. Dunajski, M., Tod, K.P.: Einstein-Weyl structures from hyper-Kähler Metrics with conformal Killing vectors. *Diff. Geom. Appl.* **14**, 39–55 (2001)
12. Fairlie, D.B., Zachos, C.: Infinite-dimensional algebras, sine brackets, and $SU(\infty)$. *Phys. Lett. B* **224**(1–2), 101–107 (1989)
13. Gaiotto, D., Moore, G., Neitzke, A.: Four-dimensional wall-crossing via three-dimensional field theory. *Comm. Math. Phys.* **299**(1), 163–224 (2010)
14. Gaiotto, D., Moore, G., Neitzke, A.: Wall-crossing, Hitchin systems, and the WKB approximation. *Adv. Math.* **234**, 239–403 (2013)
15. Gindikin, S.G.: A construction of hyper-Kähler metrics. *Funct. Anal. Appl.* **20**(3), 238–240 (1986)
16. Joyce, D.: Configurations in abelian categories. IV. Invariants and changing stability conditions, *Adv. in Math.* **217**, 125–204 (2008)
17. Joyce, D., Song, Y.: A theory of generalized Donaldson-Thomas invariants, *Mem. Amer. Math. Soc.* **217**, no. 1020, iv+199 pp (2012)
18. Joyce, D.: Holomorphic generating functions for invariants counting coherent sheaves on Calabi-Yau 3-folds. *Geom. Topol.* **11**, 667–725 (2007)
19. Kapustin, A., Kuznetsov, A., Orlov, D.: Noncommutative instantons and twistor transform. *Commun. Math. Phys.* **221**, 385–432 (2001)
20. Kontsevich, M., Soibelman, Y.: Stability structures, motivic Donaldson-Thomas invariants and cluster transformations, [arxiv 0811.2435](https://arxiv.org/abs/0811.2435)
21. LeBrun, C.: Spaces of complex null geodesics in complex-Riemannian geometry, *Trans. A.M.S.* **278**(1), 209–231 (1983)
22. Mason, L.J., Newman, E.T.: A connection between the Einstein and Yang-Mills equations. *Commun. Math. Phys.* **121**, 659–668 (1989)
23. Han Park, Q.: Self-dual gravity as a large-N limit of the 2D non-linear sigma model. *Phys. Lett.* **238B**, 287–90 (1990)
24. Penrose, R.: Non-linear gravitons and curved twistor theory. *Gen. Relativ. Gravit.* **7**, 31–52 (1976)
25. Plebański, J.F.: Some solutions of complex Einstein equations. *J. Math. Phys.* **16**, 2395–2402 (1975)
26. Plebański, J.F., Przanowski, M., Rajca, B., Tosiek, J.: The moyal deformation of the second heavenly equation. *Acta Phys. Pol. B* **26**, 889 (1995)
27. Strachan, I.A.B.: The Moyal algebra and integrable deformations of the self-dual Einstein equations. *Phys. Lett. B* **283**(1–2), 63–66 (1992)
28. Strachan, I.A.B.: A geometry for multidimensional integrable systems. *J. Geom. Phys.* **21**(3), 255–278 (1997)
29. Takasaki, K.: An infinite number of hidden variables in hyper-Kähler metrics. *J. Math. Phys.* **30**, 1515–1521 (1989)
30. Takasaki, K.: Dressing operator approach to Moyal algebraic deformations of selfdual gravity. *J. Geom. Phys.* **14**(3), 111 (1994)
31. Ward, R.S.: Integrable and solvable systems, and relations among them, *Phil. Trans. Roy. Soc. A*, (315), issue 1533, 451–457 (1985)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.