



Approximation and application of the Riesz-Caputo fractional derivative of variable order with fixed memory

Tomasz Blaszczyk · Krzysztof Bekus · Krzysztof Szajek · Wojciech Sumelka

Received: 10 November 2020 / Accepted: 19 April 2021 / Published online: 7 May 2021
© The Author(s) 2021

Abstract In this paper, the Riesz-Caputo fractional derivative of variable order with fixed memory is considered. The studied non-integer differential operator is approximated by means of modified basic rules of numerical integration. The three proposed methods are based on polynomial interpolation: piecewise constant, piecewise linear, and piecewise quadratic interpolation. The errors generated by the described methods and the experimental rate of convergence are reported. Finally, an application of the Riesz-Caputo fractional derivative of space-dependent order in continuum mechanics is depicted.

Keywords Riesz-Caputo derivative · variable order · fractional continua · numerical solution

1 Introduction

In the last decades, fractional calculus has become a powerful mathematical tool to model a lot of physical processes. Because of the nature of fractional derivatives (nonlocal operators), they are especially used in describing nonlocal and multiscale phenomena. In this regard, we refer the reader works on the following topics: viscoelastic materials [6, 23, 28], heat conduction [24, 40, 54, 56], diffusion processes in complex systems [12, 30, 32], nonlocal constitutive laws [13, 19, 46], microscopic interaction forces [18, 25], nonlocal stress-strain constitutive relations [3], non-local beams [2, 8, 10, 41, 47]. A fairly broad overview of various aspects of fractional calculus applications can be found in the review paper by Failla and Zingales [20].

Notwithstanding this branch of mathematics is not new (the creation of fractional calculus dates back to the birth of the classical theory of differential calculus [27]) there are still remain a lot of problems that require attention. In particular, the branch of fractional calculus, that is devoted to differential equations containing simultaneously both the left and right fractional derivatives, is a research area where exists a large number of an unsolved/unexplored issues [15, 21, 29]. No exact solutions, for many of the aforementioned equations, have been found till now. For this reason, many researchers focus their attention on approximate solutions of equations containing

T. Blaszczyk (✉) · K. Bekus
Czestochowa University of Technology, Department of Mathematics, al. Armii Krajowej 21, 42-200 Czestochowa, Poland
e-mail: tomasz.blaszczyk@pcz.pl

K. Bekus
e-mail: krzysztof.bekus@im.pcz.pl

K. Szajek · W. Sumelka
Poznan University of Technology, Institute of Structural Analysis, Piotrowo 5 street, 60-965 Poznan, Poland
e-mail: krzysztof.szajek@put.poznan.pl

W. Sumelka
e-mail: wojciech.sumelka@put.poznan.pl

various types of combinations of fractional derivatives. For this reason, many researchers focus their attention on approximate solutions of equations containing various types of combinations of fractional derivatives [1, 7, 11, 14, 22, 53].

On the other hand, certain types of phenomena in physics are better depicted when the order of the fractional operator is variable (it depends on the spatial or time variable) [26, 43]. Promising approaches have been presented in papers where authors studied: diffusion processes in an inhomogeneous and heterogeneous medium [16, 33, 44, 49–51, 55], control theory [17, 34, 35], mechanics [4, 31, 36, 38, 39, 42]. More details of the available applications of a variable order fractional calculus in the area of scientific and engineering modelling are presented in the following review papers: [37, 52].

In this paper we focus on the Riesz-Caputo fractional operator of variable order with fixed memory and its application in a strong form of the space-fractional continuum (1D case), which is just a straightforward extension of the results presented in [45]. Such approach allows, when adopted to the mechanical applications, for clear physical interpretation and moreover gives smooth passage to classical approach as a limit case [5, 48]. We propose three methods, based on polynomial interpolation, to approximation of the studied fractional operator. All presented numerical sachems were implemented in Python using library for real and complex floating-point arithmetic with arbitrary precision (<http://mpmath.org/>).

2 Approximation of the Riesz-Caputo fractional derivative with variable order

Let us consider the following Riesz-Caputo fractional derivative of variable order with α depending on space variable

$$\begin{aligned} {}_{x-\ell}^{RC} D_x^{\alpha(x)} f(x) &= \frac{1}{2} ({}_{x-\ell}^C D_x^{\alpha(x)} f(x) + \\ &(-1)^n {}_x^C D_{x+\ell}^{\alpha(x)} f(x)) \end{aligned} \quad (1)$$

Operators ${}_{x-\ell}^C D_x^{\alpha(x)}$ and ${}_x^C D_{x+\ell}^{\alpha(x)}$ are well known fractional Caputo derivatives, with fixed memory length ℓ and variable order $\alpha(x) > 0$, defined as

$$\begin{aligned} {}_{x-\ell}^C D_x^{\alpha(x)} f(x) &= \frac{1}{\Gamma(n-\alpha(x))} \int_{x-\ell}^x (x-\tau)^{n-\alpha(x)-1} f^{(n)}(\tau) d\tau \end{aligned} \quad (2)$$

$$\begin{aligned} {}_x^C D_{x+\ell}^{\alpha(x)} f(x) &= \frac{(-1)^n}{\Gamma(n-\alpha(x))} \int_x^{x+\ell} (\tau-x)^{n-\alpha(x)-1} f^{(n)}(\tau) d\tau \end{aligned} \quad (3)$$

where $n-1 < \alpha(x) < n$.

In following, based on our previous work [9] where the numerical schemes for differential operators of constant order α were developed, we elaborate the algorithms for the Riesz-Caputo derivative of variable order. For discretisation, we generate the grid of $(N+1)$ – equidistant nodes: $x_0 < \dots < x_{i-m} < x_{i-m+1} < \dots < x_i < \dots < x_{i+m-1} < x_{i+m} < \dots < x_N$ where $x_i = x_0 + i\Delta x$, $m = \ell/\Delta x$ and $\Delta x = (x_N - x_0)/N$. Moreover, we introduce the following notation: $f(x_i) = f_i$, $f^{(n)}(x_i) = f_i^{(n)}$, and $\alpha(x_i) = \alpha_i$.

2.1 Method I: Approximation based on piecewise constant interpolation

$$\begin{aligned} {}_{x-\ell}^{RC} D_x^{\alpha(x)} f(x) \Big|_{x=x_i} &\approx \frac{1}{2} \left(\sum_{k=i-m+1}^i f_k^{(n)} p_{i,k-1} + \sum_{k=i}^{i+m-1} f_k^{(n)} q_{i,k} \right) \end{aligned} \quad (4)$$

where

$$p_{i,k} = \frac{(\Delta x)^{n-\alpha_i}}{\Gamma(n-\alpha_i+1)} [(i-k)^{n-\alpha_i} - (i-k-1)^{n-\alpha_i}] \quad (5)$$

and

$$q_{i,k} = \frac{(\Delta x)^{n-\alpha_i}}{\Gamma(n-\alpha_i+1)} [(k-i+1)^{n-\alpha_i} - (k-i)^{n-\alpha_i}] \quad (6)$$

2.2 Method II: Approximation based on piecewise linear interpolation

$$\begin{aligned} {}_{x-\ell}^{RC} D_x^{\alpha(x)} f(x) \Big|_{x=x_i} &\approx \frac{1}{2} \left(\sum_{k=i-m}^i f_k^{(n)} r_{i,k} + \sum_{k=i}^{i+m} f_k^{(n)} s_{i,k} \right) \end{aligned} \quad (7)$$

where

$$r_{i,k} = \frac{(\Delta x)^{n-\alpha_i}}{\Gamma(n-\alpha_i+2)} \times \begin{cases} (m-1)^{n-\alpha_i+1} + (n-\alpha_i-m+1)m^{n-\alpha_i} & \text{for } k = i-m \\ (i-k+1)^{n-\alpha_i+1} - 2(i-k)^{n-\alpha_i+1} + (i-k-1)^{n-\alpha_i+1} & \text{for } k = i-m+1 \dots i-1 \\ 1 & \text{for } k = i \end{cases} \quad (8)$$

and

$$s_{i,k} = \frac{(\Delta x)^{n-\alpha_i}}{\Gamma(n-\alpha_i+2)} \times \begin{cases} (m-1)^{n-\alpha_i+1} + (n-\alpha_i-m+1)m^{n-\alpha_i} & \text{for } k = i+m \\ (k-i+1)^{n-\alpha_i+1} - 2(k-i)^{n-\alpha_i+1} + (k-i-1)^{n-\alpha_i+1} & \text{for } k = i+1 \dots i+m-1 \\ 1 & \text{for } k = i \end{cases} \quad (9)$$

2.3 Method III: Approximation based on piecewise quadratic interpolation

$$\left. {}_{x-\ell}^{RC} D_{x+\ell}^{\alpha(x)} f(x) \right|_{x=x_i} \approx \frac{1}{2} (\Delta x)^{n-\alpha_i} \times \begin{cases} \sum_{k=-1}^1 f_{i+k}^{(n)} u_k + \sum_{k=i-m}^{i-1} f_k^{(n)} w_{i,k} + \sum_{k=-1}^1 f_{i-k}^{(n)} u_k + \sum_{k=i+1}^{i+m} f_k^{(n)} v_{i,k} & \text{for odd } m \\ \sum_{k=i-m}^i f_k^{(n)} w_{i,k} + \sum_{k=i}^{i+m} f_k^{(n)} v_{i,k} & \text{for even } m \end{cases} \quad (10)$$

where

$$w_{i,k} = \begin{cases} \frac{m^{n-\alpha_i+2} - (m-2)^{n-\alpha_i+2}}{\Gamma(n-\alpha_i+3)} - \frac{3m^{n-\alpha_i+1} + (m-2)^{n-\alpha_i+1}}{2\Gamma(n-\alpha_i+2)} \\ + \frac{m^{n-\alpha_i}}{\Gamma(n-\alpha_i+1)} & \text{for } k = i-m \\ -2 \frac{(i-k+1)^{n-\alpha_i+2} - (i-k-1)^{n-\alpha_i+2}}{\Gamma(n-\alpha_i+3)} \\ +2 \frac{(i-k+1)^{n-\alpha_i+1} + (i-k-1)^{n-\alpha_i+1}}{\Gamma(n-\alpha_i+2)} & \text{for odd } k+i-m \\ \frac{(i-k+2)^{n-\alpha_i+2} - (i-k-2)^{n-\alpha_i+2}}{\Gamma(n-\alpha_i+3)} \\ - \frac{(i-k+2)^{n-\alpha_i+1} + 6(i-k)^{n-\alpha_i+1} + (i-k-2)^{n-\alpha_i+1}}{2\Gamma(n-\alpha_i+2)} & \text{for even } k+i-m \text{ and } k < i-1 \\ \frac{2^{n-\alpha_i+2}}{\Gamma(n-\alpha_i+3)} - \frac{2^{n-\alpha_i+1}}{2\Gamma(n-\alpha_i+2)} & \text{for even } m \text{ and } k = i \\ \frac{3^{n-\alpha_i+2} - 1}{\Gamma(n-\alpha_i+3)} - \frac{3^{n-\alpha_i+1} + 3}{2\Gamma(n-\alpha_i+2)} - \frac{1}{\Gamma(n-\alpha_i+1)} & \text{for odd } m \text{ and } k = i-1 \end{cases} \quad (11)$$

and




Fig. 1 Approximations of the Riesz-Caputo derivative of functions x^2 , $\exp x$, $\sin x$ and $\cos x$ for space-dependent order $\alpha(x) = \frac{1}{2} + \frac{x}{2}$ and $\alpha(x) = 1 - \frac{x}{2}$

$$v_{i,k} =$$

$$\left\{ \begin{array}{ll} \frac{m^{n-\alpha_i+2} - (m-2)^{n-\alpha_i+2}}{\Gamma(n-\alpha_i+3)} - \frac{3m^{n-\alpha_i+1} + (m-2)^{n-\alpha_i+1}}{2\Gamma(n-\alpha_i+2)} \\ \quad + \frac{m^{n-\alpha_i}}{\Gamma(n-\alpha_i+1)} & \text{for } k = i+m \\ -2 \frac{(k-i+1)^{n-\alpha_i+2} - (k-i-1)^{n-\alpha_i+2}}{\Gamma(n-\alpha_i+3)} \\ \quad + 2 \frac{(k-i+1)^{n-\alpha_i+1} + (k-i-1)^{n-\alpha_i+1}}{\Gamma(n-\alpha_i+2)} & \text{for odd } k+i+m \\ \frac{(k-i+2)^{n-\alpha_i+2} - (k-i-2)^{n-\alpha_i+2}}{\Gamma(n-\alpha_i+3)} & \\ - \frac{(k-i+2)^{n-\alpha_i+1} + 6(k-i)^{n-\alpha_i+1} + (k-i-2)^{n-\alpha_i+1}}{2\Gamma(n-\alpha_i+2)} & \text{for even } k+i+m \text{ and } k > i+1 \\ \frac{2^{n-\alpha_i+2}}{\Gamma(n-\alpha_i+3)} - \frac{2^{n-\alpha_i+1}}{2\Gamma(n-\alpha_i+2)} & \text{for even } m \text{ and } k = i \\ \frac{3^{n-\alpha_i+2} - 1}{\Gamma(n-\alpha_i+3)} - \frac{3^{n-\alpha_i+1} + 3}{2\Gamma(n-\alpha_i+2)} - \frac{1}{\Gamma(n-\alpha_i+1)} & \text{for odd } m \text{ and } k = i+1 \end{array} \right. \quad (12)$$

with

Table 1 Errors and experimental rates of convergence for the approximation of the Riesz-Caputo derivative of function $f(x) = \exp(x)$ for variable order $\alpha(x) = \exp(-x)$

x	Δx	Method I		Method II		Method III	
		Error	Rate	Error	Rate	Error	Rate
0.2	0.01	1.50965E-05	—	6.41933E-08	—	1.08056E-14	—
	0.005	7.57269E-06	1.00	1.65880E-08	1.95	6.87771E-16	3.97
	0.0025	3.79284E-06	1.00	4.26599E-09	1.96	4.36702E-17	3.98
	0.00125	1.89813E-06	1.00	1.09273E-09	1.96	2.76712E-18	3.98
0.4	0.01	2.42990E-05	—	7.68809E-08	—	9.25228E-15	—
	0.005	1.21786E-05	1.00	1.94861E-08	1.98	5.77257E-16	4.00
	0.0025	6.09687E-06	1.00	4.92442E-09	1.98	3.60283E-17	4.00
	0.00125	3.05037E-06	1.00	1.24163E-09	1.99	2.24927E-18	4.00
0.8	0.01	3.66332E-05	—	8.55870E-08	—	4.72786E-15	—
	0.005	1.83466E-05	1.00	2.14589E-08	2.00	2.92187E-16	4.02
	0.0025	9.18088E-06	1.00	5.37533E-09	2.00	1.81207E-17	4.01
	0.00125	4.59236E-06	1.00	1.34564E-09	2.00	1.12653E-18	4.01
1.2	0.01	4.88466E-05	—	9.92076E-08	—	1.41104E-15	—
	0.005	2.44551E-05	1.00	2.48227E-08	2.00	8.62417E-17	4.03
	0.0025	1.22355E-05	1.00	6.20887E-09	2.00	5.31507E-18	4.02
	0.00125	6.11977E-06	1.00	1.55271E-09	2.00	3.29302E-19	4.01
1.6	0.01	6.54281E-05	—	1.23339E-07	—	1.56315E-15	—
	0.005	3.27508E-05	1.00	3.08444E-08	2.00	9.89008E-17	3.98
	0.0025	1.63846E-05	1.00	7.71249E-09	2.00	6.22459E-18	3.99
	0.00125	8.19461E-06	1.00	1.92832E-09	2.00	3.90593E-19	3.99
1.8	0.01	7.64028E-05	—	1.40558E-07	—	3.16440E-15	—
	0.005	3.82421E-05	1.00	3.51467E-08	2.00	1.98766E-16	3.99
	0.0025	1.91312E-05	1.00	8.78768E-09	2.00	1.24576E-17	4.00
	0.00125	9.56817E-06	1.00	2.19706E-09	2.00	7.79820E-19	4.00

$$\begin{aligned} u_{-1} &= \frac{2(n - \alpha_i)^2 + 3(n - \alpha_i)}{2\Gamma(n - \alpha_i + 3)} \\ u_0 &= \frac{2 + 2(n - \alpha_i)}{\Gamma(n - \alpha_i + 3)} \\ u_1 &= \frac{-(n - \alpha_i)}{2\Gamma(n - \alpha_i + 3)} \end{aligned}$$

3 Numerical analysis: errors and experimental rates of convergence

In this subsection, we report on a few numerical examples for checking the accuracy of the presented methods. Algebraic problems given by Eqs. (4), (7), and (10) were built using original procedures developed by the authors in Python.

Figure 1 shows graphs of approximations of the Riesz-Caputo derivative of functions x^2 , $\exp x$, $\sin x$,

and $\cos x$ for space-dependent orders $\alpha(x) = \frac{1}{2} + \frac{x}{2}$ and $\alpha(x) = 1 - \frac{x}{2}$. The calculations presented on the plots have been performed for $N = 2000$ and $\ell = 0.2$.

In Tables 1, 2, and 3 we present the absolute errors of numerical calculations carried out using the proposed methods (Method I, Method II, and Method III) for the following functions:

- $f(x) = \exp(x)$ and $\alpha(x) = \exp(-x)$ (Table 1),
- $f(x) = \exp(x)$ and $\alpha(x) = \frac{1+x^2}{2}$ (Table 2),
- $f(x) = \exp(x)$ and $\alpha(x) = \sin(x) + 1$ (Table 3).

The reported errors have been determined by comparing numerical results with exact values (using series representation of the Riesz-Caputo derivative [9]). We take a particular value of the Riesz-Caputo operator and compare it with a value received from the series at the same specific point. In this case, for each node x_i the order of derivative has the form $\alpha(x_i)$ and is an

Table 2 Errors and experimental rates of convergence for the approximation of the Riesz-Caputo derivative of function $f(x) = \exp(x)$ for variable order $\alpha(x) = \frac{1+x^2}{2}$

x	Δx	Method I		Method II		Method III	
		Error	Rate	Error	Rate	Error	Rate
0.2	0.01	2.06366E-05	–	5.23628E-08	–	3.94876E-15	–
	0.005	1.03373E-05	1.00	1.31554E-08	1.99	2.44466E-16	4.01
	0.0025	5.17346E-06	1.00	3.30045E-09	1.99	1.51746E-17	4.01
	0.00125	2.58795E-06	1.00	8.27193E-10	2.00	9.43733E-19	4.01
0.4	0.01	2.52875E-05	–	6.95082E-08	–	6.46222E-15	–
	0.005	1.26695E-05	1.00	1.75087E-08	1.99	4.00927E-16	4.01
	0.0025	6.34134E-06	1.00	4.40176E-09	1.99	2.49195E-17	4.01
	0.00125	3.17235E-06	1.00	1.10504E-09	1.99	1.55101E-18	4.01
0.8	0.01	2.73888E-05	–	1.16803E-07	–	1.97114E-14	–
	0.005	1.37389E-05	1.00	3.01898E-08	1.95	1.25493E-15	3.97
	0.0025	6.88124E-06	1.00	7.76567E-09	1.96	7.96998E-17	3.98
	0.00125	3.44373E-06	1.00	1.98958E-09	1.96	5.05113E-18	3.98
1.2	0.01	4.47995E-05	–	8.55260E-08	–	6.66571E-16	–
	0.005	2.24256E-05	1.00	2.13896E-08	2.00	4.26259E-17	3.97
	0.0025	1.12193E-05	1.00	5.34859E-09	2.00	2.69926E-18	3.98
	0.00125	5.61127E-06	1.00	1.33732E-09	2.00	1.69983E-19	3.99
1.6	0.01	6.83960E-05	–	2.66922E-07	–	4.14854E-14	–
	0.005	3.43000E-05	1.00	6.85320E-08	1.96	2.62248E-15	3.98
	0.0025	1.71769E-05	1.00	1.75197E-08	1.97	1.65496E-16	3.99
	0.00125	8.59546E-06	1.00	4.46293E-09	1.97	1.04289E-17	3.99
1.8	0.01	7.20214E-05	–	1.28814E-07	–	4.46378E-15	–
	0.005	3.60466E-05	1.00	3.22071E-08	2.00	2.79546E-16	4.00
	0.0025	1.80323E-05	1.00	8.05228E-09	2.00	1.74906E-17	4.00
	0.00125	9.01840E-06	1.00	2.01314E-09	2.00	1.09381E-18	4.00

established number (the order α is constant for each node x_i). Moreover, all tables contain the experimental rates of convergence for each proposed method.

Analysing results presented in tables, one can observe that the convergence is the fastest for the piecewise quadratic interpolation (scheme 10), a little slower in case of the piecewise linear interpolation (scheme 7) and the slowest for the piecewise constant interpolation (scheme 4). Additionally, the errors generated by Method I are the biggest one, while for Method III are the smallest one.

4 Application of Riesz-Caputo derivative of variable order for the space-fractional continuum mechanics - numerical study

In this section we present the application of the Riesz-Caputo fractional derivative of variable order in continuum mechanics. We consider one-dimensional tension problem of the space-fractional continua in the following form [46]:

$$\frac{\partial}{\partial x} \left(\ell^{\alpha(x)-1} \Gamma(2-\alpha(x))_{x-\ell}^{RC} D_{x+\ell}^{\alpha(x)} f(x) \right) + \frac{b}{E} = 0 \quad (13)$$

with Dirichlet's boundary conditions

$$\begin{cases} f(0) = 0 \\ f(l) = 0.01l \end{cases} \quad (14)$$

where $\frac{b}{E} = 0.1$ and $l = 1$. In Eq. (18), x denotes spatial variable, ℓ is the length scale, f denotes the

Table 3 Errors and experimental rates of convergence for the approximation of the Riesz-Caputo derivative of function $f(x) = \exp(x)$ for variable order $\alpha(x) = \sin(x) + 1$

x	Δx	Method I		Method II		Method III	
		Error	Rate	Error	Rate	Error	Rate
0.2	0.01	1.60723E-05	—	3.02310E-08	—	4.09406E-16	—
	0.005	8.04511E-06	1.00	7.56003E-09	2.00	2.58751E-17	3.98
	0.0025	4.02481E-06	1.00	1.89034E-09	2.00	1.62750E-18	3.99
	0.00125	2.01297E-06	1.00	4.72631E-10	2.00	1.02088E-19	3.99
0.4	0.01	2.36586E-05	—	5.19790E-08	—	2.00026E-15	—
	0.005	1.18469E-05	1.00	1.30181E-08	2.00	1.23408E-16	4.02
	0.0025	5.92790E-06	1.00	3.25836E-09	2.00	7.64717E-18	4.01
	0.00125	2.96507E-06	1.00	8.15218E-10	2.00	4.75253E-19	4.01
0.8	0.01	3.44676E-05	—	1.18646E-07	—	1.60320E-14	—
	0.005	1.72791E-05	1.00	3.02127E-08	1.97	1.00487E-15	4.00
	0.0025	8.65136E-06	1.00	7.66646E-09	1.98	6.29514E-17	4.00
	0.00125	4.32873E-06	1.00	1.93990E-09	1.98	3.94203E-18	4.00
1.2	0.01	1.94921E-05	—	1.10806E-07	—	2.29987E-14	—
	0.005	9.78684E-06	0.99	2.93851E-08	1.91	1.50571E-15	3.93
	0.0025	4.90457E-06	1.00	7.74780E-09	1.92	9.81780E-17	3.94
	0.00125	2.45530E-06	1.00	2.03271E-09	1.93	6.37888E-18	3.94
1.6	0.01	2.09175E-07	—	1.45639E-09	—	3.37524E-16	—
	0.005	1.05104E-07	0.99	3.94470E-10	1.88	2.26121E-17	3.90
	0.0025	5.26960E-08	1.00	1.06208E-10	1.89	1.50804E-18	3.91
	0.00125	2.63879E-08	1.00	2.84493E-11	1.90	1.00174E-19	3.91
1.8	0.01	1.48806E-05	—	9.56471E-08	—	2.12801E-14	—
	0.005	7.47474E-06	0.99	2.56854E-08	1.90	1.41240E-15	3.91
	0.0025	3.74691E-06	1.00	6.85677E-09	1.91	9.33326E-17	3.92
	0.00125	1.87607E-06	1.00	1.82109E-09	1.91	6.14370E-18	3.93

Fig. 2 Spatial discretization for one-dimensional fractional continuum body





Fig. 3 The comparison of displacements through the length of the body for variable order $\alpha(x) = \frac{1}{2} + \frac{x}{2}$ (left side) and $\alpha(x) = 1 - \frac{x}{2}$ (right side), and $\ell \in \{0.01; 0.1; 0.2; 0.4\}$



displacements, b is the body force, and E stands for the Young modulus.

We start from introducing the spatial discretization for the analysed one-dimensional continuum body that is presented in Fig. 2.

Discrete form of the fractional operator $\frac{\partial}{\partial x} \left(\ell^{\alpha(x)-1} \Gamma(2 - \alpha(x))_{x-\ell}^{RC} D_{x+\ell}^{\alpha(x)} f(x) \right)$ can be described by the trapezoidal formula (Method II) as:

$$\begin{aligned} & \frac{\partial}{\partial x} \left(\ell^{\alpha(x)-1} \Gamma(2 - \alpha(x))_{x-\ell}^{RC} D_{x+\ell}^{\alpha(x)} f(x) \right) \Big|_{x=x_i} \\ & \approx \frac{\Gamma(2 - \alpha_i) \ell^{\alpha_i-1}}{2\Delta x} \\ & \times \left(\sum_{k=i-m}^i f'_k r_{i,k} + \sum_{k=i}^{i+m} f'_k s_{i,k} \right. \\ & \quad \left. - \sum_{k=i-m-1}^{i-1} f'_k r_{i-1,k} - \sum_{k=i-1}^{i+m-1} f'_k s_{i-1,k} \right) \end{aligned} \quad (15)$$

Next, after applying the backward difference formula, for the first order derivative, in Eq. (15) we obtain:

$$\begin{aligned} & \frac{\partial}{\partial x} \left(\ell^{\alpha(x)-1} \Gamma(2 - \alpha(x))_{x-\ell}^{RC} D_{x+\ell}^{\alpha(x)} f(x) \right) \Big|_{x=x_i} \\ & \approx \frac{\Gamma(2 - \alpha_i) \ell^{\alpha_i-1}}{2(\Delta x)^2} \\ & \times \left(\sum_{k=i-m}^i ((f_{k+1} - f_k) r_{i,k}) + \sum_{k=i}^{i+m} ((f_{k+1} - f_k) s_{i,k}) \right. \\ & \quad \left. - \sum_{k=i-m-1}^{i-1} ((f_{k+1} - f_k) r_{i-1,k}) - \sum_{k=i-1}^{i+m-1} ((f_{k+1} - f_k) s_{i-1,k}) \right) \end{aligned} \quad (16)$$

Finally, the following system of linear equations equations can be formulated:

$$\begin{cases} f_0 = 0 \\ \sum_{k=i-m}^i ((f_{k+1} - f_k) r_{i,k}) + \sum_{k=i}^{i+m} ((f_{k+1} - f_k) s_{i,k}) \\ - \sum_{k=i-m-1}^{i-1} ((f_{k+1} - f_k) r_{i-1,k}) - \sum_{k=i-1}^{i+m-1} ((f_{k+1} - f_k) s_{i-1,k}) = -\frac{b}{FE} \\ f_n = 0.01l \end{cases} \quad (17)$$

where

$$F = \frac{\Gamma(2 - \alpha_i) \ell^{\alpha_i-1}}{2(\Delta x)^2}$$

The system (17) (containing $N + 1$ equations) has been solved for two types of function α , namely $\alpha(x) = \frac{1}{2} + \frac{x}{2}$ and $\alpha(x) = 1 - \frac{x}{2}$ and the following parameters: $N = 200$ and $\ell \in \{0.01; 0.1; 0.2; 0.4\}$. The obtained results are presented in Fig. 3.

The analysis of Fig. 3 allows us to draw the conclusion that the deformation of the space-fractional continuum body solely depends on the length scale ℓ and the order α . Moreover, the introduction of variable order of fractional continua makes the model more flexible, thus in perspective validation of this model with experimental data should be easier.

5 Conclusions

The approximation of the Riesz-Caputo fractional derivative of variable order and with fixed memory can be elaborated using different approaches. Herein, three original methods are considered based on the polynomial interpolation: piecewise constant (Method I), piecewise linear (Method II), and piecewise quadratic (Method III) interpolation. As presented, the convergence is the fastest for Method III, a little slower in case of Method II and the slowest for Method I. Moreover, the errors generated by Method I are the biggest one, while for Method III are the smallest one. Finally, the presented schemes can be directly applied for modelling of scale effect in terms of space-fractional continuum mechanics, making its ability to mimic experimental results higher.

Acknowledgements This work is supported by the National Science Centre, Poland under Grant No. 2017/27/B/ST8/00351

Declarations

Conflicts of interest The authors declare that they have no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly

from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Almeida R, Malinowska AB, Morgado ML, Odzijewicz T (2017) Variational methods for the solution of fractional discrete/continuous Sturm-Liouville problems. *J Mech Mater Struct* 12(1):3–21
- Alotta G, Failla G, Zingales M (2014) Finite element method for a nonlocal Timoshenko beam model. *Finite Elem Anal Des* 89:77–92
- Atanackovic TM, Stankovic B (2009) Generalized wave equation in nonlocal elasticity. *Acta Mech* 208(1–2):1–10
- Atanackovic TM, Pilipovic S (2011) Hamilton's principle with variable order fractional derivatives. *Fract Calc Appl Anal* 14:94–109
- Aydinlik S, Kiris A, Sumelka W (2021) Nonlocal vibration analysis of microstretch plates in the framework of space-fractional mechanics-theory and validation. *Eur Phys J Plus* 136:169
- Bagley RL, Torvik PJ (1983) A theoretical basis for the application of fractional calculus to viscoelasticity. *J Rheol* 27(3):201–210
- Blaszczyk T, Ciesielski M (2017) Numerical solution of Euler-Lagrange equation with Caputo derivatives. *Adv Appl Math Mech* 9(1):173–185
- Blaszczyk T (2017) Analytical and numerical solution of the fractional Euler-Bernoulli beam equation. *J Mech Mater Struct* 12(1):23–34
- Blaszczyk T, Bekus K, Szajek K, Sumelka W (2021) On numerical approximation of the Riesz-Caputo operator with the fixed/short memory length. *J King Saud Univ Sci* 33(1):101220
- Blaszczyk T, Siedlecki J, Sun HG (2021) An exact solution of fractional Euler-Bernoulli equation for a beam with fixed-supported and fixed-free ends. *Appl Math Comput* 396:125932
- Bourdin L, Cresson J, Greff I, Inizan P (2013) Variational integrator for fractional Euler-Lagrange equations. *Appl Numer Math* 71:14–23
- Caputo M (2003) Diffusion with space memory modelled with distributed order space fractional differential equations. *Ann Geophys* 46:223–234
- Carpinteri A, Cornetti P, Sapora A (2014) Nonlocal elasticity: an approach based on fractional calculus. *Meccanica* 49(11):2551–2569
- Ciesielski M, Blaszczyk T (2015) Numerical solution of non-homogenous fractional oscillator equation in integral form. *J Theoret Appl Mech* 53(4):959–968
- Ciesielski M, Blaszczyk T (2017) The multiple composition of the left and right fractional Riemann-Liouville integrals: analytical and numerical calculations. *Filomat* 31(19):6087–6099
- Chechkin AV, Gorenflo R, Sokolov IM (2005) Fractional diffusion in inhomogeneous media. *J Phys A: Math Gen* 38:L679
- Dabiri A, Moghaddam B, Machado JT (2018) Optimal variable-order fractional PID controllers for dynamical systems. *J Comput Appl Math* 339:40–48
- Di Paola M, Zingales M (2008) Long-range cohesive interactions of non-local continuum faced by fractional calculus. *Int J Solids Struct* 45(21):5642–5659
- Drapaca CS, Sivaloganatha S (2012) A fractional model of continuum mechanics. *J Elast* 107:105–123
- Failla G, Zingales M (2020) Advanced materials modelling via fractional calculus: challenges and perspectives. *Philos Trans R Soc A* 378:20200050
- Klimek M (2009) On solutions of linear fractional differential equations of a variational type. The Publishing Office of the Czestochowa University of Technology, Czestochowa
- Klimek M, Ciesielski M, Blaszczyk T (2018) Exact and numerical solutions of the fractional Sturm-Liouville problem. *Fract Calcul Appl Anal* 21(1):45–71
- Koeller RC (1984) Application of fractional calculus to the theory of viscoelasticity. *J Appl Mech* 51:299–307
- Kukla S, Siedlecka U (2020) Time-fractional heat conduction in a finite composite cylinder with heat source. *J Appl Math Comput Mech* 19(2):85–94
- Lazopoulos KA (2006) Non-local continuum mechanics and fractional calculus. *Mech Res Commun* 33(6):753–757
- Lorenzo CF, Hartley TT (2002) Variable order and distributed order fractional operators. *Nonlinear Dyn* 29:57–98
- Machado JT, Kiryakova V, Mainardi F (2017) Recent history of fractional calculus. *Commun Nonlinear Sci* 16:1140–1153
- Mainardi F (2012) An historical perspective on fractional calculus in linear viscoelasticity. *Fract Calc Appl Anal* 15:712–717
- Malinowska AB, Odzijewicz T, Torres DFM (2015) Advanced methods in the fractional calculus of variations. Springer Briefs in Applied Sciences and Technology, Springer, Cham
- Meerschaert MM, Benson DA, Scheffler HP, Baeumer B (2002) Stochastic solution of space-time fractional diffusion equations. *Phys Rev E* 65:041103
- Meng R, Yin D, Drapaca CS (2019) Variable-order fractional description of compression deformation of amorphous glassy polymers. *Comput Mech* 64:163–171
- Metzler R, Klafter J (2000) The random walk's guide to anomalous diffusion: a fractional dynamics approach. *Phys Rep* 339:1–77
- Obembe AD, Hossain ME, Abu-Khamisin SA (2017) Variable-order derivative time fractional diffusion model for heterogeneous porous media. *J Pet Sci Eng* 152:391–405
- Orosco J, Coimbra CFM (2016) On the control and stability of variable-order mechanical systems. *Nonlinear Dyn* 86:695–710
- Ostalczyk P (2010) Stability analysis of a discrete-time system with a variable-, fractional-order controller. *Bull Pol Acad Sci Tech Sci* 58:613–619
- Patnaik S, Semperlotti F (2020) Application of variable- and distributed-order fractional operators to the dynamic analysis of nonlinear oscillators. *Nonlinear Dyn* 100:561–580
- Patnaik S, Hollkamp JP, Semperlotti F (2020) Applications of variable-order fractional operators: a review. *Proc R Soc A* 476:20190498

38. Patnaik S, Patnaik S, Sidhardh F (2021) Semperlotti, Towards a unified approach to nonlocal elasticity via fractional-order mechanics. *Int J Mech Sci* 189:105992
39. Patnaik S, Sidhardh S, Semperlotti F (2020) A Ritz-based finite element method for a fractional-order boundary value problem of nonlocal elasticity. *Int J Solids Struct* 202:398–417
40. Povstenko Y (2013) Fractional heat conduction in infinite one-dimensional composite medium. *J Therm Stresses* 36:351–363
41. Rahimi Z, Sumelka W, Yang X-J (2017) Linear and non-linear free vibration of nano beams based on a new fractional non-local theory. *Eng Comput* 34(5):1754–1770
42. Ramirez LE, Coimbra CF (2011) On the variable order dynamics of the nonlinear wake caused by a sedimenting particle. *Physica D* 240:1111–1118
43. Samko SG (1995) Fractional integration and differentiation of variable order. *Anal Math* 21:213–236
44. Santamaria F, Wils S, De Schutter E, Augustine GJ (2006) Anomalous diffusion in Purkinje cell dendrites caused by spines. *Neuron* 52:635–648
45. Sumelka W (2014) Thermoelasticity in the Framework of the Fractional Continuum Mechanics. *J Therm Stresses* 37(6):678–706
46. Sumelka W, Blaszczyk T (2014) Fractional continua for linear elasticity. *Arch Mech* 66(3):147–172
47. Sumelka W, Blaszczyk T, Liebold C (2015) Fractional Euler-bernoulli beams: theory, numerical study and experimental validation. *Eur J Mech A Solids* 54:243–251
48. Sumelka W (2016) On geometrical interpretation of the fractional strain concept. *J Theoret Appl Mech* 54(2):671–674
49. Sun HG, Chen W, Chen Y (2009) Variable-order fractional differential operators in anomalous diffusion modeling. *Phys A* 388:4586–4592
50. Sun HG, Chen W, Sheng H, Chen Y (2010) On mean square displacement behaviors of anomalous diffusions with variable and random orders. *Phys Lett A* 374:906–910
51. Sun HG, Chen W, Li C, Chen Y (2010) Fractional differential models for anomalous diffusion. *Phys A* 389:2719–2724
52. Sun HG, Chang A, Zhang Y, Chen W (2019) A review on variable-order fractional differential equations: mathematical foundations, physical models, numerical methods and applications. *Fract Calc Appl Anal* 22:27–59
53. Szajek K, Sumelka W, Blaszczyk T, Bekus K (2020) On selected aspects of space-fractional continuum mechanics model approximation. *Int J Mech Sci* 167:105287
54. Szymanek E, Blaszczyk T, Hall MR, Dehdezi PK, Leszczynski JS (2014) Modelling and analysis of heat transfer through 1D complex granular system. *Granular Matter* 16(5):687–694
55. Wu GC, Baleanu D, Xie HP, Zeng SD (2017) Lattice fractional diffusion equation of random order. *Math Methods Appl Sci* 40:6054–6060
56. Zingales M (2014) Fractional-order theory of heat transport in rigid bodies. *Commun Nonlinear Sci Numer Simul* 19:3938–3953

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.