

Targeting tuberculosis through a small focused library of 1,2,3-triazoles

Guillermo R. Labadie · Agustina de la Iglesia · Héctor R. Morbidoni

Received: 12 December 2010 / Accepted: 17 May 2011 / Published online: 2 June 2011
© Springer Science+Business Media B.V. 2011

Abstract Looking for new active molecules against *Mycobacterium tuberculosis*, a small focused library of 1,2,3-triazoles was efficiently prepared by click chemistry. Compounds were subsequently tested against different pathogenic and opportunistic mycobacteria including *M. avium* and *M. tuberculosis*. Two of them showed MIC at lower $\mu\text{g/mL}$ concentration for *M. avium* and even below that for *M. tuberculosis*, being more potent than control drugs.

Keywords Tuberculosis · Triazoles · Click chemistry

Introduction

Since the time of the identification of the causing agent of tuberculosis (*Mycobacterium tuberculosis*—Mtb) by Robert Koch more than 120 years ago all the attempts to eradicate this disease have been unsuccessful [1]. Instead of being near to a solution, the situation is so critical that in 1993 WHO declared tuberculosis “a global health emergency” [2]. That situation is caused by the combined effects of the AIDS pandemic, the advent of multidrug (MDR) [3] and extremely drug (XDR) resistant *M. tuberculosis* strains and the decline

of public health controls in some parts of the world (notably South East Asia, eastern Europe and the former Soviet Union) [4,5]. There are about 8.9–9 million new cases each year, of which 15% are children, and 1.7–2 million deaths, of which 450,000 are children. In this context, the number of cases is currently rising to 2% per year, and without a coordinated control effort, tuberculosis will infect an estimated 1 billion more people by 2020, killing 70 million [6]. Patients with a faulty immune response due to AIDS or other causes of cellular immunodepression are specially vulnerable to the tubercle bacilli replication leading to up to 31% death rate in the African region [7]. These individuals are also susceptible to infection by non-tuberculous mycobacteria (NTMs) from environmental sources. NTMs are usually cause of infection in AIDS patients as well as in individuals suffering of other chronic pulmonary pathologies, such as cystic fibrosis. Members of this mycobacterial group—specially *M. abscessus* and *M. chelonae*—have recently been linked to skin and soft tissue infections and *M. avium* is frequently a cause of disseminated infections in AIDS patients. For all the mentioned reasons—mainly because of the rise in the detection of MDR and XDR strains—there is an urgent need to develop new antimycobacterial agents with novel mechanism(s) of action that would be active on those strains [8].

Lipoic acid (Fig. 1) is an essential organosulphur compound that activates a number of protein complexes involved in key metabolic processes. Enzymes participating in lipoylation have gained attention because of their implication in bacterial pathogenicity [9]. LipB is part of the lipoic acid biosynthesis catalyzing the octanoyl-[acyl carrier protein]-protein acyltransferase and also plays an essential role in the growth of *M. tuberculosis* [10]. Recently, this enzyme has been crystallized unexpectedly presenting a decanoic acid covalently bound to the sulphur thiol of the Cys176 [11]. Another example of the critical role of fatty acids on *M. tuberculosis*

Electronic supplementary material The online version of this article (doi:10.1007/s11030-011-9319-0) contains supplementary material, which is available to authorized users.

G. R. Labadie (✉)
Instituto de Química Rosario (IQUIR-CONICET), Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Suipacha 531, S2002LRK Rosario, Argentina
e-mail: labadie@iquir-conicet.gov.ar

A. de la Iglesia · H. R. Morbidoni
Catedra de Microbiología, Virología y Parasitología, Facultad de Ciencias Médicas, Universidad Nacional de Rosario, Santa Fe 3100, S2002KTR Rosario, Argentina




Fig. 1 Antimycobacterial compounds and metabolites

metabolism was the activity of 2-alkynoic fatty acids (Fig. 1) that inhibited the bacterial growth. These compounds, at toxic concentrations, inhibit fatty acid degradation and synthesis, resulting in cell death. The mechanism of those antibacterial compounds included several pathways to achieve its effect [12]. Another interesting target for the development of new drugs against tuberculosis is the InhA enzyme, the enoyl reductase in the type II fatty acid biosynthesis pathway. This enzyme has been studied as a specific target to develop new chemotherapeutic agents against tuberculosis. In fact, InhA has been validated as a drug target by a temperature-sensitive mutation that causes a phenotypic response similar to isoniazid administration [13]. Triclosan (Fig. 1) is a broad spectrum antimicrobial compound that was long thought to have no specific cellular target until it was recently revealed as a potent inhibitor of the enoyl reductase of many organisms [14]. Different triclosan analogs have been prepared to selectively target *M. tuberculosis* InhA. The alkyl diphenyl ethers have been among the most successful inhibitors derived from triclosan having antitubercular MIC around 2–3 $\mu\text{g/mL}$, **8PP** (Fig. 1) being the most active of the series [15].

Azoles currently used as antifungal drugs (miconazole, econazole, fluconazole, clotrimazole, among others) have been tested against mycobacteria displaying a good killing activity creating a fresh interest onazole and triazole drugs as potential anti-mycobacterial scaffolds [16]. Econazole (Fig. 1) and clotrimazole (Fig. 1) have shown to be efficient antimycobacterial agents [17, 18], even in murine models [19] and in the latent form of the disease [20]. These results have moved different groups to prepare new collections of compounds having triazoles as a “new” antimycobacterial pharmacophore [21–23]. In general, new synthetic 1,2,4-triazoles have been reported as novel antimycobacterial products [24] but recently 1,2,3-isomers have been also introduced, prepared by conventional 1,3-dipolar cycloaddition [25–27]. The introduction of “click chemistry” has




Fig. 2 Strategy used for library design

significantly changed many fields of science, including drug discovery [28, 29]. The simplicity and efficiency of the copper(I)-catalyzed 1,3-dipolar Huisgen cycloaddition introduced by Sharpless [30], allowed to easily synthesize 1,2,3 triazoles. Over the last 15 years many click chemistry approaches have been successfully used to prepare diverse collections of triazoles directed to unspecific targets [31–33] even on solid support [34, 35], or focused libraries looking to selectively inhibit enzymes [36, 37], or mimicking active natural products [38, 39].

Based on the critical role displayed by fatty acid biosynthesis and metabolism and the known and renewed interest on triazoles as antimycobacterial drugs, we decided to combine them as an alternative approach for antitubercular drug design. Our strategy required the preparation of a focused library to find a simple scaffold to develop new drugs based on the known antitubercular compounds **8PP** and **2-alkynoic acid**. To do that, a series of 1,2,3-triazoles were prepared mimicking those two compounds (Fig. 2).

Results and discussion

A pool of azides and alkynes were selected and used in our synthesis to guarantee a minimum range of diversity over the products, allowing structural–activity relationship studies. Alkynes **A1–7** and azides were selected as the required building block for our synthesis (Fig. 3). Azide **Z2** were prepared by direct substitution of the bromide with sodium azide in DMF, azides **Z1** and **Z3** were also synthesized in the same way, but the corresponding bromides were prepared starting from the alcohol by substitution with phosphorous tribromide in Et_2O .

Alkynes **A1–7** were purchased from different commercial providers. With the necessary building blocks at hand, the triazoles were prepared following conditions originally reported [30] (Scheme 1).





Fig. 3 Alkynes (**A1–7**) and azides (**Z1–3**) used in triazole synthesis



Scheme 1 Triazoles synthesis by copper catalyzed Huisgen cycloaddition conditions

Reactions were conducted in a parallel solution synthesis fashion under copper(II) sulphate catalytic conditions in water:*t*-BuOH (1:1) using sodium ascorbate as reductant. A collection of 16 compounds **AZ** were obtained with an 84% average yield after purification (Table 1).

Products were completely characterized by 1D-, 2D-NMR experiments and HRMS. The products prepared have a 1,4-substitution pattern as was expected based on the original description of this methodology and by several other authors after that [31–39].

Minimum inhibitory concentration (MIC) of the compounds was determined against *M. tuberculosis* H37Rv and four non-tuberculous mycobacterial species. Of those, three (*M. fortuitum*, *M. avium* and *M. abscessus*) are increasingly becoming important opportunistic pathogens. The fourth species, *M. smegmatis* is a non-pathogenic rapid growing mycobacteria, that is usually used for rapid testing of compounds. The MIC value for Rifampicin used as control drug against Mtb, was 0.5 $\mu\text{g/mL}$ while MIC values for econazole and fluconazole were of 2.5 $\mu\text{g/mL}$ and $\geq 100 \mu\text{g/mL}$, respectively.

Compounds were tested for their activity against *M. tuberculosis* H37Rv, of those eight (compounds **A1Z1**, **A1Z2**, **A1Z3**, **A2Z2**, **A3Z1**, **A3Z2**, **A4Z1** and **A5Z1**) were not active at 100 $\mu\text{g/mL}$, three displayed activity at 50 $\mu\text{g/mL}$ (compounds **A2Z3**, **A6Z2** and **A7Z2**) and four (compounds **A4Z2**, **A4Z3**, **A5Z2** and **A6Z1**) showed activity at 25 $\mu\text{g/mL}$ (Table 1).

The remaining compound (**A7Z1**) has an MIC value of 0.50 $\mu\text{g/mL}$ (Table 1) fivefold more active than econazole. Compounds showing the highest activity against *M. avium* were **A4Z3** and **A7Z1** with MIC values of 1.6 $\mu\text{g/mL}$, and 3.1 $\mu\text{g/mL}$, respectively, while compounds **A4Z2**, **A7Z2**, showed activity at 25 $\mu\text{g/mL}$ on non-tuberculous mycobacteria. These results are promising, but will need a complete study over different strains because previous reports [40] have established that *M. avium* presents differences in cell envelope composition between strains and biotypes that will produce different responses to the same compound. None of the tested molecules displayed activity against the other mycobacterial species tested (Table 1), probably due to NTMs species having very low membrane permeability. None of these compounds showed cytotoxic effects on Vero cells, at 4.76 $\mu\text{g/mL}$, the higher concentration tested. That provides a minimum selectivity index (MIC antimycobacterial/MIC cytotoxicity) of 9.5 for compound **A7Z1** and 3.0 for compound **A4Z3**. It is important to remark that activity of the series of the aliphatic triazoles with the carboethoxy group (**A5Z1**, **A6Z1** and **A7Z1**) increase their activity as long as the number of carbon on the chain is growing. That can be explained based on the change of the lipophilicity of the compounds. To demonstrate that, log *D* of the library were calculated using adequate software (Table 1). As we predicted, the calculated values showed the expected trend: 1.77 for **A5Z1**, 2.83 for **A6Z1** and 4.42 for **A6Z1** showing a good correlation between MIC and log *D* for this small series.

Therefore, inhibitory activity of this series of compounds correlates with an increased value of the lipophilicity index in good agreement with previous observations for major antitubercular drugs and some recent reports of newly introduced active compounds [41,42].

In conclusion, we have found through a small focused collection of carefully selected 1,2,3-triazoles a very simple active scaffold against *M. tuberculosis*. The most promising candidate as antitubercular agent was compound **A7Z1** (Fig. 4) being five times more active than econazole.

Also, we have found that **A4Z3** (Fig. 4) was active against *M. avium* at 1.6 $\mu\text{g/mL}$. A complete study of the most active compounds with MRD strains is currently being conducted together with experiments to validate their mechanism of action.

Table 1 Antymycobacterial, cytotoxicity activities and yields for 1,2,3-triazoles **AZ**

Compound	<i>M. avium</i> MIC ^a	<i>M. tuberculosis</i> MIC ^a	Cytotoxicity ^b IC ₅₀	log <i>D</i> ^c pH 7.4	Yield ^d (%)
A1Z1	>100	>100	NC ^e	-0.94 ± 0.57	85
A1Z2	>100	>100	NC	0.21 ± 0.54	78
A1Z3	>100	>100	NC	1.05 ± 0.54	90
A2Z2^f	>100	>100	NC	2.23 ± 0.57	79
A2Z3	>100	50	NC	2.21 ± 0.77	70
A3Z1	>100	>100	NC	-3.19	78
A3Z2	>100	>100	NC	-2.03	72
A4Z1	>100	>100	NC	2.00 ± 0.58	94
A4Z2^g	25	25	NC	3.15 ± 0.56	86
A4Z3^g	1.6	25	NC	3.99 ± 0.55	93
A5Z1	>100	100	NC	1.77 ± 0.56	72
A5Z2	>100	25	NC	2.91 ± 0.53	78
A6Z1	>100	25	NC	2.83 ± 0.56	79
A6Z2	100	50	NC	3.97 ± 0.53	96
A7Z1	3.1	0.5	NC	4.42 ± 0.52	95
A7Z2	25	50	NC	5.57 ± 0.53	96
Rifampicin		0.5			
Econazole		2.5			
Fluconazole		≥100			

^a Minimum inhibitory concentration expressed in μg/mL

^b Vero cells, concentration expressed in μg/mL

^c Calculated using ACD labs 6.0


^d After purification

^e NC: not cytotoxic (up to the maximum dose tested

4.76 μg/mL)

^f Ethyl ester reported by Appukkuttan et al. [43]

^g Reported by Lee et al. [31] without a complete characterization

**Fig. 4** Most active compounds

Experimental

General methods

Chemical reagents were purchased from commercial suppliers and were used without further purification unless otherwise noted. All the solvents (hexane, ethyl acetate, CH₂Cl₂, Et₂O) were distilled prior to use. Yields were calculated for material judged homogenous by thin layer chromatography and nuclear magnetic resonance (NMR). Reaction progress was monitored by thin layer chromatography (TLC) performed on silica gel 60 F₂₅₄ pre-coated aluminium sheets. Spots were visualized under 254 nm UV lamp and/or by

dipping the TLC plate into a solution of *p*-anisaldehyde or phosphomolibdic acid followed by heating with a heat gun. Flash column chromatography was performed using silica gel 60 (230–400 mesh). ¹H and ¹³C NMR spectra were recorded on a Bruker Avance II 300 MHz (75.13 MHz) using CDCl₃ as solvent unless otherwise noted. Chemical shifts for proton nuclear magnetic resonance (¹H NMR) spectra are reported in parts per million relative to the signal of tetramethylsilane at 0 ppm (internal standard) and coupling constants (*J*) are reported in Hertz (Hz). Chemical shifts for carbon nuclear magnetic resonance (¹³C NMR) spectra are reported in parts per million relative to the centre line of the CDCl₃ triplet at 76.9 ppm. Assignments of proton resonances were confirmed by correlated spectroscopy. ESI-HRMS were recorded at the University of Mississippi, Department of Medicinal Chemistry on a Waters LC-QTof micro.

General procedure for azides and alkynes reaction under click chemistry conditions

Alkyne (1 eq) and the azide (1 eq) were suspended in mL/eq of ^tBuOH:H₂O (1:1) and then aqueous 1 M CuSO₄ solution and finally aqueous 1 M sodium ascorbate solution were added and the mixture stirred overnight at room temperature. Brine was added and the solution was extracted with

dichloromethane. The combined organic extracts were dried over sodium sulphate and evaporated. The resulting residue was purified by column chromatography over silica gel using an increasing AcOEt/hexane gradient to afford desired pure products.

Compound A1Z1 ethyl 2-(4-(hydroxymethyl)-1H-1,2,3-triazol-1-yl)acetate

¹H NMR: 7.62 (s, 1H), 5.07 (s, 2H), 4.64 (s, 2H), 4.15 (q, *J* = 6.9 Hz, 2H), 1.20 (t, *J* = 6.9 Hz, 3H); ¹³C NMR: 166.6 (C), 148.2 (C), 123.7 (C), 62.3 (CH₂), 55.8 (CH₂), 50.8 (CH₂), 14.0 (CH₃); ESI-HRMS Calcd for (M + H⁺) C₇H₁₂N₃O₃ 186.0879; found 186.0871.

Compound A1Z2 (1-benzyl-1H-1,2,3-triazol-4-yl)methanol

¹H NMR: 7.45 (s, 1H), 7.30 (m, 3H), 7.20 (m, 2H), 5.42 (s, 2H), 4.67 (s, 2H); ¹³C NMR: 148.3 (C), 134.6 (C), 129.0 (CH), 128.7 (CH), 128.1 (CH), 122.0 (CH), 54.1 (CH₂); ESI-HRMS Calcd for (M + H⁺) C₁₀H₁₂N₃O₂ 190.0980; found 190.0972.

Compound A1Z3 (1-(3-phenylpropyl)-1H-1,2,3-triazol-4-yl)methanol

¹H NMR: 7.36 (s, 1H), 7.10–7.30 (m, 5H), 4.74 (s, 2H), 4.28 (t, *J* = 7.2 Hz, 2H), 2.60 (t, *J* = 7.2 Hz, 2H), 2.18 (q, *J* = 7.2, 2H); ¹³C NMR: 148.0 (C), 128.6 (CH), 128.4 (CH), 126.3 (CH), 122.0 (C), 56.0 (CH₂), 49.5 (CH₂), 32.4 (CH₂), 31.6 (CH₂). HRMS Calcd for (M + H⁺) C₁₂H₁₆N₃O₂; 218.1293; found 218.1293.

Compound A2Z2 methyl 1-benzyl-1H-1,2,3-triazole-4-carboxylate

¹H NMR: 7.97 (s, 1H), 7.41 (m, 3H), 7.29 (m, 2H), 5.58 (s, 2H), 3.93 (s, 3H); ¹³C NMR: 161.1 (C), 140.3 (C), 133.7 (C), 129.4 (CH), 129.2 (CH), 128.3 (CH), 127.4 (CH), 54.5 (CH₂), 52.2 (CH₃); ESI-HRMS Calcd for (M + H⁺) C₁₁H₁₂N₃O₂ 218.0930; found 218.0931.

Compound A2Z3 (methyl 1-(3-phenylpropyl)-1H-1,2,3-triazole-4-carboxylate

¹H NMR: 8.10 (s, 1H), 7.30–7.10 (m, 5H), 4.39 (t, *J* = 7.2), 3.92 (s, 3H), 2.63 (t, *J* = 7.2, 2H), 2.25 (q, *J* = 7.2, 2H); ¹³C NMR: 161.2 (C), 139.9 (C), 139.8 (C), 128.7 (CH), 128.4 (CH), 127.5 (CH), 126.6 (CH), 52.2 (CH₃), 49.9 (CH₂), 32.3 (CH₂), 31.5 (CH₂); HRMS Calcd for (M + H⁺) C₁₃H₁₆N₃O₂ 246.1253; found 246.1243.

Compound A3Z1 1-(2-ethoxy-2-oxoethyl)-1H-1,2,3-triazole-4-carboxylic acid

¹H NMR (DMSO-*d*₆): 13.09 (bs, 1H), 8.75 (s, 1H), 7.35 (m, 5H), 5.63 (s, 2H); ¹³C NMR: 167.7 (C), 163.2 (C), 139.8 (C), 129.4 (CH), 59.6 (CH₂), 52.5 (CH₂), 13.9 (CH₃); ESI-HRMS Calcd for (M + H⁺) C₇H₁₀N₃O₄ 200.0671; found 200.0667.

Compound A3Z2 1-benzyl-1H-1,2,3-triazole-4-carboxylic acid

¹H NMR (DMSO-*d*₆): 13.09 (bs, 1H), 8.75 (s, 1H), 7.35 (m, 5H), 5.63 (s, 2H); ¹³C NMR (DMSO-*d*₆): 163.1 (C), 140.4 (C), 136.0 (C), 129.3 (CH), 129.3 (CH), 128.8 (CH), 128.5 (CH), 53.6; ESI-HRMS Calcd for (M + H⁺) C₁₀H₁₀N₃O₂ 204.0777; found 204.0777.

Compound A4Z1 ethyl 2-(4-phenyl-1H-1,2,3-triazol-1-yl)acetate

¹H NMR: 7.90 (s, 1H), 7.84 (dd, *J*₁ = 1.5, *J*₂ = 0.6, 2H), 7.50–7.30 (m, 3H), 5.21 (s, 2H), 4.29 (q, *J* = 7.2, 2H), 1.32 (t, *J* = 7.2, 3H); ¹³C NMR: 166.3 (C), 148.2 (C), 130.4 (C), 128.8 (CH), 128.3 (CH), 125.8 (CH), 121.0 (CH), 51.0 (CH₂), 62.5 (CH₂), 14.1 (CH₃); ESI-HRMS Calcd for (M + H⁺) C₁₂H₁₄N₃O₂ 232.1097; found 232.1086.

Compound A4Z2 (1-benzyl-4-phenyl-1H-1,2,3-triazole

¹H NMR: 7.80 (dd, *J*₁ = 8.4, *J*₂ = 1.5, 2H), 7.66 (s, 1H), 7.42–7.30 (m, 8H), 5.58 (s, 2H); ¹³C NMR: 148.2 (C), 134.7 (C), 130.6 (CH), 129.2 (CH), 128.8 (CH), 128.2 (CH), 128.1 (CH), 125.5 (CH), 119.5 (CH), 54.2 (CH₂); ESI-HRMS Calcd for (M + H⁺) C₁₅H₁₄N₃ 236.1188; found 236.1189.

Compound A4Z3, 4-phenyl-1-(3-phenylpropyl)-1H-1,2,3-triazole

¹H NMR: 7.82 (m, 2H); 7.70 (s, 1H); 7.46–7.16 (m, 8H); 4.83 (t, *J* = 6.9, 2H), 2.69 (t, *J* = 7.8, 2H), 2.29 (t, *J* = 6.9, 2H); ¹³C NMR: 147.6 (C), 140.1 (C), 130.6 (C), 128.7 (CH), 128.5 (CH), 128.4 (CH), 128.0 (CH), 126.3 (CH), 119.5 (CH), 49.5 (CH₂), 32.4 (CH₂), 31.4 (CH₂). ESI-HRMS Calcd for (M + H⁺) C₁₇H₁₈N₃ 264.1501; found 264.1501.

Compound A5Z1 ethyl 2-(4-propyl-1H-1,2,3-triazol-1-yl)acetate

¹H NMR: 7.41 (s, 1H), 5.11 (s, 2H), 4.25 (q, *J* = 7.2, 2H), 2.71 (t, *J* = 7.5, 2H), 1.71 (m, *J* = 7.5, 2H), 1.29 (t, *J* = 7.2, 3H), 0.97 (t, *J* = 7.2, 3H); ¹³C NMR: 166.5 (C), 148.4 (C), 122.2 (CH), 62.1 (CH₂), 50.7 (CH₂), 27.5 (CH₂), 22.5

(CH₂), 13.9 (CH₃), 13.6 (CH₃); ESI-HRMS Calcd for (M + H⁺) C₉H₁₆N₃O₂ 198.1243; found 198.1244.

Compound A5Z2 benzyl-4-proyl-1H-1,2,3-triazole

¹H NMR: 7.41–7.32 (m, 3H), 7.26–7.22 (m, 2H), 7.18 (s, 1H), 5.49 (s, 2H), 2.67 (t, *J* = 7.8, 2H), 1.67 (m, 2H), 0.95 (t, *J* = 6.9, 3H); ¹³C NMR: 148.6 (C), 135.1 (C), 129.0 (CH), 128.5 (CH), 127.9 (CH), 53.9 (CH₂), 27.7 (CH₂), 22.6 (CH₂), 13.7 (CH₃); ESI-HRMS Calcd for (M + Na⁺) C₁₂H₁₅N₃O₂Na 224.1170; found 224.1164.

Compound A6Z1 ethyl 2-(4-pentyl-1H-1,2,3-triazol-1-yl)acetate

¹H NMR: 7.41 (s, 1H), 5.12 (s, 2H), 4.26 (q, *J* = 7.3, 2H), 2.73 (t, *J* = 7.3, 2H), 1.65 (m, 4H), 1.20–1.40 (m, 10H), 1.30 (t, *J* = 7.2, 3H), 0.87 (t, *J* = 0.69, 3H); ¹³C NMR: 166.5 (C), 148.9 (C), 121.9 (CH), 62.2 (CH₂), 50.7 (CH₂), 31.8 (CH₂), 29.3 (CH₂), 29.2 (CH₂), 25.6 (CH₂), 22.6 (CH₂), 14.0 (CH₃); ESI-HRMS Calcd for (M + H⁺) C₁₁H₂₀N₃O₂ 226.1556; found 226.1554.

Compound A6Z2 benzyl-4-pentyl-1H-1,2,3-triazole

¹H NMR: 7.35 (m, 3H), 7.26 (m, 2H), 7.17 (s, 1H), 5.49 (s, 2H), 2.68 (t, *J* = 7.7, 2H), 1.67 (m, 2H), 1.31 (m, 4H), 0.88 (t, *J* = 7.0, 3H); ¹³C NMR: 149.0 (C), 135.1 (C), 129.0 (CH), 128.6 (CH), 127.9 (CH), 120.5 (CH), 53.9 (CH₂), 31.4 (CH₂), 29.1 (CH₂), 25.7 (CH₂), 22.4 (CH₂), 14.0 (CH₃); ESI-HRMS Calcd for (M + H⁺) C₁₄H₂₀N₃ 230.1657; found 230.1662.

Compound A7Z1, ethyl 2-(4-octyl-1H-1,2,3-triazol-1-yl)acetate

¹H NMR: 7.40 (s, 1H), 5.11 (s, 2H), 4.26 (q, *J* = 7.2, 2H), 2.73 (t, *J* = 7.5, 2H), 1.65 (m, 4H), 1.20–1.40 (m, 10H), 1.30 (t, *J* = 7.2, 3H), 0.87 (t, *J* = 0.69, 3H); ¹³C NMR: 166.5 (C), 148.9 (C), 121.9 (CH), 62.2 (CH₂), 50.7 (CH₂), 31.8 (CH₂), 29.3 (CH₂), 29.2 (CH₂), 25.6 (CH₂), 22.6 (CH₂), 14.0 (CH₃); ESI-HRMS Calcd for (M + H⁺) C₁₄H₂₆N₃O₂ 268.2025; found 268.2022.

Compound A7Z2 benzyl-4-octyl-1H-1,2,3-triazole

¹H NMR: 7.36 (m, 3H), 7.25 (m, 2H), 7.17 (s, 1H), 5.49 (s, 2H), 2.68 (t, *J* = 7.5, 2H), 1.63 (q, *J* = 7.5, 2H), 1.20–1.30 (m, 10H), 0.87 (t, *J* = 6.3, 3H); ¹³C NMR: 148.9 (C), 135.1 (C), 129.01 (CH), 128.5 (CH), 127.9 (CH), 120.5 (CH), 53.9 (CH₂), 31.8 (CH₂), 29.3 (CH₂), 29.2 (CH₂), 25.7 (CH₂), 22.6 (CH₂), 14.1 (CH₃); ESI-HRMS Calcd for (M + H⁺) C₁₇H₂₆N₃ 272.2127; found 272.2127.

Bacterial strains

Mycobacterium tuberculosis strain H37Rv as well as clinical isolates of *M. fortuitum*, *M. avium* subsp. *avium* and *M. abscessus* were kindly provided by Dr. L. Barrera (Instituto Nacional de Microbiología “C.G. Malbrán”, Argentina). *M. smegmatis* strain mc²155 was the kind gift from Dr. W. R. Jacobs (A. Einstein College of Medicine, NY, USA). All the mycobacterial strains were grown in Middlebrook 7H9 broth (Difco Laboratories, Detroit, MI, USA) supplemented with 1/10 v/v of ADS (a solution containing 50 g/L BSA fraction V, 20 g/L dextrose and 8.1 g/L NaCl), glycerol (1% w/v) herein designated 7H9-ADS-G for short. Tween 80 was added to prevent clumping (0.05% w/v for cultures of *M. tuberculosis* and *M. avium*, 0.2% w/v for cultures of *M. abscessus*, *M. fortuitum* and *M. smegmatis*). When needed, solid media (Middlebrook 7H11 supplemented with ADS (1/10 v/v) and glycerol (1% v/v) was used. All strains were grown at 37 °C under gentle agitation.

Anti-mycobacterial activity assay

Stock solutions for all the tested compounds were made in DMSO at 10 mg/mL. Working solutions were made by dilution in the above described 7H9-ADS-G medium at a final concentration of 400 μg/mL.

Antimycobacterial activity was determined by a twofold dilution of the compounds in Middlebrook 7H9-ADS-G medium. For this purpose 96-well plates (Falcon, Cat number 3072, Becton Dickinson, Lincoln Park, NJ) were used. The 96-well plates received 100 μL of Middlebrook 7H9 broth and a serial twofold dilution of the compounds was made directly on the plate. The initial and final drug concentrations tested were 100 and 0.8 μg/mL, respectively. Four compounds were tested in duplicate in each microtiter plate, Rifampicin (from 2 to 0.16 μg/mL; stock solution prepared as a 10 mg/mL solution in methanol) was used as control drug. For the sake of simplicity of the 96 wells plate design, control azole drugs were tested in parallel. The azole drugs used, Econozale (20 mg/mL in DMSO stock solution, 80 μg/mL working solution) and fluconazole (20 mg/mL in DMSO stock solution, 200 μg/mL working solution) were tested in the range of 20–0.16 μg/mL and 100–0.81 μg/mL, respectively. Two rows were used for growth control (medium and inoculum alone) and sterility control (medium alone). The inoculum was prepared as a 1/25 dilution of a fresh mid-log *M. tuberculosis* H37Rv suspension (O.D equivalent to Mc Farland 1.0 scale value) made in Middlebrook 7H9-ADS-G. A 100-μL aliquot (containing approximately 10⁶ Colony Forming Units) was used to inoculate the wells except for the row used for sterility testing. Plates were sealed with Parafilm and incubated at 37 °C for 5 days. After this time, 22 μL of a freshly prepared solution of

the tetrazolium dye 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide (MTT, Sigma, 2.5 mg/mL in 1:1 water; 20% (w/v) Tween 80) was added to the first growth control well plate in the growth control row and reincubated at 37 °C for 24 h. A change from yellow to purple colour indicated that the plate was ready to be fully developed in which case 22 µL of the MTT solution was added to the rest of the wells. If the colour change was not observed the second well in the growth control row was developed as describe above and incubated for a further 24 h. The operation was repeated if necessary until a colour change was clearly seen in the growth control well, at which time the remaining drug containing wells were developed. In our experience, colour change in the growth control row was seen in the first well.

Minimum inhibitory concentration (MIC) was defined as the lowest drug concentration preventing mycobacterial growth and colour change.

Cytotoxicity assay

The in vitro cytotoxicity was determined against mammalian kidney fibroblasts (VERO). The assay was performed in 96-well tissue culture-treated plates as described earlier [44]. Briefly, cells were seeded to the wells of the plate (25,000 cells/well) and incubated for 24 h. Samples were added and plates were again incubated for 48 h. The number of viable cells was determined by neutral red assay. IC₅₀ values were determined from logarithmic graphs of growth inhibition versus concentration. Doxorubicin was used as a positive control, while DMSO was used as vehicle control.

Acknowledgments Authors want to express their gratitude to L. Barrera (ANLIS Malbran, Buenos Aires, Argentina) for kind gift of mycobacterial strains. This work was supported in part by grants from National Research Council of Argentina, CONICET (PIP 5188/05); Agencia Nacional de Promoción Científica y Tecnológica, ANPCyT-Argentina (PICT-2004/25455 to GRL and PICT-2005/38198 to HRM). This investigation received financial support from the UNICEF/UNDP/WORLD BANK/WHO Special Programme for Research and Training in Tropical Diseases (TDR) to GRL. GRL is a member of the scientific staff of CONICET-Argentina. HRM is a career member of CIUNR-Argentina.

References

- Frieden TR, Sterling TR, Munsiff SS, Watt CJ, Dye C (2003) Tuberculosis. *Lancet* 362:887–899. doi:10.1016/S0140-6736(03)14333-4
- WHO (1993) Tuberculosis on the increase. *World Health Forum* 14:1–94
- Espinal MA (2003) The global situation of MDR-TB. *Tuberculosis* 83:44–51. doi:10.1016/S1472-9792(02)00058-6
- Afanas'ev M, Ikryannikova L, Il'ina E, Kuz'min A, Larionova E, Smirnova T, Chernousova L, Govorun V (2011) Molecular typing of *Mycobacterium tuberculosis* circulated in Moscow, Russian Federation. *Eur J Clin Microbiol Infect Dis* 30:181–191. doi:10.1007/s10096-010-1067-z
- Coker R (2001) Detention and mandatory treatment for tuberculosis patients in Russia. *Lancet* 358:349–350. doi:10.1016/S0140-6736(01)05587-8
- World Health Organization (2002) Global tuberculosis control: surveillance, planning, financing. WHO/CDS/TB/2002.295. World Health Organization, Geneva, Switzerland
- Sepkowitz K, Raffalli J, Riley L, Kiehn T, Armstrong D (1995) Tuberculosis in the AIDS era. *Clin Microbiol Rev* 8:180–199
- Janin YL (2007) Antituberculosis drugs: ten years of research. *Bioorg Med Chem* 15:2479–2513. doi:10.1016/j.bmc.2007.01.030
- Spalding MD, Prigge ST (2010) Lipic acid metabolism in microbial pathogens. *Microbiol Mol Biol Rev* 74:200–228. doi:10.1128/MMBR.00008-10
- Sasseti CM, Boyd DH, Rubin EJ (2003) Genes required for mycobacterial growth defined by high density mutagenesis. *Mol Microbiol* 48:77–84. doi:10.1046/j.1365-2958.2003.03425.x
- Ma Q, Zhao X, Nasser Eddine A, Geerlof A, Li X, Cronan JE, Kaufmann SH, Wilmanns M (2006) The *Mycobacterium tuberculosis* LipB enzyme functions as a cysteine/lysine dyad acyltransferase. *Proc Natl Acad Sci USA* 103:8662–8667. doi:10.1073/pnas.0510436103
- Morbidoni HR, Vilcheze C, Kremer L, Bittman R, Sacchettini JC, Jacobs WR Jr (2006) Dual inhibition of mycobacterial fatty acid biosynthesis and degradation by 2-alkynoic acids. *Chem Biol* 13:297–307. doi:10.1016/j.chembiol.2006.01.005
- Vilcheze C, Morbidoni HR, Weisbrod TR, Iwamoto H, Kuo M, Sacchettini JC, Jacobs WR Jr (2000) Inactivation of the inhA-encoded fatty acid synthase II (FASII) enoyl-acyl carrier protein reductase induces accumulation of the FASI end products and cell lysis of *Mycobacterium smegmatis*. *J Bacteriol* 182:4059–4067. doi:10.1128/JB.186.13.4051-4055.2004
- Lu H, Tonge PJ (2008) Inhibitors of FabI, an enzyme drug target in the bacterial fatty acid biosynthesis pathway. *Acc Chem Res* 41:11–20. doi:10.1021/ar700156e
- Sullivan TJ, Truglio JJ, Boyne ME, Novichenok P, Zhang X, Stratton CF, Li HJ, Kaur T, Amin A, Johnson F, Slayden RA, Kisker C, Tonge PJ (2006) High affinity InhA inhibitors with activity against drug-resistant strains of *Mycobacterium tuberculosis*. *ACS Chem Biol* 1:43–53. doi:10.1021/cb050004z
- Rivers EC, Mancera RL (2008) New anti-tuberculosis drugs with novel mechanisms of action. *Curr Med Chem* 15:1956–1967. doi:10.2174/092986708785132906
- McLean KJ, Marshall KR, Richmond A, Hunter IS, Fowler K, Kieser T, Gurucha SS, Besra GS, Munro AW (2002) Azole antifungals are potent inhibitors of cytochrome P450 mono-oxygenases and bacterial growth in mycobacteria and streptomycetes. *Microbiology* 148:2937–2949
- Ahmad Z, Sharma S, Khuller GK (2005) In vitro and ex vivo anti-mycobacterial potential of azole drugs against *Mycobacterium tuberculosis* H37Rv. *FEMS Microbiol Lett* 251:19–22. doi:10.1016/j.femsle.2005.07.022
- Ahmad Z, Sharma S, Khuller GK (2006) The potential of azole antifungals against latent/persistent tuberculosis. *FEMS Microbiol Lett* 258:200–203. doi:10.1111/j.1574-6968.2006.00224.x
- Ahmad Z, Sharma S, Khuller GK (2006) Azole antifungals as novel chemotherapeutic agents against murine tuberculosis. *FEMS Microbiol Lett* 261:181–186. doi:10.1111/j.1574-6968.2006.00350.x
- Jadhav GR, Shaikh MU, Kale RP, Shiradkar MR, Gill CH (2009) SAR study of clubbed [1,2,4]-triazolyl with fluorobenzimidazoles as antimicrobial and antituberculosis agents. *Eur J Med Chem* 44:2930–2935. doi:10.1016/j.ejmech.2008.12.001

22. Ozdemir A, Turan-Zitouni G, Kaplancikli ZA, Chevallet P (2007) Synthesis of some 4-arylideneamino-4H-1,2,4-triazole-3-thiols and their antituberculosis activity. *J Enzyme Inhib Med Chem* 22:511–516. doi:[10.1080/14756360601178424](https://doi.org/10.1080/14756360601178424)
23. Kaplancikli ZA, Turan-Zitouni G, Chevallet P (2005) Synthesis and antituberculosis activity of new 3-alkylsulfanyl-1,2,4-triazole derivatives. *J Enzyme Inhib Med Chem* 20:179–182. doi:[10.1080/14756360500043471](https://doi.org/10.1080/14756360500043471)
24. Shiradkar MR, Murahari KK, Gangadasu HR, Suresh T, Kalyan CA, Panchal D, Kaur R, Burange P, Ghogare J, Mokale V, Raut M (2007) Synthesis of new S-derivatives of clubbed triazolyl thiazole as anti-*Mycobacterium tuberculosis* agents. *Bioorg Med Chem* 15:3997–4008. doi:[10.1016/j.bmc.2007.04.003](https://doi.org/10.1016/j.bmc.2007.04.003)
25. Dabak K, Sezer O, Akar A, Anac O (2003) Synthesis and investigation of tuberculosis inhibition activities of some 1,2,3-triazole derivatives. *Eur J Med Chem* 38:215–218. doi:[10.1016/S0223-5234\(02\)01445-9](https://doi.org/10.1016/S0223-5234(02)01445-9)
26. Costa MS, Boechat N, Rangel EA, de C. da Silva F, de Souza AM, Rodrigues CR, Castro HC, Junior IN, Lourenco MC, Wardell SM, Ferreira VF (2006) Synthesis, tuberculosis inhibitory activity, and SAR study of N-substituted-phenyl-1,2,3-triazole derivatives. *Bioorg Med Chem* 14:8644–8653. doi:[10.1016/j.bmc.2006.08.019](https://doi.org/10.1016/j.bmc.2006.08.019)
27. Gill C, Jadhav G, Shaikh M, Kale R, Ghawalkar A, Nagargoje D, Shiradkar M (2008) Clubbed [1,2,3] triazoles by fluorine benzimidazole: a novel approach to H37Rv inhibitors as a potential treatment for tuberculosis. *Bioorg Med Chem Lett* 18:6244–6247. doi:[10.1016/j.bmcl.2008.09.096](https://doi.org/10.1016/j.bmcl.2008.09.096)
28. Kolb HC, Sharpless KB (2003) The growing impact of click chemistry on drug discovery. *Drug Discov Today* 8:1128–1137. doi:[10.1016/S1359-6446\(03\)02933-7](https://doi.org/10.1016/S1359-6446(03)02933-7)
29. Tron GC, Pirali T, Billington RA, Canonico PL, Sorba G, Genazzani AA (2008) Click chemistry reactions in medicinal chemistry: applications of the 1,3-dipolar cycloaddition between azides and alkynes. *Med Res Rev* 28:278–308. doi:[10.1002/med.20107](https://doi.org/10.1002/med.20107)
30. Rostovtsev VV, Green LG, Fokin VV, Sharpless KB (2002) A stepwise Huisgen cycloaddition process: copper(I)-catalyzed regioselective “ligation” of azides and terminal alkynes. *Angew Chem Int Ed Engl* 41:2596–2599. doi:[10.1002/1521-3773\(20020715\)41:14](https://doi.org/10.1002/1521-3773(20020715)41:14)
31. Lee B-Y, Park SR, Jeon HB, Kim KS (2006) A new solvent system for efficient synthesis of 1,2,3-triazoles. *Tetrahedron Lett* 47:5105–5109. doi:[10.1016/j.tetlet.2006.05.079](https://doi.org/10.1016/j.tetlet.2006.05.079)
32. Angelo NG, Arora PS (2007) Solution- and solid-phase synthesis of triazole oligomers that display protein-like functionality. *J Org Chem* 72:7963–7967. doi:[10.1021/jo701292h](https://doi.org/10.1021/jo701292h)
33. VijayaRaghava Reddy L, Venkat Reddy P, Mishra NN, Shukla PK, Yadav G, Srivastava R, Shaw AK (2010) Synthesis and biological evaluation of glycal-derived novel tetrahydrofuran 1,2,3-triazoles by ‘click’ chemistry. *Carbohydr Res* 345:1515–1521. doi:[10.1016/j.carres.2010.03.031](https://doi.org/10.1016/j.carres.2010.03.031)
34. Jlalía I, Beauvineau C, Beauvière S, Önen E, Aufort M, Beauvineau A, Khaba E, Herscovici J, Meganem F, Girard C (2010) Automated synthesis of a 96 product-sized library of triazole derivatives using a solid phase supported copper catalyst. *Molecules* 15:3087–3120. doi:[10.3390/molecules15053087](https://doi.org/10.3390/molecules15053087)
35. Turner RA, Oliver AG, Lokey RS (2007) Click chemistry as a macrocyclization tool in the solid-phase synthesis of small cyclic peptides. *Org Lett* 9:5011–5014. doi:[10.1021/ol702228u](https://doi.org/10.1021/ol702228u)
36. Colombano G, Travelli C, Galli U, Caldarelli A, Chini MG, Canonico PL, Sorba G, Bifulco G, Tron GC, Genazzani AA (2009) A novel potent nicotinamide phosphoribosyltransferase inhibitor synthesized via click chemistry. *J Med Chem* 53:616–623. doi:[10.1021/jm9010669](https://doi.org/10.1021/jm9010669)
37. Jia Z, Zhu Q (2010) ‘Click’ assembly of selective inhibitors for MAO-A. *Bioorg Med Chem Lett* 20:6222–6225. doi:[10.1016/j.bmcl.2010.08.104](https://doi.org/10.1016/j.bmcl.2010.08.104)
38. Choi KW, Brimble MA (2008) Synthesis of spiroacetal-triazoles as privileged natural product-like scaffolds using “click chemistry”. *Org Biomol Chem* 6:3518–3526. doi:[10.1039/b808454h](https://doi.org/10.1039/b808454h)
39. Bagwell CL, Moloney MG, Yaqoob M (2010) Oxazolomycins: natural product lead structures for novel antibacterials by click fragment conjugation. *Bioorg Med Chem Lett* 20:2090–2094. doi:[10.1016/j.bmcl.2010.02.066](https://doi.org/10.1016/j.bmcl.2010.02.066)
40. Philalay JS, Palermo CO, Hauge KA, Rustad TR, Cangelosi GA (2004) Genes required for intrinsic multidrug resistance in *Mycobacterium avium*. *Antimicrob Agents Chemother* 48:3412–3418. doi:[10.1128/AAC.48.9.3412-3418.2004](https://doi.org/10.1128/AAC.48.9.3412-3418.2004)
41. Barry CEIII, Slayden RA, Sampson AE, Lee RE (2000) Use of genomics and combinatorial chemistry in the development of new antimicrobial drugs. *Biochem Pharmacol* 59:221–231. doi:[10.1016/S0006-2952\(99\)00253-1](https://doi.org/10.1016/S0006-2952(99)00253-1)
42. Prado S, Ledoit H, Michel S, Koch M, Darbord JC, Cole ST, Tillequin F, Brodin P (2006) Benzofuro[3,2-f][1]benzopyrans: a new class of antitubercular agents. *Bioorg Med Chem* 14:5423–5428. doi:[10.1016/j.bmc.2006.03.033](https://doi.org/10.1016/j.bmc.2006.03.033)
43. Appukkuttan P, Dehaen W, Fokin VV, Van der Eycken E (2004) A microwave-assisted click chemistry synthesis of 1,4-disubstituted 1,2,3-triazoles via a copper(I)-catalyzed three-component reaction. *Org Lett* 6:4223–4225. doi:[10.1021/ol048341v](https://doi.org/10.1021/ol048341v)
44. Babich H, Borenfreund E (1991) Cytotoxicity of T-2 toxin and its metabolites determined with the neutral red cell viability assay. *Appl Environ Microbiol* 57:2101–2103