



# Time-time Covariance for Last Passage Percolation with Generic Initial Profile

P. L. Ferrari<sup>1</sup> · A. Occelli<sup>1</sup>

Received: 6 August 2018 / Accepted: 26 December 2018 / Published online: 5 January 2019  
© Springer Nature B.V. 2019

## Abstract

We consider time correlation for KPZ growth in 1+1 dimensions in a neighborhood of a characteristics. We prove convergence of the covariance with droplet, flat and stationary initial profile. In particular, this provides a rigorous proof of the exact formula of the covariance for the stationary case obtained in Ferrari and Spohn (2011). Furthermore, we prove the universality of the first order correction when the two observation times are close and provide a rigorous bound of the error term. This result holds also for random initial profiles which are not necessarily stationary.

**Keywords** Last passage percolation · KPZ universality class · Time-time correlations

**Mathematics Subject Classification (2010)** 60K35 · 82C22 · 82B43

## 1 Introduction

Stochastic growth models in the Kardar-Parisi-Zhang (KPZ) universality class [34] on a one-dimensional substrate are described by a height function  $h(x, t)$  with  $x$  denoting space and  $t$  time. The height function evolves microscopically according to a random and local dynamics, while on a macroscopic scale the evolution is a deterministic PDE and the limit shape is non-random. In particular, if the speed of growth as a function of the gradient of the interface is a strictly convex or concave function, then the model is in the KPZ universality class. One expects large time universality under an appropriate scaling limit.

---

✉ P. L. Ferrari  
ferrari@uni-bonn.de

A. Occelli  
occelli@iam.uni-bonn.de

<sup>1</sup> Institute for Applied Mathematics, Bonn University, Endenicher Allee 60, 53115 Bonn, Germany

By studying special models in the KPZ class, the law of the one-point fluctuations and of the spatial statistics are well-known. In particular, the fluctuations scales as  $t^{1/3}$  and the correlation length as  $t^{2/3}$  (see surveys and lecture notes [13, 17, 23, 28, 44, 46, 51]).<sup>1</sup> Furthermore, it is known that non-trivial correlations survive on the macroscopic time scale if one considers space-time points along characteristic lines of the PDE for the macroscopic evolution [19, 22]. This phenomenon is called slow-decorrelation and it indicates that non-trivial processes in a spatial  $t^{2/3}$ -neighborhood of a characteristic and for macroscopic temporal scale is to be expected. The limit process depends on the initial condition, since this is already the case for the processes at a fixed time.

The study of the time-time process started much more recently. On the experimental and numerical simulation side observables like the persistence probability or the covariance of an appropriately rescaled height function have been studied [49, 50, 53, 54]. On the analytic and rigorous side, the two-time joint distribution of the height function is known for special initial conditions: Johansson analyzed a model on full space [32, 33], while Baik and Liu considered a model on a torus [4, 5]. There are also non-rigorous works on the time-time covariance and on the upper tail of distributions using replica approach [36–38]. For general (random) initial conditions exact formulas on the joint distributions are not yet available. Also, the analysis of the covariance starting from the available formulas [4, 33] seems to be a difficult task.

In [29] Ferrari and Spohn made some predictions for the behavior of the two-time covariance for three typical initial conditions based on a last passage percolation (LPP) picture. In particular, for the stationary case, an exact formula for the covariance of two points along a characteristic has been derived. Furthermore, the behavior when the macroscopic times were either close or far from each others were provided. However, the work is not mathematically rigorous since the exchange of the large time limit and maximum over sums of Airy processes as well as justification for convergence of the covariances are not provided. The work by Corwin, Liu and Wang [20] showed the way to obtain a rigorous convergence of distribution in terms of the variational process used in [29], by lifting the finite-dimensional slow-decorrelation result of [19, 22] to a functional slow-decorrelation statement.

In this paper we consider a last passage percolation model, which can be also seen as a (version of the) polynuclear growth model. As initial condition we consider the three standard cases (called droplet, flat and stationary) as in [29], but we extend the study to random but not stationary initial profiles (see [16] for a related model). In the first three cases by the method of [20] (simplified in some aspects in [16, 27]) one knows that the limiting distributions of (rescaled) LPP times can be expressed as a variational problem in terms of some Airy processes. The first result proven in this paper is the convergence of the covariance of the LPP time to the covariance of the limiting processes, see Theorem 2.2. As a corollary, this provides a proof for the exact formula of the covariance for the stationary case of [29]. We actually extend the

---

<sup>1</sup>This holds true around point with smooth limit shape. Around shocks there are some differences, see e.g. [21, 25, 26].

result by taking points not exactly on the characteristics, but in a  $t^{2/3}$ -neighborhood of it.

Our second result concerns the behavior of the covariance when the two times are close to each other on a macroscopic scale. Physically we expect to see the signature of the stationary state as first approximation. This was noticed also in numerical experiments [50]. This is proven in Theorem 2.5 for all the initial conditions considered. We also provide a rigorous error term, which is compatible with the experiments.<sup>2</sup> To obtain the result, we need to control the spatial process at fixed time on small scales. This is achieved by comparing with stationary cases on sets of high probability. The idea goes back to Cator and Pimentel [15] for the droplet case (extended to general case in [41]). The control on the high probability sets requires bounds on exit point probabilities, which has to be obtained for each initial profile. In particular, to achieve a good control in the error term, one can not use soft bounds as in [7, 41]. Finally, for droplet initial condition we derive a result also when times are far apart, see Theorem 2.6.

A few weeks after we finished our paper, for the droplet geometry, Basu and Ganguly obtained the same exponents for the behaviour at close or far away points [8]. Unlike in our paper, they did not identify the prefactor, but on the other hand, their result are non-asymptotic as well. One input often used in their paper are the bounds on transversal fluctuations of [9].

**Outline** In Section 2 we introduce the model, state some known limiting results necessary for the rest of the paper and provide the main results. In Section 3 we recall the stationary LPP and the comparison lemmas. In Section 4 we prove Theorem 2.2 on the convergence of the covariance. In Section 5 we prove Theorem 2.5 on the close time behaviour, while in Section 6 we sketch the proof of Theorem 2.6. The appendix contains several bounds on the one-point distribution or on increments, which are used in the proofs.

## 2 Model and Results

### 2.1 LPP and Polynuclear Growth

Consider a collection of i.i.d. random variables  $\omega_{i,j}$ ,  $i, j \in \mathbb{Z}$  with exponential distribution of parameter one. An *up-right path*  $\pi = (\pi(0), \pi(1), \dots, \pi(n))$  on  $\mathbb{Z}^2$  from a point  $A$  to a point  $E$  is a sequence of points in  $\mathbb{Z}^2$  with  $\pi(k+1) - \pi(k) \in \{(0, 1), (1, 0)\}$ , with  $\pi(0) = A$  and  $\pi(n) = E$ , and  $n$  is called the length  $\ell(\pi)$  of  $\pi$ . Given a set of points  $S_A$  with some random variables (not

<sup>2</sup>The next order correction is sensitive to the scaling used to define the process. For the scaling used in this paper the error term seems to be optimal. However, if one scales the random variables to have the same one-point distribution function, then experimentally the error term is smaller: instead of an error term with exponent  $1^-$ , one gets an exponent  $\min\{5/3, 2/3 + \alpha\}$ , where  $\alpha$  is the exponent controlling the convergence of the variance of the height difference to that of the Baik-Rains distribution [52].

necessarily independent)  $h^0$  on  $S_A$ , but independent of the  $\omega$ 's, and given a point  $E$ , one defines the last passage time  $L_{S_A \rightarrow E}$  as

$$L_{S_A \rightarrow E} = \max_{\substack{\pi: A \rightarrow E \\ A \in S_A}} \left( h^0(\pi(0)) + \sum_{1 \leq k \leq n} \omega_{\pi(k)} \right). \tag{2.1}$$

Also, for two points  $P, Q$  which are not on the initial set  $S_A$ , we define  $L_{P \rightarrow Q}$  as above but without the term  $h^0(\pi(0))$ .  $\pi_{S_A \rightarrow E}^{max}$  indicates the maximizer of the last passage time. For continuous random variables, the maximizer is a.s. unique.<sup>3</sup>

LPP can be thought as a stochastic growth model, a version of the polynuclear growth model, as follows. Let  $S_A = \mathcal{L} := \{(i, j) \in \mathbb{Z}^2 \mid i + j = 0\}$  and let  $h^0$  represents a height function at time  $t = 0$ . Then one defines the height function at time  $t$  by the relation

$$h(x, t) = L_{\mathcal{L}} \rightarrow ((x + t)/2, (t - x)/2) \tag{2.2}$$

for all  $x - t$  being even numbers (and set  $h(x, t) = L_{\mathcal{L}} \rightarrow ((x + t - 1)/2, (t - x - 1)/2)$  for  $x - t$  odd). The dynamics of the height function is

$$h(x, t) = \max\{h(x - 1, t - 1), h(x, t - 1), h(x + 1, t - 1)\} + \omega_{(x+t)/2, (t-x)/2} \tag{2.3}$$

with initial conditions  $h(x, 0) = h^0(x/2, -x/2)$  (here  $\omega_{(x+t)/2, (t-x)/2} = 0$  if  $x - t$  is odd).

We are interested in the scaling limit of the height function

$$(w, \tau) \mapsto \lim_{t \rightarrow \infty} \frac{h(w2^{1/3}t^{2/3}, \tau t) - \tau t}{2^{2/3}t^{1/3}} \tag{2.4}$$

or, equivalently, setting  $E = (\tau N, \tau N) + w(2N)^{2/3}(1, -1)$ ,

$$(w, \tau) \mapsto \lim_{N \rightarrow \infty} \frac{L_{S_A \rightarrow E} - 4\tau N}{2^{4/3}N^{1/3}}, \tag{2.5}$$

for different initial conditions<sup>4</sup>

1. *Droplet case.* In this case one sets  $h^0 = 0$  and further set  $\omega(i, j) = 0$  whenever  $(i, j) \notin \mathbb{Z}_+^2$ . In terms of LPP this is equivalent to take  $S_A = (0, 0)$  and  $h^0 = 0$ .
2. *Flat with zero-slope.* This means that we take  $h^0 = 0$ .
3. *Stationary with zero-slope.* Let  $\{X_k, Y_k\}_{k \in \mathbb{Z}}$  be i.i.d. random variable  $\text{Exp}(1/2)$ -distributed. Then define

$$h^0(x, -x) = \begin{cases} \sum_{k=1}^x (X_k - Y_k), & \text{for } x \geq 1, \\ 0, & \text{for } x = 0, \\ -\sum_{k=x+1}^0 (X_k - Y_k), & \text{for } x \leq -1. \end{cases} \tag{2.6}$$

<sup>3</sup>The only exception will be if  $h^0$  is not random, since then the maximizer is unique up to the initial point, which has weight 0 and thus it is irrelevant.

<sup>4</sup>The choice of zero-slope is just for convenience as it avoids to introduce a further parameter in the scaling. However, the inputs used in the proofs are available for non-zero slopes as well.

4. *A family of random initial conditions.* We consider the case where for a given  $\sigma \geq 0$ ,  $h^0$  is given by (2.6) multiplied by  $\sigma$ . Clearly, the cases  $\sigma = 0$  and  $\sigma = 1$  correspond to the flat and to the stationary cases.

*Remark 2.1* In the setting of TASEP, a random initial condition maps to a LPP starting from a random line. Due to functional slow-decorrelation, the weight  $h^0$  should be taken to reflect the first order LPP from a point on the line to its projection onto the antidiagonal. Thus a-priori one could try to start with the random line used in [16, 27], but since in the scaling limit the result is identical to the one of our choice, we did not attempt to use this precise mapping.

### 2.1.1 Limiting Variational Formulas

For  $0 < \tau \leq 1$ , we set<sup>5</sup>  $E_\tau = (\tau N, \tau N) + (2N)^{2/3}w_\tau(1, -1)$  and define the LPP and its limit as

$$L_N^\star(\tau) = \frac{L_{S_A \rightarrow E_\tau}^\star - 4\tau N}{2^{4/3}N^{1/3}}, \quad \chi^\star(\tau) := \lim_{N \rightarrow \infty} L_N^\star(\tau), \tag{2.7}$$

where the superscript  $\star$  denotes the different configurations, point-to-point ( $\bullet$ ), point-to-line ( $\setminus$ ), stationary ( $\mathcal{B}$ ) and random ( $\sigma$ ).

The convergence in distribution of the random variables  $L_N^\star(\tau)$  are well-known. Recall that for LPP we have the identity

$$L_{S_A \rightarrow E_1}^\star = \max_{u \in \mathbb{R}} \{L_{S_A \rightarrow I(u)}^\star + L_{I(u) \rightarrow E_1}\} \tag{2.8}$$

with

$$I(u) = (\tau N, \tau N) + u(2N)^{2/3}(1, -1). \tag{2.9}$$

Provided that the limit  $N \rightarrow \infty$  and  $\max_{u \in \mathbb{R}}$  can be exchanged (which is the case in all the cases considered here, see [16, 20, 27] for related works), the limiting processes can be written in terms of Airy processes as follows.

1. *Droplet case.* Let  $\mathcal{A}_2$  and  $\tilde{\mathcal{A}}_2$  be two independent  $\text{Airy}_2$  processes. Then

$$\begin{aligned} \chi^\bullet(\tau) &= \tau^{1/3} \left[ \tilde{\mathcal{A}}_2\left(\frac{w_\tau}{\tau^{2/3}}\right) - \frac{w_\tau^2}{\tau^{4/3}} \right], \\ \chi^\bullet(1) &= \max_{u \in \mathbb{R}} \left\{ \tau^{1/3} \left[ \tilde{\mathcal{A}}_2\left(\frac{u}{\tau^{2/3}}\right) - \frac{u^2}{\tau^{4/3}} \right] + (1-\tau)^{1/3} \left[ \mathcal{A}_2\left(\frac{u-w_1}{(1-\tau)^{2/3}}\right) - \frac{(u-w_1)^2}{(1-\tau)^{4/3}} \right] \right\}, \end{aligned} \tag{2.10}$$

The  $\text{Airy}_2$  process has been discovered in a related polynuclear growth model setting [43] (see [31] for the case of geometric random variables, or [14] for a two-parameter generalization). Tightness in this setting was shown in [27], building on the approach of [15] (while for the geometric case tightness was shown already in [31]).

<sup>5</sup>Throughout the paper we do not write explicitly integer parts.

2. *Flat case.* Let  $\mathcal{A}_1$  be an  $\text{Airy}_1$  process and  $\mathcal{A}_2$  an  $\text{Airy}_2$  process, independent of each other. Then

$$\begin{aligned} \chi^\setminus(\tau) &= (2\tau)^{1/3} \mathcal{A}_1\left(\frac{w_\tau}{(2\tau)^{2/3}}\right), \\ \chi^\setminus(1) &= \max_{u \in \mathbb{R}} \left\{ (2\tau)^{1/3} \mathcal{A}_1\left(\frac{u}{(2\tau)^{2/3}}\right) + (1-\tau)^{1/3} \left[ \mathcal{A}_2\left(\frac{u-w_1}{(1-\tau)^{2/3}}\right) - \frac{(u-w_1)^2}{(1-\tau)^{4/3}} \right] \right\}. \end{aligned} \tag{2.11}$$

The  $\text{Airy}_1$  process has been discovered in the framework of the totally asymmetric simple exclusion process [11, 48], equivalent through slow-decorrelation to the LPP [18, 19, 22].

3. *Stationary case.* Let  $\mathcal{A}_2$  be an  $\text{Airy}_2$  process and  $\mathcal{A}_{\text{stat}}$  an  $\text{Airy}_{\text{stat}}$  process, independent of each other. Then

$$\begin{aligned} \chi^{\mathcal{B}}(\tau) &= \tau^{1/3} \mathcal{A}_{\text{stat}}\left(\frac{w_\tau}{\tau^{2/3}}\right), \\ \chi^{\mathcal{B}}(1) &= \max_{u \in \mathbb{R}} \left\{ \tau^{1/3} \mathcal{A}_{\text{stat}}\left(\frac{u}{\tau^{2/3}}\right) + (1-\tau)^{1/3} \left[ \mathcal{A}_2\left(\frac{u-w_1}{(1-\tau)^{2/3}}\right) - \frac{(u-w_1)^2}{(1-\tau)^{4/3}} \right] \right\}. \end{aligned} \tag{2.12}$$

The limit process  $\text{Airy}_{\text{stat}}$  (which, in spite of the name, is not stationary) was obtained in [2].

4. *Random initial conditions.* For this case, the one-point distribution is given by the following expression<sup>6</sup>

$$\mathbb{P}(\chi^\sigma(1) \leq s) = \mathbb{P}\left(\max_{u \in \mathbb{R}} \{ \mathcal{A}_2(u) - u^2 + \sqrt{2}\sigma B(u) \} \leq s\right), \tag{2.13}$$

where the  $\text{Airy}_2$  process and the two-sided standard Brownian motion  $B$  are independent of each other. Furthermore, we could write formulas similar to the one of the first three cases in terms of an Airy sheet [35]. However uniqueness in law of Airy sheet is so-far not proven [35, 40]. Therefore we state the convergence of the covariance to the covariance of its limit process only for the other cases. However, the proof could be adapted to the general  $\sigma$  as well, once uniqueness of the limit is established.

## 2.2 Main Results

### 2.2.1 Convergence of the Covariance

As our first result we give a rigorous proof of the convergence of the covariances.

---

<sup>6</sup>This was actually proven for the LPP model where instead of the random function on the antidiagonal one has a random line in [16], see also [27] for general slope. These works were based on the approach in the geometric random variables case of [20]. Adapting the proof of [27] to this setting to get the variational formula is straightforward (it is actually even slightly simpler).

**Theorem 2.2** *We have*

$$\lim_{N \rightarrow \infty} \text{Cov} (L_N^*(\tau), L_N^*(1)) = \text{Cov} (\chi^*(\tau), \chi^*(1)), \tag{2.14}$$

for  $\star \in \{\bullet, \setminus, \mathcal{B}\}$ .

*Remark 2.3* The motivation of this paper is the study of the covariance. However, by inspecting the proof, one sees that one can generalize the proof to get convergence of any joint moments of  $L_N^*(\tau)$  and  $L_N^*(1)$  without the need to new ideas and bounds.

For the stationary process  $\mathcal{A}_{\text{stat}}(w) \stackrel{(d)}{=} \max_{v \in \mathbb{R}} \{\sqrt{2}B(v) + \mathcal{A}_2(v) - (v - w)^2\}$  where the Airy<sub>2</sub> process,  $\mathcal{A}_2$ , and the two-sided standard Brownian motion,  $B(v)$ , are independent [45]. We denote

$$F_w(s) = \mathbb{P} \left( \max_{v \in \mathbb{R}} \{\sqrt{2}B(v) + \mathcal{A}_2(v) - (v - w)^2\} \leq s \right) \tag{2.15}$$

and use the notation  $\xi_{\text{stat},w}$  for a random variable distributed according to  $F_w$ . Due to stationarity one has the property [6, 42]  $\mathbb{E}(\mathcal{A}_{\text{stat}}(w)) = 0$ , which implies

$$\text{Var}(\xi_{\text{stat},w}) = \mathbb{E} \left( \max_{v \in \mathbb{R}} \{\sqrt{2}B(v) + \mathcal{A}_2(v) - (v - w)^2\} \leq s \right)^2. \tag{2.16}$$

For the stationary case, an exact expression for the covariance has been obtained in [29] for  $\tau$  in the entire interval  $[0,1]$ , in the special case  $w_\tau = w_1 = 0$ . For general values of  $w_\tau$  and  $w_1$ , we obtain

**Corollary 2.4** *For the stationary LPP, the covariance of the limiting height function for all  $\tau \in (0, 1)$  can be expressed as*

$$\begin{aligned} \text{Cov} (\chi^{\mathcal{B}}(\tau), \chi^{\mathcal{B}}(1)) &= \frac{\tau^{2/3}}{2} \text{Var} (\xi_{\text{stat},\tau^{-2/3}w_\tau}) + \frac{1}{2} \text{Var} (\xi_{\text{stat},w_1}) \\ &\quad - \frac{(1 - \tau)^{2/3}}{2} \text{Var} (\xi_{\text{stat},(1-\tau)^{-2/3}(w_1-w_\tau)}). \end{aligned} \tag{2.17}$$

**2.2.2 Universal Behavior for  $\tau \rightarrow 1$**

In [29] there is a conjecture on the behaviour of the covariance of the limit process for  $\tau \rightarrow 1$  for the other initial profiles as well. Our second goal is to provide a proof of such statements together with a rigorous error bound. We also extend the result to all initial conditions 1-4. Recall that for any random variables  $X_1, X_2$  it holds

$$\text{Cov} (X_1, X_2) = \frac{1}{2} \text{Var} (X_1) + \frac{1}{2} \text{Var} (X_2) - \frac{1}{2} \text{Var}(X_2 - X_1). \tag{2.18}$$

**Theorem 2.5** *Let us scale  $w_1 = \tilde{w}_1(1 - \tau)^{2/3}$  and  $w_\tau = \tilde{w}_\tau(1 - \tau)^{2/3}$ . Then as  $\tau \rightarrow 1$  we have<sup>7</sup>*

$$\text{Var}(\chi^\star(\tau) - \chi^\star(1)) = (1 - \tau)^{2/3} \text{Var}(\xi_{\text{Stat}, \tilde{w}_1 - \tilde{w}_\tau}) + \mathcal{O}(1 - \tau)^{1-\delta}, \tag{2.19}$$

for any  $\delta > 0$ . In particular, by (2.18), for  $\star = \{\bullet, \setminus, \mathcal{B}\}$ , we can rewrite

$$\begin{aligned} \text{Cov}(\chi^\star(\tau), \chi^\star(1)) &= \frac{1}{2} \text{Var}(\xi^\star(w_1)) + \frac{\tau^{2/3}}{2} \text{Var}(\xi^\star(w_\tau \tau^{-2/3})) \\ &\quad - \frac{(1 - \tau)^{2/3}}{2} \text{Var}(\xi_{\text{Stat}, \tilde{w}_1 - \tilde{w}_\tau}) + \mathcal{O}(1 - \tau)^{1-\delta}. \end{aligned} \tag{2.20}$$

Here  $\xi^\star(w) + w^2$  (resp.  $2^{2/3}\xi^\setminus(w)$ ) is distributed according to a GUE (resp. GOE) Tracy-Widom law and  $\xi^\mathcal{B}(w) = \xi_{\text{Stat}, w}$ .

### 2.2.3 Small $\tau$ Behavior for Droplet Initial Conditions

**Theorem 2.6** *For point-to-point LPP, let  $w_\tau = \hat{w}_\tau \tau^{2/3}$ . Then the covariance of the limiting height function for  $\tau \rightarrow 0$  can be expressed as*

$$\text{Cov}(\chi^\bullet(\tau), \chi^\bullet(1)) = \tau^{2/3} \mathbb{E}(\mathcal{A}_2(\hat{w}_\tau) \max_{u \in \mathbb{R}} \{\mathcal{A}_2(u) - u^2 + \sqrt{2}B(u)\}) + \mathcal{O}(\tau^{1-\delta}). \tag{2.21}$$

## 3 The Stationary LPP and its Comparison Lemmas

As shown in [7] the stationary situation can be realized in different ways. For the purpose of this paper, we will consider the following situations

- On  $\mathbb{Z}_+^2$ : consider the LPP from  $S_A = \{(0, 0)\}$  with

$$\omega_{i,j} = \begin{cases} 0 & \text{for } i = 0, j = 0, \\ \text{Exp}(1 - \rho) & \text{for } i \geq 1, j = 0, \\ \text{Exp}(\rho) & \text{for } i = 0, j \geq 1, \\ \text{Exp}(1) & \text{for } i \geq 1, j \geq 1. \end{cases} \tag{3.1}$$

This is called stationary LPP with density  $\rho$  since the increments of the LPP along horizontal lines are still sums of iid.  $\text{Exp}(1 - \rho)$  random variables, as a special case of Lemma 4.2 of [7]. More generically, the increments along a down-right path are sums of independent random variables,  $\text{Exp}(1 - \rho)$  for horizontal steps, and  $-\text{Exp}(\rho)$  for vertical steps.

- Consider  $S_A = \mathcal{L} = \{(i, j) \in \mathbb{Z}^2 \mid i + j = 0\}$  and with boundary terms

$$h^0(x, -x) = \begin{cases} \sum_{k=1}^x (X_k - Y_k), & \text{for } x \geq 1, \\ 0, & \text{for } x = 0, \\ -\sum_{k=x+1}^0 (X_k - Y_k), & \text{for } x \leq -1, \end{cases} \tag{3.2}$$

<sup>7</sup>One could also reformulate the result by saying that the error term is  $\mathcal{O}((1 - \tau)/\ln(1 - \tau))$ .



where  $\{X_k\}_{k \in \mathbb{Z}}$  and  $\{Y_k\}_{k \in \mathbb{Z}}$  are independent random variables with  $X_k \sim \text{Exp}(1 - \rho)$  and  $Y_k \sim \text{Exp}(\rho)$ . Then by Lemma 4.2 of [7] the increments of the LPP in this model are as in the first case.

We will call a stationary LPP model either of this two settings, depending on the cases. When we consider the point-to-point problem, we will refer to the stationary case as the first setting, while, when considering the other initial conditions, the stationary LPP will be the second setting.

To prove Theorem 2.5 we are going to use a comparison with the stationary model of density slightly higher or lower than  $1/2$ . The comparison idea was first used in [15] and then generalized in [41], with applications in [24, 27, 39, 40]. For that purpose, we need to introduce the notion of exit point, which is the location where the maximizer of the LPP exits its boundary terms. Let us define it for both stationary settings.

**Definition 3.1** • The exit point for the stationary LPP to  $(m, n)$  with boundary (3.1) is the last point on the  $x$ -axis or the  $y$ -axis of the maximizer ending at  $(m, n)$ . We introduce the random variable  $Z^\rho(m, n) \in \mathbb{Z}$  such that, if  $Z^\rho(m, n) > 0$ , then the exit point is  $(Z^\rho(m, n), 0)$ , and if  $Z^\rho(m, n) < 0$ , then the exit point is  $(0, -Z^\rho(m, n))$ .

- The exit point for the stationary LPP to  $(m, n)$  with boundary (3.2) is the starting point of the maximizer ending at  $(m, n)$ . We use the notation  $\tilde{Z}^\rho(m, n) \in \mathbb{Z}$  such that the exit point is  $(\tilde{Z}^\rho(m, n), -\tilde{Z}^\rho(m, n))$ .
- The exit point for the LPP from  $\mathcal{L}$  with initial condition  $h^0$  is the starting point of the maximizer ending at  $(m, n)$ . We use the notation  $Z_{h^0}(m, n) \in \mathbb{Z}$  such that the exit point is  $(Z_{h^0}(m, n), -Z_{h^0}(m, n))$ . For the random initial condition with parameter  $\sigma$ , we denote  $Z_{h^0} = Z_\sigma$ , and for flat initial condition  $Z_{h^0} = Z^\setminus$ .

Now we state the two comparison lemmas which we are going to use in the proof of Theorem 2.5.

**Lemma 3.2** Denote by  $L^\rho$  the LPP (3.1) and  $L^\bullet$  the LPP in the droplet case. Let  $0 \leq m_1 \leq m_2$  and  $n_1 \geq n_2 \geq 0$ . Then if  $Z^\rho(m_1, n_1) \geq 0$ , it holds

$$L^\bullet(m_2, n_2) - L^\bullet(m_1, n_1) \leq L^\rho(m_2, n_2) - L^\rho(m_1, n_1), \tag{3.3}$$

while, if  $Z^\rho(m_2, n_2) \leq 0$ , then we have

$$L^\bullet(m_2, n_2) - L^\bullet(m_1, n_1) \geq L^\rho(m_2, n_2) - L^\rho(m_1, n_1). \tag{3.4}$$

**Lemma 3.3** Denote by  $L^\rho$  the LPP (3.2) and  $L^*$  be the LPP from  $\mathcal{L}$  with boundary term  $h^0$ . Let  $0 \leq m_1 \leq m_2$  and  $n_1 \geq n_2 \geq 0$ . Then if  $\tilde{Z}^\rho(m_1, n_1) \geq \tilde{Z}_{h^0}(m_2, n_2)$ , it holds

$$L^*(m_2, n_2) - L^*(m_1, n_1) \leq L^\rho(m_2, n_2) - L^\rho(m_1, n_1), \tag{3.5}$$

while, if  $\tilde{Z}^\rho(m_2, n_2) \leq \tilde{Z}_{h^0}(m_1, n_1)$ , then we have

$$L^*(m_2, n_2) - L^*(m_1, n_1) \geq L^\rho(m_2, n_2) - L^\rho(m_1, n_1). \tag{3.6}$$

For  $n_1 = n_2$ , Lemma 3.2 is in Lemma 1 of [15], while Lemma 3.3 is Lemma 2.1 of [41]. The generalization to points on a down-right path is straightforward. It was made for instance in the LPP setting (3.2) in Lemma 3.5 of [24].

## 4 Convergence of the Covariance

### 4.1 Preliminaries and Notations

A law of large number for point-to-point LPP was proven in [47], namely, for large  $(m, n)$ ,  $L_{(0,0) \rightarrow (m,n)} \approx (\sqrt{m} + \sqrt{n})^2$ . From this we can estimate

$$\begin{aligned} L_{(0,0) \rightarrow E_\tau}^* &\approx 4\tau N - w_\tau^2 \tau^{-1} 2^{4/3} N^{1/3}, \\ L_{(0,0) \rightarrow I(u)}^* &\approx 4\tau N - u^2 \tau^{-1} 2^{4/3} (\tau N)^{1/3}, \\ L_{I(u) \rightarrow E_1} &\approx 4(1 - \tau)N - \frac{(u - w_1)^2}{1 - \tau} 2^{4/3} N^{1/3}. \end{aligned} \tag{4.1}$$

Denote the rescaled LPP by

$$L_N^*(u, \tau) := \frac{L_{S_A \rightarrow I(u)}^* - 4(1 - \tau)N}{2^{4/3} N^{1/3}}, \tag{4.2}$$

with  $I(u) = (\tau N, \tau N) + u(2N)^{2/3}(1, -1)$  and

$$L_N^{pp}(u, \tau) := \frac{L_{I(u) \rightarrow E_1} - 4(1 - \tau)N}{2^{4/3} N^{1/3}}, \tag{4.3}$$

where we recall that  $E_1 = (N, N) + w_1(2N)^{2/3}(1, -1)$ . Then, (2.7) and (2.8) become

$$\begin{aligned} L_N^*(\tau) &\equiv L_N^*(w_\tau, \tau), \\ L_N^*(1) &\equiv L_N^*(w_1, 1) = \max_{u \in \mathbb{R}} \{L_N^*(u, \tau) + L_N^{pp}(u, \tau)\}. \end{aligned} \tag{4.4}$$

Furthermore,

$$\lim_{N \rightarrow \infty} L_N^{pp}(u, \tau) = (1 - \tau)^{1/3} \left[ \mathcal{A}_2 \left( \frac{u - w_1}{(1 - \tau)^{2/3}} \right) - \frac{(u - w_1)^2}{(1 - \tau)^{4/3}} \right], \tag{4.5}$$

and

$$\lim_{N \rightarrow \infty} L_N^*(u, \tau) = \tau^{1/3} \mathcal{A}^\star \left( \frac{u}{\tau^{2/3}} \right), \quad \star \in \{\bullet, \setminus, \mathcal{B}\}, \tag{4.6}$$

where

$$\mathcal{A}^\bullet(u) = \tilde{\mathcal{A}}_2(u) - u^2, \quad \mathcal{A}^\setminus(u) = 2^{1/3} \mathcal{A}_1(u 2^{-2/3}), \quad \mathcal{A}^\mathcal{B}(u) = \mathcal{A}_{\text{stat}}(u). \tag{4.7}$$

### 4.2 Localization of the Maximizer at Time $\tau N$

The maximizer of the process  $L_N^*(u, \tau) + L_N^{pp}(u, \tau)$  is confined in the region with  $|u| \leq M$  if the following event holds

$$\Omega_M^G = \left\{ \max_{|u| \leq M} \{L_N^*(u, \tau) + L_N^{pp}(u, \tau)\} > \max_{|u| > M} \{L_N^*(u, \tau) + L_N^{pp}(u, \tau)\} \right\}. \tag{4.8}$$

Thus we need to estimate  $\mathbb{P}(\Omega_M^G)$ . For any choice of  $s \in \mathbb{R}$  we can write

$$\begin{aligned} \mathbb{P}(\Omega_M^G) &\geq \mathbb{P}\left(\max_{|u| \leq M} \{L_N^*(u, \tau) + L_N^{pp}(u, \tau)\} > s > \max_{|u| > M} \{L_N^*(u, \tau) + L_N^{pp}(u, \tau)\}\right) \\ &\geq 1 - \mathbb{P}(G_M) - \mathbb{P}(B_M), \end{aligned} \tag{4.9}$$

where we defined

$$\begin{aligned} G_M &= \{\max_{|u| \leq M} \{L_N^*(u, \tau) + L_N^{pp}(u, \tau)\} \leq s\}, \\ B_M &= \{\max_{|u| > M} \{L_N^*(u, \tau) + L_N^{pp}(u, \tau)\} > s\}. \end{aligned} \tag{4.10}$$

The right side of (4.9) is estimated using the following lemma.

**Lemma 4.1** *Let  $s = -M^2\tilde{c}$  with  $\tilde{c} = 1/(16(1 - \tau))$ . Then, there exists a finite  $M_0$  such that for any  $M \geq M_0$*

$$\begin{aligned} \mathbb{P}(G_M) &\leq Ce^{-cM^2} \\ \mathbb{P}(B_M) &\leq Ce^{-cM^2} \end{aligned} \tag{4.11}$$

for some constants  $C, c > 0$  uniform in  $N$ .

As a direct consequence we have the following localization result.

**Corollary 4.2** *For any  $M \geq M_0$ ,*

$$\mathbb{P}(\text{the maximizer of } L_N^*(u, \tau) + L_N^{pp}(u, \tau) \text{ passes by } I(u) \text{ with } |u| > M) \leq 2Ce^{-cM^2} \tag{4.12}$$

uniformly in  $N$ .

Denote by

$$\chi_M^*(1) = \max_{|u| \leq M} \left\{ \tau^{1/3} \mathcal{A}^*(\tau^{-2/3}u) + (1 - \tau)^{1/3} \left[ \mathcal{A}_2\left(\frac{u-w_1}{(1-\tau)^{2/3}}\right) - \frac{(u-w_1)^2}{(1-\tau)^{4/3}} \right] \right\} \tag{4.13}$$

and recall

$$\chi^*(\tau) = \tau^{1/3} \mathcal{A}^*(\tau^{-2/3}w_\tau). \tag{4.14}$$

**Lemma 4.3** *We have the convergence of joint distributions*

$$\begin{aligned} &\lim_{N \rightarrow \infty} \mathbb{P}\left(\max_{|u| \leq M} \{L_N^*(u, \tau) + L_N^{pp}(u, \tau)\} \leq s_1; L_N^*(w_\tau, \tau) \leq s_2\right) \\ &= \mathbb{P}(\chi_M^*(1) \leq s_1; \chi^*(\tau) \leq s_2). \end{aligned} \tag{4.15}$$

*Proof* It is enough to have weak convergence of the two rescaled process to the terms in the rhs. As mentioned above, the point-wise convergence have been already proven. So we need tightness in the space of continuous functions of  $[-M, M]$ . Tightness  $L_N^{pp}(u, \tau)$  and  $L_N^*(u, \tau)$  can be found in Corollary 4.2 of [27], for  $L_N^B(u, \tau)$

it is a direct a direct consequence of Lemma 4.2 of [7] and the standard Donsker’s theorem. Finally, tightness for  $L_N^\searrow(u, \tau)$  has been established in [41].  $\square$

### 4.2.1 Localization of the Process

Let us prove Lemma 4.1 and Corollary 4.2.

*Proof of Lemma 4.1* Recall that to prove this lemma, we take  $s = s_0$ , with the choice  $s_0 = -M^2\tilde{c} = -M^2/(16(1 - \tau))$ .

- (1) Bound on  $\mathbb{P}(G_M)$ .

We have

$$\begin{aligned} \mathbb{P}(G_M) &= \mathbb{P}\left(\max_{|u|\leq M} \{L_N^\star(u, \tau) + L_N^{\text{PP}}(u, \tau)\} \leq s_0\right) \\ &\leq \mathbb{P}\left(L_N^\bullet(0, \tau) + L_N^{\text{PP}}(0, \tau) \leq s_0\right) \\ &\leq \mathbb{P}\left(L_N^\bullet(0, \tau) \leq s_0/2\right) + \mathbb{P}\left(L_N^{\text{PP}}(0, \tau) \leq s_0/2\right). \end{aligned} \tag{4.16}$$

Now we can use standard estimates on the lower tail of the point-to-point LPP (see Proposition A.1 in Appendix A) to obtain that (4.16) is bounded by  $Ce^{-cM^3}$  uniformly in  $N$ , for some constants  $C, c$ .

- (2) Bound on  $\mathbb{P}(B_M)$ . Since similar estimates will be used to derive another result, we add an extra variable  $\hat{s} \geq 0$  in the following computations. The case  $\hat{s} = 0$  is the one relevant for the present proof.

We have

$$\begin{aligned} \mathbb{P}(B_M) &= \mathbb{P}\left(\max_{|u|>M} \{L_N^\star(u, \tau) + L_N^{\text{PP}}(u, \tau)\} > s_0 + \hat{s}\right) \\ &\leq \mathbb{P}\left(\max_{|u|>M} \left\{L_N^\star(u, \tau) - \frac{u^2}{2(1 - \tau)}\right\} > \frac{s_0 + \hat{s}}{2}\right) \\ &\quad + \mathbb{P}\left(\max_{|u|>M} \left\{L_N^{\text{PP}}(u, \tau) + \frac{u^2}{2(1 - \tau)}\right\} > \frac{s_0 + \hat{s}}{2}\right). \end{aligned} \tag{4.17}$$

We study separately the two terms of (4.17) and rename them  $\mathbb{P}(B_M^1)$  and  $\mathbb{P}(B_M^2)$  respectively. Remark that the maximum over  $u$  is actually a maximum over  $M < |u| \leq \mathcal{O}(N^{1/3})$ , since  $I(u)$  need to stay in the backward light cone of the end-point  $E_1$ . We will not write this explicitly all the time.

- (a) Estimate of  $\mathbb{P}(B_M^2)$ :

$$\mathbb{P}(B_M^2) = \mathbb{P}\left(\max_{|u|>M} \left\{L_N^{\text{PP}}(u, \tau) + \frac{u^2}{2(1 - \tau)}\right\} > \frac{s_0 + \hat{s}}{2}\right) \tag{4.18}$$

The bound can be obtained through the decay of the kernel for half-flat initial condition. The bound on the Fredholm determinant and the kernel are given as in Theorem 2.6 and Lemma 2.7 of [16] to get

$$\mathbb{P}(B_M^2) \leq Ce^{-cM^2(1-\tau)^{-4/3}} e^{-\tilde{c}\hat{s}}. \tag{4.19}$$

Alternatively, one could adapt the proof of Lemma 4.3 of [27] to get the same result.

(b) Estimate of  $\mathbb{P}(B_M^1)$ :

$$\mathbb{P}(B_M^1) = \mathbb{P}\left(\max_{|u|>M} \left\{L_N^*(u, \tau) - \frac{u^2}{2(1-\tau)}\right\} > \frac{s_0 + \hat{s}}{2}\right). \tag{4.20}$$

- Droplet initial condition: for this case, one can estimate it like we made for (4.18) (with minor changes in the terms depending on  $\tau$ ). However, since  $L_N^\bullet(u, \tau) \leq L_N^\setminus(u, \tau)$ , the droplet upper tail is simply bounded by the upper tail of the flat initial condition case.
- Flat initial condition: the bound is obtained in Lemma 4.4 below.
- Stationary initial condition: the bounds for the maximum over  $u > M$  and for  $u < -M$  are similar and thus we present the details only for the first one.

$$\begin{aligned} & \mathbb{P}\left(\max_{u>M} \left\{L_N^*(u, \tau) - \frac{u^2}{2(1-\tau)}\right\} > \frac{s_0 + \hat{s}}{2}\right) \\ & \leq \mathbb{P}\left(\max_{u>M} \left\{L_N^*(u, \tau) - L_N^*(M, \tau) - \frac{u^2}{2(1-\tau)}\right\} > s_0 + \frac{\hat{s}}{4}\right) \\ & \quad + \mathbb{P}\left(L_N^*(0, \tau) > -\frac{s_0}{4} + \frac{\hat{s}}{8}\right) + \mathbb{P}\left(L_N^*(M, \tau) - L_N^*(0, \tau) > -\frac{s_0}{4} + \frac{\hat{s}}{8}\right). \end{aligned} \tag{4.21}$$

We study separately the three terms of the last line of (4.21). The first term is bounded using (D.5), the second with (D.1) and the third one with (D.4), with the final result

$$\mathbb{P}(B_M^1) \leq C e^{-cM^2 - \tilde{c}\hat{s}} \tag{4.22}$$

for some  $c, \tilde{c}$  depending on  $\tau$ , but uniform for all  $N$  large enough.

□

*Proof of Corollary 4.2* By (4.16), (4.22), (4.19) we can conclude that

$$\mathbb{P}\left(\max_{|u|\leq M} \{L_N^*(u, \tau) + L_N^{pp}(u, \tau)\} > \max_{|u|>M} \{L_N^*(u, \tau) + L_N^{pp}(u, \tau)\}\right) \geq 1 - 2C e^{-cM^2}, \tag{4.23}$$

which implies that the probability that the maximizer of  $L_N^*(u, \tau) + L_N^{pp}(u, \tau)$  passes through  $I(u)$  with  $|u| > M$  goes to zero as  $2C e^{-cM^2}$ , for some constants  $C, c > 0$ .

□

Here for simplicity of notation we rename  $\hat{s}$  as  $s$ .

**Lemma 4.4** *For flat initial condition, there exist  $N_0, M_0$  large enough such that for all  $N \geq N_0$  and  $M \geq M_0$  it holds*

$$\mathbb{P}\left(\max_{M < |u| < \mathcal{O}(N^{1/3})} \left\{L_N^\setminus(u, \tau) - \frac{u^2}{2(1-\tau)}\right\} > \frac{s_0 + s}{2}\right) \leq C e^{-cs} e^{-\tilde{c}M^2}, \tag{4.24}$$

for some constants  $C, c, \tilde{c}$  independent of  $N$  and  $M$ .

*Proof* By symmetry we can consider only the case  $u > M$ , since the bounds for  $u < -M$  are similar. Fix an  $\varepsilon \in (0, 1/6)$ . Then,

$$\begin{aligned} & \mathbb{P}\left(\max_{M < u < \mathcal{O}(N^{1/3})} \left\{L_N^{\setminus}(u, \tau) - \frac{u^2}{2(1-\tau)}\right\} > \frac{s_0 + s}{2}\right) \\ & \leq \sum_{\ell=1}^{N^\varepsilon} \mathbb{P}\left(\max_{u \in [\ell M, (\ell+1)M]} \left\{L_N^{\setminus}(u, \tau) - \frac{u^2}{2(1-\tau)}\right\} > \frac{s_0 + s}{2}\right) \\ & + \sum_{u \in [N^\varepsilon, \mathcal{O}(N^{2/3})]} \mathbb{P}\left(L_N^{\setminus}(u, \tau) - \frac{u^2}{2(1-\tau)} > \frac{s_0 + s}{2}\right). \end{aligned} \tag{4.25}$$

Notice that  $v \mapsto L_N^{\setminus}(u + v, \tau)$  and  $v \mapsto L_N^{\setminus}(v, \tau)$  have the same law for any  $u$ . Thus, we can simply bound (using also (B.3))

$$\begin{aligned} \mathbb{P}\left(L_N^{\setminus}(u, \tau) - \frac{u^2}{2(1-\tau)} > \frac{s_0 + s}{2}\right) & \leq \mathbb{P}\left(L_N^{\setminus}(0, \tau) > \frac{s}{2} - \frac{M^2}{32(1-\tau)} + \frac{N^{2\varepsilon}}{2(1-\tau)}\right) \\ & \leq C e^{-cs/2 + cM^2/(32(1-\tau)) - cN^{2\varepsilon}/(2(1-\tau))} \\ & \leq C e^{-\tilde{c}s - \hat{c}M^2} e^{-cN^{2\varepsilon}/(4(1-\tau))}, \end{aligned} \tag{4.26}$$

for some constants  $C, c, \tilde{c}, \hat{c}$ , where the last inequality holds for all  $N \geq N_0(M)$ . From this it immediately follows that

$$\sum_{u \in [N^\varepsilon, \mathcal{O}(N^{2/3})]} \mathbb{P}\left(L_N^{\setminus}(u, \tau) - \frac{u^2}{2(1-\tau)} > \frac{s_0 + s}{2}\right) \leq C e^{-\tilde{c}s - \hat{c}M^2}. \tag{4.27}$$

Now we evaluate the first term in (4.25). Using translation-invariance in  $u$  we get

$$\begin{aligned} & \mathbb{P}\left(\max_{u \in [\ell M, (\ell+1)M]} \left\{L_N^{\setminus}(u, \tau) - \frac{u^2}{2(1-\tau)}\right\} > \frac{s_0 + s}{2}\right) \\ & \leq \mathbb{P}\left(\max_{u \in [0, M]} L_N^{\setminus}(u, \tau) > \frac{s}{2} - \frac{M^2}{32(1-\tau)} + \frac{\ell^2 M^2}{2(1-\tau)}\right) \\ & \leq \mathbb{P}\left(\max_{u \in [0, M]} L_N^{\setminus}(u, \tau) > \frac{s}{2} + \frac{\ell^2 M^2}{4(1-\tau)}\right) \end{aligned} \tag{4.28}$$

(a) The first case is  $s \geq N^{2\varepsilon}$ . We can still just use the union bound and the exponential decay to get

$$\begin{aligned} (4.28) & \leq N^{2/3} \exp\left(-c \frac{\ell^2 M^2}{4(1-\tau)} - c \frac{s}{2}\right) \leq N^{2/3} \exp\left(-c \frac{\ell^2 M^2}{4(1-\tau)} - c \frac{s}{4} - c \frac{N^{2\varepsilon}}{4}\right) \\ & \leq \exp(-c_1 \ell^2 M^2 - c_2 s), \end{aligned} \tag{4.29}$$

for some constants  $c_1, c_2 > 0$  and all  $N$  large enough.

(b) The second case is  $s \in (0, N^{2\varepsilon})$ . Since also  $\ell \leq N^\varepsilon$  we have that  $x := \frac{s}{2} + \frac{\ell^2 M^2}{8(1-\tau)} = \mathcal{O}(N^{2\varepsilon}) \ll N^{1/3}$ . The idea is to bound the process in terms

of the increments of a stationary case. For that reason we first need to get a formula including the increments of the rescaled LPP, namely we get

$$(4.50) \leq \mathbb{P} \left( L_N^{\setminus} (0, \tau) > \frac{s}{4} + \frac{\ell^2 M^2}{8(1-\tau)} \right) + \mathbb{P} \left( \max_{u \in [0, M]} \left\{ L_N^{\setminus} (u, \tau) - L_N^{\setminus} (0, \tau) \right\} > \frac{s}{2} + \frac{\ell^2 M^2}{8(1-\tau)} \right). \quad (4.30)$$

For the first term, we just use (B.3) and obtain

$$\mathbb{P} \left( L_N^{\setminus} (0, \tau) > \frac{s}{4} + \frac{\ell^2 M^2}{8(1-\tau)} \right) \leq C e^{-cs/4 - c\ell^2 M^2 / (8(1-\tau))}. \quad (4.31)$$

The sum of this bound over  $\ell \geq 1$  leads to a bound  $\tilde{C} e^{-cs/4 - cM^2 / (8(1-\tau))}$ . For the second term in (4.30), define  $\rho_+ = \frac{1}{2} + \kappa N^{-1/3}$  and the event

$$\Omega_{N, \kappa} = \{ \tilde{Z}^{\rho_+} (I(0)) > \tilde{Z}^{\setminus} (I(u)), \text{ for all } u \in [0, M] \}. \quad (4.32)$$

On this event, by Lemma 3.3, we have

$$L_N^{\setminus} (u, \tau) - L_N^{\setminus} (0, \tau) \leq L_N^{\rho_+} (u, \tau) - L_N^{\rho_+} (0, \tau), \quad (4.33)$$

which in turns gives

$$\begin{aligned} & \mathbb{P} \left( \max_{u \in [0, M]} \left\{ L_N^{\setminus} (u, \tau) - L_N^{\setminus} (0, \tau) \right\} > \frac{s}{2} + \frac{\ell^2 M^2}{8(1-\tau)} \right) \\ & \leq \mathbb{P} \left( \max_{u \in [0, M]} \left\{ L_N^{\rho_+} (u, \tau) - L_N^{\rho_+} (0, \tau) \right\} > \frac{s}{2} + \frac{\ell^2 M^2}{8(1-\tau)} \right) + \mathbb{P}(\Omega_{N, \kappa}^c). \end{aligned} \quad (4.34)$$

By stationarity of the increments we have

$$L_N^{\rho_+} (u, \tau) - L_N^{\rho_+} (0, \tau) = \frac{1}{2^{4/3} N^{1/3}} \sum_{i=1}^{\lfloor uN^{2/3} \rfloor} Z_i, \quad (4.35)$$

where  $Z_i = X_i - Y_i$  with  $X_i$ 's i.i.d.  $\text{Exp}(1 - \rho_+)$  and  $Y_i$ 's i.i.d.  $\text{Exp}(\rho_+)$  random variables. Since  $\mathcal{M}_u = \sum_{i=1}^{\lfloor uN^{2/3} \rfloor} Z_i$  is a submartingale, so it is  $\exp(t\mathcal{M}_u)$  for  $t > 0$  (at least for  $t$  small enough) and we can use Doob's inequality for submartingales,

$$\mathbb{P} \left( \max_{u \in [0, M]} \mathcal{M}_u \geq x \right) \leq \inf_{t \geq 0} \frac{\mathbb{E} [e^{t\mathcal{M}_M}] }{e^{tx}} = \inf_{t \geq 0} \frac{\mathbb{E} [e^{tZ_1}]^{M N^{2/3}} }{e^{tx}}. \quad (4.36)$$

For and  $\rho_+ = \frac{1}{2} + \kappa N^{-1/3}$ . An explicit computation gives, for  $\kappa \in (0, x/(2^{5/3}M))$ ,

$$(4.36) \leq \exp \left( -\frac{(2^{1/3}x - 4M\kappa)^2}{4M} + \mathcal{O}(x^4 N^{-2/3}, \kappa^4 N^{-2/3}) \right). \quad (4.37)$$

Thus, with the choice  $\kappa = x/(2^{8/3}M)$ , we find

$$\begin{aligned} &\mathbb{P}\left(\max_{u \in [0, M]} \{L_N^{\rho_+}(u, \tau) - L_N^{\rho_+}(0, \tau)\} > \frac{s}{2} + \frac{\ell^2 M^2}{8(1 - \tau)}\right) \\ &\leq \exp\left(-\frac{x^2}{16M}\right) \leq \exp\left(-c_1 s^2 - c_2 \ell^4 M^3\right), \end{aligned} \tag{4.38}$$

for some constants  $c_1, c_2 > 0$  and all  $N$  large enough. Summing this bound over  $\ell \geq 1$  we get  $Ce^{-c_1 s^2 - c_2 M^3}$  for some constants  $C$ , completing the proof of (4.24). □

**Lemma 4.5** *Let  $\rho_{\pm} = \frac{1}{2} \pm \kappa N^{-1/3}$ . Define the event*

$$\Omega_{N, \kappa} = \left\{ \tilde{Z}^{\rho_+}(I(0)) \geq \tilde{Z} \setminus (I(u)), \forall u \in [0, M] \right\} \cap \left\{ \tilde{Z}^{\rho_-}(I(0)) \leq \tilde{Z} \setminus (I(u)), \forall u \in [-M, 0] \right\}. \tag{4.39}$$

where the exit points are as in Definition 3.1. Then, for all  $N$  large enough and all  $\kappa > 0$  with  $\kappa = o(N^{1/3})$ ,

$$\mathbb{P}(\Omega_{N, \kappa}^c) \leq Ce^{-c\kappa^2}. \tag{4.40}$$

*Proof* We need to estimate the complement of the probabilities of the two terms in (4.39), for instance

$$\mathbb{P}(\tilde{Z} \setminus (I(u)) > \tilde{Z}^{\rho_+}(I(0))), \text{ for some } u \in [0, M]. \tag{4.41}$$

The estimates are completely analogous, thus we provide the details only for the first one.

Since  $\tilde{Z} \setminus (I(u)) \leq \tilde{Z} \setminus (I(M))$  for all  $u \in [0, M]$ , we have

$$\begin{aligned} (4.41) &\leq \mathbb{P}(\tilde{Z} \setminus (I(M)) > \tilde{Z}^{\rho_+}(I(0))) \\ &\leq \mathbb{P}(\tilde{Z} \setminus (I(M)) > \alpha(2N)^{2/3}) + \mathbb{P}(\tilde{Z}^{\rho_+}(I(0)) < \alpha(2N)^{2/3}). \end{aligned} \tag{4.42}$$

By Lemma 4.3 of [27], we have that  $\mathbb{P}(\tilde{Z} \setminus (I(M)) > \alpha(2N)^{2/3}) \leq Ce^{-c\alpha^2}$ , for some constants  $C, c \in (0, \infty)$ . Using stationarity of the increments along the antidiagonal, we have

$$\tilde{Z}^{\rho_+}(n - k, n + k) \stackrel{d}{=} \tilde{Z}^{\rho_+}(n, n) - k. \tag{4.43}$$

Thus,

$$\begin{aligned} \mathbb{P}(\tilde{Z}^{\rho_+}(I(0)) < \alpha(2N)^{2/3}) &= \mathbb{P}(\tilde{Z}^{\rho_+}(I(0)) - \alpha(2N)^{2/3} \leq 0) \\ &= \mathbb{P}(\tilde{Z}^{\rho_+}(I(-\alpha)) < 0) = \mathbb{P}(Z^{\rho_+}(I(-\alpha)) < 0). \end{aligned} \tag{4.44}$$

The last equality follows from the fact that we can construct the two models on the same randomness (define the random variables in the model (3.1) as image of the ones of (3.2) by [7]), for which  $\tilde{Z}^{\rho_+}(m, n) < 0$  iff  $Z^{\rho_+}(m, n) < 0$  by simple geometric considerations.



Setting  $(\gamma^2 n, n) = I(-\alpha)$ , and writing  $\rho_+ = 1/(1 + \gamma) + \tilde{\kappa} n^{-1/3}$ , we deduce that  $\tilde{\kappa} = \tau^{2/3} \kappa - 2^{-4/3} \alpha \tau^{-1/3} + \mathcal{O}(\kappa N^{-1/3})$ . Lemma 2.5 of [27] states<sup>8</sup> that if  $\tilde{\kappa} > 0$ , then  $\mathbb{P}(Z^{\rho_+}(\gamma^2 n, n) < 0) \leq C e^{-c\tilde{\kappa}^2}$  for some constants  $C, c > 0$ . We choose  $\alpha = 2^{1/3} \tau \kappa$ , which gives  $\tilde{\kappa} = \frac{1}{2} \kappa \tau^{2/3} (1 + \mathcal{O}(\kappa/N^{1/3}))$ . Then, for all  $N$  large enough, we obtain

$$(4.44) \leq C e^{-\tilde{c}\kappa^2} \tag{4.45}$$

for some constants  $C, c > 0$ . □

### 4.2.2 Convergence of the Covariance

To prove Theorem 2.2, first we show that the  $N \rightarrow \infty$  limit of the covariance of  $L_N^*(\tau)$  and  $\max_{|u| \leq M} \{L_N^*(u, \tau) + L_N^{pp}(u, \tau)\}$  is the covariance of  $\chi^*(\tau)$  and  $\chi_M^*(1)$ , for fixed  $M > 0$ . Now that we have proved the localization of the process, we need to show that the covariance of  $\chi^*(\tau)$  and  $\chi^*(1)$  is the  $M \rightarrow \infty$  limit of the covariance of the LPP restricted to the region  $|u| \leq M$ .

**Proposition 4.6** *For any fixed  $M > 0$ ,*

$$\lim_{N \rightarrow \infty} \text{Cov} \left( L_N^*(w_\tau, \tau), \max_{|u| \leq M} \{L_N^*(u, \tau) + L_N^{pp}(u, \tau)\} \right) = \text{Cov} (\chi^*(\tau), \chi_M^*(1)). \tag{4.46}$$

*Proof* Let us denote

$$L_{N;M}^*(1) = \max_{|u| \leq M} \{L_N^*(u, \tau) + L_N^{pp}(u, \tau)\}. \tag{4.47}$$

By Lemma 4.1 we already have the convergence of joint distributions of  $L_{N;M}^*(1)$  and  $L_N^*(\tau) \equiv L_N^*(w_\tau, \tau)$  to  $\chi_M^*(1)$  and  $\xi^*(\tau)$ . By Cauchy-Schwarz it is enough to show the convergence of the second moments of  $L_{N;M}^*(1)$  and  $L_N^*(\tau)$ .

For a random variable  $X_N$  with distribution function  $F_N(s) = \mathbb{P}(X_N \leq s)$ , we can write

$$\mathbb{E}(X_N^2) = 2 \int_{\mathbb{R}_+} s(1 - F_N(s)) ds - 2 \int_{\mathbb{R}_-} s F_N(s) ds. \tag{4.48}$$

If we know that  $X_N \rightarrow X$  in distribution, to show convergence of the second moment we need only to find  $g(s)$  independent of  $N$  such that  $1 - F_N(s) \leq g(s)$  for  $s \geq 0$ ,  $F_N(s) \leq g(s)$  for  $s < 0$  and that  $g \in L^1(\mathbb{R})$ . Then dominated convergence allows to take the limit in the integrals and obtain  $\mathbb{E}(X_N^2) \rightarrow \mathbb{E}(X^2)$ . Thus our task is to find such bounds. Since  $F_N(s) \in [0, 1]$ , it is enough to get bounds for the tails, i.e., a bound for  $1 - F_N(s)$  for  $s \geq s_0$  and for  $F_N(s)$  for  $s \leq -s_0$  for some  $s_0$ .

$$(1) \quad \lim_{N \rightarrow \infty} \mathbb{E}[(L_N^*(\tau))^2] = \mathbb{E}[(\chi^*(\tau))^2].$$

---

<sup>8</sup>By inspecting the proof of Lemma 2.5 of [27], one sees that it actually holds true not only for any given  $\kappa$ , but also for all  $\kappa \in [0, o(n^{1/3})]$ .

- bound on lower tails: due to  $L_N^*(\tau) \geq L_N^\bullet(\tau)$ , we can use for all cases the lower bound for the droplet initial condition, which is in Proposition A.1 (by appropriate change of variables).
- bound on upper tails: (a) for the droplet initial condition, this is in Proposition A.1, (b) for the flat initial condition, this is given in Proposition A.1, (c) for the stationary initial condition,<sup>9</sup> we have

$$\mathbb{P}(L_N^{\mathcal{B}}(\tau) \leq s) \leq \mathbb{P}(L_N^{\mathcal{B}}(0) \leq s/2) + \mathbb{P}(L_N^{\mathcal{B}}(\tau) - L_N^{\mathcal{B}}(0) \leq s/2). \tag{4.49}$$

The first term is bounded using Proposition A.4, while the second using Proposition A.5.

In all cases we have at least exponential decay of the both the upper and lower tails. This implies the convergence of the second moment as well.

(2)  $\lim_{N \rightarrow \infty} \mathbb{E}[(L_{N;M}^*(1))^2] = \mathbb{E}[(\chi_M^*(1))^2]$ .

- bound on lower tails: we have

$$\begin{aligned} \mathbb{P}(L_{N;M}^*(1) \leq s) &\leq \mathbb{P}(L_N^*(0, \tau) \leq s/2) + \mathbb{P}(L_N^{\text{pp}}(0, \tau) \leq s/2) \\ &\leq \mathbb{P}(L_N^\bullet(0, \tau) \leq s/2) + \mathbb{P}(L_N^{\text{pp}}(0, \tau) \leq s/2) \leq C e^{-c|s|^{3/2}} \end{aligned} \tag{4.50}$$

by Proposition A.1.

- bound on upper tails: we have  $L_{N;M}^*(1) \leq L_N^*(1)$  and thus by the estimates used already in part (1) we have

$$\mathbb{P}(L_N^*(1) \geq s) \leq C e^{-cs}. \tag{4.51}$$

These bounds implies convergence of the second moment as well.

□

What remains to prove Theorem 2.2 is a control on the contribution to the covariance from the events when the maximizer passed by  $I(u)$  for some  $|u| > M$ . We have the decomposition

$$\begin{aligned} &\text{Cov}(L_N^*(\tau), L_N^*(1)) \\ &= \text{Cov}(L_N^*(\tau), L_{N;M}^*(1)) + \text{Cov}(L_N^*(\tau), L_N^*(1) - L_{N;M}^*(1)). \end{aligned} \tag{4.52}$$

Given the convergence of the second moments for fixed  $M$  by Proposition 4.6, there is one term left to study:

$$\begin{aligned} &|\text{Cov}(L_N^*(\tau), L_N^*(1) - L_{N;M}^*(1))| \\ &= |\mathbb{E}[L_N^*(\tau)(L_N^*(1) - L_{N;M}^*(1))] - \mathbb{E}[L_N^*(\tau)]\mathbb{E}[L_N^*(1) - L_{N;M}^*(1)]| \\ &\leq 2 \left( \mathbb{E}[(L_N^*(\tau))^2] \mathbb{E}[(L_N^*(1) - L_{N;M}^*(1))^2] \right)^{1/2}, \end{aligned} \tag{4.53}$$

where we used Cauchy-Schwarz to control the second term.

---

<sup>9</sup>For stationary initial condition, the convergence of all moments was already proven in [3].

**Lemma 4.7** For any  $M > 0$ ,

$$\lim_{N \rightarrow \infty} \mathbb{E} \left[ (L_N^*(1) - L_{N;M}^*(1))^2 \right] \leq C e^{-cM^2} \tag{4.54}$$

as well as

$$\mathbb{E}[(\chi^*(1) - \chi_M^*(1))^2] \leq C e^{-cM^2}. \tag{4.55}$$

where  $C, c > 0$  are positive constants, uniformly in  $N$ .

*Proof* Let us denote

$$L_{N;M^c}^*(1) = \max_{|u| > M} \{L_N^*(u, \tau) + L_N^{pp}(u, \tau)\}. \tag{4.56}$$

Since  $L_N^*(1) = \max\{L_{N;M}^*(1), L_{N;M^c}^*(1)\}$ , we can write

$$L_N^*(1) - L_{N;M}^*(1) = \max\{0, L_{N;M^c}^*(1) - L_{N;M}^*(1)\}. \tag{4.57}$$

Integrating by parts, we obtain

$$\mathbb{E} \left[ (L_N^*(1) - L_{N;M}^*(1))^2 \right] = 2 \int_{\mathbb{R}_+} s \mathbb{P} (L_{N;M^c}^*(1) - L_{N;M}^*(1) > s). \tag{4.58}$$

The probability in the r.h.s. of (4.58) can be bounded as

$$\mathbb{P} (L_{N;M^c}^*(1) - L_{N;M}^*(1) > s) \leq \mathbb{P} \left( L_{N;M^c}^*(1) > \frac{s + \alpha}{2} \right) + \mathbb{P} \left( L_{N;M}^*(1) \leq \frac{\alpha - s}{2} \right) \tag{4.59}$$

for any choice of  $\alpha$ .

(a) We use the first inequality in (4.28) and obtain

$$\mathbb{P} \left( L_{N;M}^*(1) < \frac{\alpha - s}{2} \right) \leq \mathbb{P} \left( L_N^*(0, \tau) < \frac{\alpha - s}{4} \right) + \mathbb{P} \left( L^{pp}(0, \tau) < \frac{\alpha - s}{4} \right). \tag{4.60}$$

For any  $\alpha < 0$  and  $s \geq 0$ , by Proposition A.1 we get

$$\mathbb{P} \left( L_{N;M}^*(1) < \frac{\alpha - s}{2} \right) \leq C e^{-c(s-\alpha)^{3/2}} \tag{4.61}$$

for some constants  $C, c$ . Thus it is enough to choose  $\alpha = -\gamma M^2$  for some  $\gamma > 0$ .

(b) Next we bound  $\mathbb{P} \left( L_{N;M^c}^*(1) > \frac{s+\alpha}{2} \right)$ . Choosing  $\alpha = -\frac{M^2}{16(1-\tau)}$  and using the bounds for  $\mathbb{P}(B_M)$  in the proof of Lemma 4.1, we obtain

$$\mathbb{P} \left( L_{N;M^c}^*(1) > \frac{s + \alpha}{2} \right) \leq C e^{-cM^2} e^{-\tilde{c}s}. \tag{4.62}$$

Plugging the bounds (4.61) and (231) into (4.58) leads (4.54).

Finally, (4.55) is proven as follows. By dominated convergence we have that

$$\begin{aligned} \mathbb{E}[(\chi^*(1) - \chi_M^*(1))^2] &= 2 \int_{\mathbb{R}_+} s \mathbb{P}(\chi_{M^c}^*(1) - \chi_M^*(1) > s) \\ &= \lim_{N \rightarrow \infty} 2 \int_{\mathbb{R}_+} s \mathbb{P}(L_{N;M^c}^*(1) - L_{N;M}^*(1) > s) \leq C e^{-cM^2} \end{aligned} \tag{4.63}$$

where the last inequality follows from (4.54). □

*Proof of Theorem 2.2* We have

$$\begin{aligned} &\lim_{N \rightarrow \infty} \text{Cov}(L_N^*(\tau), L_N^*(1)) \\ &= \lim_{M \rightarrow \infty} \lim_{N \rightarrow \infty} \text{Cov}(L_N^*(\tau), L_{N,M}^*(1)) + \lim_{M \rightarrow \infty} \lim_{N \rightarrow \infty} \text{Cov}(L_N^*(\tau), L_N^*(1) - L_{N,M}^*(1)). \end{aligned} \tag{4.64}$$

By Proposition 4.6, the first term equals  $\text{Cov}(\chi^*(\tau), \chi_M^*(1))$ . By Lemma 4.7, the second term is 0. Thus what remains is to show that

$$\text{Cov}(\chi^*(\tau), \chi^*(1)) = \lim_{M \rightarrow \infty} \text{Cov}(\chi^*(\tau), \chi_M^*(1)). \tag{4.65}$$

This is obtained once we prove that

$$\lim_{M \rightarrow \infty} \mathbb{E}[(\chi^*(1) - \chi_M^*(1))^2] = 0, \tag{4.66}$$

which is also part of Lemma 4.7. □

### 4.3 Formula for the Stationary Case

Now we prove the claimed formula for the stationary case. It follows by a simple computation using the result of Theorem 2.2 for the stationary case and the identity (2.18).

*Proof of Corollary 2.4* Setting  $X_1 = \chi^{\mathcal{B}}(\tau)$  and  $X_2 = \chi^{\mathcal{B}}(1)$  in (2.18) we get

$$\text{Cov}(\chi^{\mathcal{B}}(\tau), \chi^{\mathcal{B}}(1)) = \frac{1}{2} \text{Var}(\chi^{\mathcal{B}}(\tau)) + \frac{1}{2} \text{Var}(\chi^{\mathcal{B}}(1)) - \frac{1}{2} \text{Var}(\chi^{\mathcal{B}}(1) - \chi^{\mathcal{B}}(\tau)). \tag{4.67}$$

The first two terms in (2.17) are an immediate consequence of the convergence of moments, see proof of Proposition 4.6. For the last term, we have

$$\begin{aligned} \chi^{\mathcal{B}}(1) - \chi^{\mathcal{B}}(\tau) &= \max_{u \in \mathbb{R}} \left\{ (1 - \tau)^{1/3} \left[ \mathcal{A}_2 \left( \frac{u - w_1}{(1 - \tau)^{2/3}} \right) - \frac{(u - w_1)^2}{(1 - \tau)^{4/3}} \right] \right. \\ &\quad \left. + \tau^{1/3} [\mathcal{A}_{\text{stat}}(\tau^{-2/3} u) - \mathcal{A}_{\text{stat}}(\tau^{-2/3} w_\tau)] \right\}. \end{aligned} \tag{4.68}$$

Changing the variable  $u = w_\tau + z(1 - \tau)^{2/3}$ , and calling  $\xi = \frac{w_1 - w_\tau}{(1-\tau)^{2/3}}$ , it gives

$$\begin{aligned} \chi^{\mathcal{B}}(1) - \chi^{\mathcal{B}}(\tau) &= (1 - \tau)^{1/3} \max_{z \in \mathbb{R}} \left\{ \mathcal{A}_2(z - \xi) - (z - \xi)^2 \right. \\ &\quad \left. + \frac{\tau^{1/3}}{(1 - \tau)^{1/3}} \left[ \mathcal{A}_{\text{stat}}(\tau^{-2/3}(w_\tau + z(1 - \tau)^{2/3})) - \mathcal{A}_{\text{stat}}(\tau^{-2/3}w_\tau) \right] \right\}. \end{aligned} \tag{4.69}$$

Next we use the facts: (a)  $\mathcal{A}_2(z - \xi) \stackrel{(d)}{=} \mathcal{A}_2(z)$ , (b)  $\mathcal{A}_{\text{stat}}(a + x) - \mathcal{A}_{\text{stat}}(a) \stackrel{(d)}{=} \sqrt{2}B(x)$  with  $B$  a standard Brownian motion, and (c) the scaling of Brownian motion, to get

$$\chi^{\mathcal{B}}(1) - \chi^{\mathcal{B}}(\tau) \stackrel{(d)}{=} (1 - \tau)^{1/3} \max_{z \in \mathbb{R}} \left\{ \sqrt{2}B(z) + \mathcal{A}_2(z) - (z - \xi)^2 \right\} \stackrel{(d)}{=} \mathcal{A}_{\text{stat}}(\xi)(1 - \tau)^{1/3}. \tag{4.70}$$

□

### 5 Behavior Around $\tau = 1$

What we have to prove is

$$\text{Var} [\chi^*(1) - \chi^*(\tau)] = (1 - \tau)^{2/3} \text{Var} (\xi_{\text{stat}, \tilde{w}}) + \mathcal{O}(1 - \tau)^{1-\delta}, \tag{5.1}$$

as  $\tau \rightarrow 1$  for all the initial conditions. Clearly the flat and stationary are special case of the more generic random initial conditions. Define

$$\begin{aligned} \chi_M^* &= \lim_{N \rightarrow \infty} \max_{|u| \leq (1-\tau)^{2/3}M} (L_N^*(u, \tau) + L_N^{\text{pp}}(u, \tau)) \\ &= \lim_{N \rightarrow \infty} \max_{|v| \leq M} (L_N^*((1 - \tau)^{2/3}v, \tau) + L_N^{\text{pp}}((1 - \tau)^{2/3}v, \tau)). \end{aligned} \tag{5.2}$$

In particular, for droplet, flat, stationary initial conditions, we have

$$\chi_M^* = (1 - \tau)^{1/3} \max_{|v| \leq M} \left( \left( \frac{\tau}{1-\tau} \right)^{1/3} \mathcal{A}^* \left( v \left( \frac{1-\tau}{\tau} \right)^{2/3} \right) + \mathcal{A}_2(v - \tilde{w}_1) - (v - \tilde{w}_1)^2 \right), \tag{5.3}$$

with  $\mathcal{A}^*$  being the  $\text{Airy}_2$ ,  $\text{Airy}_1$  or  $\text{Airy}_{\text{stat}}$  process for  $\star = \bullet, \setminus, \mathcal{B}$  respectively. Also, recall the notation

$$\chi^*(\tau) = \lim_{N \rightarrow \infty} L_N^*((1 - \tau)^{2/3}\tilde{w}_\tau, \tau) = (1 - \tau)^{1/3} \left( \frac{\tau}{1-\tau} \right)^{1/3} \mathcal{A}^* \left( \tilde{v}(\tau) \left( \frac{1-\tau}{\tau} \right)^{2/3} \right). \tag{5.4}$$

On short scales,  $\mathcal{A}^*$  is expected to behave similar to the stationary state, which is a two-sided Brownian motion with diffusion coefficient 2. Since the  $\text{Airy}_2$  process is stationary, for  $\tau \rightarrow 1$ ,  $\chi_M^* - \chi^*(\tau)$  should be close to the following expression

$$\xi_{M, \tilde{w}_\tau, \tilde{w}_1} := (1 - \tau)^{1/3} \max_{|v| \leq M} \left\{ \sqrt{2}B(v - \tilde{w}_\tau) + \mathcal{A}_2(v) - (v - \tilde{w}_1)^2 \right\}. \tag{5.5}$$

In this proof we set

$$\tilde{w} = \tilde{w}_1 - \tilde{w}_\tau. \tag{5.6}$$

For  $M = \infty$ , replacing  $v - \tilde{w}_\tau \rightarrow \tilde{v}$  and using the stationarity of  $\mathcal{A}_2$  we obtain

$$\xi_{\infty, \tilde{w}_\tau, \tilde{w}_1} \stackrel{(d)}{=} (1 - \tau)^{1/3} \max_{\tilde{v} \in \mathbb{R}} \{ \sqrt{2} B(\tilde{v}) + \mathcal{A}_2(\tilde{v}) - (\tilde{v} - \tilde{w})^2 \} = \xi_{\text{stat}, \tilde{w}}. \tag{5.7}$$

Note that in distribution

$$(1 - \tau)^{1/3} (\xi_{\text{GUE}} - \tilde{w}^2) \leq \xi_{M, \tilde{w}_\tau, \tilde{w}_1} \leq (1 - \tau)^{1/3} \xi_{\text{stat}, \tilde{w}} \tag{5.8}$$

and therefore we know that the  $m$ th moment of  $\xi_{M, \tilde{w}_\tau, \tilde{w}_1}$  is finite and of order  $(1 - \tau)^{m/3}$ .

To control the error term, the idea is to take  $M$  depending on  $\tau$  such that  $M \rightarrow \infty$  as  $\tau$  goes to 1. Then the task is to prove that the difference between the second moment of  $\chi_M^*(1) - \chi^*(\tau)$  and the second moment of  $\xi_{M, \tilde{w}_\tau, \tilde{w}_1}$  goes to zero as  $\tau \rightarrow 1$ .

**Lemma 5.1** *Let  $M = \frac{1}{(1-\tau)^{\delta/2}}$  with  $\delta > 0$ . Then*

$$\left| \mathbb{E} \left[ (\chi_M^*(1) - \chi^*(\tau))^2 \right] - \mathbb{E} \left[ \xi_{M, \tilde{w}_\tau, \tilde{w}_1}^2 \right] \right| = \mathcal{O}(1 - \tau)^{1-\delta}. \tag{5.9}$$

We need to control how close the increments of the process over distances of order  $(1 - \tau)^{2/3}$  at time  $\tau$  are with respect to the increments of Brownian motion.

We present a short technical lemma that will be used in the proof of Lemma 5.1. Recall Definition 3.1 of the exit point for a LPP with boundary conditions (3.1) (for the droplet case) or (3.2) (for the random case) and define  $\rho^\pm = \frac{1}{2} \pm \kappa N^{-1/3}$ . Denote by  $L^{\rho^\pm}$  its associated LPP.

**Lemma 5.2** *There is an event  $\Omega_\kappa$  with  $\mathbb{P}(\Omega_\kappa) \geq 1 - C \exp(-c\tilde{\kappa}^2)$ , with*

- (a) *for droplet initial condition,  $\tilde{\kappa} = \kappa - \frac{M(1-\tau)^{2/3}}{2^{4/3}\tau}$ ,*
- (b) *for random initial condition,  $\tilde{\kappa} = \kappa - 2 \frac{M(1-\tau)^{2/3}}{2^{4/3}\tau}$ ,*  
*and constants  $C, c, M_0 \in (0, \infty)$ , such that on  $\Omega_\kappa$  the inequalities*

$$\xi_{M, \tilde{w}_\tau, \tilde{w}_1} - \varepsilon_0 \leq \chi_M^*(1) - \chi^*(\tau) \leq \xi_{M, \tilde{w}_\tau, \tilde{w}_1} + \varepsilon_0, \tag{5.10}$$

*hold in distribution, for all  $M \geq M_0$  with  $\varepsilon_0 = \mathcal{O}(\kappa M(1 - \tau)^{2/3})$ , under the condition  $\tilde{\kappa} > 0$ .*

*Proof* Let us define

$$\Delta_N^\bullet(u) = \frac{L_{(0,0) \rightarrow I(u)} - L_{(0,0) \rightarrow E_\tau}}{2^{4/3} N^{1/3}}, \quad \Delta_N^\sigma(u) = \frac{L_{\mathcal{L}^\sigma \rightarrow I(u)} - L_{\mathcal{L}^\sigma \rightarrow E_\tau}}{2^{4/3} N^{1/3}}, \tag{5.11}$$

and recall the definitions

$$\begin{aligned} L_N^\bullet(u, \tau) &= \frac{L_{(0,0) \rightarrow I(u)} - 4\tau N}{2^{4/3} N^{1/3}}, & L_N^\sigma(u, \tau) &= \frac{L_{\mathcal{L}^\sigma \rightarrow I(u)} - 4\tau N}{2^{4/3} N^{1/3}}, \\ L_N^{\text{pp}}(u, \tau) &= \frac{L_{I(u) \rightarrow E_1} - 4(1 - \tau)N}{2^{4/3} N^{1/3}}. \end{aligned} \tag{5.12}$$

Then we have

$$L_N^\star(u, \tau) = L_N^\star(0, \tau) + \Delta_N^\star(u). \tag{5.13}$$

Also, recall that we will use the notation  $u = v(1 - \tau)^{2/3}$  and  $\tilde{M} = M(1 - \tau)^{2/3}$ .

Define the event

$$\Omega_{N,\kappa} = \begin{cases} \{Z^{\rho+}(I(-\tilde{M})) > 0, Z^{\rho-}(I(\tilde{M})) < 0\}, & \text{for droplet IC,} \\ \{\tilde{Z}^{\rho+}(I(-\tilde{M})) > Z_{\sigma}(I(\tilde{M})), \tilde{Z}^{\rho-}(I(\tilde{M})) < Z_{\sigma}(I(-\tilde{M}))\}, & \text{for random IC.} \end{cases} \tag{5.14}$$

Then, on the event  $\Omega_{N,\kappa}$  we can bound  $\Delta_N^*(u)$  with the increments of the stationary LPP with density  $\rho_{\pm}$ , defined as

$$B^{\pm}(u) = \frac{L^{\rho_{\pm}}(I(u)) - L^{\rho_{\pm}}(I(0)) - m_{\rho_{\pm}}u(2N)^{2/3}}{2^{4/3}N^{1/3}}, \tag{5.15}$$

where  $m_{\rho_{\pm}} = \frac{1}{1-\rho_{\pm}} - \frac{1}{\rho_{\pm}} = 8\kappa N^{-1/3} + \mathcal{O}(N^{-1})$ . Indeed a minimal modification of Lemma 3.2 implies, for  $-\tilde{M} \leq w < u \leq \tilde{M}$ ,

$$\begin{aligned} [B^-(u) - B^-(w)] - 4(u - w)\kappa &\leq L_N^*(u, \tau) - L_N^*(w, \tau) \\ &\leq [B^+(u) - B^+(w)] + 4(u - w)\kappa \end{aligned} \tag{5.16}$$

for  $N$  large enough. Furthermore,  $\text{Var}(B^{\pm}(u)) = u2^{1/2}(1 + \mathcal{O}(N^{-2/3}))$  and  $B^{\pm}(0) = 0$ . Thus by Donsker’s theorem,  $\lim_{N \rightarrow \infty} B^{\pm}(u) = \sqrt{2}B(u)$ , with  $B(u)$  a standard two-sided Brownian motion in the space of continuous functions on bounded sets.

Recall that

$$\begin{aligned} \chi_M^*(1) - \chi^*(\tau) &= \lim_{N \rightarrow \infty} \max_{|v| \leq M} \left\{ L_N^*(v(1 - \tau)^{2/3}, \tau) - L_N^*(\tilde{w}_{\tau}(1 - \tau)^{2/3}, \tau) \right. \\ &\quad \left. + L_N^{\text{pp}}(v(1 - \tau)^{2/3}, \tau) \right\} \end{aligned} \tag{5.17}$$

and also that  $v \mapsto L_N^{\text{pp}}(v(1 - \tau)^{2/3}, \tau)$  converges weakly to  $(1 - \tau)^{1/3}[\mathcal{A}_2(v) - (v - \tilde{w}_1)^2]$ . Thus, taking the  $N \rightarrow \infty$  limit and using the inequalities (54) we obtain

$$\begin{aligned} &\mathbb{P}\left(\chi_M^*(1) - \chi^*(\tau) \leq (1 - \tau)^{1/3}s\right) \\ &\leq \mathbb{P}\left(\max_{|v| \leq M} \left\{ \sqrt{2}B(v) - B(\tilde{w}_{\tau}) + \mathcal{A}_2(v) - (v - \tilde{w}_1)^2 - 4\kappa(v - \tilde{w}_{\tau})(1 - \tau)^{1/3} \right\} \leq s\right). \end{aligned} \tag{5.18}$$

Denoting  $\varepsilon = \max_{|v| \leq M} |4\kappa(v - w)(1 - \tau)^{1/3}| = 6\kappa M(1 - \tau)^{1/3}$  we obtain

$$\begin{aligned} &\mathbb{P}\left(\chi_M^*(1) - \chi^*(\tau) \leq (1 - \tau)^{1/3}s\right) \\ &\leq \mathbb{P}\left(\max_{|v| \leq M} \left\{ \sqrt{2}B(v - \tilde{w}_{\tau}) + \mathcal{A}_2(v) - (v - \tilde{w}_1)^2 \right\} \leq s + \varepsilon\right) \\ &= \mathbb{P}\left(\xi_{M, \tilde{w}_{\tau}, \tilde{w}_1} \leq (1 - \tau)^{1/3}s + \varepsilon_0\right) \end{aligned} \tag{5.19}$$

with  $\varepsilon_0 = (1 - \tau)^{1/3}\varepsilon$ .

Similarly for the lower bound we get

$$\mathbb{P}\left(\chi_M^*(1) - \chi^*(\tau) > (1 - \tau)^{1/3}s\right) \leq \mathbb{P}\left(\xi_{M, \tilde{w}_{\tau}, \tilde{w}_1} > (1 - \tau)^{1/3}s + \varepsilon_0\right). \tag{5.20}$$

To conclude the proof, we need to estimate  $\mathbb{P}(\Omega_{N,\kappa})$ .

(a) *Droplet initial condition:* For this case, we apply Lemma 2.5 of [27]. To estimate  $\mathbb{P}(Z^{\rho_{\pm}}(I(\mp \tilde{M})) > 0)$ , we need to set  $I(\mp \tilde{M}) = (\gamma^2 n, n)$ . This gives

$$\rho_{\pm} = \frac{1}{2} \pm \frac{\tilde{M}}{2^{4/3} \tau N^{1/3}} \pm \frac{\tilde{\kappa}}{\tau^{2/3} N^{1/3}}. \tag{5.21}$$

Then, Lemma 2.5 of [27] gives

$$\mathbb{P}(Z^{\rho_{\pm}}(I(\mp \tilde{M})) > 0) \geq 1 - Ce^{-c\tilde{\kappa}^2} = 1 - Ce^{-c(\tau^{2/3} \kappa - 2^{-4/3} \tilde{M} \tau^{-1/3})^2}. \tag{5.22}$$

The estimates are uniform for all  $N$  large enough. Renaming  $c\tau^{4/3}$  as a new constant  $c$ , and  $2C$  by  $C$ , we get

$$\mathbb{P}(\Omega_{N,\kappa}) \geq 1 - C \exp\left(-c\left(\kappa - \frac{M(1-\tau)^{2/3}}{2^{4/3}\tau}\right)^2\right). \tag{5.23}$$

(b) *Random initial condition:* We derive a bound only for  $\mathbb{P}(\tilde{Z}^{\rho_{+}}(I(-\tilde{M})) < Z_{\sigma}(I(\tilde{M})))$ , since bounding  $\mathbb{P}(\tilde{Z}^{\rho_{-}}(I(\tilde{M})) < Z_{\sigma}(I(-\tilde{M})))$  is completely analogue.

The probability we want to bound is smaller than

$$\mathbb{P}(\tilde{Z}^{\rho_{+}}(I(-\tilde{M})) \leq \alpha(2N)^{2/3}) + \mathbb{P}(Z_{\sigma}(I(\tilde{M})) > \alpha(2N)^{2/3}), \tag{5.24}$$

and we choose  $\alpha = 2^{1/3} \tau \kappa$ . Exactly as in (4.44), we have

$$\mathbb{P}(\tilde{Z}^{\rho_{+}}(I(-\tilde{M})) \leq \alpha(2N)^{2/3}) = \mathbb{P}(Z^{\rho_{+}}(I(-\tilde{M} - \alpha)) < 0) \leq Ce^{-c\tilde{\kappa}^2} \tag{5.25}$$

with  $\tilde{\kappa} = 2\tau^{2/3} \left(\kappa - \frac{(1-\tau)^{2/3} M}{2^{1/3} \tau}\right)$ , provided  $\tilde{\kappa} > 0$ .

Now we bound  $\mathbb{P}(Z_{\sigma}(I(\tilde{M})) > \alpha(2N)^{2/3})$ . Let  $J(v) = v(2N)^{2/3}(1, -1)$ , define the scaled variables

$$L_N(v) = \frac{L_{J(v) \rightarrow I(\tilde{M})} - 4(1-\tau)N}{2^{4/3} N^{1/3}}, \quad \mathcal{W}_N(v) = \frac{h^0(J(v))}{2^{4/3} N^{1/3}}. \tag{5.26}$$

Then,

$$\begin{aligned} \mathbb{P}(Z_{\sigma}(I(\tilde{M})) > \alpha(2N)^{2/3}) &\leq \mathbb{P}\left(\max_{v \leq \alpha} (L_N(v) + \mathcal{W}_N(v)) \leq -s\right) \\ &\quad + \mathbb{P}\left(\max_{v > \alpha} (L_N(v) + \mathcal{W}_N(v)) \geq -s\right). \end{aligned} \tag{5.27}$$

Since  $L_N(v) \sim -(v - \tilde{M})^2/(\tau)$ , we choose  $s = (\alpha - \tilde{M})^2/4$ .

The first term in (5.27) is bounded by

$$\mathbb{P}(L_N(\alpha) + \mathcal{W}_N(\alpha) \leq -s) \leq \mathbb{P}\left(L_N(\alpha) \leq -\frac{3}{2}s\right) + \mathbb{P}\left(\mathcal{W}_N(\alpha) \leq \frac{1}{2}s\right). \tag{5.28}$$

The first term bounded by  $C_1 e^{-c_2(\alpha - \tilde{M})^3}$  by (A.4). Since  $\mathcal{W}_N$  is a (rescaled) sum of iid. random variables, we can use the the exponential Chebyshev's inequality (see e.g. the proof of (D.4)) and obtain a bound  $C_2 e^{-c_2(\alpha - \tilde{M})^4/\alpha}$ .



The second term in (5.27) is bounded by

$$\mathbb{P}\left(\max_{v \geq \alpha} \left(L_N(v) + \frac{1}{2}(v - \tilde{M})^2\right) \geq -\frac{1}{2}s\right) + \mathbb{P}\left(\max_{v \geq \alpha} \left(\mathcal{W}_N(v) - \frac{1}{2}(v - \tilde{M})^2\right) \geq -\frac{1}{2}s\right). \tag{5.29}$$

The first term is estimated similarly to (4.18) and leads to a bound  $C_3 e^{-c_3(\alpha - \tilde{M})^2}$ . The second term is bounded using Doob’s maximal inequality (see e.g. the proof of (D.5) leading to a bound  $C_4 e^{-c_4(\alpha - \tilde{M})^4/\alpha}$ ).

Combining these bounds we get  $\mathbb{P}(Z_\sigma(I(\tilde{M})) > \alpha(2N)^{2/3}) \leq C e^{-c\tilde{\kappa}^2}$ , provided  $\tilde{\kappa} > 0$ , for some constants  $C, c \in (0, \infty)$  uniformly for all  $\tau$  in a compact subset of  $(0, 1]$ . Up to renaming  $c\tau^{4/3}$  to  $c$  and the constant  $2C$  to  $C$  we get the claimed result.  $\square$

Now we can prove Lemma 5.1.

*Proof of Lemma 5.1* By Lemma 5.2 we have, on a event  $\Omega_\kappa$  with  $\mathbb{P}(\Omega_\kappa^c) \leq C e^{-c\tilde{\kappa}^2}$ , with

$$\tilde{\kappa} = \kappa - \frac{M(1 - \tau)^{2/3}}{2^{4/3}\tau}. \tag{5.30}$$

the inequality

$$(\chi_M^*(1) - \chi^*(\tau))\mathbb{1}_{\Omega_\kappa} = \xi_{M, \tilde{w}_\tau, \tilde{w}_1}\mathbb{1}_{\Omega_\kappa} + \zeta\mathbb{1}_{\Omega_\kappa}, \tag{5.31}$$

for some random variables  $\zeta$  with  $|\zeta| \leq \varepsilon_0$ . Thus

$$\mathbb{E}[(\chi_M^*(1) - \chi^*(\tau))^2] = \mathbb{E}[(\chi_M^*(1) - \chi^*(\tau))^2\mathbb{1}_{\Omega_\kappa}] + \mathbb{E}[(\chi_M^*(1) - \chi^*(\tau))^2\mathbb{1}_{\Omega_\kappa^c}]. \tag{5.32}$$

Using (5.31) we get

$$\mathbb{E}[(\chi_M^*(1) - \chi^*(\tau))^2\mathbb{1}_{\Omega_\kappa}] = \mathbb{E}(\xi_{M, \tilde{w}_\tau, \tilde{w}_1}^2) - \mathbb{E}(\xi_{M, \tilde{w}_\tau, \tilde{w}_1}^2\mathbb{1}_{\Omega_\kappa^c}) + 2\mathbb{E}(\zeta\xi_{M, \tilde{w}_\tau, \tilde{w}_1}\mathbb{1}_{\Omega_\kappa}) + \mathbb{E}(\zeta^2\mathbb{1}_{\Omega_\kappa}). \tag{5.33}$$

Using Cauchy-Schwarz and the fact that  $|\zeta| \leq \varepsilon_0$ , we get the bounds

$$\begin{aligned} \mathbb{E}(\xi_{M, \tilde{w}_\tau, \tilde{w}_1}^2\mathbb{1}_{\Omega_\kappa^c}) &\leq \sqrt{\mathbb{E}(\xi_{M, \tilde{w}_\tau, \tilde{w}_1}^4)\mathbb{P}(\Omega_\kappa^c)} \leq C_1(1 - \tau)^{2/3}e^{-c\tilde{\kappa}^2/2}, \\ |\mathbb{E}(\zeta\xi_{M, \tilde{w}_\tau, \tilde{w}_1}\mathbb{1}_{\Omega_\kappa})| &\leq \varepsilon_0\sqrt{\mathbb{E}(\xi_{M, \tilde{w}_\tau, \tilde{w}_1}^2)} \leq C_2(1 - \tau)^{1/3}\varepsilon_0, \\ \mathbb{E}(\zeta^2\mathbb{1}_{\Omega_\kappa}) &\leq \varepsilon_0^2, \end{aligned} \tag{5.34}$$

for some constants  $C_1, C_2$  (since, as already mentioned, the  $m$ th moment of  $\xi_{M, \tilde{w}_\tau, \tilde{w}_1}$  is of order  $(1 - \tau)^{m/3}$ ).

It remains to bound the last term of (5.32). Let  $\Lambda = \{|\chi_M^*(1) - \chi^*(\tau)| \leq \lambda\}$  and decompose  $(\chi_M^*(1) - \chi^*(\tau))\mathbb{1}_{\Omega_\kappa^c}$  as  $(\chi_M^*(1) - \chi^*(\tau))\mathbb{1}_{\Omega_\kappa^c}(\mathbb{1}_\Lambda + \mathbb{1}_{\Lambda^c})$ . Then,

$$\mathbb{E}[(\chi_M^*(1) - \chi^*(\tau))^2\mathbb{1}_{\Omega_\kappa^c}] \leq \mathbb{E}[(\chi_M^*(1) - \chi^*(\tau))^2\mathbb{1}_{\Lambda^c}] + \lambda^2\mathbb{P}(\Omega_\kappa^c). \tag{5.35}$$

Integration by parts gives

$$\begin{aligned} \mathbb{E} \left[ (\chi_M^*(1) - \chi^*(\tau))^2 \mathbb{1}_{\Lambda^c} \right] &= \lambda^2 \mathbb{P}(|\chi_M^*(1) - \chi^*(\tau)| > \lambda) \\ &+ 2 \int_{\lambda}^{\infty} s \mathbb{P}(\chi_M^*(1) - \chi^*(\tau) > s) ds - 2 \int_{-\infty}^{-\lambda} s \mathbb{P}(\chi_M^*(1) - \chi^*(\tau) \leq s) ds. \end{aligned} \tag{5.36}$$

Now, for  $s > 0$ ,

$$\mathbb{P}(\chi_M^*(1) - \chi^*(\tau) > s) \leq \mathbb{P}(\chi_M^*(1) \geq s/2) + \mathbb{P}(\chi^*(\tau) \leq -s/2), \tag{5.37}$$

and for  $s < 0$ ,

$$\mathbb{P}(\chi_M^*(1) - \chi^*(\tau) \leq s) \leq \mathbb{P}(\chi_M^*(1) \leq s/2) + \mathbb{P}(\chi^*(\tau) \geq -s/2). \tag{5.38}$$

Recall that

$$\begin{aligned} \mathbb{P}(\chi_M^*(1) > s) &\leq \mathbb{P}(\chi^*(1) > s), \\ \mathbb{P}(\chi_M^*(1) \leq s) &\leq \mathbb{P}((1 - \tau)^{1/3} \tilde{\mathcal{A}}_2(0) \leq s) = F_{\text{GUE}}(s/(1 - \tau)^{1/3}). \end{aligned} \tag{5.39}$$

Since both tails of  $\chi^*(1)$  and of the GUE Tracy-Widom distributions have (at least) exponential decay (see Appendix C and D), it then follows that

$$\mathbb{E} \left[ (\chi_M^*(1) - \chi^*(\tau))^2 \mathbb{1}_{\Lambda^c} \right] \leq C \lambda^2 e^{-c\lambda/(1-\tau)^{1/3}} \tag{5.40}$$

for some constants  $C, c$ .

Summing up we have obtained

$$\begin{aligned} &\mathbb{E}[(\chi_M^*(1) - \chi^*(\tau))^2] - \mathbb{E}(\xi_{M, \tilde{w}_\tau, \tilde{w}_1}^2) \\ &= \mathcal{O} \left( (1 - \tau)^{2/3} e^{-c\tilde{\kappa}^2/2}; (1 - \tau)^{1/3} \varepsilon_0; \varepsilon_0^2; \lambda^2 e^{-c\tilde{\kappa}^2}; \lambda^2 e^{-c\lambda/(1-\tau)^{1/3}} \right), \end{aligned} \tag{5.41}$$

with  $\varepsilon_0 = \mathcal{O}(\kappa M(1 - \tau)^{2/3})$ . Now we choose  $M, \kappa, \lambda$ . Let  $\delta \in (0, 1/3)$  be any fixed number and choose

$$M = \frac{1}{(1 - \tau)^{\delta/2}}, \quad \kappa = \frac{1}{(1 - \tau)^{\delta/2}}, \quad \lambda = 1. \tag{5.42}$$

Then, the error term in (5.41) is just of order  $\mathcal{O}((1 - \tau)^{1-\delta})$ . □

Now we are ready to prove Theorem 2.5.

*Proof of Theorem 2.5* We have

$$\begin{aligned} \mathbb{E} \left[ (\chi^*(1) - \chi^*(\tau))^2 \right] &= \mathbb{E} \left[ (\chi^*(1) - \chi_M^*(1))^2 \right] + \mathbb{E} \left[ (\chi_M^*(1) - \chi^*(\tau))^2 \right] \\ &\quad + 2\mathbb{E} \left[ (\chi^*(1) - \chi_M^*(1))(\chi_M^*(1) - \chi^*(\tau)) \right] \end{aligned} \tag{5.43}$$

With the choice  $M = (1 - \tau)^{-\delta/2}$ , by Lemma 5.1 we have

$$\mathbb{E} \left[ (\chi_M^*(1) - \chi^*(\tau))^2 \right] = \mathbb{E}[\xi_{M, \tilde{w}_\tau, \tilde{w}_1}^2] + \mathcal{O}(1 - \tau)^{1-\delta}. \tag{5.44}$$

By Lemma 4.7, the Cauchy-Schwarz inequality, and  $\xi_{\infty, \hat{w}} = (1 - \tau)^{1/3} \xi_{\text{stat}, \hat{w}}$ , we obtain

$$\mathbb{E} \left[ (\chi^*(1) - \chi^*(\tau))^2 \right] = (1 - \tau)^{2/3} \mathbb{E}[\xi_{\text{stat}, \hat{w}}]^2 + \mathcal{O}((1 - \tau)^{1-\delta}). \tag{5.45}$$

Since  $\mathbb{E}[\xi_{\text{stat}, \hat{w}}] = 0$  the claimed result is proven. □

### 6 Behavior Around $\tau = 0$ for Droplet Initial Conditions

Let us finally explain the asymptotic for  $\tau \rightarrow 0$ . The details are simple modifications of what we made for the case  $\tau \rightarrow 1$ . By Theorem 1 of [41], we have local weak convergence of the  $\text{Airy}_2$  process to a Brownian motion for  $\tau \rightarrow 0$ ,

$$\lim_{\tau \rightarrow 0} \left( \frac{\tau}{1-\tau} \right)^{-1/3} \left( \tilde{\mathcal{A}}_2 \left( \left( \frac{\tau}{1-\tau} \right)^{2/3} v \right) - \tilde{\mathcal{A}}_2(0) \right) = \sqrt{2}B(v). \tag{6.1}$$

Lemmas 5.1 and 5.2 can be easily readjusted for this case. Let us call  $w_\tau = \tau^{2/3} \hat{w}_\tau$ . Then, by Theorem 2.2, renaming  $u = z\tau^{2/3}$ ,

$$\begin{aligned} & \text{Cov}(\chi^*(\tau), \chi^*(1)) \\ &= \text{Cov} \left( \tau^{1/3} [\tilde{\mathcal{A}}_2(\hat{w}_\tau) - \hat{w}_\tau^2], \tau^{1/3} \max_{z \in \mathbb{R}} \left\{ \tilde{\mathcal{A}}_2(z) - z^2 + \left( \frac{1-\tau}{\tau} \right)^{1/3} \mathcal{A}_2 \left( z \frac{\tau^{2/3}}{(1-\tau)^{2/3}} \right) - z^2 \frac{\tau}{1-\tau} \right\} \right) \\ &= \tau^{2/3} \text{Cov} \left( \tilde{\mathcal{A}}_2(\hat{w}_\tau), \max_{z \in \mathbb{R}} \left\{ \tilde{\mathcal{A}}_2(z) - z^2 + \sqrt{2}B(z) \right\} + \left( \frac{1-\tau}{\tau} \right)^{1/3} \mathcal{A}_2(0) \right) + \mathcal{O}(\tau^{1-\delta}) \\ &= \tau^{2/3} \text{Cov} \left( \tilde{\mathcal{A}}_2(\hat{w}_\tau), \max_{z \in \mathbb{R}} \left\{ \tilde{\mathcal{A}}_2(z) - z^2 + \sqrt{2}B(z) \right\} \right) + \mathcal{O}(\tau^{1-\delta}), \end{aligned} \tag{6.2}$$

for any  $\delta > 0$ , where the covariance of  $\tilde{\mathcal{A}}_2(\hat{w}_\tau)$  and  $\mathcal{A}_2(0)$  is zero, since they are independent processes. The second term in the covariance has the same distribution as  $\xi_{\text{stat}, 0}$ , which is has expected value 0. This leads to the claimed result of Theorem 2.6.

**Acknowledgments** This work is supported by the German Research Foundation in the Collaborative Research Center 1060 “The Mathematics of Emergent Effects”, project B04.

### Appendix A: Bounds on Point-to-Point LPP

In the proofs, we use known results for the point-to-point LPP with exponential random variables, which we recall here.

**Proposition A.1** For  $\eta \in (0, \infty)$  define  $\mu = (\sqrt{\eta \ell} + \sqrt{\ell})^2$ ,  $\sigma = \eta^{-1/6} (1 + \sqrt{\eta})^{4/3}$ , and the rescaled random variable

$$L_\ell^{\text{resc}} := \frac{L_{(0,0) \rightarrow (\eta \ell, \ell)} - \mu}{\sigma \ell^{1/3}}. \tag{A.1}$$

(a) *Limit law*

$$\lim_{\ell \rightarrow \infty} \mathbb{P}(L_\ell^{\text{resc}} \leq s) = F_{GUE}(s), \tag{A.2}$$

with  $F_{GUE}$  the GUE Tracy-Widom distribution function.

(b) *Bound on upper tail: there exist constants  $s_0, \ell_0, C, c$  such that*

$$\mathbb{P}(L_\ell^{\text{resc}} \geq s) \leq Ce^{-cs} \tag{A.3}$$

for all  $\ell \geq \ell_0$  and  $s \geq s_0$ .

(c) *Bound on lower tail: there exist constants  $s_0, \ell_0, C, c$  such that*

$$\mathbb{P}(L_\ell^{\text{resc}} \leq s) \leq Ce^{-c|s|^{3/2}} \tag{A.4}$$

for all  $\ell \geq \ell_0$  and  $s \leq -s_0$ .

(a) was proven in Theorem 1.6 of [30]. Using the relation with the Laguerre ensemble of random matrices (Proposition 6.1 of [1]), or to TASEP described above, the distribution is given by a Fredholm determinant. An exponential decay of its kernel leads directly to (b). See e.g. Proposition 4.2 of [25] or Lemma 1 of [3] for an explicit statement. (c) was proven in [3] (Proposition 3 together with (56)). In the present language it is reported in Proposition 4.3 of [25] as well.

### Appendix B: Bounds for Point-to-Line LPP

**Proposition B.1** *Let  $\mathcal{L} = \{(k, -k), k \in \mathbb{Z}\}$ . Consider the rescaled LPP from  $\mathcal{L}$  to  $(\ell, \ell)$  given by*

$$L_\ell^{\mathcal{L}, \text{resc}} = \frac{L_{\mathcal{L}} \rightarrow (\ell, \ell) - 4\ell}{2^{4/3} \ell^{1/3}}. \tag{B.1}$$

(a) *Limit law*

$$\lim_{\ell \rightarrow \infty} \mathbb{P}(L_\ell^{\mathcal{L}, \text{resc}} \leq s) = F_{GOE}(2^{2/3}s). \tag{B.2}$$

(b) *Bound on upper tail: there exists constants  $s_0, \ell_0, C, c$  such that*

$$\mathbb{P}(L_\ell^{\mathcal{L}, \text{resc}} \geq s) \leq Ce^{-cs} \tag{B.3}$$

for all  $\ell \geq \ell_0$  and  $s \geq s_0$ .

(c) *Bound on lower tail: there exists constants  $s_0, \ell_0, C, c$  such that*

$$\mathbb{P}(L_\ell^{\mathcal{L}, \text{resc}} \leq s) \leq Ce^{-c|s|^{3/2}} \tag{B.4}$$

for all  $\ell \geq \ell_0$  and  $s \leq -s_0$ .

(a) was obtained in [11, 48] in terms of TASEP, which can be directly rewritten in term of LPP (the complete proof is present in [10]). For general slopes of  $\mathcal{L}$  it was shown in [27]. (b) this tails follows from the asymptotic analysis on the correlation kernel made in [10]. (c) It follows from (A.4) since  $\mathbb{P}(L_{\mathcal{L}} \rightarrow (\ell, \ell) \leq x) \leq \mathbb{P}(L_{(0,0) \rightarrow (\ell, \ell)} \leq x)$ .

### Appendix C: Bounds on LPP with Random Initial Condition

**Proposition C.1** Define  $L_{\mathcal{L} \rightarrow (\ell, \ell)}^\sigma = \max_k \{L(k, -k)_{\rightarrow (\ell, \ell)} + h^0(k, -k)\}$  with  $h^0$  as in (2.6), and consider the rescaled LPP time

$$L_\ell^{\sigma, \text{resc}} = \frac{L_{\mathcal{L} \rightarrow (\ell, \ell)}^\sigma - 4\ell}{2^{4/3} \ell^{1/3}}. \tag{C.1}$$

Then, there exists constants  $s_0, \ell_0, C, c$  such that:

(a) Bound on upper tail:

$$\mathbb{P}(L_\ell^{\sigma, \text{resc}} \geq s) \leq C e^{-cs} \tag{C.2}$$

for all  $\ell \geq \ell_0$  and  $s \geq s_0$ .

(b) Tail on lower tail:

$$\mathbb{P}(L_\ell^{\sigma, \text{resc}} \leq s) \leq C e^{-c|s|^{3/2}} \tag{C.3}$$

for all  $\ell \geq \ell_0$  and  $s \leq -s_0$ .

*Proof* (a) Define  $J(u) = u(2\ell)^{2/3}(1, -1)$  and  $W_\ell(u) = h^0(J(u))/(2^{4/3} \ell^{1/3})$ . By Donsker’s theorem,  $u \mapsto W_\ell(u)$  converges weakly to a two-sided Brownian motion with diffusion coefficient  $2\sigma^2$ . Further, define

$$L_\ell^{\text{pp}}(u) := \frac{L_{J(u) \rightarrow (\ell, \ell)} - 4\ell}{2^{4/3} \ell^{1/3}}. \tag{C.4}$$

Then, we can write

$$L_\ell^{\sigma, \text{resc}} = \max_u \{L_\ell^{\text{pp}}(u) + W_\ell(u)\} \leq \max_u \{L_\ell^{\text{pp}}(u) + u^2/2\} + \max_u \{W_\ell(u) - u^2/2\}. \tag{C.5}$$

Thus,

$$\mathbb{P}(L_\ell^{\sigma, \text{resc}} \geq s) \leq \mathbb{P}(\max_u \{L_\ell^{\text{pp}}(u) + u^2/2\} \geq s/2) + \mathbb{P}(\max_u \{W_\ell(u) - u^2/2\} \geq s/2). \tag{C.6}$$

By computations based on Doob maximal inequality (used for instance in (4.36)), one obtains  $\mathbb{P}(\max_u \{W_\ell(u) - u^2/2\} \geq s/2) \leq C e^{-cs^2}$  for some constants  $C, c > 0$ . To bound the first term without new estimates, remark that for any  $M$  we can bound

$$\begin{aligned} \mathbb{P}(\max_u \{L_\ell^{\text{pp}}(u) + u^2/2\} \geq s/2) &\leq \mathbb{P}(\max_u L_\ell^{\text{pp}}(u) \geq s/4 - M^2/2) \\ &\quad + \mathbb{P}(\max_{|u| > M} \{L_\ell^{\text{pp}}(u) + u^2/2\} \geq s/4) \end{aligned} \tag{C.7}$$

The exponential decay in  $s$  for the second term is just a special case of (4.19) (set  $\tau = 0$ ) and it holds for all  $M \geq M_0$ , for some finite  $M_0$ . We fix  $M = M_0$  and then, using the fact that  $\max_u L_\ell^{\text{pp}}(u) = L_\ell^{\mathcal{L}, \text{resc}}$ , by (B.3) we have exponential decay in  $s$  for the first term as well.

(b) It follows from (A.4) since  $\mathbb{P}(L_{\mathcal{L}}^\sigma \rightarrow (\ell, \ell) \leq x) \leq \mathbb{P}(L_{(0,0) \rightarrow (\ell, \ell)} \leq x)$ . □

### Appendix D: Bounds on Stationary LPP

We now state and give a short proof of the tails of the one-point distribution in the stationary case with  $\rho = 1/2$  of the LPP to  $(\ell, \ell)$ .

**Proposition D.1** *Let  $\rho = 1/2$ . Then there exists constants  $s_0, \ell_0, C, c$  such that:*

(a) *Bound on upper tail:*

$$\mathbb{P}(L^{\mathcal{B}}_{(0,0) \rightarrow (\ell, \ell)} \geq 4\ell + 2^{4/3}s\ell^{1/3}) \leq Ce^{-cs} \tag{D.1}$$

for all  $\ell \geq \ell_0$  and  $s \geq s_0$ .

(b) *Bound on lower tail:*

$$\mathbb{P}(L^{\mathcal{B}}_{(0,0) \rightarrow (\ell, \ell)} \leq 4\ell + 2^{4/3}s\ell^{1/3}) \leq Ce^{-c|s|^{3/2}} \tag{D.2}$$

for all  $\ell \geq \ell_0$  and  $s \leq -s_0$ .

*Proof* (a) One can write  $L^{\mathcal{B}}_{(0,0) \rightarrow (\ell, \ell)} = \max\{L^{1,\rho}(\ell, \ell), L^{-,\rho}(\ell, \ell)\}$ , where  $L^{1,\rho}(\ell, \ell)$  (resp.  $L^{-,\rho}(\ell, \ell)$ ) are the LPP with one-sided perturbation only on  $i = 0$  (resp.  $j = 0$ ). Then,

$$\mathbb{P}(L^{\mathcal{B}}_{(0,0) \rightarrow (\ell, \ell)} \geq x) \leq \mathbb{P}(L^{1,\rho}(\ell, \ell) \geq x) + \mathbb{P}(L^{-,\rho}(\ell, \ell) \geq x). \tag{D.3}$$

By choosing  $x = 4\ell + s2^{4/3}\ell^{1/3}$ , Lemma 3.3 of [27] (based on the estimates on the correlation kernel in [1]) gives exponential decay in  $s$  for all  $s \geq s_0$ .

(b) It follows from (A.4), since  $\mathbb{P}(L^{\mathcal{B}}_{(0,0) \rightarrow (\ell, \ell)} \leq x) \leq \mathbb{P}(L_{(0,0) \rightarrow (\ell, \ell)} \leq x)$ . □

**Lemma D.2** *Let  $\rho = 1/2$  and define  $I(u) = (\ell - 2u\ell^{2/3}, \ell + 2u\ell^{2/3})$ . Then, for any  $\alpha > 0$ , we have*

$$\mathbb{P}(|L^{\mathcal{B}}_{(0,0) \rightarrow I(K)} - L^{\mathcal{B}}_{(0,0) \rightarrow I(0)}| \geq \alpha\ell^{1/3}) \leq 4e^{-\alpha^2/(16K)} \tag{D.4}$$

for all  $\ell$  large enough. Furthermore,

$$\mathbb{P}(\max_{u \geq K} L^{\mathcal{B}}_{(0,0) \rightarrow I(u)} - L^{\mathcal{B}}_{(0,0) \rightarrow I(K)} - \beta u^2\ell^{1/3} \geq \alpha\ell^{1/3}) \leq Ce^{-\frac{(\alpha + \beta K^2)^2}{16K}}, \tag{D.5}$$

for a constant  $C$  and for all  $\beta > 0$  and  $\alpha > -\beta K^2$  and  $\ell$  large enough.

*Proof* The process  $K \mapsto Y_K := L^{\mathcal{B}}_{(0,0) \rightarrow I(K)} - L^{\mathcal{B}}_{(0,0) \rightarrow I(0)}$  is a martingale [7] given by a sum of i.i.d.zero mean random variables  $Z_j - 2$ , with  $Z_j \sim \text{Exp}(1/2)$ . By the exponential Chebyshev inequality,

$$\begin{aligned} \mathbb{P}(|Y_K| \geq \alpha \ell^{1/3}) &\leq \mathbb{P}(Y_K \geq \alpha \ell^{1/3}) + \mathbb{P}(-Y_K \geq \alpha \ell^{1/3}) \\ &\leq \inf_{t \geq 0} e^{-t\alpha \ell^{1/3}} \mathbb{E}(e^{t(Z_1-2)})^{2K\ell^{2/3}} + \inf_{t' \geq 0} e^{-t'\alpha \ell^{1/3}} \mathbb{E}(e^{-t'(Z_1-2)})^{2K\ell^{2/3}}. \end{aligned} \tag{C.6}$$

Using  $\mathbb{E}(e^{t(Z_1-2)}) = \frac{e^{-2t}}{1-2t}$  for  $t \in (0, 1/2)$  and  $\mathbb{E}(e^{-t'(Z_1-2)}) = \frac{e^{2t'}}{1+2t'}$  for all  $t' \geq 0$ , after the minimization we obtain

$$\mathbb{P}(|Y_K| \geq \alpha \ell^{1/3}) \leq 2e^{-\alpha^2/(16K)(1+\mathcal{O}(\alpha K^{-1}\ell^{-1/3}))} \leq 4e^{-\alpha^2/(16K)} \tag{C.7}$$

for all  $\ell$  large enough.

For the second estimate, from the inequality

$$\begin{aligned} &\mathbb{P}(\max_{u \geq K} L^{\mathcal{B}}_{(0,0) \rightarrow I(u)} - L^{\mathcal{B}}_{(0,0) \rightarrow I(K)} - \beta u^2 \ell^{1/3} \geq \alpha \ell^{1/3}) \\ &\leq \sum_{m \geq 1} \mathbb{P}(\max_{u \in [Km, K(m+1)]} L^{\mathcal{B}}_{(0,0) \rightarrow I(u)} - L^{\mathcal{B}}_{(0,0) \rightarrow I(K)} \geq (\alpha + \beta K^2 m^2) \ell^{1/3}) \\ &\leq \sum_{m \geq 1} \inf_{t > 0} e^{-t(\alpha + \beta K^2 m^2) \ell^{1/3}} \mathbb{E}(e^{t(Z_1-2)})^{2Km\ell^{2/3}}. \end{aligned} \tag{C.8}$$

Maximising over  $t$  and taking the sum we finally get<sup>10</sup>

$$\mathbb{P}\left(\max_{u \geq K} L^{\mathcal{B}}_{(0,0) \rightarrow I(u)} - L^{\mathcal{B}}_{(0,0) \rightarrow I(-K)} - \beta u^2 \ell^{1/3} \geq \alpha \ell^{1/3}\right) \leq C e^{-\frac{(\alpha + \beta K^2)^2}{16K}} \tag{C.9}$$

for a constant  $C$  and for all  $\beta > 0$  and  $\alpha > -\beta K^2$  and  $\ell$  large enough. □

### Appendix E: Bounds for point-to-half line LPP

**Proposition E.1** *Let  $I(u) = (\tau N, \tau N) + u(2N)^{2/3}(1, -1)$ . Then,*

$$\mathbb{P}\left(\max_{|u| > M} L_{I(u) \rightarrow (N, N)} > 4(1 - \tau)N + 2^{4/3}(s - \gamma M^2)N^{1/3}\right) \leq C e^{-cM^2(1-\tau)^{-4/3}} e^{-\tilde{c}s(1-\tau)^{-1/3}} \tag{E.1}$$

for some constants  $C, c, \tilde{c} > 0$ , which can be taken uniform in  $N$  and uniform for  $\gamma$  in a compact subset of  $(0, 1/(1 - \tau))$ .

*Proof* By symmetry, it is enough to get the bound on the distribution of  $\max_{u < -M} L_{I(u) \rightarrow (N, N)}$ . By first shifting  $I(-M)$  to the origin, and then using the

---

<sup>10</sup>To be precise, for  $\varepsilon > 0$  small, one can bound  $\mathbb{P}(L^{\mathcal{B}}_{(0,0) \rightarrow I(u)} - L^{\mathcal{B}}_{(0,0) \rightarrow I(K)} \geq (\alpha + \beta K^2 u^2) \ell^{1/3})$  for all  $u \geq \varepsilon K \ell^{1/3}$  using (C.8) and for  $m \in \{1, \dots, \varepsilon \ell^{1/3}\}$  we can minimize over  $t$  and compute the series expansion in the exponent for large  $\ell$ .

mapping between LPP and TASEP, the distribution function is the same as the distribution of TASEP particle number  $n = t/4 + \tilde{\tau}(t/2)^{2/3}$  at time  $t = 4(1 - \tau)N + 2^{4/3}N^{1/3}(s - \gamma M^2)$ , starting at  $x_k(0) = -2k, k \geq 0$ .

From Proposition 3 of [12] we have an explicit expression in terms of Fredholm determinant. The upper tail estimate is standard. Using Hadamard's bound it is enough to have a bound on the correlation kernel. In Section 4 of [12] exponential decay of the rescaled correlation kernel has been proven. Then, simple algebraic computations give the claimed result.  $\square$

## References

- Baik, J., Ben Arous, G., P ech e, S.: Phase transition of the largest eigenvalue for non-null complex sample covariance matrices. *Ann. Probab.* **33**, 1643–1697 (2006)
- Baik, J., Ferrari, P.L., P ech e, S.: Limit process of stationary TASEP near the characteristic line. *Comm. Pure Appl. Math.* **63**, 1017–1070 (2010)
- Baik, J., Ferrari, P.L., P ech e, S.: Convergence of the two-point function of the stationary TASEP, *Singular Phenomena and Scaling in Mathematical Models*, Springer, pp. 91–110 (2014)
- Baik, J., Liu, Z.: Multi-point distribution of periodic TASEP. arXiv:1710.03284 (2017)
- Baik, J., Liu, Z.: TASEP on a ring in sub-relaxation time scale. *Comm. Pure Appl. Math.* **71**, 747–813 (2018)
- Baik, J., Rains, E.M.: Limiting distributions for a polynuclear growth model with external sources. *J. Stat. Phys.* **100**, 523–542 (2000)
- Bal azs, M., Cator, E., Sepp al ainen, T.: Cube root fluctuations for the corner growth model associated to the exclusion process. *Electron J. Probab.* **11**, 1094–1132 (2006)
- Basu, R., Ganguly, S.: Time correlation exponents in last passage percolation. arXiv:1807.09260 (2018)
- Basu, R., Sidoravicius, V., Sly, A.: Last passage percolation with a defect line and the solution of the Slow Bond Problem. arXiv:1408.346 (2014)
- Borodin, A., Ferrari, P.L.: Large time asymptotics of growth models on space-like paths I: PushASEP. *Electron J. Probab.* **13**, 1380–1418 (2008)
- Borodin, A., Ferrari, P.L., Pr ahofer, M., Sasamoto, T.: Fluctuation properties of the TASEP with periodic initial configuration. *J. Stat. Phys.* **129**, 1055–1080 (2007)
- Borodin, A., Ferrari, P.L., Sasamoto, T.: Transition between  $\text{Airy}_1$  and  $\text{Airy}_2$  processes and TASEP fluctuations. *Comm. Pure Appl. Math.* **61**, 1603–1629 (2008)
- Borodin, A., Gorin, V.: Lectures on integrable probability. arXiv:1212.3351 (2012)
- Borodin, A., P ech e, S.: Airy kernel with two sets of parameters in directed percolation and random matrix theory. *J. Stat. Phys.* **132**, 275–290 (2008)
- Cator, E., Pimentel, L.: On the local fluctuations of last-passage percolation models. *Stoch. Proc. Appl.* **125**, 879–903 (2015)
- Chhita, S., Ferrari, P.L., Spohn, H.: Limit distributions for KPZ growth models with spatially homogeneous random initial conditions. *Ann. Appl. Probab.* **28**, 1573–1603 (2018)
- Corwin, I.: The Kardar-Parisi-Zhang equation and universality class. *Random Matrices: Theory Appl.* **01**, 1130001 (2012)
- Corwin, I., Ferrari, P.L., P ech e, S.: Limit processes for TASEP with shocks and rarefaction fans. *J. Stat. Phys.* **140**, 232–267 (2010)
- Corwin, I., Ferrari, P.L., P ech e, S.: Universality of slow decorrelation in KPZ models. *Ann. Inst. H. Poincar e, Probab. Statist.* **48**, 134–150 (2012)
- Corwin, I., Liu, Z., Wang, D.: Fluctuations of TASEP and LPP with general initial data. *Ann. Appl. Probab.* **26**, 2030–2082 (2016)
- Ferrari, P.A., Fontes, L.: Shock fluctuations in the asymmetric simple exclusion process. *Probab. Theory Relat. Fields* **99**, 305–319 (1994)
- Ferrari, P.L.: Slow decorrelations in KPZ growth. *J. Stat. Mech.*, P07022 (2008)
- Ferrari, P.L.: From interacting particle systems to random matrices. *J. Stat. Mech.*, P10016 (2010)



24. Ferrari, P.L., Ghosal, P., Nejjar, P.: Limit law of a second class particle in TASEP with non-random initial condition. arXiv:[1710.02323](#) (2017)
25. Ferrari, P.L., Nejjar, P.: Anomalous shock fluctuations in TASEP and last passage percolation models. *Probab. Theory Relat. Fields* **161**, 61–109 (2015)
26. Ferrari, P.L., Nejjar, P.: Fluctuations of the competition interface in presence of shocks. *ALEA Lat. Am. J. Probab. Math. Stat.* **14**, 299–325 (2017)
27. Ferrari, P.L., Occelli, A.: Universality of the GOE Tracy-Widom distribution for TASEP with arbitrary particle density. *Electron J. Probab.* **23**(51), 1–14 (2018)
28. Ferrari, P.L., Spohn, H.: Random Growth Models, The Oxford handbook of random matrix theory. In: Akemann, G., Baik, J., Di Francesco, P. (eds.), pp. 782–801. Oxford Univ. Press, Oxford (2011)
29. Ferrari, P.L., Spohn, H.: On time correlations for KPZ growth in one dimension. *SIGMA* **12**, 074 (2016)
30. Johansson, K.: Shape fluctuations and random matrices. *Comm. Math. Phys.* **209**, 437–476 (2000)
31. Johansson, K.: Discrete polynuclear growth and determinantal processes. *Comm. Math. Phys.* **242**, 277–329 (2003)
32. Johansson, K.: Two time distribution in Brownian directed percolation. *Comm. Math. Phys.* **351**, 441–492 (2017)
33. Johansson, K.: The two-time distribution in geometric last-passage percolation. arXiv:[1802.00729](#) (2018)
34. Kardar, M., Parisi, G., Zhang, Y.Z.: Dynamic scaling of growing interfaces. *Phys. Rev. Lett.* **56**, 889–892 (1986)
35. Matetski, K., Quastel, J., Remenik, D.: The KPZ fixed point. arXiv:[1701.00018](#) (2017)
36. De Nardis, J., Le Doussal, P.: Tail of the two-time height distribution for KPZ growth in one dimension, *J. Stat. Mech* 053212 (2017)
37. De Nardis, J., Le Doussal, P.: Two-time height distribution for 1D KPZ growth: the recent exact result and its tail via replica, *J. Stat. Mech* 093203 (2018)
38. De Nardis, J., Le Doussal, P., Takeuchi, K.A.: Memory and universality in interface growth. *Phys. Rev. Lett.* **118**, 125701 (2017)
39. Nejjar, P.: Transition to shocks in TASEP and decoupling of last passage times. arXiv:[1705.08836](#) (2017)
40. Pimentel, L.P.R.: Ergodicity of the KPZ fixed point. arXiv:[1708.06006](#) (2017)
41. Pimentel, L.P.R.: Local behavior of airy processes. *J. Stat. Phys.* **173**, 1614–1638 (2018)
42. Prähofer, M., Spohn, H.: Current fluctuations for the totally asymmetric simple exclusion process In and out of equilibrium. In: Sidoravicius, V. (ed.) *Progress in Probability*, Birkhäuser (2002)
43. Prähofer, M., Spohn, H.: Scale invariance of the PNG droplet and the Airy process. *J. Stat. Phys.* **108**, 1071–1106 (2002)
44. Quastel, J.: Introduction to KPZ. *Curr. Dev. Math.*, 125–194 (2011)
45. Quastel, J., Remenik, D.: Airy processes and variational problems, Topics in percolative and disordered systems. In: Ramírez, A., Ben Arous, G., Ferrari, P., Newman, C., Sidoravicius, V., Vares, M. (eds.) Springer (2014)
46. Quastel, J., Spohn, H.: The one-dimensional KPZ equation and its universality class. *J. Stat. Phys.* **160**, 965–984 (2015)
47. Rost, H.: Non-equilibrium behavior of a many particle system: density profile and local equilibrium. *Z. Wahrsch. Verw. Gebiete* **58**, 41–53 (1981)
48. Sasamoto, T.: Spatial correlations of the 1D KPZ surface on a flat substrate. *J. Phys. A* **38**, L549–L556 (2005)
49. Takeuchi, K.A.: Statistics of circular interface fluctuations in an off-lattice Eden model. *J. Stat. Mech.*, P05007 (2012)
50. Takeuchi, K.A.: Crossover from growing to stationary interfaces in the Kardar-Parisi-Zhang class. *Phys. Rev. Lett.* **110**, 210604 (2013)
51. Takeuchi, K.A.: An appetizer to modern developments on the Kardar–Parisi–Zhang universality class. *Physica A* **504**, 77–105 (2016)
52. Takeuchi, K.A.: Private communication (2018)
53. Takeuchi, K.A., Akimoto, T.: Characteristic sign renewals of Kardar–Parisi–Zhang fluctuations. *J. Stat. Phys.* **164**, 1167–1182 (2016)
54. Takeuchi, K.A., Sano, M.: Evidence for geometry-dependent universal fluctuations of the Kardar-Parisi-Zhang interfaces in liquid-crystal turbulence. *J. Stat. Phys.* **147**, 853–890 (2012)