

Inflammatory responses of a human keratinocyte cell line to 10 nm citrate- and PEG-coated silver nanoparticles

V. Bastos · D. Brown · H. Johnston ·
A. L. Daniel-da-Silva · I. F. Duarte ·
C. Santos · H. Oliveira

Received: 15 April 2016 / Accepted: 12 July 2016 / Published online: 20 July 2016
© Springer Science+Business Media Dordrecht 2016

Abstract Silver nanoparticles (AgNPs) are among the most commonly used engineered NPs and various commercially available products are designed to come in direct contact with the skin (wound dressings, textiles, creams, among others). Currently, there is limited understanding of the influence of coatings on the toxicity of AgNPs and in particular their ability to impact on AgNP's mediated inflammatory responses. As AgNPs are often stabilized by different coatings, including citrate and polyethyleneglycol (PEG), in this study we investigate the influence of citrate (Cit10) or PEG (PEG10) coatings to 10 nm AgNP on skin, using human HaCaT keratinocytes. AgNPs cytotoxicity and inflammatory response (nuclear factor (NF)- κ B induction and cytokine production) of HaCaT were assessed

after in vitro exposure to 10 and 40 μ g/mL after 4, 24, and 48 h. Results showed that although both types of coated AgNPs decreased cell proliferation and viability, Cit10 AgNPs were more toxic. NF- κ B inhibition was observed for the highest concentration (40 μ g/mL) of PEG10 AgNPs, and the putative link to early apoptotic pathways observed in these cells is discussed. No production of IL-1 β , IL-6, IL-10, and TNF α was stimulated by AgNPs. Furthermore, Cit10 and PEG10 AgNPs decreased the release of MCP-1 by HaCaT cells after 48 h of exposure. As cytokines are vital for the immunologic regulation in the human body, and it is demonstrated that they may interfere with NPs, more research is needed to understand how different AgNPs affect the immune system.

V. Bastos · C. Santos · H. Oliveira
CESAM & Laboratory of Biotechnology and Cytomics,
University of Aveiro, 3810-193 Aveiro, Portugal

D. Brown · H. Johnston
School of Life Sciences, Heriot-Watt University,
Riccarton, Edinburgh EH14 4AS, UK

A. L. Daniel-da-Silva · I. F. Duarte
Department of Chemistry, CICECO – Aveiro Institute of
Materials, University of Aveiro, 3810-193 Aveiro,
Portugal

C. Santos (✉)
Department of Biology & Greenup/Citab -UP, Faculty of
Sciences, University of Porto, Rua do Campo, Alegre,
Porto, Portugal
e-mail: csantos@fc.up.pt

Keywords Citrate-AgNPs · PEG-AgNPs · HaCaT
cells · Inflammatory responses · Nanotoxicology ·
Skin · Environmental and health effects

Introduction

Nanotechnology-based consumer products are exponentially increasing, being nanosilver-containing products among the most commonly used (Vance et al. 2015). Silver nanoparticles (AgNPs) are widely used due to their enhanced physicochemical properties and biological activities such as their antimicrobial activity. Their applications range from medicine and industry to household and personal care products

(EPA 2010) or clothing (Abdelhalim and Jarrar 2011; Behra et al. 2013; Benn and Westerhoff 2008; Eckhardt et al. 2013; Nowack et al. 2011). The increased exploitation of AgNPs and consequent release into the environment raises concerns about their possible impacts on the environment and on human health (Nowack and Bucheli 2007). There is an array of AgNPs that are being exploited, which vary with respect to their physicochemical properties (e.g., size, shape, charge, surface coating, dispersion state) (Ahlberg et al. 2014; Boonkaew et al. 2014; Comfort et al. 2014; Kim et al. 2012; Park et al. 2011b). Existing studies have demonstrated that the physicochemical properties of AgNPs are able to influence their toxicity for different cell lines [e.g., human keratinocytes (HaCaT and primary keratinocytes), normal fibroblasts (NHF), rat adrenal pheochromocytoma (PC12), and mouse osteoblasts (MC3T3-E1), fibroblasts (L929), and macrophages (RAW 264.7)]. However, little attention has been given to the coating-dependent toxicity of AgNPs. Thus, research on the toxicity of AgNPs of varied physicochemical properties is critical in order to better predict the risks they pose.

It has been reported that nanoparticle coating, media composition, and ionic strength influence the surface chemistry, shape, aggregation state, and dissolution of AgNPs, which in turn can differently affect their cellular uptake and biological effects (Tejamaya et al. 2012). Indeed, few studies addressing the uptake (by embryonic fibroblasts NIH/3T3, keratinocytes HaCaT, and hepatoma cells Hepa-1c1c7, respectively) of different coated AgNPs and their influence on cytotoxicity have been reported (Caballero-Díaz et al. 2013; Lu et al. 2010; Pang et al. 2016). Citrate is the most commonly used reducing and stabilizing agent of AgNPs, rendering NPs with a negative surface charge and providing colloidal stability through electrostatic repulsions (Sharma et al. 2009). Among other coating agents of AgNPs, low molecular weight polyethyleneglycol (PEG), which stabilizes AgNPs through steric interactions, has been increasingly used in biomedical applications as it enhances biocompatibility and increases blood circulation time (Ginn et al. 2014; Ryan et al. 2008).

Assessment of the ability of NPs to induce inflammatory responses is commonly used as an indicator of toxicity. For example, Chalew and

Schwab (2013) studied the inflammatory effects of AgNPs, titanium dioxide (TiO₂NPs), and zinc oxide (ZnONPs) (0, 0.1, 1, 10, and 100 mg/L) on human intestinal Caco-2 and SW480 cells and found that all NPs increased IL-8 cytokine generation in both cell lines. Also, Park et al. (2011a) observed that proinflammatory cytokines (IL-1, TNF- α , and IL-6) and Th0 cytokine (IL-2) were progressively increased by day 28 after a single intratracheal instillation of AgNPs in mice. Suliman et al. (2013) found that 50 μ g/mL AgNPs exposure to human lung epithelial (A549) cells significantly increased the level of proinflammatory cytokines, namely interleukin-1 β (IL-1 β) and interleukin-6 (IL-6). Yang and collaborators also observed IL-1 β release by human blood monocytes in response to AgNPs (Yang et al. 2012). However, Wong et al. (2009) found an antiinflammatory effect of AgNPs to two mouse macrophage cell lines, RAW264.7 and J774.1, where AgNPs blocked TNF- α production. On the other hand, we could not find studies reporting the induction of antiinflammatory cytokines after exposure to NPs (Murray et al. 2013; Orłowski et al. 2013; Samberg et al. 2010). Cytokines can strongly activate inflammatory responses and cell death in various tissues, including the skin (Fujiwara and Kobayashi 2005; Graves et al. 2004). Indeed, a study on the effects of UVB radiation using HaCaT cells reported an increase of various proinflammatory cytokines—interleukin (IL)-1 β , IL-6, IL-8, interferon (IFN)- γ , granulocyte-colony stimulating factor (G-CSF), macrophage inflammatory protein (MIP)-1 β , and tumor necrosis factor (TNF)- α (Yoshizumi et al. 2008). Murray et al. (2013) found increased IL-8 and IL-6 in human epidermal keratinocytes (HEK cells) after exposure to superparamagnetic iron oxide (SPION) NPs (2.6, 5.2, 13, and 26 μ g/cm² for 24 h). In other study using HEK cells, quantum dot NPs significantly increase IL-6 at 1.25 to 10 nM, while IL-8 increased from 2.5 to 10 nM after 24 and 48 h (Zhang et al. 2008). Therefore, as products containing AgNPs can be applied to the skin (e.g., wound dressing), and as there are experimental evidences for skin penetration of 25 \pm 7 nm AgNPs (also in intact skin) (Larese et al. 2009) and 20–40 nm AgNPs (George et al. 2014), the human keratinocyte cell line HaCaT was selected as an *in vitro* model in this study. It is well known that cytokines play crucial roles in immunologic

regulation in the human body and are involved in the induction of proliferation, differentiation, and cell death in many cell types (Yarilin and Belyakov 2004). Moreover, activation of the transcription factor nuclear factor kappa B (NF- κ B) has been shown to play a central role in the enhanced expression and regulation of cytokine genes (Kelso 1998). There is also evidence that carbon NPs can activate NF- κ B in macrophages which stimulates TNF α production (Brown et al. 2004). To our knowledge, the activation of NF- κ B in keratinocytes has not been studied previously.

In our previous study (Bastos et al. 2016) we evaluated the toxicity of 30 nm AgNPs coated with citrate or PEG on HaCaT cells. Our results showed that Cit30 AgNPs were more cytotoxic than PEG30 AgNPs. Concerning cytokine release, both Cit30 and PEG30 AgNPs induced a decrease in MCP-1 production but no effect on other cytokines, namely IL-1 β , IL-6, IL-10, and TNF- α (Bastos et al. 2016).

In this study, we aimed to compare the inflammatory responses of HaCaT cells exposed to well-characterized 10 nm AgNPs coated with citrate or PEG, in order to explore the influence of smaller sizes of AgNPs on the inflammatory response. In particular, the effects on viability, expression of the proinflammatory transcription factor NF- κ B, and production of cytokines such as interleukin-1 beta (IL-1 β), IL-6, tumor necrosis factor-alpha (TNF- α), IL-10, and monocyte chemoattractant protein-1 (MCP-1) were assessed.

Materials and methods

Chemicals

Sterile, purified, and endotoxin-free silver nanoparticles (Biopure AgNPs 1.0 mg/mL in water), with a diameter of 10 nm and a citrate or polyethyleneglycol (PEG) surface, designated as Cit10 and PEG10, respectively, were purchased from Nanocomposix Europe (Prague, Czech Republic). Dulbecco's modified Eagle's medium (DMEM), fetal bovine serum (FBS), antibiotics, and phosphate buffer saline (PBS, pH 7.4) were purchased from Life Technologies (Carlsbad, CA, USA). 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT), Mowiol, and DAPI were obtained from Sigma-Aldrich (St. Louis, MO, USA).

Physicochemical characterization of AgNPs

The morphology and size of AgNPs were assessed by transmission electron microscopy (TEM) using a transmission electron microscope Hitachi H9000 NAR (Hitachi High-Technologies Europe GmbH, Germany) operating at 300 kV. Samples for TEM analysis were prepared by evaporating dilute suspensions of AgNPs on a copper grid coated with an amorphous carbon film. The hydrodynamic diameter and polydispersity index (PdI) were measured by dynamic light scattering (DLS) and the zeta potential was assessed by electrophoretic mobility, both measurements using a Zetasizer Nano ZS (Malvern Instruments, UK). Silver quantification measurements were performed by inductively coupled plasma optical emission spectrometry (ICP-OES) in an Activa M Radial spectrometer (Horiba Jobin-Yvon), employing a charge coupled device (CCD) array detector, with a wavelength range of 166–847 nm and radial plasma view. Samples for ICP-OES were prepared by the addition of 10 μ L AgNPs (1.0 mg/mL) to 990 μ L of either ultrapure water or complete culture medium, incubated for 0, 4, 24, or 48 h, then centrifuged at 40,000 rcf for 120 min at 4 °C (in accordance with the manufacturer's recommendations) to deposit the nanoparticles and separate the supernatant, which was then digested with acid (HCl:HNO₃ 2:1 v/v) before ICP-OES analysis.

Cell culture

The HaCaT cell line, a nontumorigenic immortalized human keratinocyte cell line (Boukamp et al. 1988), was obtained from Cell Lines Services (Eppelheim, Germany). Cells were grown in complete medium, i.e., Dulbecco's modified Eagle's medium, supplemented with 10 % fetal bovine serum (FBS), 2 mM L-glutamine, 100 U/mL penicillin, 100 μ g/mL streptomycin, and 250 μ g/mL fungizone at 37 °C in 5 % CO₂ humidified atmosphere. Cells were observed daily under an inverted phase-contrast Eclipse TS100 microscope (Nikon, Tokyo, Japan). For each experiment, cells were allowed to adhere for 24 h and then exposed to Cit10 or PEG10 AgNPs (dispersed through vortex in cell culture medium). For the assays cells were in passage number 45–50. Depending on the experiment, the silver ion and the coating agent per se,

dissolved in complete medium, were used as controls. The effects were measured after 4, 24, and 48 h.

Viability assay

Cell viability was determined by the colorimetric 3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyltetrazolium bromide (MTT) assay, measuring intracellular reduction of tetrazolium salts into purple formazan by viable cells (Twentyman and Luscombe 1987). Cells were seeded in 96-well plates at a concentration of 6×10^4 cells/mL. Fifty microliters of MTT (1 mg/mL) in phosphate buffered saline (PBS) were then added to each well, and incubated for 4 h at 37 °C, 5 % CO₂. Medium was then removed and 150 µL of DMSO was added to each well for solubilization of formazan crystals. The optical density of reduced MTT was measured at 570 nm in a microtiter plate reader (Synergy HT Multi-Mode, BioTeK, Winooski, VT), and cell viability was calculated as $[(\text{Sample Abs} - \text{MSO Abs})/(\text{Control Abs} - \text{DMSO Abs})] \times 100$. Three independent assays were performed with at least 2 technical replicates each and the results compared with the control (no exposure). From our previous MTT results (Bastos et al. 2016), the IC₅₀ for 30 nm citrate-coated AgNPs (the most cytotoxic) was 40 and 37.4 mg/mL at 24 and 48 h, respectively. Therefore, the concentrations of AgNPs corresponding to IC₅₀ and IC₂₀ (40 and 10 mg/mL, respectively) were selected for 10 nm AgNPs assays in order to enable comparisons between sizes.

Immunofluorescence of p65 subunit of NF-κB in human keratinocytes

After the 4 h treatments with 10 and 40 µg/mL of Cit10 or PEG 10AgNPs, coverslips were washed with PBS and permeabilized with 0.2 % Triton X-100 for 15 min followed by three washes with PBS. Cells were treated with PBS containing BSA at a concentration of 1 mg/mL as a blocking agent for 1 h. Cells were then washed three times with PBS and treated with antihuman NF-κB antibody (p65 subunit, Santa Cruz Biotechnology, Inc. Dallas, Texas USA) diluted 1/200 in PBS plus 0.5 % BSA for 1 h at room temperature. After three washes with PBS, coverslips were treated with a second antibody, Alexa fluor 488 antirabbit IgG diluted 1:200 in PBS plus 0.5 % BSA for 1 h at room temperature. After three washes with PBS, coverslips were treated with 0.5 µg/ml DAPI in

PBS plus 0.5 % BSA for 20 s, washed in PBS, and mounted on glass microscope slides using Mowiol. Cells were imaged using confocal microscopy.

Cytokine estimation using cytometric bead array

Cytokine production was assessed using Bioplex kits. Briefly, the supernatants (collected from cell viability studies, centrifuged and frozen at -80 °C) were used to estimate the release of the following cytokines from treated cells: interleukin-1 beta (IL-1β), IL-6, tumor necrosis factor-alpha (TNF-α), IL-10, and monocyte chemoattractant protein-1 (MCP-1). Bead array kits were obtained from Beckton Dickinson (Oxford, UK) and a master mix prepared according to the manufacturer's instructions. The master mix was incubated with each of the test supernatants for 1 h, followed by the addition of detection beads and incubated for a further 2 h at room temperature. The beads were then washed in wash buffer and analyzed using a BD FACSAarray™ flow cytometer which had previously been set up and calibrated using standard beads for each cytokine under investigation.

Statistical analysis

The results are reported as mean ± standard deviation (SD) of 2 technical replicates in each of the 3 independent experiments. For MTT assay, the statistical significance between control and exposed cells was performed by one-way ANOVA, followed by Dunnett and Dunn's method (as parametric and nonparametric test, respectively), using Sigma Plot 12.5 software (Systat Software Inc.). For the other assays, results were compared using two-way ANOVA, followed by Holm-Sidak test using also Sigma Plot 12.5 software (Systat Software Inc.). The differences were considered statistically significant for $p < 0.05$.

Results

Physicochemical characterization of AgNPs

A summary of the physicochemical properties of the NMs is provided in Table 1. The spherical shape and diameter of the AgNPs were verified by transmission electron microscopy, TEM (Fig. 1) and found to agree

Table 1 Physicochemical properties of AgNPs

	Cit10	PEG10
D (nm) ^a	9.3 ± 1.8	9.8 ± 2.0
D (nm) ^b	11.0 ± 2.4	9.4 ± 2.8
D_h (nm) ^c	25.1 ± 3.6	40.6 ± 3.8
PdI ^c	0.42–0.52	0.26–0.37
ζ (mV) ^d	−34.5 ± 3.3	−14.3 ± 1.0
λ_{\max}^e (nm)	389	396
%Ag+ in water 0 h ^f	0.93 ± 0.02	n.a.
%Ag+ in DMEM 0 h ^f	3.7 ± 0.05	9.1 ± 0.13
%Ag+ in DMEM 4 h ^f	14.3 ± 0.22	6.0 ± 0.12
%Ag+ in DMEM 24 h ^f	13.6 ± 0.35	11.2 ± 0.33
%Ag+ in DMEM 48 h ^f	12.9 ± 0.16	11.2 ± 0.34

Standard deviations calculated for D_h , ζ , and %Ag+ correspond to 3 replicate measurements

n.a. Not available

^a Diameter indicated by the manufacturer

^b Diameter measured by TEM

^c Hydrodynamic diameter (in cell culture medium) and polydispersity index (PdI) measured by DLS

^d Zeta potential assessed by electrophoretic mobility (in cell culture medium)

^e Wavelength of maximum absorbance peak in the UV–Vis spectrum

^f Percentage of ionic silver in the AgNP suspension (10 µg/mL)

with the manufacturer information (Table 1). The wavelength of the maximum absorbance peak in the UV–Vis spectra also matched the expected values.

Regarding the DLS assessment of hydrodynamic diameters (D_h), polydispersity indexes (PdI > 0.3) indicated large variability in particle size, especially for the Cit10 NPs, hence the Z-average sizes may lack accuracy. PEG10 NPs showed higher D_h than Cit10, as expected based on the larger size of PEG compared to citrate. The zeta potential values confirmed Cit10 AgNPs to have a negative surface charge (ζ −34 mV), which is expected as citrate is used as a coating to prevent agglomeration/aggregation through electrostatic repulsions, whereas PEG10 NPs, which are also designed to stabilize NPs through steric interactions, showed a less negative surface (ζ −14 mV). We have also assessed the amount of ionic silver (Ag^+) released from AgNPs, which was found to be low in water (<1 %), but significantly increased when the NPs were incubated in culture medium. Dissolved Ag^+ reached 14 % in Cit10 suspensions after 4 h and 11 % in PEG10 suspensions after 24 and 48 h, this lower value likely relating to a more efficient protection of PEG coating against NP surface oxidation.

Effects on cell growth and viability

HaCaT cells in control conditions (exposed to cell culture medium) showed typical morphology (Figs. 2a, 3a). When cells were exposed to Cit10 and PEG10 AgNPs for 24 h (Fig. 2b–e), their confluence decreased, especially at the highest concentration tested (40 µg/mL). The decrease in cell confluence was more visible after 48 h (Fig. 3b–e). Morphologically, exposed cells (to both Cit10 and PEG10 NPs)




Fig. 1 Transmission electron microscopy (TEM) images of Cit10 and PEG10 AgNPs used in this work




Fig. 2 Light microscopy images (100 \times) of HaCaT cells exposed to 10 nm citrate-AgNPs or PEG-AgNPs (Cit10 or PEG10) for 24 h. **a** 0 (control); **b** Cit10, 10 $\mu\text{g}/\text{mL}$; **c** Cit10, 40 $\mu\text{g}/\text{mL}$; **d** PEG10, 10 $\mu\text{g}/\text{mL}$; and **e** PEG10, 40 $\mu\text{g}/\text{mL}$. Bar corresponds to 100 μm




Fig. 3 Light microscopy images (100 \times) of HaCaT cells exposed to 10 nm citrate-AgNPs or PEG-AgNPs (Cit10 or PEG10) for 48 h. **a** 0 (control); **b** Cit10, 10 $\mu\text{g}/\text{mL}$; **c** Cit10, 40 $\mu\text{g}/\text{mL}$; **d** PEG10, 10 $\mu\text{g}/\text{mL}$; and **e** PEG10, 40 $\mu\text{g}/\text{mL}$. Bar corresponds to 100 μm

showed large precipitates/aggregates of AgNPs in the medium, and confluence appeared to be, on average, lower for Cit10 exposed cells.

The viability of HaCaT cells was negatively affected by both types of AgNP investigated in this study (Fig. 4). Relative to controls, the viability of




Fig. 4 Relative cell viability (%) of HaCaT cells after exposure to 10 and 40 µg/mL of 10 nm citrate-AgNPs or PEG-AgNPs (Cit10 or PEG10), measured by MTT assay, at 4, 24, and 48 h post exposure. Data expressed as mean and standard deviation ($n = 3$). Asterisk indicate significant differences between control at $p < 0.05$

exposed cells was significantly reduced ($p < 0.05$) upon exposure to Cit10 AgNPs at 10 and 40 µg/mL after 4, 24, and 48 h. Following PEG10 AgNP exposure, the viability of cells following exposure at a concentration of 10 µg/mL was not affected at 4 h but a significant reduction in cell viability was observed at 24 and 48 h at this concentration. At a concentration of 40 µg/mL, PEG10 NPs significantly decreased cell viability at all time points (4, 24, and 48 h).

NF-κB activation and inflammatory cytokine release

Activation of NF-κB in HaCaT cells by AgNPs was evaluated by immunofluorescence; in its inactive state NF-κB is located in the cytoplasm, and in its active state is localized in the nucleus. Figure 5a–c shows a positive control with 240 µM of H₂O₂ where there is a great intensity of p65 staining in the nucleus. Also, controls (including: cells only (no staining), cells stained with only the primary antibody and samples stained with the second antibody only) were done to check the autofluorescence in cells (data not shown). In control cells, most NF-κB staining was localized in the cell cytoplasm, with occasional occurrence in the nucleus. Regarding AgNP-exposed cells, there was no evidence of NF-κB activation (i.e., no increase in the intensity of staining in the nucleus) (Fig. 5 e–g). A decrease in p65 staining in the nucleus after exposure

to 40 µg/mL PEG10 AgNPs was observed compared to control cells (Fig. 5h). To confirm the decrease in p65 staining observed in Fig. 5h, a quantification of the nucleus fluorescence intensity of HaCaT microscopy images were done using the ImageJ software (Fig. 5i).

The release of cytokines by HaCaT cells treated with AgNPs is shown in Fig. 6. Lipopolysaccharide (LPS) stimulated a significant increase in MCP-1 release at 48 h, when compared to the control. MCP-1 production significantly decreased following exposure of HaCaT cells to both AgNP types and was most pronounced at a concentration of 40 µg/mL, compared to negative and positive controls ($p < 0.001$). No effects were observed on the other cytokines studied, IL-1β, IL-6, IL-10, and TNF-α, after exposure to both AgNPs at both times (data not shown).

Discussion

Citrate-coated AgNPs are among the most widely used AgNPs in multiple industrial applications (Tolaymat et al. 2010), while the less used PEG-coated AgNPs have gained increasing attention over recent years by, for instance, the biomedical industry due to their high stability and reduced reactivity (Brandenberger et al. 2010; Ginn et al. 2014; Ryan et al. 2008; Thorley and Tetley 2013). Therefore, there is a need to better understand the influence of coatings on the biological effects of NPs, and in particular the effects on the inflammatory responses. As skin represents one of the major organs in contact with AgNPs, we compared the viability and inflammatory responses induced by citrate- or PEG-coated AgNPs with a core diameter of 10 nm (Cit10 and PEG10 AgNPs) in human epidermal keratinocytes (HaCaT cells).

From the cytotoxicity results, Cit10 AgNPs were more toxic than PEG10 NPs. The higher toxicity of Cit10 AgNPs was particularly relevant for low doses, since a lower concentration (10 µg/mL) of Cit0 NPs induced a statistically significant decrease in cell viability 4 h post exposure, that was not evident for PEG-AgNPs. At higher doses (40 µg/mL) and exposure periods, PEG10 and Cit10 AgNPs reduced viability in similar ways. These data suggest that, for these skin cells, the influence of coating is more important at low AgNP concentrations, whereas by increasing concentration, the influence of coating




Fig. 5 Microscopy images of HaCaT cells. **a–c** are fluorescence microscopy images (400 \times) of immunofluorescence of HaCaT cells treated with anti NF- κ B p65 antibody after stimulation with 240 μ M H₂O₂ for 10 min prior fixation; **a** p65 subunit of NF- κ B (alexa fluor 488 antirabbit IgG); **b** DAPI staining the nucleus; **c** Overlap of a and b images. **d–h** are confocal microscopy images of immunofluorescence of p65 subunits of NF- κ B in human HaCaT keratinocytes exposed

to AgNPs for 4 h: **d** 0 (control); **e** 10 μ g/mL of Cit10; **f** 40 μ g/mL of Cit10; **g** 10 μ g/mL of PEG10; **h** 40 μ g/mL of PEG10. Bar corresponds to 100 μ m. **i** is the quantification of the nucleus fluorescence intensity of HaCaT microscopy images presented in Fig. 4h using the ImageJ software. Data expressed the mean and standard deviation. *Asterisks* indicate significant differences between control at $p < 0.01$

seems to be less relevant. A significant decrease in BEAS-2B (bronchial epithelial) cell viability upon exposure to 20 nm citrate-coated AgNPs at 6.25–50 μ g/mL after 24 h has been observed previously (Wang et al. 2014). Also, Song et al. (2012) showed a decrease in cell viability in human liver cell

line—HL-7702 after exposure to PEG-coated AgNPs in dose- and time-dependent manner at doses from 6.25 μ g/mL. Future studies could assess the sensitivity of different cell types to the AgNPs used in this study.

A complex interplay between environmental and genetic factors control immune system responses and




Fig. 6 Cytokine release by HaCaT cells after 24 and 48 h exposure to 10 nm citrate- or PEG- AgNPs (Cit10 or PEG10). Control+ is a positive control by adding LPS to cells. Data represent the mean \pm standard deviation ($n = 3$) of the concentration (pg/ml) of MCP-1 cytokine released from the cells after NPs treatment

when a deregulation of immune homeostasis occurs, host defense can be impaired and at the same time cause excessive and potential harmful inflammatory responses, which could be responsible for several immune disorders (Bieber 2008; Morar et al. 2006). The ability of NPs to elicit proinflammatory responses is frequently assessed in *in vitro* and *in vivo* studies as a marker of their toxicity [e.g., Schoemaker et al. (2002)]. Thus, understanding NP-dependent regulation of cytokine production is essential, since this process conditions shifts from acute to chronic phases of allergic inflammation (Rossi and Zlotnik 2000). NF- κ B pathways have been traditionally associated to increases in the production of inflammatory cytokines which could be implicated in the development of a variety of diseases (Driscoll et al. 1997; Mossman and Churg 1998). AgNPs did not activate NF- κ B in HaCaT cells in this study. In fact, NF- κ B may be inhibited after exposure to the higher concentration (40 μ g/mL) of PEG10 AgNPs. It is described in literature that the inhibition or absence of NF- κ B activation induces apoptosis or sensitizes cells to apoptosis (Schoemaker et al. 2002).

Murray et al. (2013) found that a coexposure of human epidermal keratinocytes (HEK cells) to superparamagnetic iron oxide (SPION) nanoparticles and UVB, induced NF- κ B activation and release of inflammatory mediators such as the cytokines IL-6

and IL-8. Carbon black NPs have also been demonstrated to induce NF- κ B activation in macrophages to stimulate TNF α production (Brown et al. 2004). However, it has also been recognized that NF- κ B signaling has important functions in the maintenance of physiological immune homeostasis, particularly in epithelial cells (Wullaert et al. 2011). In a previous work (Bastos et al. 2016), we have demonstrated, by Annexin-V/PI assay and expression of genes involved in apoptosis, that Cit30 AgNPs induced preferably late apoptotic/necrotic pathways, while cells exposed to PEG30 AgNPs stimulated increases of cells in earlier phase of apoptosis (increasing the expression of genes involved in apoptosis) and no apoptosis/necrosis, supporting that the coating of these AgNPs influence differently the cell apoptosis/necrosis pathways. A major role of NF- κ B pathways involve the regulation of antiapoptotic genes, by NF- κ B directly binding and inhibiting CASP3, -7, and -9 which seems to be happening in citrate-AgNPs exposed cells (Schoemaker et al. 2002). Considering that only PEG10 AgNPs inhibit NF- κ B, we therefore hypothesize that this may be involved in the induction of apoptosis by activating CASP3 found in PEG30-exposed cells versus Cit30-exposed ones. In the future we suggest that NF- κ B activation/inhibition may be used to a greater extent when assessing the hazard of coating and AgNPs to better understand the mechanisms (i.e., cellular and molecular events) underlying their toxicity. On the other hand, Brown et al. (2004) showed that ultrafine carbon black particles (UfCB) induced nuclear translocation of NF- κ B in human monocytes which occurred through ROS-mediated mechanism. Indeed, in our previous study 30 nm citrate- and PEG-AgNPs induced a significant increase in the production of ROS by HaCaT cells at the highest dose tested (40 μ g/mL), compared to control cells. However, the ROS levels were similar for both NP types which do not explain the NF- κ B inhibition by PEG10 AgNPs. Thus, for further studies we also suggest the quantification of NF- κ B activation (e.g., by western blotting) and determination of ROS production.

Concerning cytokine release, neither Cit10 nor PEG10 AgNPs induced IL-1 β , IL-6, IL-10, TNF- α , and MCP-1 production by HaCaT cells. Instead, compared to control cells, they decreased MCP-1 production after 48 h exposure, this reduction being more pronounced at the higher concentration (40 μ g/mL). MCP-1 production in control cells increased with

time in culture, as already described by Takahashi et al. (1995) who reported values close to ours for monocytes and endothelial cells in vitro. The information regarding the influence of AgNPs on the stimulation of these cytokines from keratinocytes release is scarce, and sometimes contradictory, as different exposure conditions, concentrations, coating, cell type, NP size, and synthesis have been considered in the literature (Chalew and Schwab 2013; Giovanni et al. 2015; Miethling-Graff et al. 2014; Orłowski et al. 2013; Samberg et al. 2010; Suliman et al. 2013; Wong et al. 2009; Yen et al. 2009). For instance, Orłowski et al. (2013) found an increase of MCP-1 production in murine keratinocytes (murine 291.03C) and by monocytes (RAW 264.7) after exposure to unmodified AgNPs. Also, human umbilical vein endothelial (HUVEC) cells showed a significant increase of IL-6, IL-8, and MCP-1 at doses higher than 1 mg/L AgNPs (Shi et al. 2014). Moreover, confirming the inflammatory potential of AgNPs, several interleukins and TNF- α were reported to increase upon exposure of HEK cells (Samberg et al. 2010) and macrophages (Yen et al. 2009) to AgNPs. On the other hand, several authors reported an undetectable stimulation of cytokines in response to metal NPs, as observed in the present study. For example, Murray et al. (2013) demonstrated that HEK cells exposed to superparamagnetic iron oxide nanoparticles maintained the IL-1 β , IL-10, and TNF- α below detectable levels (while increasing IL-6); and also that mouse epidermal cells (JB6 P+) maintained INF- γ and IL-12 below the detectable levels after exposure to the same nanoparticles. Similarly, Samberg et al. (2010) did not find detectable levels of IL-10 in HEK cells exposed to unwashed AgNPs. In mice peritoneal tissues and in RAW 264.7 cells, Wong et al. (2009) demonstrated that AgNPs have an antiinflammatory effect decreasing TNF- α , and INF- γ . Also, Parnsamut and Brimson (2015) found that AgNPs inhibited TNF- α expression in leukemic cell lines. It is known that cytokines can adsorb onto the surface of particles, which may compromise their detection (Brown et al. 2010). Thus, it should not be excluded that AgNPs may induce cytokine production by keratinocytes but that the cytokines bind to the AgNP surface to prevent their detection. How proteins bind to nanoparticles is currently an important topic of debate. For example, Deng et al. (2013) showed that human plasma proteins differently bind to positively and negatively charged

polymer-coated gold NPs, which elicited different biological responses, and that only the negatively charged nanoparticles induced cytokine release from THP-1 cells. While proteins can bind to different nanoparticles, the biological outcome may not be the same. Selection of cytokines for assessment in this study was prioritized based on the outcome of a literature search which identified cytokines that are commonly produced following exposure of cells to NPs. Future studies could therefore assess a wider panel of cytokines.

In summary, our study demonstrated that while citrate- and PEG-coated AgNPs decreased the viability of HaCaT cells. Citrate-coated AgNPs were more cytotoxic than PEG-coated NPs, particularly at low concentrations and shorter incubation times. At higher AgNPs concentration, the influence of coating became less relevant. Also, we demonstrated that, independent of the coating, AgNPs did not induce cytokine production, and decreased MCP-1 release. Finally, PEG10 AgNPs at high concentrations inactivated the transcription factor NF- κ B, and putative correlation with antiinflammatory and antiapoptotic homeostasis should be further explored.

Acknowledgments This work was developed in the scope of the projects CICECO-Aveiro Institute of Materials (Ref. FCT UID/CTM/50011/2013) and CESAM (Ref. FCT UID/AMB/50017/2013), financed by national funds through the FCT/MEC and when applicable cofinanced by the European Regional Development Fund (FEDER) under the PT2020 Partnership Agreement. Funding to the project FCOMP-01-0124-FEDER-021456 (Ref. FCT PTDC/SAU-TOX/120953/2010) by FEDER through COMPETE and by national funds through FCT, and the FCT-awarded grants (SFRH/BD/81792/2011; SFRH/BPD/111736/2015) are acknowledged. I.F.D and A.L.D.S. acknowledge FCT/MCTES for the research contracts under the Program ‘Investigador FCT’ 2014.

Compliance with ethical standards

Conflict of interest The authors declare that they have no competing interests.

References

- Abdelhalim MAK, Jarrar BM (2011) Renal tissue alterations were size-dependent with smaller ones induced more effects and related with time exposure of gold nanoparticles. *Lipids Health Dis* 10(1):1
- Ahlberg S, Meinke MC, Werner L, Epple M, Diendorf J, Blume-Peytavi U, Lademann J, Vogt A, Rancan F (2014) Comparison of silver nanoparticles stored under air or argon

- with respect to the induction of intracellular free radicals and toxic effects toward keratinocytes. *Eur J Pharm Biopharm* 88(3):651–657
- Bastos V, de Oliveira JF, Brown D, Jonhston H, Malheiro E, Daniel-da-Silva AL, Duarte IF, Santos C, Oliveira H (2016) The influence of Citrate or PEG coating on silver nanoparticle toxicity to a human keratinocyte cell line. *Toxicol Lett* 249:29–41
- Behra R, Sigg L, Clift M, Herzog F, Minghetti M, Johnston B, Petri-Fink A, Rothen-Rutishauser B (2013) Bioavailability of silver nanoparticles and ions: from a chemical and biochemical perspective. *J R Soc Interface* 10(87):20130396
- Benn T, Westerhoff P (2008) Nanoparticle silver released into water from commercially available sock fabrics. *Environ Sci Technol* 42(11):4133–4139
- Bieber T (2008) Atopic dermatitis. *N Engl J Med* 358(14):1483–1494
- Boonkaew B, Kempf M, Kimble R, Cuttle L (2014) Cytotoxicity testing of silver-containing burn treatments using primary and immortal skin cells. *Burns* 40(8):1562–1569
- Boukamp P, Petrussevska R, Breitkreutz D, Hornung J, Markham A, Fusenig N (1988) Normal keratinization in a spontaneously immortalized aneuploid human keratinocyte cell line. *J Cell Biol* 106:761–771
- Brandenberger C, Mühlfeld C, Ali Z, Lenz A-GG, Schmid O, Parak WJ, Gehr P, Rothen-Rutishauser B (2010) Quantitative evaluation of cellular uptake and trafficking of plain and polyethylene glycol-coated gold nanoparticles. *Small* 6(15):1669–1678
- Brown DM, Donaldson K, Borm PJ, Schins RP, Dehnhardt M, Gilmour P, Jimenez LA, Stone V (2004) Calcium and ROS-mediated activation of transcription factors and TNF- α cytokine gene expression in macrophages exposed to ultrafine particles. *Am J Physiol Lung Cell Mol Physiol* 286(2):L344–L353
- Brown DM, Dickson C, Duncan P, Al-Attili F, Stone V (2010) Interaction between nanoparticles and cytokine proteins: impact on protein and particle functionality. *Nanotechnology* 21(21):215104
- Caballero-Díaz E, Pfeiffer C, Kastl L, Gil P, Simonet B, Valcárcel M, Lamana J, Laborda F, Parak WJ (2013) The toxicity of silver nanoparticles depends on their uptake by cells and thus on their surface chemistry. *Part Part Syst Charact* 30(12):1079–1085
- Chalew TEA, Schwab KJ (2013) Toxicity of commercially available engineered nanoparticles to Caco-2 and SW480 human intestinal epithelial cells. *Cell Biol Toxicol* 29(2):101–116
- Comfort KK, Maurer EI, Hussain SM (2014) Slow release of ions from internalized silver nanoparticles modifies the epidermal growth factor signaling response. *Colloids Surf B* 123:136–142
- Deng ZJ, Liang M, Toth I, Monteiro M, Minchin RF (2013) Plasma protein binding of positively and negatively charged polymer-coated gold nanoparticles elicits different biological responses. *Nanotoxicology* 7(3):314–322
- Driscoll KE, Carter JM, Hassenbein DG (1997) Cytokines and particle-induced inflammatory cell recruitment. *Environ Health Perspect* 105(Suppl 5):1159
- Eckhardt S, Brunetto P, Gagnon J, Priebe M, Giese B, Fromm K (2013) Nanobio silver: its interactions with peptides and bacteria, and its uses in medicine. *Chem Rev* 113(7):4708–4754
- EPA (2010) State of the science literature review: everything nanosilver and more. Scientific, Technical, Research, Engineering and modeling support final report
- Fujiwara N, Kobayashi K (2005) Macrophages in inflammation. *Curr Drug Targets Inflamm Allergy* 4(3):281–286
- George R, Merten S, Wang TT, Kennedy P, Maitz P (2014) In vivo analysis of dermal and systemic absorption of silver nanoparticles through healthy human skin. *Australas J Dermatol* 55(3):185–190
- Ginn C, Khalili H, Lever R, Brocchini S (2014) PEGylation and its impact on the design of new protein-based medicines. *Future Med Chem* 6(16):1829–1846
- Giovanni M, Yue J, Zhang L, Xie J, Ong CN (2015) Pro-Inflammatory Responses of RAW264. 7 Macrophages when Treated with Ultralow Concentrations of Silver, Titanium Dioxide, and Zinc Oxide Nanoparticles. *J Hazard Mater* 297:146–152
- Graves JD, Craxton A, Clark EA (2004) Modulation and function of caspase pathways in B lymphocytes. *Immunol Rev* 197(1):129–146
- Kelso A (1998) Cytokines: principles and prospects. *Immunol Cell Biol* 76(4):300–317
- Kim TH, Kim M, Park HS, Shin US, Gong MS, Kim HW (2012) Size-dependent cellular toxicity of silver nanoparticles. *J Biomed Mater Res A* 100(4):1033–1043
- Larese FF, D'Agostin F, Crosera M, Adami G, Renzi N, Bovenzi M, Maina G (2009) Human skin penetration of silver nanoparticles through intact and damaged skin. *Toxicology* 255(1):33–37
- Lu W, Senapati D, Wang S, Tovmachenko O, Singh A, Yu H, Ray P (2010) Effect of surface coating on the toxicity of silver nanomaterials on human skin keratinocytes. *Chem Phys Lett* 487(1):92–96
- Miethling-Graff R, Rumpker R, Richter M, Verano-Braga T, Kjeldsen F, Brewer J, Hoyland J, Rubahn H-GG, Erdmann H (2014) Exposure to silver nanoparticles induces size- and dose-dependent oxidative stress and cytotoxicity in human colon carcinoma cells. *Toxicol In Vitro* 28(7):1280–1289
- Morar N, Willis-Owen SAG, Moffatt MF (2006) The genetics of atopic dermatitis. *J Allergy Clin Immunol* 118(1):24–34
- Mossman BT, Churg A (1998) Mechanisms in the pathogenesis of asbestosis and silicosis. *Am J Respir Crit Care Med* 157(5):1666–1680
- Murray AR, Kisin E, Inman A, Young S-H, Muhammed M, Burks T, Uheida A, Tkach A, Waltz M, Castranova V (2013) Oxidative stress and dermal toxicity of iron oxide nanoparticles in vitro. *Cell Biochem Biophys* 67(2):461–476
- Nowack B, Bucheli T (2007) Occurrence, behavior and effects of nanoparticles in the environment. *Environ Pollut* 150(1):5–22
- Nowack B, Krug H, Height M (2011) 120 years of nanosilver history: implications for policy makers. *Environ Sci Technol* 45(4):1177–1183
- Orlowski P, Krzyzowska M, Zdanowski R, Winnicka A, Nowakowska J, Stankiewicz W, Tomaszewska E, Celichowski G, Grobelny J (2013) Assessment of in vitro cellular responses of monocytes and keratinocytes to tannic acid modified silver nanoparticles. *Toxicol In Vitro* 27(6):1798–1808

- Pang C, Brunelli A, Zhu C, Hristozov D, Liu Y, Semenzin E, Wang W, Tao W, Liang J, Marcomini A, et al (2016) Demonstrating approaches to chemically modify the surface of Ag nanoparticles in order to influence their cytotoxicity and biodistribution after single dose acute intravenous administration. *Nanotoxicology* 10(2):129–139
- Park E-JJ, Choi K, Park K (2011a) Induction of inflammatory responses and gene expression by intratracheal instillation of silver nanoparticles in mice. *Arch Pharm Res* 34(2):299–307
- Park J, Lim D-H, Lim H-J, Kwon T, J-S Choi, Jeong S, Choi I-H, Cheon J (2011b) Size dependent macrophage responses and toxicological effects of Ag nanoparticles. *Chem Commun (Camb)* 47(15):4382–4384
- Parnsamut C, Brimson S (2015) Effects of silver nanoparticles and gold nanoparticles on IL-2, IL-6, and TNF- α production via MAPK pathway in leukemic cell lines. *Genet Mol Res* 14(2):3650
- Rossi D, Zlotnik A (2000) The biology of chemokines and their receptors. *Annu Rev Immunol* 18(1):217–242
- Ryan SMM, Mantovani G, Wang X, Haddleton DM, Brayden DJ (2008) Advances in PEGylation of important biotech molecules: delivery aspects. *Expert Opin Drug Deliv* 5(4):371–383
- Samberg ME, Oldenburg SJ, Monteiro-Riviere NA (2010) Evaluation of silver nanoparticle toxicity in skin in vivo and keratinocytes in vitro. *Environ Health Perspect* 118(3):407
- Schoemaker MH, Ros JE, Homan M, Trautwein C (2002) Cytokine regulation of pro- and anti-apoptotic genes in rat hepatocytes: NF- κ B-regulated inhibitor of apoptosis protein 2 (cIAP2) prevents apoptosis. *J Hepatol* 36(6):742–750
- Sharma V, Yngard R, Lin Y (2009) Silver nanoparticles: green synthesis and their antimicrobial activities. *Adv Colloid Interface Sci* 145(1):83–96
- Shi J, Sun X, Lin Y, Zou X, Li Z, Liao Y, Du M, Zhang H (2014) Endothelial cell injury and dysfunction induced by silver nanoparticles through oxidative stress via IKK/NF- κ B pathways. *Biomaterials* 35(24):6657–6666
- Song X-l, Li B, Xu K, Liu J, Ju W, Wang J, Liu X-d, Li J, Qi Y-f (2012) Cytotoxicity of water-soluble mPEG-SH-coated silver nanoparticles in HL-7702 cells. *Cell Biol Toxicol* 28:225–237
- Suliman Y, Omar A, Ali D, Alarifi S, Harrath AH, Mansour L, Alwasel S (2013) Evaluation of cytotoxic, oxidative stress, proinflammatory and genotoxic effect of silver nanoparticles in human lung epithelial cells. *Environ Toxicol* 30(2):149–160
- Takahashi M, Masuyama JI, Ikeda U, Kasahara T, Kitagawa SI, Takahashi YI, Shimada K, Kano S (1995) Induction of monocyte chemoattractant protein-1 synthesis in human monocytes during transendothelial migration in vitro. *Circ Res* 76(5):750–757
- Tejamaya M, Römer I, Merrifield R, Lead J (2012) Stability of citrate, PVP, and PEG coated silver nanoparticles in ecotoxicology media. *Environ Sci Technol* 46(13):7011–7017
- Thorley AJ, Tetley TD (2013) New perspectives in nanomedicine. *Pharmacol Ther* 140(2):176–185
- Tolaymat TM, El Badawy AM, Genaidy A, Scheckel KG, Luxton TP, Suidan M (2010) An evidence-based environmental perspective of manufactured silver nanoparticle in syntheses and applications: a systematic review and critical appraisal of peer-reviewed scientific papers. *Sci Total Environ* 408(5):999–1006
- Twentyman P, Luscombe M (1987) A study of some variables in a tetrazolium dye (MTT) based assay for cell growth and chemosensitivity. *Br J Cancer* 56(3):279
- Vance ME, Kuiken T, Vejerano EP, McGinnis SP, Hochella MF Jr, Rejeski D, Hull MS (2015) Nanotechnology in the real world: redeveloping the nanomaterial consumer products inventory. *Beilstein J Nanotechnol* 6(1):1769–1780
- Wang X, Ji Z, Chang C, Zhang H, Wang M, Liao Y-P, Lin S, Meng H, Li R, Sun B et al (2014) Use of coated silver nanoparticles to understand the relationship of particle dissolution and bioavailability to cell and lung toxicological potential. *Small* 10(2):385–398
- Wong KK, Cheung SO, Huang L, Niu J, Tao C, Ho C-MM, Che C-MM, Tam PK (2009) Further evidence of the anti-inflammatory effects of silver nanoparticles. *ChemMedChem* 4(7):1129–1135
- Wullaert A, Bonnet MC, Pasparakis M (2011) NF- κ B in the regulation of epithelial homeostasis and inflammation. *Cell Res* 21(1):146–158
- Yang E-JJ, Jang J, Lim D-HH, Choi I-HH (2012) Enzyme-linked immunosorbent assay of IL-8 production in response to silver nanoparticles. *Methods Mol Biol* 926:131–139
- Yarilin AA, Belyakov IM (2004) Cytokines in the thymus: production and biological effects. *Curr Med Chem* 11(4):447–464
- Yen H-J, Hsu S-H, Tsai C-L (2009) Cytotoxicity and immunological response of gold and silver nanoparticles of different sizes. *Small* 5(13):1553–1561
- Yoshizumi M, Nakamura T, Kato M, Ishioka T, Kozawa K, Wakamatsu K, Kimura H (2008) Release of cytokines/chemokines and cell death in UVB-irradiated human keratinocytes, HaCaT. *Cell Biol Int* 32(11):1405–1411
- Zhang LW, Yu WW, Colvin VL, Monteiro-Riviere NA (2008) Biological interactions of quantum dot nanoparticles in skin and in human epidermal keratinocytes. *Toxicol Appl Pharmacol* 228(2):200–211