

Lump solutions to dimensionally reduced p -gKP and p -gBKP equations

Wen Xiu Ma · Zhenyun Qin · Xing Lü

Received: 26 September 2015 / Accepted: 28 November 2015 / Published online: 14 December 2015
© Springer Science+Business Media Dordrecht 2015

Abstract Based on generalized bilinear forms, lump solutions, rationally localized in all directions in the space, to dimensionally reduced p -gKP and p -gBKP equations in (2+1)-dimensions are computed through symbolic computation with Maple. The sufficient and necessary conditions to guarantee analyticity and rational localization of the solutions are presented. The resulting lump solutions contain six parameters, two of which are totally free, due to the translation invari-

ance, and the other four of which only need to satisfy the presented sufficient and necessary conditions. Their three-dimensional plots with particular choices of the involved parameters are made to show energy distribution.

Keywords Lump solution · Generalized bilinear derivative · (3+1)-dimensional bilinear p -gKP and p -gBKP equations

Mathematics Subject Classification 35Q51 · 35Q53 · 37K40

W. X. Ma
College of Mathematics and Physics, Shanghai University
of Electric Power, Shanghai 200090, People's Republic of
China

W. X. Ma
International Institute for Symmetry Analysis and
Mathematical Modelling, Department of Mathematical
Sciences, North-West University, Mafikeng Campus,
Private Bag X2046, Mmabatho 2735, South Africa

Z. Qin
School of Mathematics and Key Lab for Nonlinear
Mathematical Models and Methods, Fudan University,
Shanghai 200433, People's Republic of China
e-mail: zyqin@fudan.edu.cn

X. Lü
Department of Mathematics, Beijing Jiao Tong University,
Beijing 100044, People's Republic of China
e-mail: xlv@bjtu.edu.cn

W. X. Ma (✉)
Department of Mathematics and Statistics, University of
South Florida, Tampa, FL 33620-5700, USA
e-mail: mawx@cas.usf.edu

1 Introduction

Integrable equations possess soliton solutions—exponentially localized solutions in certain directions [1]. They can also possess positon solutions—a kind of periodic solutions [2]—and complexiton solutions—combinations of solitons and positons [3]. The Hirota bilinear forms [4] play an important role in presenting solitons, positons and complexitons.

In contrast to soliton solutions, lump solutions are a kind of rational function solutions, localized in all directions in the space. Particular examples of lump solutions are found for many integrable equations such as the KPI equation [5–8], the three-dimensional three-wave resonant interaction equation [9], the B-KP equation [10], the Davey–Stewartson II equation [7] and the Ishimori I equation [11]. There are general searches

for rational function solutions to the KdV equation, the Boussinesq equation and the Toda lattice equation (see, e.g., [12–14]), systematically through the Wronskian and Casoratian determinant techniques for integrable equations [15, 16]. Generalized bilinear forms are also used to compute rational function solutions to the generalized KdV, KP and Boussinesq equations [17–19]. A natural question arises what kind of lump solutions can exist for nonlinear partial differential equations which possess generalized bilinear forms.

The (3+1)-dimensional generalized KP and BKP (gKP and gBKP) equations are as follows [20]:

$$u_{xxx}y + 3(u_x u_y)_x + u_{tx} + u_{ty} - u_{zz} = 0, \tag{1.1}$$

and

$$u_{ty} - u_{xxx}y - 3(u_x u_y)_x + 3u_{xz} = 0. \tag{1.2}$$

Under the transformation $u = 2(\ln f)_x$, they become the Hirota bilinear equations

$$(D_x^3 D_y + D_t D_x + D_t D_y - D_z^2) f \cdot f = 0, \tag{1.3}$$

and

$$(D_t D_y - D_x^3 D_y + 3D_x D_z) f \cdot f = 0, \tag{1.4}$$

respectively. Here, D_t, D_x, D_y and D_z are Hirota bilinear derivatives [4], which have connections with Kac–Moody algebras and quantum field theory [21]. For the above bilinear gKP and gBKP equations, resonant solitons are presented, forming linear subspaces of solutions [20], and three-wave solutions are computed by using the multiple exp-function method [22].

In this paper, we would like to consider the following two generalized bilinear equations in (3+1)-dimensions, called the (3+1)-dimensional bilinear p -gKP and p -gBKP equations:

$$(D_{p,x}^3 D_{p,y} + D_{p,t} D_{p,x} + D_{p,t} D_{p,y} - D_{p,z}^2) f \cdot f = 0, \tag{1.5}$$

and

$$(D_{p,t} D_{p,y} - D_{p,x}^3 D_{p,y} + 3D_{p,x} D_{p,z}) f \cdot f = 0, \tag{1.6}$$

with p being an arbitrarily given natural number, often a prime number, and the generalized bilinear operators being defined by [23]:

$$\begin{aligned} & (D_{p,x_1}^{n_1} \cdots D_{p,x_M}^{n_M} f \cdot g)(x_1, \dots, x_M) \\ &= \prod_{i=1}^M \left(\frac{\partial}{\partial x_i} + \alpha \frac{\partial}{\partial x_i'} \right)^{n_i} f(x_1, \dots, x_M) g(x_1', \dots, x_M') \\ & \quad \times \Big|_{x_1'=x_1, \dots, x_M'=x_M} \\ &= \prod_{i=1}^M \sum_{l_i=0}^{n_i} \alpha^{l_i} \binom{n_i}{l_i} \frac{\partial^{n_i-l_i}}{\partial x_i^{n_i-l_i}} f(x_1, \dots, x_M) \\ & \quad \times \frac{\partial^{l_i}}{\partial x_i^{l_i}} g(x_1, \dots, x_M), \end{aligned} \tag{1.7}$$

where n_1, \dots, n_M are arbitrary nonnegative integers, and for an integer m , the m -th power of α is computed as follows:

$$\begin{aligned} \alpha^m &= (-1)^{r(m)}, \text{ if } m \equiv r(m) \pmod p \\ & \text{with } 0 \leq r(m) < p. \end{aligned} \tag{1.8}$$

The choices for powers in (1.8) just give a rule to take the signs: +1 or –1. When $p = 2k, k \in \mathbb{N}$, the two generalized bilinear Eqs. (1.5) and (1.6) simplify to the two Hirota bilinear Eqs. (1.3) and (1.4), respectively.

With symbolic computation with Maple, we will do a search for positive quadratic function solutions to the dimensionally reduced bilinear p -gKP and p -gBKP equations from taking $z = x$ or $z = y$ in Eqs. (1.5) and (1.6). To search for quadratic function solutions, we begin with

$$\begin{aligned} f &= g^2 + h^2 + a_9, \quad g = a_1x + a_2y + a_3t + a_4, \\ h &= a_5x + a_6y + a_7t + a_8, \end{aligned} \tag{1.9}$$

where $a_i, 1 \leq i \leq 9$, are real parameters to be determined. In the two-dimensional space, a sum involving one square does not generate exact solutions which are rationally localized in all directions in the space, through the dependent variable transformations $u = 2(\ln f)_x$ or $u = 2(\ln f)_{xx}$. Noting that the generalized bilinear equations

$$P(D_{p,x}, D_{p,y}, D_{p,z}, D_{p,t}) f \cdot f = 0$$

with a given polynomial P but different values of $p \geq 2$ have the same set of quadratic function solutions, and the resulting quadratic function solutions will generate the same set of lump solutions to the corresponding nonlinear p -gKP and p -gBKP equations with different values of p . Because of the same set of solutions, our discussion will focus on the case of $p = 3$. The sufficient and necessary conditions to guarantee analyticity and localization of the corresponding rational function solutions will be explicitly presented. A few concluding remarks will be given at the end of the paper.

2 Lump solutions to the reduced p -gKP equations

2.1 Reduction with $z = x$

When $p = 3$, the (3+1)-dimensional bilinear p -gKP Eq. (1.5) reduces to the following generalized bilinear equation in (2+1)-dimensions:

$$\begin{aligned}
 B_{p\text{-gKP}_x}(f) &:= (D_{3,x}^3 D_{3,y} + D_{3,t} D_{3,x} \\
 &\quad + D_{3,t} D_{3,y} - D_{3,x}^2) f \cdot f \\
 &= 2(3f_{xx} f_{xy} - f_t f_x + f_{tx} f + f_{ty} f - f_t f_y \\
 &\quad + f_x^2 - f_{xx} f) = 0,
 \end{aligned}
 \tag{2.1}$$

under $z = x$. Through the link between f and u :

$$u = 2(\ln f)_x, \tag{2.2}$$

the reduced bilinear p -gKP Eq. (2.1) is transformed into

$$\begin{aligned}
 P_{p\text{-gKP}_x}(u) &= \frac{9}{8}u^2 u_x v + \frac{3}{8}u^3 u_y + \frac{3}{4}u u_{xx} v \\
 &\quad + \frac{3}{4}u_x^2 v + \frac{3}{4}u^2 u_{xy} + \frac{9}{4}u u_x u_y + \frac{3}{2}u_{xx} u_y \\
 &\quad + \frac{3}{2}u_x u_{xy} - u_{xx} + u_{tx} + u_{ty} = 0,
 \end{aligned}
 \tag{2.3}$$

where $u_y = v_x$. The transformation (2.2) is also a characteristic one in establishing Bell polynomial theories of integrable equations [24, 25], and the actual relation between the reduced p -gKP Eq. (2.3) and the reduced bilinear

p -gKP Eq. (2.1) reads

$$P_{p\text{-gKP}_x}(u) = \left[\frac{B_{p\text{-gKP}_x}(f)}{f^2} \right]_x. \tag{2.4}$$

Therefore, if f solves the reduced bilinear p -gKP Eq. (2.1), then $u = 2(\ln f)_x$ will solve the reduced p -gKP Eq. (2.3).

For Eq. (2.1), a direct symbolic computation with f in (1.9) leads to the following set of constraining equations for the parameters:

$$\begin{cases}
 a_1 = a_1, \\
 a_2 = \frac{a_1^2 a_3 - a_1 a_3^2 + 2 a_1 a_5 a_7 - a_1 a_7^2 - a_3 a_5^2}{a_3^2 + a_7^2}, \\
 a_3 = a_3, \quad a_4 = a_4, \quad a_5 = a_5, \\
 a_6 = -\frac{a_1^2 a_7 - 2 a_1 a_3 a_5 + a_3^2 a_5 - a_5^2 a_7 + a_5 a_7^2}{a_3^2 + a_7^2},
 \end{cases}$$

$$\left. \begin{aligned}
 a_7 &= a_7, \quad a_8 = a_8, \\
 a_9 &= \frac{3(a_1^2 + a_5^2)^2 [(a_1 - a_3)a_3 + (a_5 - a_7)a_7]}{(a_1 a_7 - a_3 a_5)^2} \end{aligned} \right\}, \tag{2.5}$$

which needs to satisfy the conditions

$$a_3 a_7 \neq 0, \quad a_1 a_7 - a_3 a_5 \neq 0, \tag{2.6}$$

to make the corresponding solutions f to be well defined. The condition

$$(a_1 - a_3)a_3 + (a_5 - a_7)a_7 > 0, \tag{2.7}$$

guarantees the positiveness of f , and the condition

$$(a_1^2 - a_5^2)(a_1 a_7 + a_3 a_5) - 2 a_1 a_5 (a_1 a_3 - a_5 a_7) \neq 0, \tag{2.8}$$

guarantees the localization of u in all directions in the (x, y) -plane. The parameters in the set (2.5) generate a class of positive quadratic function solutions to the reduced bilinear p -gKP Eq. (2.1):

$$\begin{aligned}
 f &= \left(a_1 x + \frac{a_1^2 a_3 - a_1 a_3^2 + 2 a_1 a_5 a_7 - a_1 a_7^2 - a_3 a_5^2}{a_3^2 + a_7^2} \right. \\
 &\quad \left. y + a_3 t + a_4 \right)^2 \\
 &\quad + \left(a_5 x - \frac{a_1^2 a_7 - 2 a_1 a_3 a_5 + a_3^2 a_5 - a_5^2 a_7 + a_5 a_7^2}{a_3^2 + a_7^2} \right. \\
 &\quad \left. y + a_7 t + a_8 \right)^2 \\
 &\quad + \frac{3(a_1^2 + a_5^2)^2 [(a_1 - a_3)a_3 + (a_5 - a_7)a_7]}{(a_1 a_7 - a_3 a_5)^2},
 \end{aligned}
 \tag{2.9}$$

and the resulting class of quadratic function solutions, in turn, yields a class of lump solutions to the reduced p -gKP Eq. (2.3) through the transformation (2.2):

$$u = \frac{4(a_1 g + a_5 h)}{f}, \tag{2.10}$$

where the function f is defined by (2.9), and the functions of g and h are given as follows:

$$g = a_1x + \frac{a_1^2a_3 - a_1a_3^2 + 2a_1a_5a_7 - a_1a_7^2 - a_3a_5^2}{a_3^2 + a_7^2}y + a_3t + a_4, \tag{2.11}$$

$$h = a_5x - \frac{a_1^2a_7 - 2a_1a_3a_5 + a_3^2a_5 - a_5^2a_7 + a_5a_7^2}{a_3^2 + a_7^2}y + a_7t + a_8. \tag{2.12}$$

In this class of lump solutions, all six involved parameters of a_1, a_3, a_4, a_5, a_7 and a_8 are arbitrary, provided that the three conditions (2.6), (2.7) and (2.8) are satisfied, which guarantee the definedness, positiveness and localization in all directions in the space for the solutions, respectively.

2.2 Reduction with $z = y$

When $p = 3$, the (3+1)-dimensional bilinear p -gKP Eq. (1.5) reduces the following generalized bilinear equation:

$$B_{p\text{-gKP}_y}(f) := (D_{3,x}^3D_{3,y} + D_{3,t}D_{3,x} + D_{3,t}D_{3,y} - D_{3,y}^2)f \cdot f = 2(3f_{xx}f_{xy} - f_t f_x + f_{tx}f + f_{ty}f - f_t f_y + f_y^2 - f_{yy}f) = 0, \tag{2.13}$$

under $z = y$. Through the link between f and u defined by (2.2), the reduced bilinear p -gKP Eq. (2.13) is transformed into

$$P_{p\text{-gKP}_y}(u) = \frac{9}{8}u^2u_xv + \frac{3}{8}u^3u_y + \frac{3}{4}uu_{xx}v + \frac{3}{4}u_x^2v + \frac{3}{4}u^2u_{xy} + \frac{9}{4}uu_xu_y + \frac{3}{2}u_{xx}u_y + \frac{3}{2}u_xu_{xy} - u_{yy} + u_{tx} + u_{ty} = 0, \tag{2.14}$$

where $u_y = v_x$. The actual relation between the reduced p -gKP Eq. (2.14) and the reduced bilinear p -gKP Eq. (2.13) reads

$$P_{p\text{-gKP}_y}(u) = \left[\frac{B_{p\text{-gKP}_y}(f)}{f^2} \right]_x. \tag{2.15}$$

Therefore, if f solves the reduced bilinear p -gKP Eq. (2.13), then $u = 2(\ln f)_x$ will solve the reduced p -gKP Eq. (2.14).

For Eq. (2.13), a direct symbolic computation with f defined by (1.9) yields the following set of constraining equations for the parameters:

$$\left\{ \begin{aligned} a_1 &= \frac{a_2^2a_3 - a_2a_3^2 + 2a_2a_6a_7 - a_2a_7^2 - a_3a_6^2}{a_3^2 + a_7^2}, \\ a_2 &= a_2, \quad a_3 = a_3, \quad a_4 = a_4, \\ a_5 &= -\frac{a_2^2a_7 - 2a_2a_3a_6 + a_3^2a_6 - a_6^2a_7 + a_6a_7^2}{a_3^2 + a_7^2}, \\ a_6 &= a_6, \quad a_7 = a_7, \quad a_8 = a_8, \\ a_9 &= \frac{3(a_1^2 + a_5^2)(a_1a_2 + a_5a_6)(a_3^2 + a_7^2)}{(a_2a_7 - a_3a_6)^2} \end{aligned} \right\}, \tag{2.16}$$

which needs to satisfy

$$a_3a_7 \neq 0, a_2a_7 - a_3a_6 \neq 0. \tag{2.17}$$

Noting the expression of a_9 in (2.16), the positiveness of f needs $a_1a_2 + a_5a_6 > 0$, which is equivalent to

$$(a_2 - a_3)a_3 + (a_6 - a_7)a_7 > 0, \tag{2.18}$$

thanks to (2.17). The localization of f needs $a_1a_6 - a_2a_5 \neq 0$, which equivalently requires

$$(a_2^2 - a_6^2)(a_2a_7 + a_3a_6) - 2a_2a_6(a_2a_3 - a_6a_7) \neq 0. \tag{2.19}$$

The parameters in the set (2.16) lead to a class of positive quadratic function solutions to the reduced bilinear p -gKP Eq. (2.13):

$$f = \left(\frac{a_2^2a_3 - a_2a_3^2 + 2a_2a_6a_7 - a_2a_7^2 - a_3a_6^2}{a_3^2 + a_7^2} x + a_2y + a_3t + a_4 \right)^2 + \left(-\frac{a_2^2a_7 - 2a_2a_3a_6 + a_3^2a_6 - a_6^2a_7 + a_6a_7^2}{a_3^2 + a_7^2} x + a_6y + a_7t + a_8 \right)^2 + \frac{3(a_1^2 + a_5^2)(a_1a_2 + a_5a_6)(a_3^2 + a_7^2)}{(a_2a_7 - a_3a_6)^2}, \tag{2.20}$$

where a_1 and a_5 are defined as in (2.16), and the resulting class of quadratic function solutions, in turn,

yields a class of lump solutions to the reduced p -gKP Eq. (2.14) through the transformation (2.2):

$$u = \frac{4(a_1g + a_5h)}{f}, \tag{2.21}$$

where the function f is defined by (2.20), and the functions of g and h are given as follows:

$$g = \frac{a_2^2a_3 - a_2a_3^2 + 2a_2a_6a_7 - a_2a_7^2 - a_3a_6^2}{a_3^2 + a_7^2}x + a_2y + a_3t + a_4, \tag{2.22}$$

$$h = -\frac{a_2^2a_7 - 2a_2a_3a_6 + a_3^2a_6 - a_6^2a_7 + a_6a_7^2}{a_3^2 + a_7^2}x + a_6y + a_7t + a_8. \tag{2.23}$$

In this class of lump solutions, all six involved parameters of a_2, a_3, a_4, a_6, a_7 and a_8 are arbitrary, provided that the conditions in (2.17), (2.18) and (2.19) are satisfied.

$$P_{p\text{-gBKP}_x}(u) := -\frac{9}{8}u^2u_xv - \frac{3}{8}u^3u_y - \frac{3}{4}uu_{xx}v - \frac{3}{4}u_x^2v - \frac{3}{4}u^2u_{xy} - \frac{9}{4}uu_xu_y - \frac{3}{2}u_{xx}u_y - \frac{3}{2}u_xu_{xy} + 3u_{xx} + u_{ty} = 0, \tag{3.2}$$

where $u_y = v_x$. The actual relation between the reduced p -gKP equation and the reduced bilinear p -gKP equation reads

$$P_{p\text{-gBKP}_x}(u) = \left[\frac{B_{p\text{-gBKP}_x}(f)}{f^2} \right]_x. \tag{3.3}$$

Therefore, if f solves the reduced bilinear p -gBKP Eq. (3.1), then $u = 2(\ln f)_x$ will solve the reduced p -gBKP Eq. (3.2).

For Eq. (3.1), a direct symbolic computation with f in (1.9) yields the following set of constraining equations for the parameters:

$$\left\{ \begin{aligned} a_1 &= a_1, a_2 = a_2, a_3 = -\frac{3(a_1^2a_2 + 2a_1a_5a_6 - a_2a_5^2)}{a_2^2 + a_6^2}, a_4 = a_4, \\ a_5 &= a_5, a_6 = a_6, a_7 = \frac{3(a_1^2a_6 - 2a_1a_2a_5 - a_6a_5^2)}{a_2^2 + a_6^2}, a_8 = a_8, \\ a_9 &= \frac{a_1^3a_2^3 + (a_1^2a_6^2 + a_1a_2a_5a_6 + a_2^2a_5^2)(a_1a_2 + a_5a_6) + a_5^3a_6^3}{(a_1a_6 - a_2a_5)^2} \end{aligned} \right\}, \tag{3.4}$$

3 Lump solutions to the reduced p -gBKP equations

3.1 Reduction with $z = x$

When $p = 3$, the (3+1)-dimensional bilinear p -gBKP Eq. (1.6) reduces to the following generalized bilinear equation:

$$B_{p\text{-gBKP}_x}(f) := (D_{3,t}D_{3,y} - D_{3,x}^3D_{3,y} + 3D_{3,x}^2)f \cdot f - 2(f_{ty}f - f_t f_y - 3f_{xx}f_{xy} - 3f_x^2 + 3f_{xx}f) = 0, \tag{3.1}$$

under $z = x$. Through the link between f and u defined by (2.2), the reduced bilinear gBKP Eq. (3.1) is transformed into

which needs to satisfy a determinant condition

$$a_1a_6 - a_2a_5 = \begin{vmatrix} a_1 & a_2 \\ a_5 & a_6 \end{vmatrix} \neq 0. \tag{3.5}$$

When $a_9 > 0$, i.e.,

$$a_1^3a_2^3 + (a_1^2a_6^2 + a_1a_2a_5a_6 + a_2^2a_5^2)(a_1a_2 + a_5a_6) + a_5^3a_6^3 > 0, \tag{3.6}$$

the corresponding quadratic function f , defined by (1.9), is positive. Now the parameters in the set (3.4) generate a class of positive quadratic function solutions to the reduced bilinear p -gBKP Eq. (3.1):

$$f = \left[a_1x + a_2y - \frac{3(a_1^2a_2 + 2a_1a_5a_6 - a_2a_5^2)}{a_2^2 + a_6^2}t + a_4 \right]^2 + \left[a_5x + a_6y + \frac{3(a_1^2a_6 - 2a_1a_2a_5 - a_6a_5^2)}{a_2^2 + a_6^2}t + a_8 \right]^2 + \frac{a_1^3a_2^3 + (a_1^2a_6^2 + a_1a_2a_5a_6 + a_2^2a_5^2)(a_1a_2 + a_5a_6) + a_5^3a_6^3}{(a_1a_6 - a_2a_5)^2}, \tag{3.7}$$

and the resulting class of quadratic function solutions, in turn, yields a class of lump solutions to the reduced p -gBKP equation in (3.2) through the transformation (2.2):

$$u = \frac{4(a_1g + a_5h)}{f}, \tag{3.8}$$

where the function f is defined by (3.7), and the functions of g and h are given as follows:

$$g = a_1x + a_2y - \frac{3(a_1^2a_2 + 2a_1a_5a_6 - a_2a_5^2)}{a_2^2 + a_6^2}t + a_4, \tag{3.9}$$

$$h = a_5x + a_6y + \frac{3(a_1^2a_6 - 2a_1a_2a_5 - a_6a_5^2)}{a_2^2 + a_6^2}t + a_8. \tag{3.10}$$

In this class of lump solutions, all six involved parameters of a_1, a_2, a_4, a_5, a_6 and a_8 are arbitrary, provided that the solutions are well defined and positive, i.e., if the conditions in (3.5) and (3.6) are satisfied. That determinant condition (3.5) precisely means that two directions (a_1, a_2) and (a_5, a_6) in the (x, y) -plane are not parallel, which is essential in formulating lump solutions in $(2+1)$ -dimensions by using a sum involving two squares.

3.2 Reduction with $z = y$

When $p = 3$, the $(3+1)$ -dimensional bilinear p -gBKP Eq. (1.6) reduces the following generalized bilinear equation:

$$B_{p\text{-gBKP}_y}(f) := (D_{3,t}D_{3,y} - D_{3,x}^3D_{3,y} + 3D_{3,x}D_{3,y})f \cdot f = 2(f_t y f - f_t f_y - 3f_{xx} f_{xy} - 3f_x f_y + 3f_{xy} f) = 0, \tag{3.11}$$

under $z = y$. Through the link between f and u defined by (2.2), the reduced bilinear p -gBKP Eq. (3.11) is transformed into

$$P_{p\text{-gBKP}_y}(u) := -\frac{9}{8}u^2u_xv - \frac{3}{8}u^3u_y - \frac{3}{4}uu_{xx}v - \frac{3}{4}u_x^2v - \frac{3}{4}u^2u_{xy} - \frac{9}{4}uu_xu_y - \frac{3}{2}u_{xx}u_y - \frac{3}{2}u_xu_{xy} + 3u_{xy} + u_{ty} = 0, \tag{3.12}$$

where $u_y = v_x$. The actual relation between the reduced p -gBKP Eq. (3.12) and the reduced bilinear p -gBKP Eq. (3.11) reads

$$P_{p\text{-gBKP}_y}(u) = \left[\frac{B_{p\text{-gBKP}_y}(f)}{f^2} \right]_x. \tag{3.13}$$

Therefore, if f solves the reduced bilinear p -gBKP Eq. (3.11), then $u = 2(\ln f)_x$ will solve the reduced p -gBKP Eq. (3.12).

For Eq. (3.11), a direct symbolic computation with f defined by (1.9) yields the following set of constraining equations for the parameters:

$$\left\{ \begin{aligned} a_1 &= -\frac{a_5a_6}{a_2}, \quad a_2 = a_2, \quad a_3 = \frac{3a_5a_6}{a_2}, \quad a_4 = a_4, \\ a_5 &= a_5, \quad a_6 = a_6, \quad a_7 = -3a_5, \quad a_8 = a_8, \quad a_9 = a_9 \end{aligned} \right\}, \tag{3.14}$$

where

$$a_2 \neq 0 \tag{3.15}$$

and all the other parameters are arbitrary. The parameters in this set lead to a class of positive quadratic function solutions to the reduced bilinear p -gBKP Eq. (3.1):

$$f = \left(-\frac{a_5a_6}{a_2}x + a_2y + \frac{3a_5a_6}{a_2}t + a_4 \right)^2 + (a_5x + a_6y - 3a_5t + a_8)^2 + a_9, \tag{3.16}$$

and the resulting class of quadratic function solutions, in turn, yields a class of lump solutions to the reduced p -gBKP equation in (3.12) through the transformation (2.2):

$$u = \frac{4(a_1g + a_5h)}{f}, \tag{3.17}$$




Fig. 1 The 3d plots of the lump solutions via (2.10). Parameters adopted here are: $a_1 = 4, a_2 = -12/5, a_3 = 2, a_4 = 0, a_5 = 6, a_6 = 86/5, a_7 = 1, a_8 = 0$ and $a_9 = 4563/4$

where the function f is defined by (3.16), and the functions of g and h are given as follows:

$$g = -\frac{a_5 a_6}{a_2} x + a_2 y + \frac{3 a_5 a_6}{a_2} t + a_4, \tag{3.18}$$

$$h = a_5 x + a_6 y - 3 a_5 t + a_8. \tag{3.19}$$

In this class of rational function solutions, all six involved parameters of a_2, a_4, a_5, a_6, a_8 and a_9 are arbitrary, provided that the solutions are well defined, i.e., if the condition (3.15) is satisfied. Under the condition (3.15), the determinant condition, which guarantees that two directions (a_1, a_2) and (a_5, a_6) in the (x, y) -plane are not parallel, is equivalent to

$$a_5 \neq 0. \tag{3.20}$$

Therefore, the conditions on the parameters




Fig. 2 The 3d plots of the lump solutions via (2.21). Parameters adopted here are: $a_1 = -68/13, a_2 = 4, a_3 = 3, a_4 = 0, a_5 = 184/13, a_6 = 8, a_7 = 2, a_8 = 0$ and $a_9 = 41625/13$

$$a_2 a_5 \neq 0, a_9 > 0, \tag{3.21}$$

will guarantee analyticity and localization of the solutions in (3.17) and thus present lump solutions to the reduced p -gBKP equation in (3.12).

4 Concluding remarks

Based on the generalized bilinear formulation and Maple symbolic computation, we presented positive quadratic functions solutions to the (2+1)-dimensional reduced bilinear p -gKP and p -gBKP equations, and thus, lump solutions to the (2+1)-dimensional reduced p -gKP and p -gBKP equations associated with $p = 3$.




Fig. 3 The 3d plots of the lump solutions via (3.8). Parameters adopted here are: $a_1 = 1, a_2 = 3, a_3 = 96/25, a_4 = 0, a_5 = 5, a_6 = 4, a_7 = -378/25, a_8 = 0$ and $a_9 = 14950/121$

The results actually work for all other values of $p \geq 2$ as well [26]. The representatives of the considered reduced generalized bilinear equations and their corresponding nonlinear differential equations with $p = 3$ were computed explicitly as in the set of Eqs. (2.1), (2.13), (3.1) and (3.11) and the set of Eqs. (2.3), (2.14), (3.2) and (3.12), respectively. The 3d plots of the presented lump solutions with some special choices of the involved parameters can be found in Figs. 1, 2, 3 and 4, which show energy distribution.

We point out that resonant solutions, in terms of exponential functions, to generalized trilinear differential equations have been systematically analyzed [27]. It would be very interesting to determine when there exist positive polynomial solutions including quadratic




Fig. 4 The 3d plots of the lump solutions via (3.17). Parameters adopted here are: $a_1 = -3, a_2 = 2, a_3 = 9, a_4 = 0, a_5 = 6, a_6 = 1, a_7 = -18, a_8 = 0$ and $a_9 = 4$

function solutions to generalized multi-linear equations. This kind of polynomial solutions will generate lump solutions to the corresponding nonlinear equations through $u = m(\ln f)_x$ or $u = m(\ln f)_{xx}$, where m is a constant related to the multi-linearity of the associated multi-linear equations. Rogue wave solutions could be generated as well in terms of positive polynomial solutions, being a particularly interesting class of exact solutions with rational function amplitudes. Such wave solutions are used to describe significant nonlinear wave phenomena in both oceanography [28] and nonlinear optics [29], which received a great deal

of recent attention in the mathematical physics community. To explore more soliton phenomena, it would be very interesting to consider multi-component and higher-order extensions of lump solutions, more importantly in (3+1)-dimensional cases and fully discrete cases.

Acknowledgments The work was supported in part by NNSFC under the Grants 11371326 and 11271008, the distinguished professorship of the Shanghai University of Electric Power, the Fundamental Research Funds for the Central Universities (2013XK03), the Natural Science Foundation of Shandong Province (Grant No. ZR2013AL016), Zhejiang Innovation Project of China (Grant No. T200905), the First-class Discipline of Universities in Shanghai and the Shanghai Univ. Leading Academic Discipline Project (No. A.13-0101-12-004). ZY was supported by the NNSFC under the Grant 11571079, Shanghai Pujiang Program (No. 14PJD007) and the Natural Science Foundation of Shanghai (No. 14ZR1403500), and the Young Teachers Foundation (No. 1411018) of Fudan university. XL was supported by NNSFC under the Grant 61308018 and the Fundamental Research Funds for the Central Universities (2014RC019 and 2015JBM111).

References

1. Ablowitz, M.J., Clarkson, P.A.: Solitons, Nonlinear Evolution Equations and Inverse Scattering. Cambridge University Press, Cambridge (1991)
2. Matveev, V.B.: Generalized Wronskian formula for solutions of the KdV equations: first applications. Phys. Lett. A **166**(3–4), 205–208 (2002)
3. Ma, W.X.: Complexiton solutions to the Korteweg–de Vries equation. Phys. Lett. A **301**(1–2), 35–44 (2002)
4. Hirota, R.: The Direct Method in Soliton Theory. Cambridge University Press, New York (2004)
5. Manakov, S.V., Zakharov, V.E., Bortog, L.A., Matveev, V.B.: Two-dimensional solitons of the Kadomtsev–Petviashvili equation and their interaction. Phys. Lett. A **63**(3), 205–206 (1977)
6. Johnson, R.S., Thompson, S.: A solution of the inverse scattering problem for the Kadomtsev–Petviashvili equation by the method of separation of variables. Phys. Lett. A **66**(4), 279–281 (1978)
7. Satsuma, J., Ablowitz, M.J.: Two-dimensional lumps in nonlinear dispersive systems. J. Math. Phys. **20**(7), 1496–1503 (1979)
8. Ma, W.X.: Lump solutions to the Kadomtsev–Petviashvili equation. Phys. Lett. A **379**(36), 1975–1978 (2015)
9. Kaup, D.J.: The lump solutions and the Bäcklund transformation for the three-dimensional three-wave resonant interaction. J. Math. Phys. **22**(6), 1176–1181 (1981)
10. Gilson, C.R., Nimmo, J.J.C.: Lump solutions of the BKP equation. Phys. Lett. A **147**(8–9), 472–476 (1990)
11. Imai, K.: Dromion and lump solutions of the Ishimori-I equation. Prog. Theor. Phys. **98**(5), 1013–1023 (1997)
12. Ma, W.X., You, Y.: Solving the Korteweg–de Vries equation by its bilinear form: Wronskian solutions. Trans. Am. Math. Soc. **357**(5), 1753–1778 (2005)
13. Ma, W.X., Li, C.X., He, J.S.: A second Wronskian formulation of the KP equation. Nonlinear Anal. TMA **70**(12), 4245–4258 (2009)
14. Ma, W.X., You, Y.: Rational solutions of the Toda lattice equation in Casoratian form. Chaos Solitons Fractals **22**(2), 395–406 (2004)
15. Freeman, N.C., Nimmo, J.J.C.: Soliton solutions of the Korteweg–de Vries and Kadomtsev–Petviashvili equations: the Wronskian technique. Phys. Lett. A **95**(1), 1–3 (1983)
16. Ma, W.X.: The Casoratian technique for integrable lattice equations. Dyn. Contin. Discrete Impuls. Syst. Ser. A Math. Anal. **16**(Differential Equations and Dynamical Systems, suppl. S1), 201–207 (2009)
17. Zhang, Y., Ma, W.X.: Rational solutions to a KdV-like equation. Appl. Math. Comput. **256**, 252–256 (2015)
18. Zhang, Y.F., Ma, W.X.: A Study on rational solutions to a KP-like equation. Z. Naturforsch. A **70**(4), 263–268 (2015)
19. Shi, C.G., Zhao, B.Z., Ma, W.X.: Exact rational solutions to a Boussinesq-like equation in (1+1)-dimensions. Appl. Math. Lett. **48**, 170–176 (2015)
20. Ma, W.X., Fan, E.G.: Linear superposition principle applying to Hirota bilinear equations. Comput. Math. Appl. **61**(4), 950–959 (2011)
21. Curry, J.M.: Soliton solutions of integrable systems and Hirota's method. Harvard College. Math. Rev. **2**(1), 43–59 (2008)
22. Ma, W.X., Zhu, Z.N.: Solving the (3+1)-dimensional generalized KP and BKP equations by the multiple exp-function algorithm. Appl. Math. Comput. **218**(24), 11871–11879 (2012)
23. Ma, W.X.: Generalized bilinear differential equations. Stud. Nonlinear Sci. **2**(4), 140–144 (2011)
24. Gilson, C., Lambert, F., Nimmo, J., Willox, R.: On the combinatorics of the Hirota D-operators. Proc. R. Soc. Lond. Ser. A **452**(1945), 223–234 (1996)
25. Ma, W.X.: Bilinear equations, Bell polynomials and linear superposition principle. J. Phys. Conf. Ser. **411**, 012021 (2013)
26. Ma, W.X., Zhou, Y.: Lump-type solutions to nonlinear differential equations derived from generalized bilinear equations. Mod. Phys. Lett. B (2015)
27. Ma, W.X.: Trilinear equations, Bell polynomials, and resonant solutions. Front. Math. China **8**(5), 1139–1156 (2013)
28. Müller, P., Garrett, C., Osborne, A.: Rogue waves. Oceanography **18**(3), 66–75 (2005)
29. Solli, D.R., Ropers, C., Koonath, P., Jalali, B.: Optical rogue waves. Nature **450**, 1054–1057 (2007)