

Iteration functions for p th roots of complex numbers

João Ribeiro Cardoso · Ana F. Loureiro

Received: 12 April 2010 / Accepted: 4 October 2010 /
Published online: 15 October 2010
© Springer Science+Business Media, LLC. 2010

Abstract A novel way of generating higher-order iteration functions for the computation of p th roots of complex numbers is the main contribution of the present work. The behavior of some of these iteration functions will be analyzed and the conditions on the starting values that guarantee the convergence will be stated. The illustration of the basins of attractions of the p th roots will be carried out by some computer generated plots. In order to compare the performance of the iterations some numerical examples will be considered.

Keywords Basins of attraction · Higher-order convergence · Iteration function · p th root · Residual

1 Introduction

Let $p \geq 2$ be a positive integer. Given a complex number w which do not belong to the closed negative real axis, there is a unique complex z such that

J. R. Cardoso (✉) · A. F. Loureiro
Instituto Superior de Engenharia de Coimbra,
Rua Pedro Nunes, 3030-199 Coimbra, Portugal
e-mail: jocar@isec.pt

J. R. Cardoso
Institute of Systems and Robotics, University of Coimbra,
Pólo II, 3030-290 Coimbra, Portugal

A. F. Loureiro
Centro de Matemática da Universidade do Porto,
Rua do Campo Alegre, 687, 4169-007 Porto, Portugal
e-mail: anafsl@fc.up.pt

$z^p = w$ and $-\pi/p < \arg(z) < \pi/p$, where $\arg(z)$ denotes the argument of z . This unique z is called the *principal* p th root of the complex w and will be denoted by $w^{1/p}$. All the p th roots of w can be related with $w^{1/p}$ by

$$\sqrt[p]{w} = e^{\frac{2\ell\pi i}{p}} w^{1/p},$$

with $\ell = 0, 1, \dots, p - 1$. The p th roots of w are exactly the real or complex roots of the polynomial equation

$$z^p - w = 0. \quad (1.1)$$

Recently, the iterative methods for computing the principal p th root of a complex number have received particular interest. This has been motivated, in part, by the relevance of this topic in the computation of p th roots of matrices [2, 9, 10, 12, 13, 16], which involves the p th roots of their complex or real eigenvalues. Moreover, the scalar iteration functions can be extended to matrices in a natural way.

The Newton's and Halley's iterative methods are the ones most widely used for solving nonlinear equations and, in particular, polynomial equations, whose orders of local convergence are, respectively, 2 and 3. For more details we refer the reader to [1, 4, 6, 19, 23] and the references therein. The Newton iteration function associated to the polynomial equation (1.1) is given by

$$N(z) = \frac{(p-1)z + wz^{1-p}}{p} \quad (1.2)$$

whereas the Halley iteration function is

$$H(z) = z \frac{(p-1)z^p + (p+1)w}{(p+1)z^p + (p-1)w}. \quad (1.3)$$

Since both iteration functions are rational, their dynamical behavior can be analyzed using results from the classical theory developed by Fatou and Julia [3, 8, 14].

Let λ be a fixed point of a rational iteration function $\psi(z)$, that is, $\psi(\lambda) = \lambda$. The set of initial values for which the sequence generated by $z_{k+1} = \psi(z_k)$ converge to λ is called the *basin of attraction* of λ , while the connected component of this set that contains λ is the *immediate basin of attraction*. It is well known that the boundaries of the basins of attraction are Julia sets and so exhibit a fractal behavior. In general, when the initial guess is close to such sets, the iterative methods are particularly sensitive, which means that the sequence $\{z_k\}_{k=0}^{\infty}$ can converge to a complex number distinct from λ or not converge at all.

A fixed point λ is called *attracting*, *repelling* or *indifferent* whether $|\psi'(\lambda)|$ is less than, greater than or equal to 1, respectively. If $\psi'(\lambda) = 0$ then λ is called *superattracting*. In the case of Newton iteration $N(z)$ the only existing fixed points are the p th roots of w , which are superattracting. For Halley iteration $H(z)$, the p th roots of w are also superattracting fixed points, but it has the extra fixed point $\lambda = 0$, which is repelling.

A rational iteration function $\psi(z)$ converges to λ with order of convergence j if there exists a real constant $c > 0$ such that $|z_{k+1} - \lambda| \leq c|z_k - \lambda|^j$, with $z_{k+1} = \psi(z_k)$, for all sufficiently large k . A classical convergence result for fixed point iterations is stated below.

Theorem 1.1 [11, 22] *Let $\psi(z)$ be an analytic function in a neighborhood of λ such that*

$$\psi(\lambda) = \lambda, \quad \psi'(\lambda) = \dots = \psi^{(j-1)}(\lambda) = 0, \quad \psi^{(j)}(\lambda) \neq 0.$$

Then there is z_0 sufficiently close to λ such that the sequence $\{z_k\}_{k=0}^{\infty}$ generated by $z_{k+1} = \psi(z_k)$ converges to λ with order j .

In this paper we present a novel way of generating infinitely many iteration functions for computing p th roots of complex numbers, with prefixed orders of convergence. Many known iteration functions, including Newton and Halley as well as the Schröder iteration functions, will appear as particular cases. As pointed out in [23], higher-order iteration functions may have additional fixed points that are not p th roots of w . Although in general these extraneous fixed points are repelling, they may complicate the structure of the basins of attractions of the p th roots by inserting new sets of “petals”. This phenomenon will be discussed for some higher-order iterations and will be illustrated by computer generated plots of the basins of attractions of the p th roots.

One important issue of our work is the computation of the principal p th root of a complex number by an iterative method. It is widely known that the success of these kind of methods depends on the choice of the initial guess. In [9, 10, 12, 13] the authors derived practical conditions for choosing an appropriate initial guess that guarantees the convergence of Newton’s and Halley’s method to the principal p th root. We extend their results to other iteration functions, by proving that if the initial approximation satisfy a certain condition then the sequence generated by the iterations converges toward the right p th root with the expected order.

This paper is organized as follows. In Section 2 an important result (Theorem 2.1), which can be used to generate several families of iterative methods with higher-order convergence, is derived and a definition for the residual of the terms of a sequence converging to a p th root is proposed. We also give examples of some families of iteration functions and find connections with families already existing in the literature. Sections 3, 4 and 5 are devoted to the study of particular families of iterations, where some convergence results are stated and, in particular, conditions for choosing an initial guess in terms of the residuals are derived. The results we bring into discussion are illustrated by computer generated basins of attraction. Numerical examples that permit to compare the performance of several iteration functions are given in Section 6. Finally, in the last section, after presenting a brief conclusion, we point out some issues that need further research.

Unless otherwise stated, throughout the text we will systematically consider $p \geq 2$ and $j \geq 1$ as integer numbers and w as a complex number not belonging to the closed negative real axis.

2 Generation of iteration functions

The following theorem allows the generation of infinitely many families of higher-order iteration functions that converge to the p th roots of a complex number.

Theorem 2.1 Consider the complex function f defined by $f(z) = [\alpha(z)]^{1/p}$, where $\alpha(\cdot)$ represents an analytic, one-to-one complex-valued function defined in an open set containing zero such that $\alpha(0) = 1$. If $T_j(z)$ denotes the Taylor polynomial of degree j of $f(z)$ at zero, then the principal p th root $w^{1/p}$ of w is a superattracting fixed point of the function

$$F_j(z) := z T_{j-1}(\alpha^{-1}(wz^{-p})), \tag{2.1}$$

with $j = 2, 3, \dots$ and $\alpha^{-1}(\cdot)$ representing the inverse function of α : $(\alpha^{-1} \circ \alpha)(t) = t$. Moreover, the iteration functions $F_j(z)$ are of orders at least j .

Proof Consider

$$R(z) := \alpha^{-1}(wz^{-p}). \tag{2.2}$$

The k -th order derivative of the composite function $(f \circ R)(z)$ at the point $z = w^{1/p}$ can be easily computed and is given by

$$\begin{aligned} \frac{d^k}{dz^k}(f \circ R)(z) \Big|_{z=w^{1/p}} &= \frac{d^n}{dz^n}[w z^{-p}]^{1/p} \Big|_{z=w^{1/p}} \\ &= \frac{(-1)^k k!}{w^{k/p}}, \quad k = 1, 2, 3, \dots, j - 1, \dots \end{aligned} \tag{2.3}$$

As we are dealing with derivatives of the composite function, this latter may also be computed based on the *Faà di Bruno's formula* ([5, pp. 137–138], [7]), and, in this case we have:

$$\begin{aligned} \frac{d^k}{dz^k}(f \circ R)(z) \Big|_{z=w^{1/p}} &= \sum_{\mu=1}^k \frac{d^\mu f(t)}{dt^\mu} \Big|_{t=0} B_{k,\mu}(r_1, r_2, \dots, r_{k-\mu+1}), \\ k &= 1, 2, 3, \dots, j - 1, \dots \end{aligned} \tag{2.4}$$

where $B_{k,\mu}(r_1, r_2, \dots, r_{k-\mu+1})$ denote the *Bell polynomials* ([5, pp. 133–136], [18, Chapter 5]) and

$$\begin{aligned} r_\sigma &= \frac{d^\sigma R(z)}{dz^\sigma} \Big|_{z=w^{1/p}} := \frac{d^\sigma}{dz^\sigma}(\alpha^{-1}(wz^{-p})) \Big|_{z=w^{1/p}}, \quad \text{for} \\ \sigma &= 1, \dots, k - \mu + 1 \text{ and } k = 1, 2, \dots \end{aligned} \tag{2.5}$$

Likewise, the formula of *Faà di Bruno* permits to obtain an expression for the k -th order derivative of the composite function $(T_{j-1} \circ R)(z)$ at the point $z = w^{1/p}$ and we have:

$$\frac{d^k}{dz^k}(T_{j-1} \circ R)(z) \Big|_{z=w^{1/p}} = \sum_{\mu=1}^k \frac{d^\mu T_{j-1}(t)}{dt^\mu} \Big|_{t=0} B_{k,\mu}(r_1, r_2, \dots, r_{k-\mu+1}),$$

$$k = 1, 2, 3, \dots, j - 1, \dots \tag{2.6}$$

where r_σ (with $\sigma = 1, \dots, k - \mu + 1$) are given by (2.5). Since

$$T_{j-1}(z) = \sum_{v=0}^{j-1} \left(\frac{d^v f(t)}{dt^v} \Big|_{t=0} \right) \frac{z^v}{v!}$$

we then have

$$\frac{d^\mu T_{j-1}(t)}{dt^\mu} \Big|_{t=0} = \begin{cases} \frac{d^\mu f(t)}{dt^\mu} \Big|_{t=0}, & 0 \leq \mu \leq j - 1; \\ 0, & \mu \geq j \end{cases};$$

consequently (2.6) becomes

$$\frac{d^k}{dz^k}(T_{j-1} \circ R)(z) \Big|_{z=w^{1/p}} = \sum_{\mu=1}^{\min(k, j-1)} \frac{d^\mu f(t)}{dt^\mu} \Big|_{t=0} B_{k,\mu}(r_1, r_2, \dots, r_{k-\mu+1}),$$

$$k = 1, 2, 3, \dots, j - 1, \dots \tag{2.7}$$

The comparison between (2.4) and (2.7) ensures the equality

$$\frac{d^k}{dz^k}(T_{j-1} \circ R)(z) \Big|_{z=w^{1/p}} = \frac{d^k}{dz^k}(f \circ R)(z) \Big|_{z=w^{1/p}} \quad \text{for any } k = 1, 2, \dots, j - 1$$

which, after (2.3), provides

$$\frac{d^k}{dz^k}(T_{j-1} \circ R)(z) \Big|_{z=w^{1/p}} = \frac{(-1)^k k!}{w^{k/p}} \quad \text{for any } k = 1, 2, \dots, j - 1$$

implying

$$\frac{d^k}{dz^k} F_j(z) \Big|_{z=w^{1/p}} = 0 \quad \text{for any } k = 1, 2, \dots, j - 1,$$

inasmuch as, according to the Leibniz formula for the product derivation, it holds

$$\frac{d^k}{dz^k} F_j(z) \Big|_{z=w^{1/p}} = \left(z \frac{d^k(T_{j-1} \circ R)}{dz^k}(z) + k \frac{d^{k-1}(T_{j-1} \circ R)}{dz^{k-1}}(z) \right) \Big|_{z=w^{1/p}}.$$

Finally, Theorem 1.1 allows to complete the proof. □

Using similar arguments, it is straightforward that Theorem 2.1 can be extended to the non principal p th roots of w . One important consequence

of the previous result is that for any function $\alpha(z)$ satisfying the required conditions, the sequence $\{z_k\}_{k=0}^\infty$ defined by

$$z_{k+1} = z_k T_{j-1} \left[\alpha^{-1}(wz_k^{-p}) \right], \tag{2.8}$$

converges, with order at least j , to a p th root of w , provided that an initial guess z_0 sufficiently close to that root is taken. Note that there are an infinity of possibilities for choosing $\alpha(z)$. For instance, for each $a, b, c \in \mathbb{C}$, the Möbius transformation

$$\alpha(z) = \frac{az + b}{cz + b}$$

(with $ab - bc \neq 0$) provides an important source of iteration functions. Some particular Möbius transformations will be analyzed throughout the text. A list of illustrating examples of iteration functions thus generated is discussed below, but first it shall be recalled that for any two complex numbers a and x the function $(a + z)^{-x}$ has a formal (binomial) series [5, p. 37]:

$$(a + z)^{-x} = \sum_{n \geq 0} (-1)^n (x)_n \frac{a^{-x-n} z^n}{n!}, \tag{2.9}$$

where the symbol $(x)_n$ denotes the *Pochhammer symbol* (also known as *rising factorial*) which is defined by $(x)_0 = 1$ and $(x)_n := \prod_{v=0}^{n-1} (x + v) = x(x + 1) \dots (x + n - 1)$ for any integer $n \geq 1$.

Example 1 Under the same assumptions and notations of Theorem 2.1 with $\alpha(z) = 1 - z$, one has the following family of iteration functions of order j :

$$\begin{aligned} N_j(z) &= z T_{j-1}(1 - wz^{-p}) \\ &= z \left(a_0 + a_1(1 - wz^{-p}) + \dots + a_{j-1}(1 - wz^{-p})^{j-1} \right), \end{aligned} \tag{2.10}$$

where $j = 2, 3, \dots$ and $a_n = \frac{1}{n!} \left(-\frac{1}{p}\right)_n$, for any $n = 0, \dots, j - 1$. For the particular cases $j = 2, 3$,

$$N_2(z) = z \left(1 - \frac{1}{p}(1 - wz^{-p}) \right)$$

and

$$N_3(z) = z \left(1 - \frac{1}{p}(1 - wz^{-p}) + \frac{1-p}{2!p^2}(1 - wz^{-p})^2 \right),$$

which have orders 2 and 3, respectively. It is to check that $N_2(z) \equiv N(z)$ given in (1.2), that is, N_2 is exactly the Newton iteration for the p th root of w . Moreover, as it will be seen in Section 3, N_j coincides with the Schröder iteration of order j applied to the polynomial equation (1.1).

Example 2 Consider the assumptions and notations of Theorem 2.1 with $\alpha(z) = 1/(1 - z)$. Now we have the following family of iteration functions:

$$L_j(z) = zT_{j-1}(1 - w^{-1}z^p) = z \left(a_0 + a_1(1 - w^{-1}z^p) + \dots + a_{j-1}(1 - w^{-1}z^p)^{j-1} \right), \quad (2.11)$$

where $a_n = \frac{1}{n!} \left(\frac{1}{p} \right)_n$, for $n = 0, 1, \dots, j - 1$. These iteration functions have already appeared in the literature. For instance, the particular case $j = 2$, L_2 , is the Newton’s iteration associated to the equation $\frac{1}{z^p} - \frac{1}{w} = 0$. Moreover, replacing w^{-1} by w in (2.11), we can observe that the iterations L_j coincide with the iteration functions addressed firstly in [21], and later on in [15]. L_j is also a particular case of the Padé family of iterations studied in [16] (see more details in Section 4).

For a given $\alpha(z)$ as in Theorem 2.1, the complex valued function $R(\cdot)$ defined in (2.2) will play an important role in our analysis. If we consider the sequence $\{z_k\}_{k=0}^\infty$ defined in (2.8), $R(z_k)$ can be seen as a residual, in the sense that it controls the error of the approximation

$$wz_k^{-p} \approx 1.$$

Indeed, assuming that, for a sufficiently large k , $R(z_k) = \alpha^{-1}(wz_k^{-p}) \approx 0$, the continuity of α^{-1} ensures that

$$wz_k^{-p} \approx \alpha(0) = 1.$$

Residuals of iteration functions for p th roots have already appeared in the literature; see, for instance, [15, 21] and [10]. Nevertheless, they have been treated separately as being associated to a particular iteration function. Our approach here is new in the sense that we can give an unified definition that includes those residuals as particular cases. The same holds for the recurrence relationship stated in the next lemma.

Lemma 2.1 *Under the same assumptions and notations of Theorem 2.1 and for each $k \in \mathbb{N}$, the residual $R(z_k) = \alpha^{-1}(wz_k^{-p})$ fulfills*

$$R(z_{k+1}) = \alpha^{-1}(\alpha(R(z_k)) [T_{j-1}(R(z_k))]^{-p}).$$

Proof The result follows from the identities

$$\begin{aligned} R(z_{k+1}) &= \alpha^{-1}(wz_{k+1}^{-p}) = \alpha^{-1} \left(w \left[z_k T_{j-1} \left(\alpha^{-1}(wz_k^{-p}) \right) \right]^{-p} \right) \\ &= \alpha^{-1} \left(wz_k^{-p} \left[T_{j-1} \left(\alpha^{-1}(wz_k^{-p}) \right) \right]^{-p} \right) \\ &= \alpha^{-1} \left(\alpha(R(z_k)) \left[T_{j-1}(R(z_k)) \right]^{-p} \right). \end{aligned}$$

□

3 The iteration functions N_j

In this section we focus our attention on the iteration functions N_j defined in (2.10). We start by proving that N_j is a Schröder iteration function associated to the polynomial equation $z^p - w = 0$ (Lemma 3.1) and for $j = 2, 3$ we define conditions over the initial guesses that guarantee the convergence to the right p th root.

In [20] Schröder constructed a family of prescribed order iteration functions for finding the simple roots of a nonlinear equation $g(z) = 0$. These functions are defined by the expression

$$S_j(z) = z + \sum_{n=1}^{j-1} c_n(z)[-g(z)]^n,$$

where

$$c_n(z) = \frac{1}{n!} \left[\frac{1}{g'(z)} \frac{d}{dz} \right]^{n-1} \frac{1}{g'(z)} \tag{3.1}$$

and $\left[\frac{1}{g'(z)} \frac{d}{dz} \right]^k := \frac{1}{g'(z)} \frac{d}{dz} \left(\left[\frac{1}{g'(z)} \frac{d}{dz} \right]^{k-1} \right)$ for any $k = 1, 2, \dots$

Throughout this section, the assumptions and notations of Theorem 2.1 will be considered but with

$$\alpha(z) = 1 - z,$$

which brings us to the same notations of Example 1.

Lemma 3.1 *Let N_j be defined as in (2.10) and let S_j be the Schröder iteration function associated to the polynomial equation (1.1). Then*

$$N_j(z) = S_j(z),$$

for all $z \in \mathbb{C} \setminus \{0\}$ and $j = 2, 3, \dots$

Proof Let $R(z)$ be the function given by (2.2), i.e.,

$$R(z) = \alpha^{-1}(wz^{-p}) = 1 - wz^{-p}, \tag{3.2}$$

denote the residual associated to $N_j(z)$. If $g(z) = z^p - w$, then $R(z) = g(z)/z^p$. In this case $N_j(z)$ becomes

$$\begin{aligned} N_j(z) &:= z T_{j-1}(R(z)) \\ &= z \left(a_0z + \frac{a_1z}{z^p}g(z) + \frac{a_2z}{(z^p)^2}[g(z)]^2 + \dots + \frac{a_{j-1}z}{(z^p)^{j-1}}[g(z)]^{j-1} \right) \end{aligned} \tag{3.3}$$

with $a_n = \frac{1}{n!} \left(-\frac{1}{p}\right)_n$, for any $n = 0, \dots, j - 1$. On the other hand, proceeding by finite induction, one has

$$c_n(z) = \frac{(-1)^n}{n!} \left(-\frac{1}{p}\right)_n \frac{z}{(z^p)^n}, \quad n = 0, 1, \dots, j - 1.$$

The comparison between (3.3) and the expression of S_j yields the equalities

$$c_n(z) = (-1)^n \frac{a_n z}{(z^p)^n}, \quad n = 0, 1, \dots, j - 1,$$

whence the result. □

Theorem 2.1 and also Lemma 3.1 confirm that, for each j , the iteration function $N_j(z)$ converges locally to a root of the polynomial equation (1.1) with order j . Now, we analyze the residuals of the sequence generated by N_j :

$$z_{k+1} = N_j(z_k). \tag{3.4}$$

For each $k = 0, 1, 2, \dots$, the residuals of (3.4) are given by

$$R(z_k) = 1 - w z_k^{-p},$$

and, according to Lemma 2.1, we have

$$R(z_{k+1}) = 1 - (1 - R(z_k)) [T_{j-1}(R(z_k))]^{-p}.$$

In order to find a condition on the residual $R(z_0)$ that ensures the convergence of (3.4) to a p th root of w , we seek an upper bound of $|R(z_{k+1})|$ in terms of $|R(z_k)|$, which allows us to relate the absolute values of the residuals of the successive terms of the sequence. This has already been done for the particular case of Newton’s method ($j = 2$), as we recall in the next theorem (see also [16, Theorem 4.5] for a similar result).

Theorem 3.1 [10] *Consider the Newton sequence $z_{k+1} = N_2(z_k)$ and the corresponding residuals $R(z_k)$. If*

$$|R(z_0)| = |1 - w z_0^{-p}| < 1, \tag{3.5}$$

then, for each k ,

$$|R(z_{k+1})| \leq |R(z_k)|^2 \leq |R(z_0)|^{2^k}.$$

Our goal is to extend this result for $j \geq 3$. A proof for the case $j = 3$ is based on Theorem 3.2 and will be presented in Corollary 3.1. However, we did not succeed to reach a general proof for all $j \geq 4$. The main reason for this is connected with the requirement of the computation of the coefficients of the Taylor series expansion for the residual function

$$\tilde{R}_j(z) := 1 - (1 - z) [T_{j-1}(z)]^{-p}, \tag{3.6}$$

whose expressions become more complicated as j increases. For $j = 2$ there is a nice and simple expression for the Taylor coefficients of \tilde{R}_2 (see [10]),

while the case $j = 3$ requires more laborious arguments, as it can be followed in Theorem 3.2.

The roots of the 2-degree Taylor polynomial

$$T_2(z) = 1 - \frac{1}{p} z - \frac{p-1}{2p^2} z^2 \tag{3.7}$$

associated to the function $f(z) = (1 - z)^{1/p}$ are

$$z_1 = -\frac{p(1 + \sqrt{2p-1})}{p-1} \quad \text{and} \quad z_2 = -\frac{p(1 - \sqrt{2p-1})}{p-1}$$

which are real and lie outside the unit circle. Consequently, the rational functions $(T_2(z))^{-p}$ and the residual function $\tilde{R}_3(z)$ defined in (3.6) are analytic inside the unit circle (i.e., for any z such that $|z| < 1$).

Theorem 3.2 *Consider the power series expansion of the two rational functions $(T_2(\cdot))^{-p}$ and $\tilde{R}_3(\cdot)$:*

$$(T_2(z))^{-p} = \sum_{v=0}^{\infty} c_v z^v \quad , \quad \tilde{R}_3(z) = \sum_{v=0}^{\infty} d_v z^v .$$

The following statements hold:

- (i) *The coefficients $d_v = 0$ whenever $v = 0, 1, 2$, and $d_v > 0$ for any $v = 3, 4, \dots$*
- (ii) *The series $\sum_{v=0}^{\infty} c_v$ is convergent.*
- (iii) *$\sum_{v=0}^{\infty} d_v$ is convergent and $\sum_{v=0}^{\infty} d_v = 1$.*
- (iv) *The series $\sum_{v=0}^{\infty} c_v z^v$ is absolutely convergent for any z such that $|z| < 1$.*
- (v) *The series $\sum_{v=0}^{\infty} d_v z^v$ is absolutely convergent for any z such that $|z| < 1$.*

Proof We begin by deriving explicit expressions for c_v and $d_v, v \in \mathbb{N}$, in order to prove statements (i) and (ii). At last we will show that (ii) provides (iii). The statements (iv) and (v) are a direct consequence of (ii) and (iii), on the basis of Abel’s convergence theorem for formal power series.

From its definition, the polynomial $T_2(z)$ may be written as $T_2(z) = T_1(z) + b_2 z^2$ with $b_2 = \frac{-(p-1)}{2p^2}$, and on account of (2.9), we successively have

$$\begin{aligned} (T_2(z))^{-p} &= (T_1(z) + b_2 z^2)^{-p} = \sum_{n \geq 0} \frac{(-1)^n (p)_n}{n!} (T_1(z))^{-p-n} (b_2 z^2)^n \\ &= \sum_{n \geq 0} \frac{(p-1)^n (p)_n}{n! 2^n p^{2n}} z^{2n} (T_1(z))^{-p-n} . \end{aligned}$$

According to (2.9), it follows

$$\begin{aligned} (T_1(z))^{-p-n} &= \left(1 - \frac{1}{p}z\right)^{-p-n} = \sum_{v \geq 0} \frac{(-1)^v (p+n)_v}{v!} (-1/p)^v z^v \\ &= \sum_{v \geq 0} \frac{(p+n)_v}{v! p^v} z^v \end{aligned}$$

whence

$$(T_2(z))^{-p} = \sum_{n \geq 0} \frac{(p-1)^n (p)_n}{n! 2^n p^{2n}} \sum_{v \geq 0} \frac{(p+n)_v}{v! p^v} z^{v+2n}. \tag{3.8}$$

Now, based on the equality,

$$\sum_{n \geq 0} \sum_{v \geq 0} A(n, v) = \sum_{v \geq 0} \sum_{n \geq 0} A(n, v) = \sum_{v \geq 0} \sum_{n=0}^{\lfloor v/2 \rfloor} A(n, v-2n),$$

with $\lfloor x \rfloor$ denoting the floor function of x , that is, the highest integer lower than or equal to x , the relation (3.8) becomes like

$$\begin{aligned} (T_2(z))^{-p} &= \sum_{v \geq 0} \sum_{n=0}^{\lfloor v/2 \rfloor} \frac{(p)_n (p+n)_{v-2n} (p-1)^n}{n!(v-2n)! p^v 2^n} z^v \\ &= \sum_{v \geq 0} \sum_{n=0}^{\lfloor v/2 \rfloor} \frac{(p)_{v-n} (p-1)^n}{n!(v-2n)! p^v 2^n} z^v, \end{aligned}$$

yielding

$$c_v = \sum_{n=0}^{\lfloor v/2 \rfloor} \frac{(p)_{v-n} (p-1)^n}{n!(v-2n)! p^v 2^n}. \tag{3.9}$$

Clearly, we have $c_0 = c_1 = c_2 = 1$. Inasmuch as $d_0 = 0$ and

$$d_{v+1} = c_v - c_{v+1}, \quad v \geq 2 \tag{3.10}$$

it subsequently follows that $d_1 = d_2 = 0$. With a few more computations, we derive the expression for d_v whenever $v \geq 3$:

$$\begin{aligned} d_{v+1} &= \sum_{n=0}^{\lfloor \frac{v}{2} \rfloor} \frac{(p)_{v-n} (p-1)^n}{n!(v-2n)! p^v 2^n} - \sum_{n=0}^{\lfloor \frac{v+1}{2} \rfloor} \frac{(p)_{v+1-n} (p-1)^n}{n!(v+1-2n)! p^{v+1} 2^n} \\ &= \sum_{n=0}^{\lfloor \frac{v+1}{2} \rfloor} \frac{(p)_{v+1-n} (p-1)^n}{n!(v+1-2n)! p^{v+1} 2^n} \frac{((v-2n+1)p - (p+v-n))}{(p+v-n)} \end{aligned}$$

whence

$$d_{v+1} = \sum_{n=0}^{\lfloor \frac{v+1}{2} \rfloor} \frac{(p)_{v+1-n} (p-1)^n}{n!(v+1-2n)! p^{v+1} 2^n} \frac{((v-2n)(p-1) - n)}{(p+v-n)}, \quad v \geq 2. \tag{3.11}$$

From (3.11) and on account of the fact that $p \geq 2$, we successively have

$$\begin{aligned}
 d_v &\geq \frac{1}{p^v} \sum_{n=0}^{\lfloor \frac{v}{2} \rfloor} \frac{(1)_{v-n}}{n! (v-2n)!} \frac{v-1-3n}{2^n p+v+1-n} \\
 &= \frac{1}{p^v} \sum_{n=0}^{\lfloor \frac{v}{2} \rfloor} \binom{v-n}{n} \frac{(v-1-3n)}{2^n (p+v+1-n)} \\
 &\geq \frac{1}{p^v (p+v+1)} \sum_{n=0}^{\lfloor \frac{v}{2} \rfloor} \frac{(v-1-3n)}{2^n} \\
 &= \frac{1}{p^v (p+v+1)} [2v-8+2^{-k}(-v+3k+7)]_{k=\lfloor \frac{v}{2} \rfloor}, \quad v \geq 3,
 \end{aligned}$$

whence (ii) holds.

On the other hand, and again based on the fact that $p \geq 2$, we are able to establish the inequality

$$\frac{(p)_{v-n} (p-1)^n}{p^v} = \left(\prod_{\mu=0}^{v-n-1} \frac{p+\mu}{p} \right) \left(\frac{p-1}{p} \right)^n \leq \prod_{\mu=0}^{v-n-1} \left(1 + \frac{\mu}{2} \right) = \frac{(v+1-n)!}{2^{v-n}}$$

and from the definition of (3.9), it follows

$$0 < c_v \leq \frac{(v+2)(v+4)}{2^{v+3}}$$

because

$$c_v \leq \sum_{n=0}^{\lfloor v/2 \rfloor} \frac{(v+1-n)!}{n!(v-2n)!} \frac{1}{2^v} = \frac{1}{2^v} \sum_{n=0}^{\lfloor v/2 \rfloor} \binom{v+1-n}{n+1} (n+1) \leq \frac{1}{2^v} \sum_{n=0}^{\lfloor v/2 \rfloor} (n+1).$$

Thus, the series of nonnegative terms $\sum_{v=0}^{\infty} c_v$ is convergent because of the convergence of the series $\sum_{v=0}^{\infty} \frac{(v+2)(v+4)}{2^{v+3}}$.

The statement (ii) provides statement (iii). Indeed, from (3.10), the sequence of the partial sums of the series $\sum_{v=3}^{\infty} d_v$ is given by

$$D_N = \sum_{v=3}^N d_v = c_2 - c_N = 1 - c_N,$$

which is convergent to 1 insofar as the convergence of the series $\sum_{v=0}^{\infty} c_v$ provides

$$\lim_{N \rightarrow \infty} c_N = 0.$$

Finally, on account of the well known Abel’s theorem on power series — if the series $\sum_{n=0}^{\infty} a_n$ converges, then the power series $\sum_{n=0}^{\infty} a_n z^n$ converges absolutely for $|z| < 1$ — the statements (iv) and (v) come as consequence of statements (ii) and (iii), respectively. \square

Corollary 3.1 Consider the residual function $R(\cdot)$ given by (3.2) and let z_k be defined by

$$z_{k+1} = N_3(z_k).$$

If $|R(z_0)| = |1 - wz_0^{-p}| < 1$, then

$$|R(z_{k+1})| \leq |R(z_k)|^3 \leq |R(z_0)|^{3^k},$$

for all $k = 1, 2, \dots$

Proof From Theorem 3.2 we have

$$\begin{aligned} |R(z_1)| &= \left| 1 - (1 - R(z_0))(T_2(R(z_0)))^{-p} \right| = \left| \sum_{v=0}^{\infty} d_v (R(z_0))^v \right| \\ &\leq |R(z_0)|^3 \sum_{v=0}^{\infty} d_{v+3} |R(z_0)|^v \leq |R(z_0)|^3 \sum_{v=0}^{\infty} d_{v+3} = |R(z_0)|^3. \end{aligned}$$

Now the result follows by induction. \square

From Theorem 2.1 we know that if z_0 is sufficiently close to a p th root of w , then the sequence (3.4) converges to that p th root with order j . The following result states a stronger convergence result for $j = 2, 3$, because it gives a specific condition on the initial guess.

Lemma 3.2 If z_0 is such that $|R(z_0)| = |1 - wz_0^{-p}| < 1$, with $R(\cdot)$ given by (3.2), then, for $j = 2, 3$, the sequence (3.4) converges to a p th root of w with order of convergence j .

Proof Let s be a p th root of w and assume that the sequence (3.4) converges to s . We know that $R(z_k) - R(s) = R(z_k)$ because $R(s) = 0$. The derivative of the residual at s can be written as

$$\mu := R'(s) = \lim_{k \rightarrow \infty} \frac{R(z_k) - R(s)}{z_k - s}.$$

Thus for each $\epsilon > 0$ there exists a positive integer k_0 such that for any $k \geq k_0$,

$$\left| \frac{R(z_k) - R(s)}{z_k - s} - \mu \right| < \epsilon$$

and then

$$|R(z_k) - R(s)| - |\mu(z_k - s)| \leq |R(z_k) - R(s) - \mu(z_k - s)| < \epsilon |z_k - s|.$$

Hence the following inequality holds:

$$|R(z_k) - R(s)| < (\epsilon + |\mu|)|z_k - s|,$$

or equivalently

$$|R(z_k) - R(s)| < \eta_1|z_k - s|, \tag{3.12}$$

with $\eta_1 = \epsilon + |\mu|$. Since

$$\frac{1}{R'(s)} = \lim_{k \rightarrow \infty} \frac{z_k - s}{R(z_k) - R(s)},$$

a similar argument lead us to conclude that for $k \geq k_1$, for some k_1 , there is $\eta_2 > 0$ such that

$$\eta_2|z_k - s| < |R(z_k) - R(s)|. \tag{3.13}$$

From (3.12), (3.13) and Corollary 3.1 it follows that

$$\begin{aligned} |z_{k+1} - s| &\leq \frac{1}{\eta_2} |R(z_{k+1}) - R(s)| = \frac{1}{\eta_2} |R(z_{k+1})| \leq \frac{1}{\eta_2} |R(z_k)|^j \\ &\leq \frac{1}{\eta_2} (\eta_1 |z_k - s|)^j = \frac{\eta_1^j}{\eta_2} |z_k - s|^j \end{aligned}$$

which shows that (3.4), with $j = 2, 3$ is of order j . □

Lemma 3.3 *Let $\theta = \arg(w) \in] - \pi, \pi[$ and let $\mathcal{S} = \{z \in \mathbb{C} : |1 - wz^{-p}| < 1\}$. Then*

$$\mathcal{S} = \mathcal{S}_0 \cup \mathcal{S}_1 \cup \dots \cup \mathcal{S}_{p-1} \tag{3.14}$$

where, for each $n = 0, 1, \dots, p - 1$,

$$\mathcal{S}_n = \left\{ z \in \mathcal{S} : \frac{(2n - 1)\pi + \theta}{p} < \arg(z) < \frac{(2n + 1)\pi + \theta}{p} \right\}$$

is a connected set and $\mathcal{S}_n \cap \mathcal{S}_\ell = \emptyset$, for $\ell \neq n$.

Proof Firstly we prove that, for each n , the points of the ray emerging from the origin

$$\mathcal{R}_n = \left\{ z \in \mathbb{C} : \arg(z) = \frac{(2n - 1)\pi + \theta}{p} \right\} \tag{3.15}$$

are not in \mathcal{S} . If $z \in \mathcal{R}_n$ then

$$z = \rho_1 e^{\frac{(2n-1)\pi+\theta}{p}i},$$

for some nonnegative ρ_1 . Let $w = \rho e^{\theta i}$ be the polar decomposition of w . Since

$$|1 - wz^{-p}| = \left| 1 - \rho e^{\theta i} \left(\rho_1 e^{\frac{(2n-1)\pi+\theta}{p}i} \right)^{-p} \right| = |1 + \rho \rho_1^{-p}| \geq 1,$$

it follows that $z \notin \mathcal{S}$, which shows that (3.14) holds. The sets \mathcal{S}_n are disjoint by definition. Now we show that each \mathcal{S}_n is a connected set. Let $P_n(z)$ be the complex valued function which assigns to each $z \in \mathbb{C} \setminus \mathbb{R}_0^-$ the unique p th root of z that satisfies

$$\frac{(2n - 1)\pi}{p} < \arg(z) < \frac{(2n + 1)\pi}{p}.$$

By definition, we have $[P_n(z)]^p = z$. Consider the function

$$g_n(z) = \frac{w^{1/p}}{P_n(z)}$$

which is continuous in $\mathbb{C} \setminus \mathbb{R}_0^-$. If

$$C = \{z \in \mathbb{C} : |1 - z| < 1\},$$

it is not hard to show that, for each n , $g_n(C \setminus \{0\}) = \mathcal{S}_n$. Therefore \mathcal{S}_n is a connected set because it is the image of the connected set $C \setminus \{0\}$ by a continuous function. □

Theorem 3.3 *Let \mathcal{S}_n be as in Lemma 3.3. If $z_0 \in \mathcal{S}_n$ then for $j = 2, 3$ the sequence (3.4) converges with order j to the unique p th root of w which lies on the wedge*

$$\mathcal{W}_n = \left\{ z \in \mathbb{C} : \frac{(2n - 1)\pi + \theta}{p} < \arg(z) < \frac{(2n + 1)\pi + \theta}{p} \right\}. \tag{3.16}$$

Proof Let $j = 2, 3$. By Theorem 3.1 and Corollary 3.1, N_j takes \mathcal{S} into \mathcal{S} . Moreover $N_j(\mathcal{S}_n) \subset \mathcal{S}_\ell$, for some ℓ , because all the sets in the decomposition (3.14) are disjoint and each \mathcal{S}_n is connected. Let w_n be the p th root of w which belongs to \mathcal{W}_n . It is easy to see that $w_n \in \mathcal{S}_n$. Since $N_j(w_n) = w_n \in \mathcal{S}_n$, the set $N_j(\mathcal{S}_n)$ must lie entirely in \mathcal{S}_n , for all n , and consequently, for $z_0 \in \mathcal{S}_n$, $z_k \in \mathcal{S}_n$ for all n . By Corollary 3.1, z_k converges to w_n . Thanks to Theorem 3.1 and Lemma 3.2, the convergence is of order j . □

Corollary 3.2 *Let \mathcal{S}_0 be the connected component of \mathcal{S} as in (3.14). If $z_0 \in \mathcal{S}_0$ for $j = 2, 3$ the sequence (3.4) converges to the principal p th root of w , $w^{1/p}$, with order j .*

Proof According to Theorem 3.3, one needs to show that $w^{1/p}$ lies in \mathcal{S}_0 . We know that $|1 - w(w^{1/p})^{-p}| = 0 < 1$ and that the argument of the principal p th root of w is θ/p which satisfies

$$-\frac{\pi}{p} < \frac{\theta}{p} < \frac{\pi}{p}.$$

From these inequalities we conclude that

$$\frac{-\pi + \theta}{p} < \frac{-\pi + \theta}{2p} < \frac{\theta}{p} < \frac{\pi + \theta}{2p} < \frac{\pi + \theta}{p},$$

which shows that $w^{1/p} \in \mathcal{S}_0$. □

If one is interested in taking real numbers as starting values, next corollary states the conditions that need to be satisfied.

Corollary 3.3 *Assume that w is such that $\Re(w) > 0$, where $\Re(w)$ denotes the real part of w , and let z_0 be a positive real number such that*

$$z_0 > \left(\frac{|w|^2}{2\Re(w)} \right)^{1/p}.$$

Then for $j = 2, 3$ the sequence (3.4) converges with order j to the principal p th root of w .

Proof The condition $|1 - wz_0^{-p}| < 1$ can be rewritten in the form

$$|z_0^p - w| < z_0^p.$$

This means that the positive real number z_0^p must lie on the open half plane that contains w and which is defined by the perpendicular bisector of the segment joining w with the origin. Since $\Re(w) > 0$, some calculations lead us to conclude that this half plane contains the part of real axis corresponding to real numbers greater than $\frac{|w|^2}{2\Re(w)}$. Thus

$$z_0^p > \frac{|w|^2}{2\Re(w)},$$


and so the set

$$\left\{ z_0 \in \mathbb{R} : z_0 > \left(\frac{|w|^2}{2\Re(w)} \right)^{1/p} \right\}$$

is entirely contained in \mathcal{S}_0 defined in (3.14). □

In Fig. 1, the top-left plot displays the sets $\mathcal{S}_0, \dots, \mathcal{S}_4$ with different colors together with the rays \mathcal{R}_n defined in (3.15), for $w = 1 + i$ and $p = 5$. For each n , \mathcal{S}_n corresponds to a specific 5th root of w and the previous results guarantee that, at least for $j = 2, 3$, it is contained in the basin of attraction of that p th root. This latter fact is illustrated in the remaining plots, where the basins of attraction of each 5th root of w were determined experimentally in Matlab. More precisely, a point z_0 in the rectangle $[-2, 2] \times [-2, 2]$ is marked with the same color of the 5th root $\sqrt[5]{w}$ of w , whenever $|z_{50} - \sqrt[5]{w}| < 10^{-4}$. We have also overlapped the boundaries of the regions of convergence on the basins of attraction for N_2, N_3 and N_4 . Black color is assigned to points that are not in

Fig. 1 Regions of convergence defined by the residuals of N_j (top left) and basins of attraction of the 5th roots of $w = 1 + i$ for N_2, N_3 and N_4



\mathcal{S}_n (top-left) or that do not belong to the basins of attraction of the p th roots of w (remaining plots).

While the only fixed points of N_2 are the p th roots of w , the iterations N_3 and N_4 have fixed points other than the p th roots of w . Since these extra fixed points are in general repelling, they do belong to the Julia sets of N_j and so contribute to change the shape of the boundaries. In the plots for N_3 and N_4 , we can see new sets of petals which are typical in iteration functions of orders greater than 2. We refer the reader to [23] for more details about the Julia sets of N_j .

The plot corresponding to N_4 (bottom right) suggests that the condition

$$|R(z_k)| = |1 - wz_0^{-p}| < 1$$

also guarantees the convergence of the sequence

$$z_{k+1} = N_4(z_k)$$

to a p th root of w . So assuming that results similar to the ones in Theorem 3.1 and Corollary 3.1 are stated for $j \geq 4$, it is easy to conclude that Lemmas 3.2 and 3.3, Theorem 3.3 and Corollaries 3.2 and 3.3 also hold for $j \geq 4$.

4 The iteration functions L_j

The analysis of the iterations functions L_j defined in (2.11) is the topic of this section. We proceed as in the previous section. For the particular cases $j = 2, 3, 4$, L_j is defined by

$$L_2(z) = z \left(1 + \frac{1}{p} (1 - w^{-1}z^p) \right)$$

$$\begin{aligned}
 L_3(z) &= z \left(1 + \frac{1}{p}(1 - w^{-1}z^p) + \frac{1+p}{2!p^2}(1 - w^{-1}z^p)^2 \right) \\
 L_4(z) &= z \left(1 + \frac{1}{p}(1 - w^{-1}z^p) + \frac{1+p}{2!p^2}(1 - w^{-1}z^p)^2 \right. \\
 &\quad \left. + \frac{(1+p)(1+2p)}{3!p^3}(1 - w^{-1}z^p)^3 \right).
 \end{aligned}$$

The residuals associated to the sequence

$$z_{k+1} = L_j(z_k) \tag{4.1}$$

are given by $R(z_k) = 1 - w^{-1}z_k^p$. By Lemma 2.1 these residuals satisfy the following relationship:

$$R(z_{k+1}) = 1 - (1 - R(z_k)) [T_{j-1}(R(z_k))]^p,$$

where T_{j-1} is given as in (2.11) (see Example 2). An important convergence result involving the sequence (4.1) and its residuals is stated in the following theorem. The proof can be found in [15, 21].

Lemma 4.1 *Let $R(z_k)$ denotes the residual of z_k defined in (4.1). If $|R(z_0)| = |1 - w^{-1}z_0^p| < 1$, then*

$$|R(z_{k+1})| \leq |R(z_k)|^j \leq |R(z_0)|^{j^k},$$

for all k and j . Moreover, the sequence (4.1) converges to a p th root of w .

The iterations L_j are also a particular case of the Padé family of iterations analyzed recently in [16]. More precisely, L_j is the iteration generated by the $[j - 1/0]$ Padé approximant to the function $f(z) = 1/(1 - z)^{1/p}$. The convergence of the Padé family associated to the matrix sector function was a topic that has been addressed in that paper, where some convergence results and conjectures were presented. Since the p -sector function and p th roots are closely related, a suitable change of variable can convert iterative methods for the p -sector function into iterative methods for p th roots and *vice-versa*. The details can be found in [16, Section 5].

Below, we state for L_j some convergence results that are similar to the ones presented in Section 3 for N_j . The proofs will be omitted, because they can be derived from the work developed in [16] (Lemma 4.2, Theorem 4.1 and Corollary 4.1) or are similar to the ones presented in Section 3.

Lemma 4.2 *If z_0 is such that $|R(z_0)| = |1 - w^{-1}z_0^p| < 1$, then the sequence (4.1) converges to a p th root of w with order of convergence j .*

Lemma 4.3 *Let $\theta = \arg(w) \in] - \pi, \pi[$ and let $\mathcal{T} = \{z \in \mathbb{C} : |1 - w^{-1}z^p| < 1\}$. Then*

$$\mathcal{T} = \mathcal{T}_0 \cup \mathcal{T}_1 \cup \dots \cup \mathcal{T}_{p-1} \tag{4.2}$$

where, for each $n = 0, 1, \dots, p - 1$,

$$\mathcal{T}_n = \left\{ z \in \mathcal{T} : \frac{(2n - 1)\pi + \theta}{p} < \arg(z) < \frac{(2n + 1)\pi + \theta}{p} \right\}$$

is a connected set and $\mathcal{T}_n \cap \mathcal{T}_\ell = \emptyset$, for $\ell \neq n$.

Theorem 4.1 *Let \mathcal{T}_n be as in Lemma 4.3. If $z_0 \in \mathcal{T}_n$ then the sequence (4.1) converges with order j to the unique p th root of w which lies on the wedge*

$$\mathcal{W}_n = \left\{ z \in \mathbb{C} : \frac{(2n - 1)\pi + \theta}{p} < \arg(z) < \frac{(2n + 1)\pi + \theta}{p} \right\}.$$

Corollary 4.1 *Let \mathcal{T}_0 be the connected component of \mathcal{T} as in (4.2). If $z_0 \in \mathcal{T}_0$ then z_k converges with order j to the principal p th root of w , $w^{1/p}$.*

Corollary 4.2 *Assume that w is such that $\Re(w) > 0$ and let z_0 be a positive real number such that*


$$z_0 < (2\Re(w))^{1/p}.$$

Then z_k converges with order j to the principal p th root of w .

Figure 2 displays Matlab generated plots to compare the domains of convergence defined by the sets \mathcal{T}_n in (4.2) with the basins of attractions of the 5th roots of $w = 1 + i$. The top-left plot shows the sets \mathcal{T}_n together with the rays \mathcal{R}_n defined in (3.15). The remaining plots include the basins of attraction of the 5th roots of w with respect to the iteration functions L_2, L_3 and L_4 combined with the boundaries of the sets \mathcal{T}_n . Compare the plots for L_2, L_3 and L_4 with the ones in the first column of [16, Fig. 1]. We can observe that the iteration functions L_j have much smaller domains of convergence than N_j . Although this can be seen as a disadvantage, it turns out that iterations L_j have been implemented successfully for computing matrix p th roots [9, 15]. The main reason lies on the fact that each iterate of (4.1) does not involve the computation of the inverse of z_k . When working with matrices this contributes to reduce the computational cost of the method and eventually to improve the accuracy.

Similarly to the iteration functions N_j , it happens that for $j \geq 3$, L_j has extra fixed points that are not p th roots of w . In general these fixed points are repelling and are responsible for the complicated structure of the Julia sets of L_j , by inserting new sets of petals.

Fig. 2 Regions of convergence defined by the residuals of L_j (top left) and basins of attraction of the 5th roots of $w = 1 + i$ for L_2, L_3 and L_4



5 More iteration functions

In this section we consider more iteration functions generated by the functions of the form

$$f(z) = \left(\frac{az + b}{cz + b} \right)^{1/p}, \tag{5.1}$$

with $ab - bc \neq 0$. Let T_j denotes the j -degree Taylor polynomial of f at zero. For each j the iteration function

$$F_j(z) = zT_{j-1} \left(\frac{b(wz^{-p} - 1)}{a - cwz^{-p}} \right)$$

has order j . The 2-degree Taylor polynomial of f is

$$T_2(z) = 1 + \frac{a - c}{bp}z + \frac{(a - c)^2 + p(c^2 - a^2)}{2b^2p^2}z^2.$$

If a and c are related by

$$\frac{a}{c} = \frac{1 + p}{1 - p}, \tag{5.2}$$

and $b \neq 0$, then the coefficient of z^2 in T_2 vanishes. This means that we have an iteration of order 3 defined upon a 1-degree Taylor polynomial, which has a simpler expression. Thus assuming, for instance, that $a = 1 + p$, $b = 1$ and $c = 1 - p$, the function

$$h(z) = \left[\frac{(1 + (p + 1)z)}{(1 + (1 - p)z)} \right]^{1/p},$$

gives rise to the following 3-order iteration function

$$\begin{aligned}
 H_3(z) &= zT_1 \left(\frac{w - z^p}{(p + 1)z^p + (p - 1)w} \right) \\
 &= z \frac{(p - 1)z^p + (p + 1)w}{(p + 1)z^p + (p - 1)w}
 \end{aligned}
 \tag{5.3}$$

which is exactly the Halley iteration $H(z)$ mentioned in (1.3). We note that other choices for a , b and c satisfying (5.2) lead to the same iteration function. This nice property of Halley iteration is not shared by other iterations generated by functions of the form (5.1), because the coefficient of z^2 in the corresponding Taylor polynomial does not vanish. However, several tests we carried out showed that the corresponding iteration functions converge faster as the numerator of the coefficient of z^2 becomes close to zero, that is, when

$$\frac{a}{c} \approx \frac{1 + p}{1 - p}.
 \tag{5.4}$$

Another function of the form (5.1) that will be addressed in this section is

$$m(z) = \left(\frac{1 + z}{1 - z} \right)^{1/p},$$

which satisfies (5.4) for p sufficiently large. The corresponding iteration functions are

$$M_j(z) = zT_{j-1} \left(\frac{w - z^p}{w + z^p} \right)
 \tag{5.5}$$

For $j = 2$, we have

$$M_2(z) = z \left(\frac{(p - 2)z^p + (p + 2)w}{pz^p + pw} \right).$$

Consider also the iteration functions generated by h ,

$$H_j(z) = zT_{j-1} \left(\frac{p(w - z^p)}{(p + 1)z^p + (p - 1)w} \right).
 \tag{5.6}$$

and the sequences generated by the recursions

$$z_{k+1} = M_j(z_k),
 \tag{5.7}$$

and

$$z_{k+1} = H_j(z_k).
 \tag{5.8}$$

Let us denote the residuals associated to the iterations functions M_j and H_j by R_M and R_H , respectively. These residuals are defined by (see (2.2))

$$R_M(z) = \frac{w - z^p}{w + z^p}$$

and

$$R_H(z) = \frac{p(w - z^p)}{(p + 1)z^p + (p - 1)w}.$$

According to Lemma 2.1, for z_k defined by (5.7) the following residual recurrence hold:

$$R_M(z_{k+1}) = \frac{(1 + R_M(z_k)) - (1 - R_M(z_k)) [T_{j-1}(R_M(z_k))]^p}{(1 + R_M(z_k)) + (1 - R_M(z_k)) [T_{j-1}(R_M(z_k))]^p}; \tag{5.9}$$

also for z_k given in (5.8) we have

$$\begin{aligned} &R_H(z_{k+1}) \\ &= \frac{(1 + (p + 1)R_H(z_k)) - ((1 - p)R_H(z_k) + 1) [T_{j-1}(R_H(z_k))]^p}{(p - 1)(1 + (p + 1)R_H(z_k)) + (p + 1)(1 + (1 - p)R_H(z_k)) [T_{j-1}(R_H(z_k))]^p}. \end{aligned} \tag{5.10}$$

Consider the following regions:

$$\mathcal{M} = \left\{ z_0 \in \mathbb{C} : |R_M(z_0)| = \left| \frac{w - z_0^p}{w + z_0^p} \right| < 1 \right\}$$

and

$$\mathcal{H} = \left\{ z_0 \in \mathbb{C} : |R_H(z_0)| = \left| \frac{w - z_0^p}{(p - 1)w + (p + 1)z_0^p} \right| < 1 \right\}.$$

A question that arises now is to know if \mathcal{M} (resp. \mathcal{H}) defines a region of convergence for (5.7) (resp. (5.8)), that is, if $z_0 \in \mathcal{M}$ (resp., $z_0 \in \mathcal{H}$) does the sequence (5.7) (resp. (5.8)) converge to a p th root of w with order j ? This seems to be a very hard issue because of the complicated expressions of the residuals recurrences (5.9) and (5.10). This means that we cannot expect to state nice convergence results like the ones in Corollary 3.1 and Lemma 4.1. However, we proved the following convergence result which concerns to the weaker case when (5.7) is restricted to real numbers and $j = 2$.

Theorem 5.1 Consider the real version of the iterative formula (5.7), with $j = 2$,

$$z_{k+1} = \frac{z_k}{p} \left(\frac{(p - 2)z_k^p + (p + 2)a}{z_k^p + a} \right),$$

where a is a positive real number. If z_0 is a positive real number then the sequence $\{z_k\}_{k=0}^\infty$ converges to $a^{1/p}$ with order of convergence at least 2.

Proof Firstly, we define the real function of real variable (see (5.9))

$$g(x) = \frac{(1 + x) - (1 - x)(1 + 2/p)^p}{(1 + x) + (1 - x)(1 + 2/p)^p}$$

and show that

$$|g(x)| < x^2,$$

for all $0 < x < 1$, which is equivalent to

$$g(x) < x^2 \tag{5.11}$$

and

$$-x^2 < g(x). \tag{5.12}$$

Let $0 < x < 1$. Since

$$(1 + 2/p)^p = a_0 + a_1x + a_2x^2 + \dots + a_px^p,$$

where $a_n = \binom{p}{p-n} \left(\frac{2}{p}\right)^n$, performing some calculations it is straightforward to conclude that the following equivalences hold:

$$\begin{aligned} g(x) < x^2 &\Leftrightarrow (1 + x)^2 - (1 + x^2)(1 + 2/p)^p < 0 \\ &\Leftrightarrow -a_2x^2 - \left(\sum_{n=3}^p (a_n + a_{n-2})x^n\right) \\ &\quad - a_{p-1}x^{p+1} - a_px^{p+2} < 0. \end{aligned} \tag{5.13}$$

We can easily observe that (5.13) holds and so (5.11) follows. To prove (5.12), we note that

$$\begin{aligned} -x^2 < g(x) &\Leftrightarrow (1 + x^2) - (1 - x)^2(1 + 2/p)^p > 0 \\ &\Leftrightarrow (2a_1 - a_2)x^2 + (-a_3 + 2a_2 - a_1)x^3 \\ &\quad + \left(\sum_{n=4}^p (-a_n + 2a_{n-1} - a_{n-2})x^n\right) \\ &\quad + (2a_p - a_{p-1})x^{p+1} - a_px^{p+2} > 0. \end{aligned} \tag{5.14}$$

Let b_n denote the coefficient of x^n ($n = 2, 3, \dots, p + 2$) in (5.14). It is easy to show that b_2 and b_3 are nonnegative, that is, $b_2 = 2a_1 - a_2 \geq 0$ and $b_3 = -a_3 + 2a_2 - a_1 \geq 0$. Next we show that the remaining coefficients are negative. Indeed, for $n = 4, \dots, p$, we have that

$$\begin{aligned} b_n &= -a_n + 2a_{n-1} - a_{n-2} \\ &= \left(\frac{2}{p}\right)^{n-2} \frac{(p-1) \dots (p-n+3)}{pn(n-1)(n-2)!} \\ &\quad \times ((5n - n^2 - 4)p^2 + (-4n^2 + 16n - 12)p + (-4n^2 + 12n - 8)) < 0, \end{aligned}$$

$$b_{p+1} = 2a_p - a_{p-1} < 0$$

$$b_{p+2} = -a_p < 0.$$

Since

$$b_2 + b_3 + \dots + b_{p+1} + b_{p+2} = a_1 > 0,$$

it follows that

$$b_2 + b_3 > -b_4 - \dots - b_{p+1} - b_{p+2}.$$

Thus

$$\begin{aligned} b_2x^2 + b_3x^3 &> b_2x^3 + b_3x^3 \\ &> -b_4x^3 - \dots - b_{p+1}x^3 - b_{p+2}x^3 \\ &> -b_4x^4 - \dots - b_{p+1}x^{p+1} - b_{p+2}x^{p+2} \end{aligned}$$

and so


$$b_2x^2 + b_3x^3 + b_4x^4 + \dots + b_{p+1}x^{p+1} + b_{p+2}x^{p+2} > 0,$$

which proves (5.14) and consequently (5.12). Since $|R_M(z_0)| = |(a - z_0^p)/(a + z_0^p)| < 1$ for each positive z_0 , proceeding as in the proof of Corollary 3.1 the result follows. □

We have compared experimentally the region \mathcal{M} (resp. \mathcal{H}) with the basins of attraction of the 5th roots of $w = 1 + i$ corresponding to the iteration function M_2 (resp., H_3). The pictures are displayed in Fig. 3. Comparing top-left and top-right plots we can observe that the region \mathcal{M} apparently defines a convergence region for M_2 , where starting values belonging to a given sector lead to convergence for the 5th root lying exactly on that sector. However, this does not seem to happen with the region \mathcal{H} for H_3 , which suggests that the residuals considered for H_j may not be the most appropriate. We recall that in [10] (and also in [17]) a different residual was proposed to study the convergence of Halley’s method.

Another interesting fact we can see in Fig. 3 is the similarity between the patterns exhibited by the basins of attraction for the iterations M_2 and H_3 .

Fig. 3 Region \mathcal{M} (top left), basins of attractions of the 5th roots of $w = 1 + i$ for M_2 (top right), region \mathcal{H} (bottom left) and basins of attractions of the 5th roots of w for H_3 (bottom right)



Contrarily to the basins of attraction for N_j and L_j , there are no black regions and so it may happen that both M_2 and H_3 always converge to a root of w , for any starting value not belonging to the Julia sets. We note that both iteration functions have 0 and the p th roots of w as fixed points, though 0 is a repelling fixed point.

6 Numerical examples

We have implemented the iteration functions N_2, L_2, M_2 , with order of convergence 2, and N_3, L_3 and H_3 with order of convergence 3 to illustrate the computation of the p th roots of some particular complex numbers. For any k , the error $|z_k - w^{1/p}|$ is displayed in each entry of the tables. All the tests were performed in Matlab using variable precision arithmetic with 1,000 significant decimal digits. Table 1 reports on the principal 5th root of $w = 1 + i$, taking $z_0 = 1$ as starting value. The residuals of z_0 are

$$|1 - wz_0^{-5}| = 1$$

for iterations N_2 and N_3 and

$$|1 - w^{-1}z_0^5| \approx 0.7$$

for L_2 and L_3 . Although the residuals of N_2 and N_3 do not satisfy the requirements of Corollary 3.2, the convergence to the principal 5th root of w occurs. Residuals of the starting value for M_2 and H_3 are respectively

$$\left| \frac{w - z_0^5}{w + z_0^5} \right| \approx 0.4$$

and

$$\left| \frac{5(w - z_0^5)}{6z_0^5 + 4w} \right| \approx 0.5.$$

Table 2 illustrates the computation of a non principal 8th root of $w = -1 + 2i$. We have taken $z_0 = -1 + 0.4i$ as the initial guess and computed the 8th root of w , $-0.9535... + 0.5600...i$, that lie on the wedge \mathscr{W}_4 defined in (3.16). The residuals of z_0 have absolute values less than 1 for the iterations M_2 and H_3 , and between 1 and 2 for the other iterations. Finally, Table 3 shows the application of the iterations to the particular case of the real p th root of a positive real number, with $p = 11, w = 29$ and $z_0 = 1.3$. All the residuals of z_0 have absolute values less than 1.

Our numerical examples show that, among the iteration functions of order 2, M_2 apparently converges faster than N_2 and L_2 . They also confirm that Halley’s method is a very good choice for computing p th roots.

Table 1 Values of the error $|z_k - (1 + i)^{1/5}|$, with $z_0 = 1$

k	Error N_2	Error L_2	Error M_2	Error N_3	Error L_3	Error H_3
1	$0.669... \times 10^{-1}$	$0.793... \times 10^{-1}$	$0.227... \times 10^{-1}$	$0.387... \times 10^{-1}$	$0.421... \times 10^{-1}$	$0.114... \times 10^{-1}$
2	$0.923... \times 10^{-2}$	$0.182... \times 10^{-1}$	$0.221... \times 10^{-3}$	$0.272... \times 10^{-3}$	$0.800... \times 10^{-3}$	$0.255... \times 10^{-5}$
3	$0.158... \times 10^{-3}$	$0.952... \times 10^{-3}$	$0.229... \times 10^{-7}$	$0.106... \times 10^{-9}$	$0.491... \times 10^{-8}$	$0.290... \times 10^{-16}$
4	$0.469... \times 10^{-7}$	$0.253... \times 10^{-5}$	$0.245... \times 10^{-15}$	$0.628... \times 10^{-29}$	$0.113... \times 10^{-23}$	$0.426... \times 10^{-49}$
5	$0.411... \times 10^{-14}$	$0.180... \times 10^{-10}$	$0.280... \times 10^{-31}$	$0.129... \times 10^{-86}$	$0.140... \times 10^{-70}$	$0.134... \times 10^{-147}$

Table 2 Values of the error $|z_k - (-0.9535... + 0.5600...i)|$, with $p = 8$, $w = 1 - 2i$ and $z_0 = -1 + 0.4i$

k	Error N_2	Error L_2	Error M_2	Error N_3	Error L_3	Error H_3
1	$0.220... \times 10^0$	$0.240... \times 10^0$	$0.144... \times 10^0$	$0.250... \times 10^0$	$0.260... \times 10^0$	$0.133... \times 10^0$
2	$0.306... \times 10^0$	$0.309... \times 10^0$	$0.646... \times 10^{-2}$	$0.885... \times 10^0$	$0.129... \times 10^1$	$0.100... \times 10^{-1}$
3	$0.171... \times 10^0$	$0.650... \times 10^0$	$0.199... \times 10^{-4}$	$0.531... \times 10^0$	$0.729... \times 10^2$	$0.434... \times 10^{-5}$
4	$0.755... \times 10^{-1}$	$0.236... \times 10^1$	$0.180... \times 10^{-9}$	$0.254... \times 10^0$	$0.676... \times 10^{46}$	$0.353... \times 10^{-15}$
5	$0.196... \times 10^{-1}$	$0.184... \times 10^3$	$0.147... \times 10^{-19}$	$0.684... \times 10^{-1}$	inf	$0.189... \times 10^{-45}$

Table 3 Values of the error $|z_k - 29^{1/11}|$, with $z_0 = 1.3$

k	Error N_2	Error L_2	Error M_2	Error N_3	Error L_3	Error H_3
1	$0.149... \times 10^{-1}$	$0.129... \times 10^{-1}$	$0.233... \times 10^{-2}$	$0.561... \times 10^{-2}$	$0.358... \times 10^{-2}$	$0.111... \times 10^{-2}$
2	$0.784... \times 10^{-3}$	$0.722... \times 10^{-3}$	$0.207... \times 10^{-5}$	$0.348... \times 10^{-5}$	$0.112... \times 10^{-5}$	$0.745... \times 10^{-8}$
3	$0.225... \times 10^{-5}$	$0.230... \times 10^{-5}$	$0.159... \times 10^{-11}$	$0.803... \times 10^{-15}$	$0.359... \times 10^{-16}$	$0.224... \times 10^{-23}$
4	$0.187... \times 10^{-10}$	$0.234... \times 10^{-10}$	$0.933... \times 10^{-24}$	$0.985... \times 10^{-44}$	$0.115... \times 10^{-47}$	$0.616... \times 10^{-70}$
5	$0.129... \times 10^{-20}$	$0.242... \times 10^{-20}$	$0.320... \times 10^{-48}$	$0.181... \times 10^{-130}$	$0.384... \times 10^{-142}$	$0.126... \times 10^{-209}$

7 Conclusion

We have presented a new method to generate higher-order iteration functions for computing the p th roots of a complex number. Among the infinitely many iteration functions that can be derived by this method, we can find well known iteration functions such as Newton, Halley and Schröder iterations. Some iteration functions not addressed previously in the literature can also be derived. Important convergence results that shed light on the problem of choosing an appropriated initial guess were proved as well. Our work raises a considerable number of questions that still need to be answered. For example, one needs to investigate if Corollary 3.1 can be extended to any j and to study the convergence behavior of the sequences (5.7). The complexity of the associated calculations compelled us to leave such study for a future work.

Acknowledgements The authors thank the referees for their valuable and insightful comments and suggestions. The second author acknowledges support from FCT (Portugal) through CMUP.

References

1. Amat, S., Bermúdez, C., Busquier, S., Leauthier, P., Plaza, S.: On the dynamics of some Newton's type iterative functions. *Int. J. Comput. Math.* **86**(4), 631–639 (2009)
2. Bini, D.A., Higham, N.J., Meini, B.: Algorithms for the matrix p th root. *Numer. Algorithms* **39**, 349–378 (2005)
3. Blanchard, P.: Complex analytic dynamics on the Riemann sphere. *Bull. AMS* **11**, 85–141 (1984)
4. Buff, X., Henriksen, C.: On König's root-finding algorithms. *Nonlinearity* **16**, 989–1015 (2003)
5. Comtet, L.: *Advanced Combinatorics: The Art of Finite and Infinite Expansions*. Reidel, Dordrecht (1974)
6. Curry, J.H., Garnett, L., Sullivan, D.: On the iteration of a rational function: computer experiments with Newton's method. *Commun. Math. Phys.* **91**, 267–277 (1983)
7. di Bruno, F.: *Traité de l'Élimination*. Paris (1859)
8. Fatou, P.: Sur les équations fonctionnelles. *Bull. Soc. Math. Fr.* **47**, 161–271 (1919); *Bull. Soc. Math. Fr.* **48**, 33–94, 208–314 (1920)
9. Guo, C.-H., Higham, N.J.: A Schur–Newton method for the matrix p th root and its inverse. *SIAM J. Matrix Anal. Appl.* **28**, 788–804 (2006)
10. Guo, C.-H.: On Newton's method and Halley's method for the principal p th root of a matrix. *Linear Algebra Appl.* **432**, 1905–1922 (2010)
11. Householder, A.S.: *The Numerical Treatment of a Single Nonlinear Equation*. McGraw-Hill, New York (1970)
12. Iannazzo, B.: On the Newton method for the matrix p th root. *SIAM J. Matrix Anal. Appl.* **28**, 503–523 (2006)
13. Iannazzo, B.: A family of rational iterations and its applications to the computation of the matrix p th root. *SIAM J. Matrix Anal. Appl.* **30**, 1445–1462 (2008)
14. Julia, G.: Memoire sur l'iteration des fonctions rationnelles. *J. Math. Pures Appl.* **8**, 47–245 (1918)
15. Lakić, S.: On the computation of the matrix k -th root. *Z. Angew. Math. Mech.* **78**, 167–172 (1998)
16. Laszkiewicz, B., Ziętak, K.: A Padé family of iterations for the matrix sector function and the matrix p th root. *Numer. Linear Algebra Appl.* **16**, 951–970 (2009)
17. Lin, M.: A residual recurrence for Halley's method for the matrix p th. *Linear Algebra Appl.* **432**, 2928–2930 (2010)
18. Riordan, J.: *Combinatorial Identities*. Wiley, New York (1968)

19. Roberts, G.E., Kobelski, J.H.: Newton's versus Halley's method: an approach via complex dynamics. *Int. J. Bifurc. Chaos* **14**, 1–17 (2004)
20. Schröder, E.: Ueber unendlich viele Algorithmen zur Auflösung der Gleichungen. *Math. Ann.* **2**, 317–365 (1870). English translation by G.W. Stewart: E. Schröder, On infinitely many algorithms for solving equations (1993)
21. Smith, R.A.: Infinite product expansions for matrix n -th roots. *J. Aust. Math. Soc.* **8**, 242–249 (1968)
22. Traub, J.F.: *Iterative Methods for Solution of Equations*. Prentice-Hall, Englewood Cliffs (1964)
23. Vrscay, E.R.: Julia sets and Mandelbrot-like sets associated with higher order Schröder rational iterative maps: a computer assisted study. *Math. Comput.* **46**, 151–169 (1986)