

Maximal Condorcet Domains

Vladimir I. Danilov · Gleb A. Koshevoy

Received: 29 April 2010 / Accepted: 8 September 2011 / Published online: 5 October 2011
© Springer Science+Business Media B.V. 2011

Abstract A Condorcet domain is a subset of the set of linear orders on a finite set of candidates (alternatives to vote), such that if voters preferences are linear orders belonging to this subset, then the simple majority rule does not yield cycles. It is well-known that the set of linear orders is the Bruhat lattice. We prove that a maximal Condorcet domain is a distributive sublattice in the Bruhat lattice. An explicit lattice formula for the simple majority rule is given. We introduce the notion of a symmetric Condorcet domain and characterize symmetric Condorcet domains of maximal size.

Keywords Weak Bruhat order · Distributive lattice · Binary plane tree · Simple majority rule

1 Introduction

A Condorcet domain is a set of linear orders on a finite set of candidates (alternatives to vote), such that if voters preferences are linear orders belonging to this set, then the simple majority rule does not yield cycles. We use the abbreviation CD for a Condorcet domain. A CD is *maximal* if it is not possible to add a linear order outside of the CD such that the expanded set is a CD.

In this paper we study normal CDs on the set $[n] = \{1, \dots, n\}$. A CD is *normal* if it contains the natural linear order $\alpha = (1 < 2 < \dots < n)$ and the opposite to it, the ‘anti-natural’ linear order $\omega = (n < n - 1 < \dots < 1)$.

V. I. Danilov (✉) · G. A. Koshevoy
Central Institute of Economics and Mathematics of RAS,
47, Nakhimovskii Prospect, 117418, Moscow, Russia
e-mail: vdanilov43@mail.ru, danilov@cemi.rssi.ru

G. A. Koshevoy
e-mail: koshevoy@cemi.rssi.ru

The first interesting example of a normal CD was constructed by Black [3]. Namely he showed that the set of single-peaked linear orders forms a normal CD. For other examples of CDs see [1, 4, 9]; Monjardet [11] gives a state of art of the theory. In [7] and then in [1, 4, 6], a connection between normal CDs and the lattice structure on the set \mathcal{LO} , so-called the Bruhat lattice, was revealed. Specifically, Chameni–Nembua in [4] shows that if a CD is a sublattice of the Bruhat lattice then it is a distributive sublattice.

In Theorem 1 we prove that any maximal normal CD is a sublattice of the Bruhat lattice. Next, we provide in Theorem 2 an explicit lattice polynomial formula for the simple majority rule when preferences of voters belong to a normal CD. In particular, this formula shows that the aggregated (due to the majority rule) linear order belongs to the sublattice generated by the preferences of voters. Hence, if preferences of all voters belong to a maximal normal Condorcet domain \mathcal{D} then the majority rule ordering belongs to \mathcal{D} as well.

We apply these results to study symmetric CDs. A CD is symmetric if it contains with any linear order the opposite linear order as well. We show that any symmetric CD on $[n]$ has at most 2^{n-1} elements. We obtain in Theorem 3 a complete description of symmetric CDs of maximal size 2^{n-1} . Describing the structure of maximal symmetric CDs of smaller size is an open problem. We show that, for every $2 \leq m < n$, there exists a maximal symmetric CD of size 2^m .

In Section 2 we recall the notion of weak Bruhat order on the set \mathcal{LO}_n of linear orders on $[n]$. In Section 3 we define a key notion of compatibility of linear orders, for the first time considered by Chameni–Nembua in [4]. We prove that any maximal clique (a maximal collection of pairwise compatible linear orders) is a distributive sublattice of the Bruhat lattice. In Section 4 an explicit lattice formula for the simple majority rule is given. As a consequence we obtain that a maximal normal CD is a distributive lattice. We also get a bijection between normal CDs and cliques established in [4]. Section 5 is devoted to general facts on symmetric CDs. A classification of maximal size symmetric CDs in term of plane binary trees is given in Section 6. In the last section we construct maximal symmetric CDs of size 4 for every $n \geq 3$.

2 Bruhat Lattice

Fix a natural number n and denote by $[n]$ the set $\{1, \dots, n\}$. A (strict) *linear order* on $[n]$ is a complete irreflexive transitive binary relation $<$ on $[n]$. One can identify a linear order $x_1 < x_2 < \dots < x_n$ with the word $x_1x_2\dots x_n$ of n different symbols from $[n]$. We denote by \mathcal{LO}_n or \mathcal{LO} the set of linear orders on $[n]$. Elements of \mathcal{LO} will be denoted by Greek letters like σ, τ and so on; we use also the notation $<_\sigma$. For a linear order σ we denote by σ° the opposite linear order, that is $x <_\sigma y$ if and only if $y <_{\sigma^\circ} x$. The set $[n]$ possesses two distinguished linear orders: the natural linear order $1 < 2 < \dots < n$, denoted as α , and the opposite to α , denoted as ω .

Let $\Omega = \Omega_n = \{(i, j), 1 \leq i < j \leq n\}$. A pair (i, j) from Ω is called an *inversion* of a linear order σ if $j <_\sigma i$, that is if σ inverts the natural relation $i < j$. The set of all inversions of σ is denoted $Inv(\sigma)$; it is a subset of Ω . For example, $Inv(\alpha)$ is the empty subset, whereas $Inv(\omega)$ is the whole Ω . In general, $Inv(\sigma^\circ) = \Omega - Inv(\sigma)$.

The mapping Inv embeds the set \mathcal{LO} into the set of subsets of Ω . To describe the image of this mapping we need some notions.

Definition 1 A subset S of Ω is *transitive* if $(i, j) \in S$ and $(j, k) \in S$ implies $(i, k) \in S$. A set $S \subset \Omega$ is *co-transitive* if $(i, k) \in S$ and $i < j < k$ implies either $(i, j) \in S$ or $(j, k) \in S$ (or both).

Note that the union of co-transitive sets is co-transitive, and the intersection of transitive sets is transitive.

Lemma 1 [13] *For a subset S of Ω the following assertions are equivalent:*

- (i) S is transitive and co-transitive (for short, *TcoT*);
- (ii) $S = \text{Inv}(\sigma)$ for some linear order σ .

The implication (ii) \Rightarrow (i) is obvious. The inverse implication is proved by an explicit construction of the corresponding linear order σ . Namely, we set $i <_{\sigma} j$ if either $i < j$ and $(i, j) \notin S$ or $j < i$ and $(j, i) \in S$. We leave to the reader to prove that σ is a linear order.

Thus, the set \mathcal{LO} of linear orders on $[n]$ can be identified with the set of TcoT-subsets in Ω . Because of this, one can compare linear orders via the inclusion of their inversion sets.

Definition 2 For linear orders σ and τ , we write $\sigma \ll \tau$ if $\text{Inv}(\sigma) \subseteq \text{Inv}(\tau)$. The relation \ll is called the *weak Bruhat order*.

The linear order α is the minimal element of the Bruhat poset (\mathcal{LO}, \ll) whereas ω is the maximal one. The poset (\mathcal{LO}, \ll) is a lattice indeed, see, for example [2, 8, 13]. It is convenient to use here the following lemma which remained unproved in [13].

Lemma 2 *Let S be a co-transitive subset in Ω . Then the transitive closure of S is co-transitive as well.*


Proof Let R be the transitive closure of S . By definition, a pair (i, i') belongs to R if there exists a chain $i = i_0 < i_1 < \dots < i_p = i'$ such that every neighbor pair (i_s, i_{s+1}) belongs to S .

Suppose now that $(i, k) \in R$ and $i < j < k$. We have to prove that either $(i, j) \in R$ or $(j, k) \in R$.

Let i_0, \dots, i_p be a chain as above which connects i and k . If j is one of the nodes of this chain then (i, j) and (j, k) belong to R . If not then j lies inside some link of the chain, $i_s < j < i_{s+1}$. The pair (i_s, i_{s+1}) belongs to S . Due to the co-transitivity of S , we have either $(i_s, j) \in S$ or $(j, i_{s+1}) \in S$. In the first case we obtain the chain $i = i_0 < \dots < j_s < j$ connecting i with j and giving $(i, j) \in R$. In the second case we have a chain from j to k and $(j, k) \in R$. □

For linear orders σ and τ , denote $S = \text{Inv}(\sigma)$ and $T = \text{Inv}(\tau)$. Let R be the transitive closure of $S \cup T$. Since $S \cup T$ is co-transitive, the set R is transitive and co-transitive due to Lemma 2. Therefore R is the inversion set for some linear order ρ . Obviously, $\sigma \ll \rho$ and $\tau \ll \rho$. Now, suppose that ρ' is a linear order such that $\sigma, \tau \ll \rho'$. Let $R' = \text{Inv}(\rho')$. Due to $\sigma \ll \rho'$ we have $S \subseteq R'$; similarly $T \subseteq R'$, hence $S \cup T \subseteq R'$. Since R' is a transitive set, R' contains the transitive closure of $S \cup T$ whence $\rho \ll \rho'$.

Fig. 1 The Bruhat lattice \mathcal{LO}_4



This proves the existence of the join $\sigma \vee \tau$ in the poset (\mathcal{LO}, \ll) . The existence of the meet follows by the formula $\sigma \wedge \tau = (\sigma^\circ \vee \tau^\circ)^\circ$.

Thus, the poset (\mathcal{LO}, \ll) is a lattice called the *Bruhat lattice* (Fig. 1). The operation $\sigma \mapsto \sigma^\circ$ is an ortho-complementation of the lattice, that is an antitone involution such that $\sigma \vee \sigma^\circ = \omega$ for every σ .

3 Chameni–Nembua Relation

In general case, the set $Inv(\sigma \vee \tau)$ is strictly larger than $Inv(\sigma) \cup Inv(\tau)$. Similarly the set $Inv(\sigma \wedge \tau)$ can be strictly smaller than $Inv(\sigma) \cap Inv(\tau)$.

Example 1 Let $n = 3, \sigma = 213, \tau = 132$. The linear order σ has one inversion $(1, 2)$, τ also has one inversion $(2, 3)$. However the set $\{(1, 2), (2, 3)\}$ is not transitive since it does not contain the pair $(1, 3)$. If we add this pair, we obtain $\sigma \vee \tau = \omega$. Similarly, $Inv(\sigma^\circ) \cap Inv(\tau^\circ)$ consists of the single pair $(1, 3)$ and is not co-transitive.

Nevertheless, sometimes the set $Inv(\sigma) \cup Inv(\tau)$ is transitive. In such a case we denote by $\sigma \cup \tau$ the join $\sigma \vee \tau$. Similarly, in the case of co-transitivity of the set $Inv(\sigma) \cap Inv(\tau)$, we denote by $\sigma \cap \tau$ the meet $\sigma \wedge \tau$. The following notion was introduced by Chameni–Nembua [4].

Definition 3 Linear orders σ and τ are *compatible* if the set $Inv(\sigma) \cup Inv(\tau)$ is transitive and the set $Inv(\sigma) \cap Inv(\tau)$ is co-transitive. The same terminology is applied to TcoT-sets as well.

For instance, σ and τ are compatible if $\sigma \ll \tau$. In particular, α and ω are compatible with any linear order. For any σ , the linear orders σ and σ° are compatible. Linear orders σ and τ are compatible if and only if σ° and τ° are compatible.

Lemma 3 *Let three linear orders ρ , σ , and τ be pairwise compatible. Then ρ is compatible with $\sigma \cup \tau$ and $\sigma \cap \tau$ and $\rho \cap (\sigma \cup \tau) = (\rho \cap \sigma) \cup (\rho \cap \tau)$.*

Proof We prove the compatibility of ρ and $\sigma \cup \tau$; the compatibility ρ and $\sigma \cap \tau$ is proved similarly. Let R , S , and T be the inversion sets for ρ , σ and τ . We have to check that the sets $R \cup (S \cup T)$ and $R \cap (S \cup T)$ are transitive and co-transitive.

The co-transitivity of $R \cup S \cup T$ is obvious. The transitivity follows from a simple observation. If two pairs (i, j) and (j, k) belong to $R \cup S \cup T$ then together they belong to at most two of the sets R , S or T , for instance, to R and T . But the union of R and T is transitive, hence $(i, k) \in R \cup T \subseteq R \cup S \cup T$.

The transitivity of the intersection $R \cap (S \cup T)$ follows from transitivity of R and $S \cup T$. The set $R \cap (S \cup T) = (R \cap S) \cup (R \cap T)$ is co-transitive as the union of two co-transitive sets $R \cap S$ and $R \cap T$. □

A *clique* is a subset in \mathcal{LO} which consists of pairwise compatible linear orders.

Because of Lemma 3, one can expand any clique by adding joins and meets of elements of the clique. In particular, if a clique is maximal (by inclusion) then it is a sublattice, moreover, a distributive lattice (since it is implemented by means of the intersection and union operations). This proves the following

Theorem 1 *Let \mathcal{C} be a maximal clique. Then \mathcal{C} is a distributive sublattice of the Bruhat lattice (\mathcal{LO}, \ll) .*

Example 2 Let us list all maximal cliques in the case $n = 3$. It was shown in Example 1 that there are two pairs of non-compatible linear orders: 213 and 132, and 231 and 312. All other pairs are compatible. Therefore a maximal clique must contain exactly one element from the first pair and one from the second pair. We obtain four maximal cliques:

- 1) 123, 132, 312, and 321. The alternative 2 is never the worst. This is the *peak* domain $\mathcal{D}_3(\cap)$.
- 2) 123, 213, 231, and 321. Here 2 is never the best; it is the *pit* domain $\mathcal{D}_3(\cup)$.
- 3) 123, 213, 312, and 321. Here 1 and 2 appear a tight group separated from the alternative 3. We denote this domain as $\mathcal{D}(\rightarrow)$.
- 4) 123, 132, 231, and 321. Here 1 is separated from 2 and 3, this is the domain $\mathcal{D}(\leftarrow)$.

For what follows, it is useful to establish properties of the compatibility relation under restrictions to subsets.

Let $\varphi : [m] \rightarrow [n]$ be a strictly increasing (hence, injective) mapping. Then the restriction mapping

$$\varphi^* : \mathcal{LO}_n \rightarrow \mathcal{LO}_m$$

is monotone with respect to weak Bruhat orders. Indeed, if σ is a linear order on $[n]$ then $Inv(\varphi^*(\sigma)) = \varphi^*(Inv(\sigma))$. Further, φ^* commutes with the orthocomplementations. Moreover, φ^* sends α_n in α_m and ω_n in ω_m . However, in general, φ^* does not commute with joins and meets.

Example 3 Let $\sigma = 213, \tau = 132$; then $\sigma \vee \tau = 321$. Under the restriction to the subset $\{1, 3\}$, σ is sent into 13, τ is sent into 31, and their join 321 differs from the restriction of 321 to $\{1, 3\}$ which is equal to 31. Similarly for the meet of 231 and 312.

However, there are at least two cases when the restriction φ^* commutes with \vee and \wedge . The first case is when φ is the natural inclusion of $[n - 1]$ into $[n]$. The second case is when σ and τ are compatible.

Proposition 1 *Let σ and τ be compatible linear orders on $[n]$. Then, for any strictly increasing mapping $\varphi : [m] \rightarrow [n]$, the linear orders $\varphi^*(\sigma)$ and $\varphi^*(\tau)$ are compatible. Moreover, $\varphi^*(\sigma \cup \tau) = \varphi^*(\sigma) \cup \varphi^*(\tau)$ and $\varphi^*(\sigma \cap \tau) = \varphi^*(\sigma) \cap \varphi^*(\tau)$.*

Proposition 1 follows from a simple remark: if $S \subseteq \Omega_n$ is a transitive (or co-transitive) set then $\varphi^*(S)$ is a transitive (respectively, co-transitive) set in Ω_m .

The reverse to this remark is partly true. Namely, if the intersection of a set $S \subset \Omega_n$ with every triple $\{i, j, k\}$ is transitive (co-transitive) then S is transitive (co-transitive). In particular, linear orders σ and τ are compatible if the restriction of σ and τ to every triple are compatible.

4 Cliques and Condorcet Domains

Recall that a Condorcet domain (CD) is a subset $\mathcal{D} \subseteq \mathcal{L}\mathcal{O}_n$ such that the simple majority rule does not yield cycles. Let us say this more precisely. A \mathcal{D} -opinion is a mapping $\nu : \mathcal{D} \rightarrow \mathbb{N} = \{0, 1, 2, \dots\}$. Intuitively, this means that $\nu(\sigma)$ voters have the preference σ on the set $[n]$ of alternatives. The number $|\nu| = \sum_{\sigma \in \mathcal{D}} \nu(\sigma)$ is equal to total number of voters. Let $sm(\nu)$ denote the binary relation (the ‘social preference’) on $[n]$ defined by the simple majority rule:

$$ism(\nu)j \text{ if and only if the number of voters which prefer } i \text{ to } j \text{ is strictly larger than the number of voters having opposite preference.}$$

If (obviously, asymmetric) relation $sm(\nu)$ is acyclic for every \mathcal{D} -opinion ν , the set \mathcal{D} is called a *Condorcet domain* (or an *acyclic set of linear orders*).

It is well known (see, for example, [11]) that in this definition one can consider only \mathcal{D} -opinions with odd number of voters. In this case the relation $sm(\nu)$ is complete (and is a linear order provided that \mathcal{D} is a CD). Below we give an explicit lattice polynomial expression for this linear order.

Let \mathcal{C} be a set of linear orders and ν be a \mathcal{C} -opinion. We say that a subset $A \subseteq \mathcal{C}$ is ν -admissible if $\nu(A) := \sum_{\sigma \in A} \nu(\sigma) > |\nu|/2$.

Theorem 2 *Let \mathcal{C} be a clique and let $|\nu|$ be odd. Then*

$$sm(\nu) = \bigvee_A \left(\bigwedge_{\sigma \in A} \sigma \right), \tag{1}$$

where A runs over ν -admissible subsets of \mathcal{C} .

Proof It suffices to check that the left and the right hand sides of Eq. 1 coincide under the restriction to arbitrary two-element subsets in $[n]$. Let $\varphi : [2] \rightarrow [n]$ be

a strictly increasing mapping. By definition, $\varphi^*(sm(v)) = sm(\varphi^*(v))$ (where $\varphi^*(v)$ is the corresponding recalculated opinion). Due to Proposition 1, $\varphi^*(\bigvee_A(\bigwedge A)) = \bigvee_A(\bigwedge \varphi^*(A))$. Thus, we have to check the equality 1 in the case $n = 2$. In this case there are only two linear orders: $\alpha = 12$ and $\omega = 21$. And an opinion v is given by two numbers: $v(\alpha)$ and $v(\omega)$, $|v| = v(\alpha) + v(\omega)$. Exactly one of these numbers is larger than $|v|/2$.

Suppose that $v(\alpha) > |v|/2$. Then $sm(v) = \alpha$. On the other hand, any v -admissible set A contains α , so that $\bigwedge A = \alpha$ and the join of all $\bigwedge A$ is equal to α as well.

Suppose now that $v(\omega) > |v|/2$. Then $sm(v) = \omega$. In this case the set $A_0 = \{\omega\}$ is admissible, and $\bigwedge A_0 = \omega$. Hence the join (over all admissible A) is equal to ω as well. □

Remark 1 For an arbitrary opinion $v : \mathcal{LO} \rightarrow \mathbb{N}$ (with odd $|v|$), the right hand side of the formula 1 defines an aggregation rule with values in \mathcal{LO} . This rule is a simple majority rule if the set $v^{-1}(\mathbb{N} \setminus \{0\})$ is a clique.

Corollary *Let \mathcal{C} be a maximal clique. Then, for every \mathcal{C} -opinion v with odd $|v|$, the social preference $sm(v)$ belongs to \mathcal{C} .*

Indeed, due to Theorem 1, \mathcal{C} is a sublattice of the Bruhat lattice. Therefore the right hand side of the above formula belongs to \mathcal{C} .

Remark 2 The simple majority rule is an anonymous rule in the case of aggregation of binary relations. For a so-called majority system \mathcal{F} (see [10]), one can define more general aggregation rule $a_{\mathcal{F}}: a_{\mathcal{F}}(R_1, \dots, R_N) = \bigcup_{A \in \mathcal{F}} (\bigcap_{i \in A} R_i)$. This rule has a form resembling the right hand side of Eq. 1. The same reasons as above show that if preferences $R_i, i = 1, \dots, N$, are linear orders and belong to a maximal clique \mathcal{C} then, for the aggregated preference $a_{\mathcal{F}}((R_i))$, there hold: i) it is a linear order, and ii) it belongs to \mathcal{C} . The assertion i) was proved in [10].

Recall that a subset $\mathcal{D} \subset \mathcal{LO}$ is *normal* if it contains α and ω . It follows from Theorems 2 that any maximal clique is a normal CD. Thus we get a half of the proof of the following proposition from [4].

Proposition 2 *Let \mathcal{C} be a subset of \mathcal{LO} containing α and ω . Then \mathcal{C} is a CD if and only if \mathcal{C} is a clique.*

Proof It remains to show that a normal CD is a clique. It is well-known (see, for example, [11]) that \mathcal{D} is a CD if and only if the restrictions of \mathcal{D} to every triple ijk ($i < j < k$) get into one of the domains $\mathcal{D}_3(\cap), \mathcal{D}_3(\cup), \mathcal{D}_3(\rightarrow)$ or $\mathcal{D}_3(\leftarrow)$ from Example 2. Because of the remark at the end of Section 3, the proposition follows. □

If a CD does not contain α and ω , then it might be not a clique, see Example 4. From Proposition 2 and Theorem 1 we obtain the following strengthening of a result of [1].

Corollary *Let \mathcal{D} be a maximal normal Condorcet domain. Then it is a distributive sublattice of the Bruhat lattice.*

We say that a Condorcet domain \mathcal{D} is *normalizable* if there exists a pair of opposite linear orders σ and σ° such that the set $\mathcal{D} \cup \{\sigma, \sigma^\circ\}$ is a CD as well. For normalizable CD all above results are valid for the reordered set $\{\sigma(1), \dots, \sigma(n)\}$. Thus, without loss of generality, one can assume that $\sigma = \alpha$ and consider normal CDs.

The class of normalizable Condorcet domains does not exhaust all Condorcet domains. Namely, the following example shows the existence of a non-normalizable CD.

Example 4 Let the domain \mathcal{D} in \mathcal{LO}_4 consists of four linear orders $\alpha = 1234$, $\beta = 2314$, $\gamma = 2413$ $\delta = 2143$. One can directly check that \mathcal{D} is a CD. We assert that this domain is not normalizable. In other words, for any pair σ, σ° of opposite linear orders, the domain $\mathcal{D} \cup \{\sigma, \sigma^\circ\}$ is not a CD.

For the proof we consider the restriction to the subset $\{1, 3, 4\}$, that is we delete the alternative 2. We obtain four orders 134, 314, 413 and 143. These orders form a maximal CD on the set $\{1, 3, 4\}$. From this observation, one can see that, after deleting 2, the pair σ, σ° has to be the pair 314, 413. Wlog we assume that the restriction of σ is 314. Thus, we have to examine four possibility for σ : 2314, 3214, 3124 and 3142.

- 1) $\sigma = 2314$ or 3214 . Delete the alternative 3 (and denote the restriction by $'$). Then $\sigma'^\circ = 412, \alpha' = 124, \delta' = 241$ form a cyclic triple. Therefore $\mathcal{D} \cup \{\sigma^\circ\}$ is not a CD.
- 2) $\sigma = 3124$ or 3142 . Deleting the alternative 4 we obtain the following cyclic triple $\sigma' = 312, \alpha' = 123, \beta' = 231$. Therefore $\mathcal{D} \cup \{\sigma^\circ\}$ is not a CD.

5 Symmetric Condorcet Domains

A large class of the so-called tiling type (or peak-pit) CDs was considered in [5] (see also [1, 6, 9]). A tiling type CD is a normal CD and it does not contains other pairs of opposite linear orders, except α and ω . Here we present a kind of a complementary class of normal CDs (Fig. 2).

Definition 4 A Condorcet domain \mathcal{D} is called *symmetric* if, for every linear order $\sigma \in \mathcal{D}$, the opposite linear order σ° belongs to the domain \mathcal{D} as well.




Fig. 2 Black circles depict three CDs in the lattice \mathcal{LO}_4 (see Fig. 1). On the left is a tiling CD, in the middle is the symmetric CD $((1 * 2) * 3) * 4$ from Section 6, on the right is the maximal CD of size 4 from Section 7

In other words, $\mathcal{D}^\circ = \mathcal{D}$. Domains $\mathcal{D}_3(\rightarrow)$ and $\mathcal{D}_3(\leftarrow)$ are examples of symmetric CDs. Of course, a symmetric CD is normalizable. In what follows it is convenient to consider normal symmetric CDs. We do not loss generality under this assumption.

Suppose that \mathcal{D} is a normal symmetric CD, that is a symmetric clique (by virtue of Proposition 2). If a linear order σ is compatible with every elements of \mathcal{D} then its opposite order σ° is also compatible with every element of \mathcal{D} (and with σ). Therefore any symmetric CD can be extended to a maximal and symmetric CD.

How to construct symmetric CDs? Suppose we have a partition $\Omega = S_1 \sqcup \dots \sqcup S_t$ (\sqcup denotes disjoint union). Denote by $\mathcal{B}(S_1, \dots, S_t)$ the Boolean algebra of subsets of Ω generated by S_1, \dots, S_t . It consists of $S(I) = \cup_{i \in I} S_i$, where I runs over subsets of $[t]$.

Lemma 4 *In these notations suppose that the sets $S_i, i = 1, \dots, t$, are co-transitive. Then any element $S(I)$ of the Boolean algebra $\mathcal{B}(S_1, \dots, S_t)$ is a TcoT-set.*

Indeed, any set $S(I)$ is co-transitive as the union of co-transitive sets S_i and is transitive as the complement (in Ω) to the co-transitive set $S([t] - I)$.

In particular, the Boolean algebra $\mathcal{B}(S_1, \dots, S_t)$ consists of pairwise compatible sets and, due to Proposition 2, defines a CD, which is denoted by the same symbol. Since $\mathcal{B}(S_1, \dots, S_t)$ is stable with respect to $^\circ$, this CD is symmetric. It can be non-maximal if there exists a finer partition. But any maximal symmetric CD \mathcal{D} has such a form. Indeed, due to Theorem 1 and Proposition 2, \mathcal{D} is a distributive sublattice of the Bruhat lattice. By virtue of symmetry, \mathcal{D} is ortho-complemented. Therefore \mathcal{D} is a Boolean lattice. Let $\sigma_1, \dots, \sigma_t$ be atoms of the Boolean lattice, and $S_i = Inv(\sigma_i), i = 1, \dots, t$, be the corresponding inversion sets. Obviously, the S_i 's do not intersect and moreover they cover Ω .

Proposition 3 *The size of a symmetric CD does not exceed 2^{n-1} .*

Proof We can assume that the symmetric CD \mathcal{D} is maximal. Therefore it has the form $\mathcal{B}(S_1, \dots, S_t)$ for some partition of Ω by TcoT-sets, and its size is equal to 2^t . It remains to remark that every (non-empty) co-transitive set $S \subseteq \Omega$ contains a ‘short arrow’, that is a pair of the form $(i, i + 1)$. Therefore every set S_r contains a ‘short arrow’. Since there are $n - 1$ ‘short arrows’, we conclude that $t \leq n - 1$. □

6 Structure of Symmetric Condorcet Domains of Maximal Size

From Proposition 3, we know that the size of a symmetric CD is less or equal to 2^{n-1} . Therefore any symmetric CD of size 2^{n-1} is maximal. Here we give a complete description of such CDs. This description uses the following simple construction (a particular case of the more general block construction from [9]).

Let $1 < m < n$, and suppose we have two normal CDs, \mathcal{D} on the set $[m]$ and \mathcal{D}' on the set $[m'] = [n - m]$. Denote by $\mathcal{D} * \mathcal{D}'$ the set of linear orders on $[n]$ of the form $\sigma\sigma'$ or $\sigma'\sigma$, where $\sigma \in \mathcal{D}, \sigma' \in \mathcal{D}'$. Here we consider σ and σ' as words in the alphabet $1, \dots, m$ and $m + 1, \dots, n$ respectively. We regard the set $[m']$ as the set of the last $n - m$ symbols in $[n]$.

Lemma 5 $\mathcal{D} * \mathcal{D}'$ is a normal CD on $[n]$.

Intuitively, it is quite clear. Suppose we have a $\mathcal{D} * \mathcal{D}'$ -opinion. At first we decide whether to put the first m alternatives above or below the last m' alternatives. Then we rank the first m alternatives (because \mathcal{D} is a CD) and similarly for the last m' alternatives (because \mathcal{D}' is a CD).

One can argue more formally using Proposition 2. We have to check that every two linear orders from $\mathcal{D} * \mathcal{D}'$ are compatible. Here four cases are possible. For instance, $\sigma\sigma'$ and $\tau\tau'$, or $\sigma\sigma'$ and $\tau'\tau$. In any case, it is useful to consider the inversion sets for $\sigma\sigma'$ and $\sigma'\sigma$. Let $S = \text{Inv}(\sigma)$, $S' = \text{Inv}(\sigma')$. Then $\text{Inv}(\sigma\sigma') = S \cup S'$ whereas $\text{Inv}(\sigma'\sigma) = S \cup S' \cup P$, where $P = \{(i, j), i \leq m, j \geq m + 1\}$. Consider, for example, the join of $\sigma\sigma'$ and $\tau'\tau$ (setting $T = \text{Inv}(\tau)$ and $T' = \text{Inv}(\tau')$). Let us prove that $(S \cup S') \cup (T \cup T' \cup P)$ is transitive (the co-transitivity is obvious). The set $S \cup T$ is transitive due to compatibility of σ and τ ; $S' \cup T'$ is transitive due to compatibility of σ' and τ' . The union of these transitive sets is transitive because, for any pair of arrows from S and T , the end of the arrow from S cannot be the origin of the arrow from T , and vice versa. Adding P does not violate the transitivity. Similarly with the other cases.

Lemma 6 Let \mathcal{D} and \mathcal{D}' be symmetric CDs. Then $\mathcal{D} * \mathcal{D}'$ is a symmetric CD.

This follows from $(\sigma\sigma')^\circ = \sigma' \circ \sigma^\circ$.

Proposition 4 Let \mathcal{D} and \mathcal{D}' be maximal normal CDs. Then $\mathcal{D} * \mathcal{D}'$ is a maximal normal CD.

Proof Suppose that a linear order ρ is compatible with all elements from $\mathcal{D} * \mathcal{D}'$. Then it is compatible with the linear order $\omega_m\omega_{m'}$ whose inversion set is equal to $\Omega_m \cup \Omega_{m'}$. In particular, $\text{Inv}(\rho) \cup (\Omega_m \cup \Omega_{m'})$ must be transitive. Further, due to compatibility of ρ and $\alpha_{m'}\alpha_m$, the intersection $\text{Inv}(\rho) \cap P$ must be co-transitive. We assert that it is possible only in two cases: either $\text{Inv}(\rho)$ contains P or $\text{Inv}(\rho)$ does not intersect P .

Namely, suppose that $R = \text{Inv}(\rho)$ contains a pair of the form (i, j) , where $i \leq m$, $j \geq m + 1$. Since the set $R \cap P$ is co-transitive, it contains the short arrow $(m, m + 1)$. The set Ω_m contains any arrow (i, m) ($i \leq m$), the set $\Omega_{m'}$ contains any arrow $(m + 1, j)$ ($j \geq m + 1$). Due to the transitivity, $R \cup \Omega_m \cup \Omega_{m'}$ contains all arrow of the form (i, j) , that is $R \supseteq P$.

Thus, either P does not intersect R or $P \subseteq R$. In the first case ρ has the form $\tau\tau'$, where τ is a linear order on $[m]$ and τ' is a linear order on $[m']$. In the second case ρ has the form $\tau'\tau$. Since τ is compatible with every element of \mathcal{D} , $\tau \in \mathcal{D}$ due to maximality of \mathcal{D} . Similarly $\tau' \in \mathcal{D}'$. \square

Let us consider a particular case of the construction when $m = n - 1$. In this case we take \mathcal{D}' as the set of linear orders on the singleton $\{n\}$. This set contains a single element denoted by n (we hope that this misuse of language does not yield

a confusion). In this case the set $\mathcal{D} * \mathcal{D}' = \mathcal{D} * n$ consists of linear orders of the form σn or $n\sigma$, where $\sigma \in \mathcal{D}$. If \mathcal{D} is a symmetric (respectively, maximal) CD on $[n - 1]$ then $\mathcal{D} * n$ is a symmetric (respectively, maximal) CD on $[n]$ and doubles the size of \mathcal{D} .

This construction can be iterated. In particular, one can define by induction a CD for any binary parenthesization of the expression $1 * 2 * \dots * n$. For instance, $(1 * (2 * 3)) * (4 * 5)$ is one of 14 possible parenthesizations of the set [5]. Here the symbol i is considered as a unique linear order on the set $\{i\}$. (It is well-known that the set of parenthesizations is bijective to the set of plane binary tree with n leaves. Exercise 6.19 in Stanley’s book [12] contains yet 64 interpretations of parenthesizations.)

Example 5 Obviously, $1 * 2 = \mathcal{LO}_2$. The domain $(1 * 2) * 3$ is exactly the Condorcet domain on the set [3], in which the alternative 3 is never middle, that is the domain $\mathcal{D}_3(\rightarrow)$ from Example 2. Similarly $1 * (2 * 3)$ is the domain $\mathcal{D}_3(\leftarrow)$.

Example 6 Let us consider in detail the case $(\dots((1 * 2) * 3)\dots) * n$. Linear orders from this domain are characterized by the following property: every alternative i either is better than any element from the set $[i - 1]$ or is worse than any element from it. N.S.Kukushkin proposed the following interpretation of such linear orders. Let us consider alternatives as ‘reform projects’, increasingly ordered by their degree of ‘radicalism’. Then every project is perceived either better than all of the project that preceded it, or worse than all of them. This domain is somewhat similar to the Black’s domain of single-peaked preferences. As for the single-peaked domain, there is a simple inductive procedure for aggregation of these preferences. At first, we compare the alternatives n and $n - 1$. If n wins in the comparison, it becomes the best in the group sense. In the opposite case it is worst. Next we compare $n - 1$ and $n - 2$, and so on.

Similarly one can aggregate the preferences for any CD, produced by a parenthesization of $1 * 2 * \dots * n$.

Let \mathcal{P} be a binary parenthesization of $1 * 2 * \dots * n$, and $\mathcal{D}(\mathcal{P})$ be the corresponding CD. Then $\mathcal{D}(\mathcal{P})$ is a symmetric CD of size 2^{n-1} , hence is a maximal CD. We assert that the converse is also true.

Theorem 3 *Let \mathcal{D} be a symmetric Condorcet domain of size 2^{n-1} . Then it has the form $\mathcal{D}(\mathcal{P})$ for some parenthesization \mathcal{P} of $1 * 2 * \dots * n$.*

We need the following

Lemma 7 *Let P be a co-transitive set in Ω , which contains the ‘long arrow’ $(1, n)$ and a unique short arrow $(m, m + 1)$. Then $P = \{(i, j), i \leq m < j\}$.*

Proof First of all, $P \subseteq \{(i, j), i \leq m < j\}$. Indeed, if P contains an arrow (i, j) with $i, j \leq m$ then it contains a short arrow of the same form. This contradicts the uniqueness of the short arrow $(m, m + 1)$. Similarly for $i, j > m$.

On the other hand, every arrow (i, j) with $i \leq m, j \geq m + 1$ belongs to P . Indeed, applying the definition of co-transitivity to the arrow $(1, n)$ and the symbol i , we obtain that either $(1, i) \in P$ or $(i, n) \in P$. The first case is impossible, since $P \subseteq \{(i, j), i \leq m < j\}$. Therefore $(i, n) \in P$. Applying again the definition of co-transitivity to the arrow (i, n) and the symbol j , we obtain $(i, j) \in P$. \square

Proof of Theorem 3 It follows from Section 5 that \mathcal{D} is defined by a partition $\Omega = S_1 \sqcup \dots \sqcup S_{n-1}$ consisting of $n - 1$ TcoT-sets S_1, \dots, S_{n-1} . One of them (say, S_1) contains the long arrow $(1, n)$. Moreover, S_1 (as well as all other S_r) contains a unique short arrow $(m, m + 1)$. Due to Lemma 7, $S_1 = \{(i, j), i \leq m < j\}$. This gives us the first decomposition of $[n]$ on two sub-intervals: $[m]$ and $[m + 1..n]$, and this corresponds to parenthesization $(1\dots m)(m + 1\dots n)$. Now we repeat the previous arguments to the interval $[m]$ and decompose it into two sub-intervals. Similarly for the interval $[m + 1..n]$, and so on. This process leads us to a parenthesization of $1 * 2 * \dots * n$ and completes the proof. \square

7 One More Example


Here we show that (for any $n \geq 3$) there exists a maximal symmetric CD $\mathcal{Q}_n \subseteq \mathcal{LO}_n$ of size 4. It consists of four linear orders α, ω, σ and σ° , where σ has the form $24\dots(2k)1(2k \pm 1)\dots 53$ (here $2k \pm 1$ is equal to $2k + 1 = n$, if n is odd, and $2k - 1 = n - 1$ if n is even). In other words, at first even symbols go in increasing order, further the symbol 1 stays, and then odd symbols (up to 3) go in decreasing order. For example, $\sigma = 2461753$ for $n = 7$, whereas $\sigma = 24681753$ for $n = 8$. The opposite order σ° is arranged similarly: at first odd symbols (beginning with 3) go, further 1, and next even symbols go in decreasing order.

Proposition 5 *The domain $\mathcal{Q}_n = \{\alpha, \sigma, \sigma^\circ, \omega\}$ is a maximal Condorcet domain.*

Proof Let $S = \text{Inv}(\sigma)$; this set is depicted by black circles in the Fig. 3.

Suppose that the domain \mathcal{Q}_n is not maximal. Then there exists a TcoT-set T (corresponding to a linear order which can be added to \mathcal{Q}_n) which is compatible with S and $\Omega - S$ (and differs from them). Passing to $T \cap S$ or to $T \cap (\Omega - S)$, we can

Fig. 3 The picture is slightly different for even and odd n . Here $n = 8$



suppose that T is contained in S or in $\Omega - S$. We consider in detail the case when the nonempty TcoT-set T is contained in $\Omega - S$ and $T \cup S$ is transitive (the other case is considered similarly); we have to show that T coincides with $\Omega - S$.

We will argue by induction. The induction base, $n = 3$, is obviously true. Therefore we assume $n \geq 4$. We will denote the reduction of the symbol n (or restriction on $[n - 1]$) by a prime. By induction the domain $\mathcal{D}' = \{\alpha', \omega', \sigma', (\sigma')^\circ\}$ is maximal.

We claim that T' is non-empty as well. Indeed, if T' is empty then T consists only of arrows of the form (i, n) . Moreover, T is co-transitive and, hence, contains the short arrow $(n - 1, n)$. Since $T \subseteq \Omega - S$, this is possible only at odd n . (In this case i has to be an even symbol or the symbol 1.) Further, $(1, n - 1) \in S$; due to transitivity of $T \cup S$ we obtain that $(1, n) \in T \cup S$. Since $(1, n) \notin S$, we have $(1, n) \in T$. Since T is co-transitive, we obtain $(3, n) \in T$, in contradiction with $(3, n) \in S$.

Thus, T' is non-empty. By the inductive assumption T' coincides with $\Omega' - S$. Therefore $T \cup S$ contains whole $\Omega' = \Omega_{n-1}$. In other words, every arrow (i, j) belongs to $T \cup S$ provided that $j \neq n$. Moreover, the set $T \cup S$ is transitive. Recall we should show that $T = \Omega - S$. We consider the cases of even and odd n separately.

1. n is odd. In this case $(1, 3) \in \Omega' \subseteq T \cup S$ and $(3, n) \in S$; due to transitivity of $T \cup S$ we obtain $(1, n) \in T \cup S$. But $(1, n) \notin S$ and hence $(1, n) \in T$. Let now i be an even symbol. It lies between 1 and n ; due to co-transitivity of T we obtain that either $(1, i) \in T$ or $(i, n) \in T$. The first is impossible, since then $(1, i) \in \Omega - S$ that is not the case. Therefore, for every even i we have $(i, n) \in T$. Together with $(1, n) \in T$ this gives the equality $T = \Omega - S$.
2. n is even. In this case $(2, 3) \in \Omega' \subseteq T \cup S$ and $(3, n) \in S$; the transitivity of $T \cup S$ implies $(2, n) \in T \cup S$. But $(2, n) \notin S$, hence $(2, n) \in T$. Again, let i be an even symbol (greater than 2). It lies between 2 and n ; due to the co-transitivity of T , we have either $(2, i) \in T$ or $(i, n) \in T$. The first is impossible since then $(2, i) \in T \subseteq \Omega - S$, that is not the case. Thus, for every even symbol i we have $(i, n) \in T$, which gives $T = \Omega - S$.

□

Now one can easily construct a maximal (and symmetric) CD of the size 2^m for any m , $2 \leq m < n$. For this, one takes a domain of the form $\mathcal{Q}_{n-m+2} * ((\dots(n - m + 3) * \dots) * n)$. For instance, at $n = 5$ the CD $\mathcal{Q}_4 * 5$ has the size 8. It would be interesting to find a structure which describes maximal symmetric CDs of any size.

Acknowledgements We thank the referees and the action-editor for useful remarks and suggestions. This work is partially supported by the Russian Foundation of Basic Research and the Ministry of National Education of France (project RFBR 10-01-9311-CNRS-L-a).

References

1. Abello, J.M.: The weak Bruhat order on S_n , consistent sets, and Catalan numbers. *SIAM J. Discrete Math.* **4**, 1–16 (1991)
2. Björner, A., Brenti, F.: *Combinatorics of Coxeter Groups*. Graduate Texts in Mathematics, vol. 231. Springer (2005)
3. Black, D.: On the rationale of group decision-making. *J. Polit. Econ.* **56**, 23–34 (1948)
4. Chameni-Nembua, C.: Règle majoritaire et distributivité dans le permutoèdre. *Math. Inform. Sci. Hum.* **108**, 5–22 (1989)

5. Danilov, V., Karzanov, A., Koshevoy, G.: Condorcet domains and rhombus parquets. *Èkon. Mat. Metody* **46**(4), 55–68 (2010, in Russian)
6. Galambos, A., Reiner, V.: Acyclic sets of linear orders via the Bruhat order. *Soc. Choice Welf.* **30**, 245–264 (2008)
7. Guilbaud, G.-Th.: Les théories de l'intérêt général et le problème logique de l'agrégation. *Écon. Appliquée* **5**, 501–584 (1952)
8. Guilbaud, G.-Th., Rosenstiehl, P.: Analyse algébrique d'un scrutin. *Math. Sci. Hum.* **4**, 9–33 (1963)
9. Fishburn, P.: Acyclic sets of linear orders. *Soc. Choice Welf.* **14**, 113–124 (1997)
10. Monjardet, B.: An axiomatic theory of tournament aggregation. *Math. Oper. Res.* **3**, 334–351 (1978)
11. Monjardet, B.: Acyclic domains of linear orders: a survey. In: Brams, S., Gehrlein, W., Roberts, F. (eds.) *The Mathematics of Preference, Choice and Order*, pp. 136–160. Springer (2009)
12. Stanley, R.: *Enumerative Combinatorics*, vol. 2. Cambridge University Press (1999)
13. Yanagimoto, T., Okamoto, M.: Partial orderings of permutations and monotonicity of a rank correlation statistic. *Ann. Inst. Stat. Math.* **21**, 489–506 (1969)