

Rutin Suppresses Palmitic Acids-Triggered Inflammation in Macrophages and Blocks High Fat Diet-Induced Obesity and Fatty Liver in Mice

Mingming Gao · Yongjie Ma · Dexi Liu

Received: 23 March 2013 / Accepted: 12 June 2013 / Published online: 20 June 2013
© Springer Science+Business Media New York 2013

ABSTRACT

Purpose To elucidate the mechanism of rutin in blocking macrophage-mediated inflammation and high fat diet-induced obesity and fatty liver.

Methods Both *in vitro* and *in vivo* approaches were taken in evaluating the effects of rutin on palmitic acids-triggered inflammation in cultured macrophages, and on weight gain and development of fatty liver of mice fed a high fat diet.

Results Palmitic acids increase mRNA levels of pro-inflammatory cytokines, and elevate the production of TNF α in cultured macrophages. Pre-exposure of rutin to cells greatly suppressed these elevations. The suppressed inflammation by rutin was correlated with a decrease in transcription of genes responsible for ER stress and production of reactive oxygen species. *In vivo*, rutin protects mice from high fat diet-induced obesity, fatty liver and insulin resistance. The protective effects were associated with lack of hypertrophy and crown-like structures in the white adipose tissue, decreased mRNA levels of marker genes for macrophages including *F4/80*, *Cd11c* and *Cd68*, and repressed transcription of genes involved in chronic inflammation such as *Mcp1* and *Tnfa* in white adipose tissue. In addition, rutin increases the expression of genes responsible for energy expenditure in brown adipose tissue including *Pgc1 α* and *Dio2*. Furthermore, rutin suppresses transcription of *Srebp1c* and *Cd36* in the liver, leading to a blockade of fatty liver development.

Conclusion These results suggest that supplementation of rutin is a promising strategy for blocking macrophage-mediated inflammation and inflammation-induced obesity and its associated complications.

KEY WORDS anti-oxydant · chronic inflammation · fatty liver · high fat diet · insulin resistance · obesity · rutin

INTRODUCTION

Along with changes in life style and diet structure, the prevalence of obesity is increasing globally. Many strategies have been attempted to control obesity and to prevent complications associated with it, including fatty liver, insulin resistance and cardiovascular diseases. Among strategies, supplementation of natural anti-oxidants is a promising one. For example, rutin, a potent natural anti-oxidant, has been demonstrated being capable of producing beneficial effects on several metabolic disorders (1–4). It has been reported that rutin suppresses high fat diet (HFD)-induced dyslipidemia in rats (1). Panchal *et al.* showed that rutin protects rats from the metabolic changes and cardiovascular remodeling caused by a high-fat and high-carbohydrate diet (2). Additionally, Hao *et al.* and Kamalakkannan *et al.* demonstrated that rutin improves the metabolic status of diabetic rats (3,4). Similarly, chronic inflammation has been indicated a critical role in development of metabolic disorders (5,6). For example, Henao-Mejia *et al.* reported that chronic inflammation is essential for progression of obesity and non-alcoholic fatty liver (5). A more recent study by Stienstra *et al.* identified a crucial function of inflammasome in pathogenesis of obesity and insulin resistance (6). Importantly, Hu *et al.* demonstrated that rutin ameliorates inflammasome activation, leading to suppressed inflammation in the kidney of rats (7). Similarly, Javed *et al.* reported the activity of rutin in preventing cognitive impairments in rats (8). Results from these studies prompt us to hypothesize that the beneficial effect of rutin in obesity is achieved through suppression of chronic inflammation.

Employing both *in vitro* and *in vivo* systems, we demonstrate in this study that rutin suppresses palmitic acid-triggered

M. Gao · Y. Ma · D. Liu (✉)

Department of Pharmaceutical and Biomedical Sciences
College of Pharmacy, University of Georgia, 450 Pharmacy South
250 West Green Street, Athens, Georgia 30602, USA
e-mail: dliu@uga.edu

inflammation in macrophages, and twice weekly injections of rutin protect mice from development of HFD-induced obesity, fatty liver and insulin resistance. These beneficial effects are correlated with blockade of macrophage infiltration and chronic inflammation in adipose tissue. Our results offer new and critical information regarding the beneficial effects of rutin in the management of metabolic disorders.

MATERIALS AND METHODS

Animals and Treatments

The animal protocol was approved by the Institutional Animal Care and Use Committee of the University of Georgia (Protocol Number, A2011 07-Y2-A3). Male C57BL/6 mice purchased from Charles River (Wilmington, MA) were housed under standard conditions with a 12 h light–dark cycle. Mice were divided into 3 groups ($n=5$), and animals in group one were intraperitoneally (*i.p.*) injected twice weekly with rutin (50 mg/kg) solubilized in DMSO and fed a HFD (60% calories from fat, 20% from carbohydrates, and 20% from proteins). Mice in group two and three were injected (*i.p.*) with carrier solution fed a HFD or regular chow (20% calories from fat, 60% from carbohydrates, and 20% from proteins), respectively. Body weight and food-intake were measured weekly. Body composition analysis using EchoMRI-100 (Echo Medical Systems, Houston, TX) was conducted at the end of treatment.

Cell Culture

Murine macrophage cell line (RAW 264.7, TIB-71 at ATCC) was cultured at 37°C in a 5% CO₂-humidified incubator and grown in Dulbecco's minimum essential medium (DMEM) supplemented with 10% fetal bovine serum, penicillin (100 U/ml) and streptomycin (100 µg/ml). Cell viability was measured by MTT assay. For TNF α determination in supernatants, cells were pre-treated with various concentrations of rutin for 4 h and further incubated with 400 µM palmitic acids with or without rutin for 12 h. Supernatants were collected and TNF α levels were determined using an enzyme-linked immunosorbent assay (ELISA) kit (#88-7324-88) purchased from eBioscience (San Diego, CA). In the gene expression analysis, cells were pre-treated with or without 200 µM rutin for 4 h and further incubated for 12 h with 400 µM palmitic acids with or without the presence of rutin (200 µM).

Measurement of Reactive Oxygen Species (ROS)

ROS was determined using the Total ROS Detection Kit (ENZ-51011) from Enzo Life Sciences (Farmingdale, NY).

Cells were pre-treated with or without 200 µM rutin for 4 h and further incubated with 400 µM palmitic acids with or without 200 µM rutin for 24 h. Cells were washed with phosphate buffered saline (pH=7.2) and incubated with the detection reagent from the kit for 30 min at 37°C. Cells were examined under a fluorescent microscope (ECLIPSE Ti, Nikon) and fluorescent intensity from at least 5 randomly selected fields was measured and normalized to cell numbers using the NIS-Elements imaging platform from Nikon Instruments Inc. (Melville, NY).

Intra-Peritoneal Glucose Tolerance Test (IPGTT) and Insulin Tolerance Test (ITT)

Mice were fasted for 6 h before IPGTT. Glucose solubilized in phosphate buffered saline was injected (*i.p.*) at 1.5 g/kg, and the time-point was set as 0 min. Blood glucose levels at 0, 30, 60 and 120 min were measured using glucose test strips and glucose meters. For ITT, mice were fasted for 4 h before the injection of insulin (Humulin, 0.75 U/kg) from Eli Lilly (Indianapolis, IN), and blood glucose levels were measured at 0, 30, 60 and 120 min after insulin injection.

Gene Expression Analysis

Total mRNA from the liver was isolated using the TRIZOL reagent from Invitrogen (Grand Island, NY). A commercial kit (#74804) from QIAGEN (Valencia, CA) was used for mRNA isolation from adipose tissue. Reverse transcription polymerase chain reaction (RT-PCR) was performed using a Superscript RT III enzyme kit (#11752-050) from Invitrogen. Quantitative real time PCR (qPCR) was conducted using SYBR Green as detection reagent and the data were analyzed using the $\Delta\Delta C_t$ method (9). GAPDH mRNA level was used as an internal control, and all primer sequences used are listed in Table I. Primers were synthesized at Sigma (St. Louis, MO). Melting curve analysis of all real time PCR products were conducted and showed a single DNA duplex.

Determination of Blood Concentrations of Insulin, Aspartate Aminotransferase (AST) and Alanine Aminotransferase (ALT)

Blood samples were collected from heart cavities immediately after euthanizing mice. Serum was isolated by centrifugation at 4,000 rpm for 5 min. Insulin levels in the blood were measured using a commercial ELISA kit (10-1113-01, Mercodia Developing Diagnostics). AST (TR70121) and ALT (TR71121) concentrations were determined following manufacture's instruction (Thermo Scientific, Middletown, VA).

Table 1 PCR Primer Sequences

Gene name	Forward primer sequence (F)	Reverse primer sequence (R)
<i>Acc</i>	F: ATGGGCGAATGGTCTCTTTC	R: TGGGGACCTTGTCTTCATCAT
<i>Arg1</i>	F: CTCCAAGCCAAAGTCCTTAGAG	R: AGGAGCTGTCATTAGGGACATC
<i>Cd11c</i>	F: CTGGATAGCCTTTCTTCTGCTG	R: GCACACTGTGTCCGAACCTCA
<i>Cd36</i>	F: ATGGGCTGTGATCGGAACT	R: GTCTTCCCAATAAGCATGTCTCC
<i>Cd68</i>	F: TGTCTGATCTTGCTAGGACCG	R: GAGAGTAACGGCCTTTTTGTGA
<i>Cd206</i>	F: CTCTGTTTCAGCTATTGGACGC	R: CGGAATTTCTGGGATTGAGCTTC
<i>Chop</i>	F: CTGGAAGCCTGGTATGAGGAT	R: CAGGGTCAAGAGTAGTGAAGGT
<i>Cidea</i>	F: TGACATTTCATGGGATTGCAGAC	R: GGCCAGTTGTGATGACTAAGAC
<i>Cpt1α</i>	F: CTCCGCCTGAGCCATGAAG	R: CACCAGTGATGATGCCATTCT
<i>Cpt1β</i>	F: GCACACCAGGCAGTAGCTTT	R: CAGGAGTTGATTCCAGACAGGT
<i>Dio2</i>	F: AATTATGCCTCGGAGAAGACCG	R: GGCAGTTGCCTAGTGAAGGT
<i>Edem1</i>	F: AGTCAAATGTGGATATGCTACGC	R: ACAGATATGATATGGCCCTCAGT
<i>Elovl3</i>	F: TTCTCACGCGGGTAAAAATGG	R: GAGCAACAGATAGACGACCAC
<i>F4/80</i>	F: TGACTCACCTTGTGGTCCTAA	R: CTTCCCAGAAATCCAGTCTTTCC
<i>Fas</i>	F: GGAGGTGGTG ATAGCCGGTAT	R: TGGGTAATCCATAGAGCCCAG
<i>G6p</i>	F: CGACTCGCTATCTCCAAGTGA	R: GTTGAACCACTCTCCGACCA
<i>Glut4</i>	F: GTGACTGGAACACTGGTCTTA	R: CCAGCCACGTTGCATTGTAG
<i>Gpr78</i>	F: CAGATCTTCTCCACGGCTTC	R: AGCAGGAGGAATTCCAGTCA
<i>Gpr94</i>	F: ATGCTCGGGACAATCAACACC	R: GATGTGGACGAACACCCAGAG
<i>Herp</i>	F: GCAGTTGGAGTGTGAGTCG	R: TCTGTGGATTGAGCACCCCTTT
<i>Ifnγ</i>	F: ATGAACGCTACACACTGCATC	R: CCATCTTTTGGCAGTTCCTC
<i>Il1β</i>	F: GCAACTGTTCTGAACTCAACT	R: ATCTTTTGGGGTCCGCTCAACT
<i>Il6</i>	F: TAGTCCTTCTACCCCAATTTCC	R: TTGGTCCTTAGCCACTCCTTC
<i>Mcp1</i>	F: TTAAAAACCTGGATCGGAACCAA	R: GCATTAGCTTCAGATTTACGGGT
<i>Pepck</i>	F: CTGCATAACGGTCTGGACTTC	R: CAGCAACTGCCCCGTACTCC
<i>Pdk4</i>	F: AGGGAGGTCGAGCTGTTCTC	R: GGAGTGTTCACTAAGCGGTCA
<i>Pgc1α</i>	F: TATGGAGTGACATAGAGTGTGCT	R: CCACTTCA ATCCACCCAGAAAG
<i>Pgc1β</i>	F: TCCTGTAAAAGCCCGAGTAT	R: GCTCTGGTAGGGGACAGTGA
<i>Prdm16</i>	F: CCAAGGCAAGGGCGAAGAA	R: AGTCTGGTGGGATTGGAATGT
<i>Scd1</i>	F: TTCTTGCGATACACTCTGGTGC	R: CGGGATTGAATGTTCTTGTCTG
<i>Srebp-1c</i>	F: GCAGCCACCATCTA GCCTG	R: CAGCAGTGAGTCTGCCTTGAT
<i>Tnfa</i>	F: CCCTCACACTCAGATCATCTTCT	R: GCTACGACGTGGGCTACAG
<i>Ucp-1</i>	F: AGGCTTCCAGTACCATTAGGT	R: CTGAGTGAGGCAAAGCTGATTT
<i>Ucp-2</i>	F: ATGGTTGGTTTCAAGGCCACA	R: CGGTATCCAGAGGGAAAGTGAT
<i>Ucp-3</i>	F: CTGCACCGCCAGATGAGTTT	R: ATCATGGCTTGAATCCGGACC
<i>Xbp1</i>	F: GACAGAGAGTCAAACCTAACGTGG	R: GTCCAGCAGGCAAGAAGGT

Determination of Triglyceride Level in the Livers

The measurement was performed according to a previously reported procedure (10). Briefly, the liver was freshly cut at ~200 mg per piece and homogenized using a solution consisting of chloroform and methanol (2:1). The homogenates were incubated at 4°C overnight and centrifuged at 12,000 rpm for 20 min. Supernatants were collected, dried and re-dissolved in 5% Triton-X100. The triglyceride levels were determined using a commercial kit (TR22203) from Thermo-Scientific (Middletown, VA).

H&E Staining

Tissues were freshly collected after euthanizing mice. Tissue samples were fixed, dehydrated and embedded into paraffin. Tissue sections were cut at 6 μ m in thickness and dried at 37°C for 1 h. H&E staining was performed following the instruction of a commercial kit (#3500) from BBC Biochemical (Atlanta, GA). Slides were examined using an optical microscope and size measurements of adipocytes were taken using the NIS-Elements imaging platform purchased from Nikon Instruments Inc. (Melville, NY).

Oil-Red O Staining

Liver samples were freshly collected and immediately frozen in liquid nitrogen. Tissue sections were cut at 8 μm in thickness using a Cryostat. Sections were placed on slides and fixed using neutral buffered formalin for 30 min. The sections were washed with 60% isopropanol before being stained with freshly prepared Oil-red O working solution (#26079-05, Electron Microscopy Sciences) and counterstained with haematoxylin.

Statistics

Statistical analysis was conducted using one-way analysis of variance, and a P value below 0.05 ($P < 0.05$) is considered significantly different. The data were expressed as the mean \pm SD.

RESULTS

Rutin Suppressed the Palmitic Acid-Triggered Production of TNF α by Macrophages

We first determined the suppressive effect of rutin (Fig. 1a) on palmitic acid-triggered release of TNF α from macrophages. Palmitic acids induced TNF α release in a concentration-dependent manner. The TNF α level in culture medium was

900 pg/ml at palmitic acid concentration of 400 μM without affecting the viability of cells (Fig. 1b). Pre-treatment with rutin at 400 μM did not affect the viability of cells but suppressed the elevated production of TNF α by $\sim 33\%$ (Fig. 1c).

Rutin Decreased Transcription of Genes for Inflammation and ER Stress in Macrophages

We next determined the mRNA levels of genes involved in inflammation and ER stress in macrophages. Palmitic acids increased the expression of *Tnf α* by ~ 2.3 -fold and pre-treatment with rutin repressed the elevation by $\sim 64.7\%$ (Fig. 2a). Similarly, palmitic acids elevated the transcription of genes critical for inflammation including *Iffy*, *Il1 β* , *Il6* but not *Mcp1*, and pre-treatment with rutin greatly decreased these elevations (Fig. 2b). Moreover, palmitic acids increased the expression of *Gpr94*, a gene important for ER stress, by ~ 4.2 -fold and pre-treatment with rutin suppressed the elevation by $\sim 85.2\%$ (Fig. 2c). Additionally, palmitic acids also slightly increased the expression of other genes involved in ER stress such as *Xbp1*, *Ede1* and *Herp*, and the elevations were completely blocked by pre-treatment with rutin (Fig. 2d).

Rutin Repressed ROS Production in Macrophages

To explore the possible mechanism, we next examined the production of ROS in macrophages. Palmitic acids markedly increased the intracellular ROS in macrophages, and pre-





Fig. 1 Suppression effect of rutin on palmitic acid-triggered release of TNF α by macrophages. **(a)** Molecular structure of rutin. **(b)** Palmitic acid concentration dependent TNF α release. RAW 264.7 macrophages cultured in DMEM were incubated with different concentrations of palmitic acid for 12 h and supernatants were collected for TNF α determinations using ELISA kit. **(c)** Rutin suppressed the palmitic acid-induced release of TNF α in macrophages. RAW 264.7 macrophages were pre-treated with various concentrations of rutin for 4 h and then incubated with 400 μM palmitic acids with or without rutin for 12 h. Supernatants were collected for determination of TNF α concentration. ** $P < 0.01$ vs control; ## $P < 0.01$ vs cells treated with palmitic acids without rutin. Data were from 3 independent experiments.

Fig. 2 Suppression effect of rutin on palmitic acid-induced elevation of gene expression involved in inflammation and ER stress. RAW 264.7 macrophages were pre-treated with or without 200 μ M rutin for 4 h and then incubated with 400 μ M palmitic acid and/or 200 μ M rutin for 12 h. **(a)** Relative mRNA levels of *Tnfa*. **(b)** Relative mRNA levels of genes involved in inflammation. **(c)** Relative mRNA levels of *Gpr94*. **(d)** Relative mRNA levels of genes involved in ER stress. ** $P < 0.01$ vs control; ## $P < 0.01$ vs cells treated with palmitic acid without rutin. Data were from 3 independent experiments.



treatment with rutin significantly suppressed this elevation (Fig. 3a). Treatment of cells with rutin alone did not have any effect. Quantitative analysis shows that palmitic acids increased the signals of ROS by ~5.2-fold, and pre-treatment with rutin reduced this elevation by ~68% (Fig. 3b).

Rutin Protected Mice from HFD-Induced Obesity

We evaluated the anti-obese activity of rutin in male C57BL/6 mice fed a HFD. Compared to mice fed a regular chow, Fig. 4a shows HFD-fed mice had a significantly high body weight at ~39.2 g. Twice weekly intra-peritoneal injections of rutin completely blocked HFD-induced weight gain. The difference between HFD-fed mice with or without rutin injection was ~9.8 g and can be recognized visually (Fig. 4b). To investigate

the impact of the treatment on body composition, we measured lean and fat mass. Rutin greatly suppressed the increase of fat mass with no significant impact on lean mass (Fig. 4c). In addition, treatment with rutin did not show significant impact on food-intake in mice on HFD (Fig. 4d).

Rutin Prevented HFD-Induced Hypertrophy in Adipocytes

HFD feeding for 8 weeks increased the weight of the epididymal white adipose tissue (WAT) by ~4.1-fold, and rutin completely blocked the increase (Fig. 5a). Interestingly, neither HFD feeding nor rutin treatment showed impact on brown adipose tissue (BAT) (Fig. 5a). H&E staining of adipose tissues show that HFD-feeding caused apparent hypertrophy in





Fig. 3 Suppression effect of rutin on palmitic acid-caused production of ROS in macrophages. RAW 264.7 macrophages were pre-treated with or without 200 μ M rutin for 4 h and further incubated with 400 μ M palmitic acid and/or 200 μ M rutin for 24 h. Cells were washed with PBS and incubated with the detecting reagent from the Total ROS Detection Kit for 30 min at 37°C. **(a)** Representative pictures of ROS detection. **(b)** Quantitative analysis of ROS production (* $P < 0.05$ vs control; ** $P < 0.01$ vs control; ## $P < 0.01$ vs cells treated with palmitic acids without rutin).

Fig. 4 Effect of rutin on body weight and food-intake of animals fed a HFD. **(a)** Growth curve ($n=5$, $**P<0.01$ vs mice fed with regular chow; $##P<0.01$ vs mice fed with HFD). **(b)** Representative pictures of mice (Bar length = 1 cm). **(c)** Body composition ($n=5$, $**P<0.01$ vs mice fed with regular chow; $##P<0.01$ vs mice fed with HFD). **(d)** Average food intake.




adipocytes of WAT and also increased fat accumulation in BAT (Fig. 5b). The hypertrophy in WAT was associated with an increase in crown-like structures (arrows in Fig. 5b), a sign of macrophage activation. Rutin injection completely blocked the developments of hypertrophy and formation of crown-like structures in WAT (Fig. 5b). Quantitative analysis of white adipocytes shows the same conclusion (Fig. 5c).

Rutin Suppressed the Transcription of Genes Involved in Chronic Inflammation in WAT

To explore the possible mechanism, we measured the mRNA levels of genes that are pivotal for chronic inflammation in WAT. HFD-feeding dramatically increased the expression of *F4/80*, *Cd11c* and *Cd68* by ~29.2-fold, ~26.3-fold and ~15.4-

Fig. 5 Rutin-mediated blockade of hypertrophy and crown-like structures in the WAT of mice fed a HFD. **(a)** Weight of adipose tissues ($n=5$, $**P<0.01$ vs mice fed with regular chow; $##P<0.01$ vs mice fed with HFD). **(b)** Representative pictures of H&E staining for adipose tissues (Arrowheads points to crown-like structures). **(c)** Results of quantitative analysis ($**P<0.01$ vs mice fed with regular chow; $##P<0.01$ vs mice fed with HFD).



fold, respectively (Fig. 6a). Rutin completely blocked the elevated transcription of *F4/80* and *Cd11c*, and greatly reduced the mRNA level of *Cd68* by 78.4% (Fig. 6a). Interestingly, HFD-feeding also increased the mRNA levels of *Cd206* and *Arg1*, two marker genes for alternatively activated macrophages, by ~3.8-fold and ~1.9-fold, respectively (Fig. 6b). However, rutin partly repressed *Cd206* and increased *Arg1* by ~7.5-fold (Fig. 6b). Moreover, HFD-feeding elevated the mRNA levels of *Tnfa* and *Mcp1* in WAT (Fig. 6c). Rutin greatly repressed these elevations (Fig. 6c). Additionally, neither HFD-feeding nor rutin treatment significantly changed the transcription levels of genes involved in energy expenditure in WAT, including *Pgc1a*, *Pgc1b*, *Ucp1*, *Ucp2* and *Ucp3* (Fig. 6d).

Rutin Up-Regulated the Transcription of Genes Involved in Energy Expenditure in BAT

In BAT, rutin significantly increased the mRNA levels of *Dio2* and *Elovl3* by ~6.9-fold and ~5.1-fold, respectively (Fig. 7a). Moreover, rutin also increased the transcription of *Pgc1a* by ~1.9-fold (Fig. 7b). Additionally, we determined the mRNA levels of other genes involved in energy expenditure and nutrient metabolism in BAT, including *Ucp1*, *Ucp2*, *Ucp3*, *Pgc1b*, *Cpt1a*, *Cpt1b*, *Pdk4* and *Glut4*. No mRNA levels of these genes were significantly changed by HFD-feeding or rutin treatment (Fig. 7c and d)

Rutin Blocked HFD-Induced Fatty Liver

HFD feeding for 8 weeks markedly increased the weight of livers by ~31.3%, and rutin completely blocked the increase (Fig. 8a). Consistently, HFD-feeding caused apparent fatty liver, as evidenced by the vacuoles in H&E staining and

further confirmed by Oil-red O staining, and rutin completely blocked the development of HFD-induced fatty liver (Fig. 8b). Quantitative determination of triglycerides in the liver leads to the same conclusion (Fig. 8c). Both AST and ALT levels were in normal ranges for all animals fed with either HFD or regular chow (Fig. 8d).

Rutin Modulated the Transcription of Genes Involved in Inflammation and Nutrient Metabolism in the Liver

Results in Fig. 9a indicate a slight increase in mRNA levels of several inflammatory genes including *F4/80*, *Tnfa* and *Ifny* in animals fed a HFD, but the difference was not statistically significant. Treatment with rutin did not show a notable difference. However, HFD-feeding increased the mRNA level of *Srebp1c* by ~2.8-fold, but had no significant impact on *Acc*, *Fas* and *Scd1* (Fig. 9b). Rutin suppressed the elevated transcription of *Srebp1c* by ~78.9% (Fig. 9b). Additionally, rutin increased the transcription of *Pdk4* and decreased the HFD-induced transcription increase of *Cd36*, *Pepck* and *G6p* in the livers (Fig. 9c).

Rutin Maintained Glucose Sensitivity in Mice fed a HFD

Mice fed a HFD for 8 weeks showed significantly impaired glucose sensitivity, and treatment with rutin maintained such sensitivity (Fig. 10a). AUC calculations make this point clearer (Fig. 10b). To assess insulin sensitivity in these mice, we performed ITT. HFD-feeding caused an apparent insulin resistance, and treatment with rutin completely blocked this resistance (Fig. 10c). Blood insulin determinations show that HFD-feeding increased the levels of insulin by ~4.6-fold and

Fig. 6 Rutin-mediated transcription inhibition of the genes involved in chronic inflammation in WAT of mice fed a HFD. **(a)** Relative mRNA levels of marker genes for classically activated macrophages ($n=4$, ** $P<0.01$ vs mice fed with regular chow; ## $P<0.01$ vs mice fed with HFD). **(b)** Relative mRNA levels of marker genes for alternatively activated macrophages ($n=4$, ** $P<0.01$ vs mice fed with regular chow; ## $P<0.01$ vs mice fed with HFD). **(c)** Relative mRNA levels of pro-inflammatory cytokines ($n=4$, ** $P<0.01$ vs mice fed a regular chow; ## $P<0.01$ vs mice fed with HFD). **(d)** Relative mRNA level of genes involved in energy metabolism.





Fig. 7 Rutin-mediated modulation of gene expression in BAT of mice fed a HFD. **(a)** Relative mRNA level of BAT signature genes ($n=4$, ** $P<0.01$ vs mice fed with regular chow; ## $P<0.01$ vs mice fed with HFD). **(b–c)** Relative mRNA level of genes involved in energy expenditure ($n=4$, ** $P<0.01$ vs mice fed a regular chow; ## $P<0.01$ vs mice fed a HFD). **(d)** Relative mRNA level of genes involving glucose and lipids metabolism.



treatment with rutin greatly reduced the elevation by ~71.2% (Fig. 10d).

DISCUSSION

In this study, we demonstrate that rutin suppresses palmitic acid-triggered inflammation in macrophages (Fig. 1) and blocks HFD-induced obesity in mice (Fig. 4). The anti-inflammatory effect of rutin in macrophages is correlated with its activity in suppressing ER stress and ROS production (Figs. 2 and 3). The anti-obese effect of rutin in HFD-

feeding mice was associated with its action in blocking macrophage infiltration (Fig. 5) and chronic inflammation in adipose tissue (Fig. 6).

Recent studies by Henao-Mejia *et al.* and Stienstra *et al.* show that chronic inflammation plays critical roles in the pathogenesis of obesity and its complications (5,6). Chronic inflammation in obesity is primarily mediated by macrophages (11,12). Emerging evidence shows that macrophages can be activated by saturated fatty acids (13,14). Moreover, recent studies by Cruz *et al.* and Bulua *et al.* demonstrate that the promoted inflammation is correlated with increased ROS production (15,16). In addition, recent studies also

Fig. 8 Rutin-mediated blockade of HFD-induced fatty liver. **(a)** Liver weight ($n=5$, ** $P<0.01$ vs mice fed a regular chow; ## $P<0.01$ vs mice fed a HFD). **(b)** Representative pictures of H&E staining and Oil-red O staining. **(c)** Liver triglyceride levels ($n=5$, ** $P<0.01$ vs mice fed a regular chow; ## $P<0.01$ vs mice fed a HFD). **(d)** Blood level of AST and ALT.





Fig. 9 Effect of rutin on hepatic gene expression **(a)** Relative mRNA levels of genes involving inflammation ($^{##} P < 0.01$ vs mice fed a HFD). **(b)** Relative mRNA levels of genes for lipogenesis ($n = 4$, $^{**} P < 0.01$ vs mice fed a regular chow; $^{##} P < 0.01$ vs mice fed a HFD). **(c)** Relative mRNA levels of genes for glucose and lipids metabolism ($n = 4$, $^{**} P < 0.01$ vs mice fed with regular chow; $^{##} P < 0.01$ vs mice fed with HFD).




show that ER stress may play important roles in promoting inflammation as well (17,18). Our results are in concurrence with these earlier studies in support that rutin suppresses the macrophage activation by decreasing ROS production and ER stress (Figs. 2 and 3).

Macrophage infiltration into adipose tissue is one typical feature of obesity (19,20). The original observation was reported by Weisberg *et al.* (19) and Xu *et al.* (20), and subsequently confirmed and expanded by others using different models (21–23). Although the detailed molecular mechanism

for this process needs additional investigation, it is largely accepted that hypertrophy and necrosis of adipocytes play critical roles (24). Hypertrophy is a central cause for adult obesity and is usually associated with adipocyte death (19,24). Moreover, Cinti and colleagues proposed that necrosis of adipocytes, driven by hypertrophy, is one of the predominate signals regulating macrophage infiltration (21,25). The infiltrated macrophages aggregate around necrotic adipocytes, leading to development of crown-like structures and chronic inflammation (21,25). Consistent with

Fig. 10 Rutin-mediated inhibition of HFD-induced glucose tolerance and insulin resistance. **(a)** Glucose profiles of IPGTT ($n = 5$, $^{**} P < 0.01$ vs mice fed a regular chow; $^{##} P < 0.01$ vs mice fed a HFD). **(b)** Results of AUC calculations ($n = 5$, $^{**} P < 0.01$ vs mice fed a regular chow; $^{##} P < 0.01$ vs mice fed a HFD). **(c)** Results of ITT ($n = 5$, $^{**} P < 0.01$ vs mice fed a regular chow; $^{##} P < 0.01$ vs mice fed a HFD). **(d)** Insulin level in blood ($n = 5$, $^{**} P < 0.01$ vs mice fed a regular chow; $^{##} P < 0.01$ vs mice fed a HFD).



these studies, our data show that HFD-feeding causes apparent hypertrophy of adipocytes, resulting in an increased number of crown-like structures and elevated chronic inflammation in adipose tissue (Figs. 5 and 6). Treatment with rutin significantly suppressed this process, suggesting that the anti-obese effect of this anti-oxidant may be achieved, at least in part, through blocking necrosis and chronic inflammation in adipose tissues. Additionally, rutin also has a direct effect on hepatocytes and adipocytes. A previous study by Wu *et al.* using hepatocarcinoma cells demonstrated that rutin can inhibit oleic acid-induced lipid accumulation in hepatocytes *in vitro*, probably *via* reducing lipogenesis and oxidative stress (26). In addition, using 3 T3-L1 cells Choi *et al.* demonstrated that rutin is capable of suppressing differentiation of pre-adipocytes *in vitro* (27). These direct effects may be related to the anti-oxidant activity of rutin.

Insulin resistance is a major complication associated with obesity, and its underlying mechanism remains unclear (28,29). Shulman and colleagues proposed that ectopic accumulation of lipids and dysregulation of intracellular signal pathways may play central roles in the etiology of insulin resistance (30,31). On the other hand, accumulating evidence links chronic inflammation to the induction of insulin resistance, suggesting that macrophage-related inflammation may also contribute to the pathogenesis of obesity-induced insulin resistance (20, 32,33). Our data show that treatment with rutin blocks development of insulin resistance and maintains glucose sensitivity in mice fed a HFD (Fig. 10). These effects were correlated with the blocked ectopic fat accumulation in livers and the suppressed chronic inflammation in adipose tissues (Figs. 6 and 8).

As a potent natural anti-oxidant, rutin can be found in many fruits and vegetables. Previous studies have suggested that rutin be used as a dietary supplement to modulate blood level of lipids and glucose. Additionally, it has also been applied in clinic for treatment of many diseases including poor circulation, high blood pressure and capillary fragility (34). The underlying mechanism, however, was considered complicated because no specific target has been identified. Our current study shows that rutin blocks macrophage activation and represses chronic inflammation in adipose tissue, providing new information for better understanding its beneficial effects and for using rutin to manage metabolic disorders. Evidently, further studies at molecular levels are still needed to see whether other pathways are also involved.

Rutin altered the expression levels of multiple genes in mice fed a HFD. *Arg1* is a marker for alternatively activated macrophages (M2 macrophages), and elevated expression of *Arg1* is usually associated with a higher ratio of M2 macrophages that are essential to maintain the homeostasis of adipose tissue (35). *Pgc1 α* , *Dio2* and *Elovl3* are three critical genes tightly involved in energy expenditure in BAT (36). Rutin up-regulated expression levels of these three genes,

indicating that this natural anti-oxidant may be able to increase energy expenditure. *Pdk4* encodes pyruvate dehydrogenase kinase isozyme 4 which is an enzyme located in the matrix of the mitochondria and regulates the balance between glucose and lipid metabolism (37). Rutin increased transcription of *Pdk4* in liver, suggesting that it may elevate lipid oxidation in this organ. Further studies are warranted to elucidate how rutin treatment altered expression levels of these genes *in vivo*.

In conclusion, our results show that rutin is able to block HFD-induced obesity, fatty liver and insulin resistance in mice, and these beneficial effects were correlated with a blockade of macrophage infiltration and chronic inflammation in adipose tissue. Our results confirm that supplementation of rutin is a promising strategy for preventing HFD-induced obesity and its associated complications.

ACKNOWLEDGMENTS AND DISCLOSURES

The study was supported in part by grants from NIH (RO1EB007357 and RO1HL098295). We thank Ms. Ryan Fugett for English editing. The authors claim no conflicts of interest.

REFERENCES

- Hsu CL, Wu CH, Huang SL, Yen GC. Phenolic compounds rutin and o-coumaric acid ameliorate obesity induced by high-fat diet in rats. *J Agric Food Chem*. 2009;57(2):425–31.
- Panchal SK, Poudyal H, Arumugam TV, Brown L. Rutin attenuates metabolic changes, nonalcoholic steatohepatitis, and cardiovascular remodeling in high-carbohydrate, high-fat diet-fed rats. *J Nutr*. 2011;141(6):1062–9.
- Hao HH, Shao ZM, Tang DQ, Lu Q, Chen X, Yin XX, *et al.* Preventive effects of rutin on the development of experimental diabetic nephropathy in rats. *Life Sci*. 2012;91(19–20):959–67.
- Kamalakkannan N, Stanely Mainzen Prince P. The influence of rutin on the extracellular matrix in streptozotocin-induced diabetic rat kidney. *J Pharm Pharmacol*. 2006;58(8):1091–8.
- Henaoui-Mejia J, Elinav E, Jin C, Hao L, Mehal WZ, Strowig T, *et al.* Inflammasome-mediated dysbiosis regulates progression of NAFLD and obesity. *Nature*. 2012;482(7384):179–85.
- Stienstra R, van Diepen JA, Tack CJ, Zaki MH, van de Veerdonk FL, Perera D, *et al.* Inflammasome is a central player in the induction of obesity and insulin resistance. *Proc Natl Acad Sci U S A*. 2011;108(37):15324–9.
- Hu QH, Zhang X, Pan Y, Li YC, Kong LD. Allopurinol, quercetin and rutin ameliorate renal NLRP3 inflammasome activation and lipid accumulation in fructose-fed rats. *Biochem Pharmacol*. 2012;84(1):113–25.
- Javed H, Khan MM, Ahmad A, Vaibhav K, Ahmad ME, Khan A, *et al.* Rutin prevents cognitive impairments by ameliorating oxidative stress and neuroinflammation in rat model of sporadic dementia of Alzheimer type. *Neuroscience*. 2012;210:340–52.
- Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta C(T)}$ Method. *Methods*. 2001;25(4):402–8.

10. Hara A, Radin NS. Lipid extraction of tissues with a low-toxicity solvent. *Anal Biochem.* 1978;90(1):420–6.
11. Monteiro R, Azevedo I. Chronic inflammation in obesity and the metabolic syndrome. *Mediators Inflamm.* 2010. doi:10.1155/2010/289645.
12. Nijhuis J, Rensen SS, Slaats Y, van Dielen FM, Buurman WA, Greve JW. Neutrophil activation in morbid obesity, chronic activation of acute inflammation. *Obesity (Silver Spring).* 2009;17(11):2014–8.
13. Huang S, Rutkowski JM, Snodgrass RG, Ono-Moore KD, Schneider DA, Newman JW, *et al.* Saturated fatty acids activate TLR-mediated proinflammatory signaling pathways. *J Lipid Res.* 2012;53(9):2002–13.
14. Shi H, Kokoeva MV, Inouye K, Tzameli I, Yin H, Flier JS. TLR4 links innate immunity and fatty acid-induced insulin resistance. *J Clin Invest.* 2006;116(11):3015–25.
15. Cruz CM, Rinna A, Forman HJ, Ventura AL, Persechini PM, Ojcius DM. ATP activates a reactive oxygen species-dependent oxidative stress response and secretion of proinflammatory cytokines in macrophages. *J Biol Chem.* 2007;282(5):2871–9.
16. Bulua AC, Simon A, Maddipati R, Pelletier M, Park H, Kim KY, *et al.* Mitochondrial reactive oxygen species promote production of proinflammatory cytokines and are elevated in TNFR1-associated periodic syndrome (TRAPS). *J Exp Med.* 2011;208(3):519–33.
17. Goodall JC, Wu C, Zhang Y, McNeill L, Ellis L, Saudek V, *et al.* Endoplasmic reticulum stress-induced transcription factor, CHOP, is crucial for dendritic cell IL-23 expression. *Proc Natl Acad Sci U S A.* 2010;107(41):17698–703.
18. Wheeler MC, Rizzi M, Sasik R, Almanza G, Hardiman G, Zanetti M. KDEL-retained antigen in B lymphocytes induces a proinflammatory response: a possible role for endoplasmic reticulum stress in adaptive T cell immunity. *J Immunol.* 2008;181(1):256–64.
19. Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, Ferrante AW, Jr. Obesity is associated with macrophage accumulation in adipose tissue. *J Clin Invest.* 2003;112(12):1796–808.
20. Xu H, Barnes GT, Yang Q, Tan G, Yang D, Chou CJ, *et al.* Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance. *J Clin Invest.* 2003;112(12):1821–30.
21. Cinti S, Mitchell G, Barbatelli G, Murano I, Ceresi E, Faloia E, *et al.* Adipocyte death defines macrophage localization and function in adipose tissue of obese mice and humans. *J Lipid Res.* 2005;46(11):2347–55.
22. Kanda H, Tateya S, Tamori Y, Kotani K, Hiasa K, Kitazawa R, *et al.* MCP-1 contributes to macrophage infiltration into adipose tissue, insulin resistance, and hepatic steatosis in obesity. *J Clin Invest.* 2006;116(6):1494–505.
23. Canello R, Tordjman J, Poitou C, Guilhem G, Bouillot JL, Hugol D, *et al.* Increased infiltration of macrophages in omental adipose tissue is associated with marked hepatic lesions in morbid human obesity. *Diabetes.* 2006;55(6):1554–61.
24. Sun K, Kusminski CM, Scherer PE. Adipose tissue remodeling and obesity. *J Clin Invest.* 2011;121(6):2094–101.
25. Murano I, Barbatelli G, Parisani V, Latini C, Muzzonigro G, Castellucci M, *et al.* Dead adipocytes, detected as crown-like structures, are prevalent in visceral fat depots of genetically obese mice. *J Lipid Res.* 2008;49(7):1562–8.
26. Wu CH, Lin MC, Wang HC, Yang MY, Jou MJ, Wang CJ. Rutin inhibits oleic acid induced lipid accumulation *via* reducing lipogenesis and oxidative stress in hepatocarcinoma cells. *J Food Sci.* 2011;76(2):T65–72.
27. Choi I, Park Y, Choi H, Lee EH. Anti-adipogenic activity of rutin in 3T3-L1 cells and mice fed with high-fat diet. *Biofactors.* 2006;26(4):273–81.
28. Qatanani M, Lazar MA. Mechanisms of obesity-associated insulin resistance: many choices on the menu. *Genes Dev.* 2007;21(12):1443–55.
29. Samuel VT, Petersen KF, Shulman GI. Lipid-induced insulin resistance: unravelling the mechanism. *Lancet.* 2010;375(9733):2267–77.
30. Shulman GI. Cellular mechanisms of insulin resistance. *J Clin Invest.* 2000;106(2):171–6.
31. Petersen KF, Shulman GI. Etiology of insulin resistance. *Am J Med.* 2006;119(5 Suppl 1):S10–6.
32. Verschuren L, Kooistra T, Bernhagen J, Voshol PJ, Ouwens DM, van Erk M, *et al.* MIF deficiency reduces chronic inflammation in white adipose tissue and impairs the development of insulin resistance, glucose intolerance, and associated atherosclerotic disease. *Circ Res.* 2009;105(1):99–107.
33. Shoelson SE, Lee J, Goldfine AB. Inflammation and insulin resistance. *J Clin Invest.* 2006;116(7):1793–801.
34. Pescosolido N, Librando A. Oral administration of an association of forskolin, rutin and vitamins B1 and B2 potentiates the hypotonising effects of pharmacological treatments in POAG patients. *Clin Ter.* 2010;161(3):e81–5.
35. Murray PJ, Wynn TA. Obstacles and opportunities for understanding macrophage polarization. *J Leukoc Biol.* 2011;89(4):557–63.
36. Bostrom P, Wu J, Jedrychowski MP, Korde A, Ye L, Lo JC, *et al.* A PGC1-alpha-dependent myokine that drives brown-fat-like development of white fat and thermogenesis. *Nature.* 2012;481(7382):463–8.
37. Berggren JR, Boyle KE, Chapman WH, Houmard JA. Skeletal muscle lipid oxidation and obesity: influence of weight loss and exercise. *Am J Physiol Endocrinol Metab.* 2008;294(4):E726–32.