

Zinc-enriched fertilisers as a potential public health intervention in Africa

Edward J. M. Joy · Alexander J. Stein ·
Scott D. Young · E. Louise Ander · Michael J. Watts ·
Martin R. Broadley

Received: 28 November 2014 / Accepted: 23 February 2015 / Published online: 8 March 2015
© Springer International Publishing Switzerland 2015

Abstract

Background In this review, we examine the potential of Zn-enriched fertilisers to alleviate human dietary Zn deficiency. The focus is on ten African countries where dietary Zn supply is low and where fertiliser subsidies are routinely deployed on cereal crops.

Scope Dietary Zn supply and deficiency prevalence were quantified from food supply and composition data. Typical effects of soil (granular) and foliar Zn applications on Zn concentrations in maize (*Zea mays* L.), rice (*Oryza sativa* L.) and wheat (*Triticum aestivum* L.) grains were based on a systematic literature review. Reductions in disease burdens attributable to Zn deficiency and cost-effectiveness were estimated using a disability-adjusted life years (DALYs) approach.

Conclusions Baseline Zn supply in 2009 ranged from 7.1 (Zambia) to 11.9 (Mali) mg *capita*⁻¹ day⁻¹; prevalence of Zn deficiency ranged from 24 (Nigeria) to 66 %

(Zambia). In reviewed studies, soil Zn application led to an increase in median Zn concentration in maize, rice and wheat grains of 23, 7 and 19 %; foliar application led to increases of 30, 25 and 63 %. Enriching granular fertilisers within current subsidy schemes would be most effective in Malawi, reducing DALYs lost due to Zn deficiency by 10 %. The cost per DALY saved ranged from US\$ 624 to 5893 via granular fertilisers and from US\$ 46 to 347 via foliar fertilisers. Foliar applications are likely to be more cost effective than soil applications due to fixation of Zn in the soil but may be more difficult to deploy. Zinc fertilisation is likely to be less cost-effective than breeding in the longer term although other micronutrients such as selenium could be incorporated.

Keywords Agronomic biofortification · Fertiliser · HarvestPlus · Micronutrient deficiency · Phytic acid · Zinc

Responsible Editor: Philippe Hinsinger.

Electronic supplementary material The online version of this article (doi:10.1007/s11104-015-2430-8) contains supplementary material, which is available to authorized users.

E. J. M. Joy · S. D. Young · M. R. Broadley (✉)
School of Biosciences, University of Nottingham, Sutton
Bonington Campus, Loughborough LE12 5RD, UK
e-mail: martin.broadley@nottingham.ac.uk

E. J. M. Joy · E. L. Ander · M. J. Watts
Centre for Environmental Geochemistry, British Geological
Survey, Keyworth, Nottingham NG12 5GG, UK

A. J. Stein
Agricultural Economist, Brussels, Belgium

Introduction

Zinc (Zn) is an integral component of thousands of proteins for all organisms. Adult human bodies contain 1.5–2.5 g of Zn with a daily intake requirement of 10–14 mg (WHO and FAO 2004). The estimated prevalence of inadequate dietary Zn intake is >25 % in sub-Saharan Africa (Wessells and Brown 2012; Wessells et al. 2012; Joy et al. 2014). Dietary Zn deficiency can have a range of health impacts including increased risk of child mortality due to diarrhoeal disease and stunting

(Salgueiro et al. 2002) and imposes considerable individual suffering as well as social and economic costs (Stein 2010, 2014). An estimated 0.7 % of the global disease burden is attributable to Zn deficiency, rising to 1.5 % in low income countries (WHO 2009). Factors contributing to Zn deficiency in humans include low consumption of animal products, high phytate intakes that inhibit Zn absorption and low concentrations of Zn in crops grown on Zn deficient soils (Cakmak et al. 1999; Sandstead 2000; Gibson 2012). Phytate refers to mixed salts of phytic acid (PA), the principal form of phosphorus (P) in cereal grains and a potent inhibitor of Zn absorption in the human gut. A PA:Zn molar ratio of >15 is commonly used to classify diets having low levels of bioavailable Zn.

In crop plants, a leaf Zn concentration of 15–20 mg Zn kg⁻¹ dry weight (DW) is typically required for adequate growth (Broadley et al. 2007). However, Zn deficiency in crops is widespread globally, in particular, due to low phytoavailability of Zn in soils. Such soils are commonly defined as having a Zn concentration extractable by ethylene diamine tetra-acetic acid (EDTA) or diethyl triamine penta-acetic acid (DTPA) less than 1.5 and 1.0 mg kg⁻¹ DW of soil, respectively (Trierweiler and Lindsay 1969; Lindsay and Norvell 1978). Low phytoavailability of Zn can result from low soil Zn concentrations or the influence of soil characteristics that limit Zn solubility such as high pH values or large concentrations of available phosphate or CaCO₃ (Brümmer et al. 1983; Graham et al. 1992; White and Zasoski 1999; Cakmak 2002, 2004; Alloway 2008; Lu et al. 2011). Deficiency of Zn on cultivated soils is widespread, affecting >50 % of soils in India, Pakistan and Turkey, >30 % of soils in China and most soils in Western Australia and Africa (Alloway 2008).

Zinc fertilisers are widely used to improve crop yields where soil Zn phytoavailability is low (Ahmad et al. 2012). The first reports of using Zn fertilisers to ameliorate crop Zn deficiency were in peach, pecan and pineapple orchards (Hoagland 1948). However, major increases in arable crop production due to Zn fertiliser use are now well-established. For example, wheat grain yield increases of >600 % were reported in Central Anatolia in Turkey from the mid-1990s which returned economic benefits of ca. US\$ 100 million annually in the following decade (Cakmak 2004). More recently, there has been research exploring the use of Zn fertilisers to increase Zn concentrations beyond that which is needed for maximum yield, to enrich the edible

portions of crops for human health benefits (Rengel et al. 1999; Genc et al. 2005; Ortiz-Monasterio et al. 2007; Broadley et al. 2007; Cakmak 2008; White and Broadley 2009, 2011). However, the cost-effectiveness of this approach has not previously been determined.

Several Zn forms have been used in fertilisers, including Zn-sulphate (ZnSO₄) and Zn-oxide. Such forms can be delivered either in combination with granular nitrogen (N) fertilisers or as a foliar spray. An advantage of enriching granular fertilisers is that farmers already using fertilisers can be reached with no extra labour or machinery required at the farm level, nor additional distribution infrastructure. However, plant uptake of soil-applied Zn is limited by a low availability or diffusion of Zn in certain soils, particularly those with high pH, organic matter or CaCO₃ contents (Tye et al. 2003; Zhao et al. 2014). For example, recovery of soil-applied Zn may be <1 % in calcareous soils (Lu et al. 2012). Thus, a more effective strategy for increasing grain Zn concentrations might be via foliar sprays. With foliar application, Zn is absorbed by the leaf epidermis, remobilized and transferred to the grain through the phloem (Fernández and Eichert 2009; White and Broadley 2011).

Common soil fertility management practices can also affect soil Zn status and concentrations of Zn in the grain. For example, N application can increase Zn uptake, xylem transport and remobilization via the phloem, and hence the concentration of Zn in the grain (Erenoglu et al. 2002, 2011; Kutman et al. 2010, 2011; Xue et al. 2012), while excessive P fertilisation can reduce availability of Zn in the soil (Marschner 1993; Lu et al. 2011). Manzeke et al. (2012) reported that farmer fields in Zimbabwe receiving cattle manure or leaf litter in combination with NPK had greater concentrations of EDTA-extractable Zn in soils and greater concentrations of Zn in maize grain compared to unfertilised fields or those receiving only NPK, while rotation with legumes was also reported to increase concentration of Zn in maize grain.

The use of Zn-containing fertilisers to increase dietary Zn supply is one of several strategies to address dietary Zn deficiency. These include dietary diversification, provision of supplements and addition of Zn during food processing (Gibson et al. 2000; Shrimpton et al. 2005). Other agricultural strategies to increase the concentration of Zn in harvested grain include crop breeding for high-Zn varieties (Cakmak 2008; White and Broadley 2009, 2011; Bouis and Welch 2010), while

soaking or ‘priming’ of seeds in ZnSO₄ solution might be more efficient than soil applications and confer yield benefits (Slaton et al. 2001; Harris et al. 2007; Harris et al. 2008) although increased Zn concentration in progeny grain is not consistently reported (Johnson et al. 2005). It may also be possible to breed lower PA concentrations into the grains of staple crops and the benefits of even marginal reductions in grain PA concentration on Zn bioavailability could be large at population scales (Joy et al. 2014). Interestingly, it has been reported that Zn-enriched fertilisers can decrease concentrations of PA in cereal grains while Zn deficiency may lead to increased P uptake and accumulation in plants (Loneragan et al. 1982; Erdal et al. 2002).

The impact of public health interventions can be measured using a disability-adjusted life-years (DALYs) approach. A DALY is equivalent to a lost year of ‘healthy life’ and is the sum of years of life lost due to premature mortality and years of life lost due to a disability (YLD; Murray 1994). The YLD is the product of the number of incident cases, average duration of the disease and a disability weight to reflect the severity of the disease, which ranges between 0 (i.e., full health) and 1 (i.e., death). As there are limited resources available to invest in public health, a DALY approach allows direct comparison between different public health interventions to guide policy making and to increase the efficient use of scarce funds. Previously, Stein et al. (2006) estimated that biofortification via breeding for high-Zn rice and wheat varieties could save up to 55 % of the 2.8 million DALYs lost annually due to Zn deficiency in India at a cost of US\$ 0.68 and 8.80 *per* DALY saved, for optimistic and pessimistic scenarios respectively. Fielder et al. (2013) estimated that fortifying maize meal with a premix containing vitamin A, iron and Zn at large-scale mills in Zambia could save 5657 DALYs annually of which 1757 were due to Zn deficiency, at a cost of US\$ 401 *per* DALY saved. Similar studies have not yet been conducted for fertilisers and so the aim of this review is to assess the cost-effectiveness of Zn fertilisers in reducing disease burdens due to dietary Zn deficiency.

The focus of the review is sub-Saharan Africa because of the high incidence of Zn deficiency relative to other regions of the World (Lim et al. 2012; Wessells and Brown 2012). Specifically, we have focussed on ten countries which routinely deploy fertiliser subsidy schemes (Burkina Faso, Kenya, Ghana, Mali, Malawi, Nigeria, Senegal, Tanzania and Zambia) or which have

government control of imports (Ethiopia) as this offers a mechanism for mandating Zn-enrichment of fertilisers (Jayne and Rashid 2013; Wanzala-Mlobela et al. 2013).

The first aspect of this review quantifies dietary Zn supply and deficiency prevalence for the ten focus countries using food supply and food composition data, the latter adjusted according to cereal processing methods. The second aspect is a meta-analysis of published field experiments that investigate the effect of soil- and foliar-applied Zn fertilisers on Zn and PA concentrations in the grain of three staple crops: maize (*Zea mays* L.), rice (*Oryza sativa* L.) and wheat (*Triticum aestivum* L.). The third aspect of this study models the effect of enriching fertilisers with Zn on dietary Zn supplies, deficiency and associated disease burdens using a DALY framework. We model the effect of enriching subsidised fertiliser with Zn and compare it to a scenario in which subsidised and non-subsidised fertilisers are enriched.

Materials and methods

Baseline dietary Zn supplies and deficiency prevalence

Baseline national dietary Zn supplies and deficiencies were estimated for ten countries on the basis of food supply and composition data: Burkina Faso, Ethiopia, Ghana, Kenya, Malawi, Mali, Nigeria, Senegal, Tanzania and Zambia. Food supply and population data were downloaded from United Nations datasets (UNDSEA 2013; FAO 2014a). Food Balance Sheets (FBSs) compiled by the FAO record estimates of food supply for up to 92 edible items at a national level, representing net *per capita* food supply calculated from national production, trade, transport losses, storage, non-food uses, livestock feed etc., but with no adjustment for household waste or inter- and intra-household variation in access to food (FAO 2001). Data are now available for 2011, but 2009 was chosen as the reference year to match with available fertiliser usage statistics (Supplementary Table 1). The Institute for Health Metrics and Evaluation’s (IHME) estimates of DALYs lost due to Zn deficiency are based on the study of Wessells and Brown (2012) (Lim et al. 2012). Thus, food composition data compiled by Wessells and Brown (2012) are used in the present study, including adjustments made due to processing of staple foods, such as milling and fermentation of cereals and cassava

(*Manihot esculenta* Crantz; Supplementary Table 2). Food supply and composition data were combined in order to generate estimates of dietary Zn and PA supplies by food item (Wessells and Brown 2012; Supplementary Tables 2 and 3). National mean supplies of Zn and PA were estimated and the amount of absorbable Zn in the diet was calculated using the Miller equation (Miller et al. 2007; Supplementary Table 4). As assumed previously (Wessells and Brown 2012; Wessells et al. 2012), variation in individual intakes was captured through a coefficient of variation in absorbable Zn intake of 25 %. We adopted the approach employed in the Estimated Average Requirement (EAR) cut-point method, in that the proportion of the population below the mean national physiological requirement for Zn was assumed to be deficient.

Effect of Zn-enriched fertilisers on concentrations of Zn and PA in the grains of maize, rice and wheat

A systematic literature review was conducted in order to assess the impact of Zn fertilisers on Zn concentrations in major grains. The terms ‘zinc OR Zn AND biofortification’, ‘zinc OR Zn AND fertili*’, ‘zinc OR Zn AND application’, ‘zinc OR Zn AND concentration’, ‘zinc OR Zn AND response’, ‘zinc OR Zn AND uptake’, ‘zinc OR Zn AND soil’ and ‘zinc OR Zn AND foliar’ were queried in the search engines Web of Science (Thomson Reuters, New York, U.S.A.), Science Direct (Elsevier, Philadelphia, U.S.A.) and Google Scholar (Google Inc., California, U.S.A.). Further studies were identified by searching reference sections of review and research papers found using the search terms stated. Criteria for inclusion were that studies were published in a peer-reviewed journal, that Zn was added via either soil or foliar applications under field conditions, and that concentrations of Zn in the grain were reported for treatments and controls. Both rainfed and irrigated plots were included. A number of studies assessed Zn applications in combination with varying N or P application rates. In such instances, the treatment closest to 100 kg ha⁻¹ year⁻¹ of N and 25 kg ha⁻¹ year⁻¹ of P was included for consistency. The effect of Zn fertiliser was determined for maize, wheat and rice; insufficient studies were identified to allow a similar systematic analysis of data on other crops. The most commonly used form of Zn in both granular and

foliar fertilisers is ZnSO₄·7H₂O. Other forms of Zn including Zn-bonded amino acids may be more effective at increasing grain Zn concentrations (Ghasemi et al. 2013), however insufficient studies were identified to include alternatives to ZnSO₄ in this review.

The effect of Zn fertiliser was quantified as concentration of Zn in the grain at harvest as a percent of control and the median effect was used across studies by taking study site, crop type, cultivar, application method (soil or foliar) and application rate (kg Zn ha⁻¹) as factors. Mean effect over seasons was taken for multi-season trials. Some studies examined the residual effect of Zn fertilisers but these data were not included; this is revisited in the Discussion. Studies examining the effect of different application timings of foliar sprays with later applications (i.e., post-flowering) appeared to have a greater impact on grain Zn concentration but possibly a lower impact on grain yield (Cakmak et al. 2010; Mabesa et al. 2013); in such instances, the treatment when Zn was applied at flowering or heading stages was taken for consistency. Results from 26 journal articles were included in the literature review: four studies of maize, six of rice, 15 of wheat and one of maize and wheat (Table 1). Fourteen and four studies, respectively, reported effects of soil and foliar applications while eight reported effects of both soil and foliar applications. No studies were identified that reported the effect of Zn application via soil on PA concentration in the grain of rice, nor via foliar spray in maize. Preliminary analysis of results from all studies revealed that Zn application via soil and foliar routes tended to increase concentrations of Zn in the grain. However, larger application rates did not appear to increase concentrations more than smaller rates when comparing trials. This is likely to be due to the variety of soil characteristics encountered. The majority of studies did not report variance of Zn or PA concentration in the grain so it was not possible to perform a standard meta-analysis in which variance is used to weight the contribution of effect size (Field and Gillett 2010). For these reasons, results were pooled by crop and application method (soil or foliar) with the median effect on crop Zn and PA concentrations taken.

Milling of maize, rice and wheat grains is common practice. It was assumed that the relative

Table 1 Summary of studies included in the meta-analysis. Application methods of zinc (Zn) are soil (S) or foliar (F). ‘n’ is the number of data points contributing to the meta-analysis in which

individual studies were stratified by crop, cultivar, location and Zn application rate and pooled by application method

Crop	Varieties	Application via (n)	Country	Reference
Wheat		S (4)	Pakistan	Abid et al. 2013
Wheat	Bread and durum	S (1), F (3)	Turkey	Cakmak et al. 2010
Wheat		S (12)	Turkey	Ekiz et al. 1998
Wheat	20 cultivars	S (20)	Turkey	Erdal et al. 2002
Wheat	2 Zn-deficiency tolerant cultivars	F (2)	Iran	Ghasemi et al. 2013
Maize		S (2)	Pakistan	Harris et al. 2007
Maize	2 cultivars	S (6)	Pakistan	Kanwal et al. 2010
Wheat	30 cultivars	S (60)	Iran	Khoshgofarmanesh et al. 2012
Wheat		S (1)	China	Lu et al. 2011
Wheat	2 winter wheat cultivars	S (8)	China	Lu et al. 2012
Rice	10 biofortification breeding line genotypes	F (19)	Philippines	Mabesa et al. 2013
Maize		S (1)	Zambia	Manzeke et al. 2014
Maize		S (2)	Spain	Martin-Ortiz et al. 2009
Rice	Cultivars commonly used	S (10), F (10)	China, India, Lao PDR, Thailand, Turkey	Phattarakul et al. 2012
Rice		S (4)	India	Shivay et al. 2008
Rice		S (1)	India	Srivastava et al. 2009
Maize, wheat		S (1,1), F (1,1)	China	Wang et al. 2012
Rice	3 cultivars	F (3)	China	Wei et al. 2012
Rice	5 ‘high-’;5 ‘low-’ Zn genotypes	S (15)	Philippines	Wissuwa et al. 2008
Wheat	10 cultivars	S (15), F (6)	China	Yang et al. 2011a
Wheat		F (1)	China	Yang et al. 2011b
Wheat	3 bread, 1 durum	S (4), F (4)	Turkey	Yilmaz et al. 1997
Wheat	Common cultivars	S (1), F (7)	China	Zhang et al. 2012
Wheat		S (4)	China	Zhao et al. 2011
Wheat	5 cultivars. (Results not presented by cultivar)	S (1), F (1)	China	Zhao et al. 2014
Wheat	11 cultivars	S (14), F (13)	China, India, Kazakhstan, Mexico, Pakistan, Turkey, Zambia	Zou et al. 2012

increase in Zn concentration in the whole grain and the endosperm fraction were equivalent (the assumption is revisited in the [Discussion](#)). Thus, studies that reported whole grain data were assumed to give a good prediction of the effect of Zn fertilisation on the Zn concentration of edible portions. Rice is most commonly eaten in its polished form; data for polished rice were used

where available, otherwise, brown rice data were used.

Effect of Zn fertilisers on Zn deficiency prevalence

Four scenarios were modelled to quantify the impacts of agronomic biofortification with Zn on dietary Zn and PA supplies at national levels in the ten focus countries:

- the first scenario modelled a policy to enrich *subsidised* fertiliser, which would be easiest to implement given the pre-existing government involvement;
- the second scenario modelled a policy to enrich *subsidised and non-subsidised* fertiliser;
- the third scenario modelled a policy to introduce foliar fertilisation of cereals with a target of 50 % coverage; and
- the fourth scenario modelled a policy to introduce foliar fertilisation of cereals with a target of 75 % coverage.

All scenarios assumed that maize, rice and wheat crops were targeted. In Ethiopia, teff (*Eragrostis tef* (Zucc.) Trotter) was also included as this grain accounts for almost one-fifth of national energy consumption from cereals and *ca.* 40 % of national inorganic fertiliser consumption (CSA 2011; FAO 2014a). Demand for fertiliser was also assumed to arise from millet (*Eleusine coracana* L. and *Pennisetum glaucum* L.), sorghum (various spp.), cocoa (*Theobroma cacao* L.), coffee (*Coffea* spp.), cotton (*Gossypium* spp.), palm oil (*Elaeis guineensis* Jacq.), sugarcane (*Saccharum officinarum* L.) and tobacco (*Nicotiana* spp.). Scenarios one and two assumed that fertiliser used as basal dressing was enriched using $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ to give a N:Zn mass ratio of 2:1. Assuming recommended application rates of N (see below) this would provide 23, 12, 20 and 12 $\text{kg ha}^{-1} \text{ year}^{-1}$ of Zn for hybrid maize, local maize, rice and wheat, respectively. A compliance of 90 % was used to account for enforcement problems. Scenarios three and four assumed that 600 L ha^{-1} of 0.5 % (w/v) aqueous $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ solution was sprayed twice annually on maize, rice and wheat crops, supplying 1.36 kg ha^{-1} of Zn.

The efficacy and costs of scenarios one and two are partly determined by the quantity of fertiliser enriched and the proportion of maize, rice, wheat and teff covered by this fertiliser. Fertiliser usage data were derived from the International Fertilizer Development Center (IFDC) for subsidised fertiliser consumption (IFDC 2013a) and total national fertiliser consumption (IFDC 2011; 2012a, b, c, d, e, f; 2013b, c; Supplementary Table 5) from which the supply of N was calculated. Fertiliser was assumed to contain 23 % N by mass where product information was not available. ‘Demand’ for N was

calculated as the product of crop-specific fertiliser requirements and cropping areas of maize, rice, wheat and cash crops (Supplementary Table 5). Cropping areas were derived from FAO production data (FAO 2014b; Supplementary Table 5).¹ Half of maize production area was assumed to be hybrid varieties and half local varieties in all countries. Crop-specific recommended fertiliser application rates were identified only for Malawi, (Malawi Government Ministry of Agriculture and Food Security, date unknown, accessed 2014; Supplementary Table 5) and were applied to all countries for most crops. Teff was assumed to require the same N rate as wheat. Requirements for tea, coffee, palm oil and cocoa were identified through a literature search (FAO 1984; Grice 1990; Makono and Chanika 2008).

Thus, the proportion of crops receiving Zn-enriched fertiliser in scenarios one and two was calculated as supply of N divided by demand for N (Eq. 1).

$$P_{\text{Zn}} = F_{s,w} \times q / \sum (C_{a-l} \times A_{a-l}) \quad (1)$$

Where:

- P_{Zn} proportion of crop receiving Zn-enriched fertiliser
 $F_{s,w}$ national N usage (metric tonnes, t) via subsidised fertiliser (s) or subsidised and non-subsidised fertiliser (w)
 C_{a-l} Cropping area of maize (a), rice (b), wheat (c), teff (d), millet (e), sorghum (f), cocoa (g), coffee (h), cotton (i), palm oil (j), sugarcane (k) and tobacco (l)
 A_{a-l} Recommended N application rate ($\text{t ha}^{-1} \text{ year}^{-1}$) for maize (a), rice (b), wheat (c), teff (d), millet (e), sorghum (f), cocoa (g), coffee (h), cotton (i), palm oil (j), sugarcane (k) and tobacco (l)
 q compliance factor (0 to 1).

The effect of Zn fertilisers was modelled through changes to the concentrations of Zn and PA in the grains of

¹ Total cereal production area in Ethiopia in 2009 was 9.2×10^6 ha and teff appears to be included in the production sub-category ‘Cereals, nes’ (production area 2.6×10^6 ha) rather than the sub-category ‘Millet’ (production area 0.4×10^6 ha), contrary to the FAO’s own production definitions (FAO 2014c). ‘Cereals, nes’ production area was assumed to be solely teff.

maize, rice and wheat. The proportion of each crop receiving fertiliser was multiplied by the median effects of applied Zn on grain Zn and PA concentrations to generate new composition data for maize, rice, wheat and teff. Teff grain was assumed to have the same response to Zn-enriched fertiliser as wheat grain. National dietary Zn and PA supplies, quantity of absorbable Zn in the diet and estimated prevalence of Zn deficiency were re-calculated using the new composition data, assuming that composition of all other food items and quantity of food supply had not changed. The proportion of DALYs saved was assumed to equal the reduction in proportion of deficiency prevalence.

Estimating the cost-effectiveness of an agronomic biofortification approach to addressing Zn deficiency in sub-Saharan Africa

Baseline disease burdens attributable to Zn deficiency for the ten focus countries were derived from the Global Burden of Disease Study in which a proportion of ‘diarrheal diseases’, ‘typhoid and paratyphoid fevers’ and ‘lower respiratory infections’ are attributed to Zn deficiency and assigned a ‘disability weight’ (Lim et al. 2012; IHME 2014).

The cost of enriching fertilisers with Zn was estimated assuming a wholesale retail price of $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ of US\$ 500 t^{-1} . Only the fertiliser used by maize, rice, wheat and teff was assumed to be enriched and, for soil applications, only the proportion applied as a basal dressing (Supplementary Table 6). The proportion of crops receiving basal fertiliser was assumed to equal the proportion of crops receiving fertiliser (Eq. 1). The cost of supplying knapsack sprayers was estimated for the foliar scenarios by assuming that knapsack sprayer sets cost US\$ 150 *per* unit and that each would last 10 years and cover 20 ha of cropland annually through sharing among farmers (Supplementary Table 6). Other implementation costs including agricultural extension services and the distribution of equipment and fertiliser were not considered.

Results

Baseline dietary Zn supplies and deficiency prevalence

Dietary Zn deficiency is likely to be widespread in sub-Saharan Africa. Using national-level food supply and

food composition data adjusted by common processing methods, we estimate a high prevalence of Zn deficiency in the ten focus countries, ranging from 24 % in Nigeria to 66 % in Zambia (Table 2; Supplementary Table 4). The estimated large disease burden attributable to Zn deficiency is consistent with previous work. For example, IHME (2014) estimated that the burden of Zn deficiency in the ten focus countries ranged from 161 to 1219 DALYs lost 100 k population⁻¹ in Ghana and Burkina Faso, respectively. This is two orders of magnitude greater than in the UK where it is estimated that <4 DALYs 100 k population⁻¹ are lost due to Zn deficiency (IHME 2014).

Effect of Zn-enriched fertilisers on concentrations of Zn and PA in the grains of maize, rice and wheat

A summary of the studies included in the literature review is presented in Table 1. The trials combined a range of soil types and cultivars (Table 3; Supplementary Tables 7 and 8). In control plots (i.e., without application of Zn), concentration of Zn in maize grain ranged from 14.9 mg kg⁻¹ DW on calcareous, Zn-deficient silty-clay-loam soils in Pakistan (Harris et al. 2007) to 22.5 mg kg⁻¹ DW on Zn-deficient sandy-loam in Pakistan (Kanwal et al. 2010); concentration of Zn in brown rice (i.e., husk removed but grain unpolished) ranged from 9.9 mg kg⁻¹ DW in Thailand (Phattarakul et al. 2012) to 41.6 mg kg⁻¹ DW in ‘high-Zn’, upland soil in the Philippines (Wissuwa et al. 2008); concentration of Zn in polished rice grain ranged from 12.3 mg kg⁻¹ DW in pH 7.7 soils in Turkey (Phattarakul et al. 2012) to 28.0 mg kg⁻¹ DW in Zn-adequate soils in China (Wei et al. 2012); and concentration of Zn in wheat grain ranged from 6.6 mg kg⁻¹ DW in borderline Zn-deficient soils with pH 7.8 in Iran (Khoshgoftarmansh et al. 2012) to 40.2 mg kg⁻¹ DW in Zn-adequate soils in India (Zou et al. 2012).

The median increases in Zn concentration in the grains of maize, rice and wheat were, respectively, 28, 11 and 18 % for soil application and 30, 24 and 63 % for foliar application (Table 4; Fig. 1; Supplementary Tables 7 and 8). The result for maize with foliar-applied Zn is based on only one data point (Wang et al. 2012). Application of Zn-enriched fertilisers via the soil decreased PA concentration in the grain of wheat by 11 % while foliar application decreased concentration in the grain of rice and wheat by 1 and 13 %, respectively. No studies were identified that reported

Table 2 Baseline national-level estimates of zinc (Zn) deficiency and associated disease burden in comparison to published studies for the ten focus countries of this review

Country	Estimated risk of inadequate Zn supply			Stunting	DALYs lost due to Zn deficiency
	% total population			% children 0–59 months	100 k population ⁻¹
Rank of country for each study in (brackets)					
	Present study	Wessells and Brown (2012)	Joy et al. (2014)	UNICEF (2013)	IHME (2014)
Reference year	2009	2003–7	2009	2007–11	2012
Burkina Faso	49.6 (3)	39.4 (3)	5.5 (9)	35 (7)	1,219 (1)
Ethiopia	31.6 (6)	11.0 (10)	81.5 (1)	44 (3)	344 (7)
Ghana	27.1 (7)	21.6 (8)	36.4 (5)	28 (9)	161 (10)
Kenya	27.0 (8)	25.3 (5)	60.6 (4)	35 (7)	281 (9)
Malawi	54.8 (2)	40.6 (2)	32.8 (6)	47 (1)	769 (2)
Mali	25.5 (9)	22.3 (7)	5.2 (10)	38 (6)	448 (5)
Nigeria	24.1 (10)	20.6 (9)	8.7 (8)	41 (5)	408 (6)
Senegal	36.0 (5)	24.6 (6)	11.7 (7)	27 (10)	489 (4)
Tanzania	41.1 (4)	34.1 (4)	64.4 (3)	42 (4)	341 (8)
Zambia	65.7 (1)	44.9 (1)	72.4 (2)	45 (2)	665 (3)

the effect of Zn application via soil on PA concentration in the grain of rice, nor via foliar spray in maize.

Yield data are important to rule out the ‘concentration effect’ whereby lower yields may lead to greater Zn concentrations in the grain as the Zn taken up by the plant is distributed to fewer or smaller grains. This is particularly so for foliar applications as high concentrations of Zn in the spray solution could damage leaf cuticles (Eichert and Fernández 2012). Yield was reported for only 122 out of 273 plots included in the systematic review so a consistent approach to excluding plots with low yield relative to control was not possible, although this is re-visited in the Discussion. Of the studies that reported yield data, soil application led to a median 10–11 % increase in grain yield for maize, rice and wheat, whereas foliar application had no obvious effect (Table 4; Supplementary Table 7). In addition, of the studies that did not report yield data by plot, Mabesa et al. (2013) found no significant difference in yield of rice due to foliar application of Zn, but also reported a significant negative correlation among different varieties between grain yield and grain Zn concentration, a relationship also reported by Wissuwa et al. (2008). Very high yield responses of >150 % of control were found in ten and four wheat data points for soil and

foliar-applied Zn, respectively (Supplementary Table 7). All of these data points also exhibited high Zn concentrations in grains relative to controls, with Q1, median and Q3 of 170, 189 and 257 % for the soil-applied treatments and 252, 295 and 316 % for the foliar-applied treatments.

Effect of Zn fertilisers on Zn deficiency prevalence

The effectiveness of the biofortification approach is greatly dependent on the coverage of fertilisers, i.e., the proportion of crops that would receive fertilisers enriched with Zn. Scenario one modelled the potential impact of Zn-enrichment of granular fertiliser currently distributed under national subsidy schemes. The Ethiopian Government control fertiliser imports so all fertiliser usage was considered under the ‘subsidised’ bracket. The percentage of cereal production receiving subsidised fertiliser was: Burkina Faso (2), Ethiopia (18), Ghana (6), Kenya (8), Malawi (24), Nigeria (10), Senegal (9), Tanzania (6) and Zambia (20) (Supplementary Table 5). Data for Mali were not available. The estimated reduction in DALYs lost due to Zn deficiency was lowest in Burkina Faso (<1 %) and greatest in Malawi (10 %) where there would be a 3 %

Table 3 Summary of baseline soil properties and zinc (Zn) application rates in maize, rice and wheat crops. 'n' is the number of data points contributing to the meta-analysis in which individual studies were stratified by crop, cultivar, location and Zn application rate and pooled by application method. Q1 and Q3 are first and third quartiles, respectively

Crop	Method	Baseline soil properties										Zn application rate								
		DTPA-extractable Zn					pH (H ₂ O)					n			Mean			SD		
		n	Mean mg kg ⁻¹	SD	Q1	Median	Q3	n	Mean	SD	Q1	Median	Q3	n	Mean kg ha ⁻¹	SD	Q1	Median	Q3	
Maize [†]	Soil	11	0.69	0.12	0.68	0.72	0.72	11	8.05	0.11	7.98	7.98	8.20	12	15.72	18.75	3.44	8.50	18.00	
Rice		29	1.75	1.52	0.79	0.97	2.70	30	6.89	1.13	5.70	6.85	7.80	30	11.77	4.18	11.00	13.00	15.00	
Wheat		158	0.82	0.79	0.10	0.67	1.00	137	7.73	0.32	7.55	7.80	7.98	158	25.36	13.55	11.00	23.00	50.00	
Maize	Foliar	1	0.56	-	-	0.56	-	1	8.24	-	-	8.24	-	1	0.91	-	-	0.91	-	
Rice		33	1.30	1.74	0.28	0.36	2.10	34	6.59	0.76	6.35	6.40	6.95	34	3.18	1.51	1.25	4.00	4.00	
Wheat		38	0.80	1.13	0.32	0.52	0.71	37	7.69	0.50	7.53	7.80	7.98	38	1.62	0.73	1.29	1.59	1.63	

[†] Manzeke et al. (2014) measured pH in CaCl₂ and extractable Zn using EDTA so these data were excluded

increase in the mean amount of absorbable Zn in the diet (Table 5; Supplementary Tables 9 and 10).

Scenario two modelled the potential impact of Zn-enrichment of all fertiliser currently used. The percentage of cereal production receiving both subsidised and non-subsidised fertiliser was: Ghana (24), Kenya (39), Malawi (39), Mali (14), Nigeria (22), Senegal (21), Tanzania (13) and Zambia (33) (Supplementary Table 5). Data for Burkina Faso were not available. The estimated reduction in DALYs lost due to Zn deficiency was lowest in Mali (3 %) and greatest in Malawi (15 %) where there would be a 5 % increase in the national mean amount of absorbable Zn in the diet (Table 5; Supplementary Tables 9 and 10).

Scenarios three and four modelled the potential impact of foliar Zn application, covering 50 and 75 %, respectively, of maize, rice and wheat production. The response of grain Zn concentration to foliar Zn application was greater in wheat than in maize or rice and it was assumed that Zn concentration in teff grain responded as in wheat. Wheat and teff consumption combined was greatest in Ethiopia, (161 g *capita*⁻¹ day⁻¹; Supplementary Table 1), where the increase in mean amount of absorbable Zn in the diet for scenarios three and four, respectively, was 13 and 19 % and the estimate of DALYs lost due to Zn deficiency decreased by 41 and 54 %, respectively (Table 5; Supplementary Tables 9 and 10). Response of grain PA concentration to foliar Zn application was greatest for rice, and dietary PA supply in scenario four decreased by 2.8, 1.5 and 1.4 % in Senegal, Ghana and Mali where rice consumption was 188, 157 and 72 g *capita*⁻¹ day⁻¹, respectively (Table 5; Supplementary Tables 1 and 9).

Estimating the cost-effectiveness of an agronomic biofortification approach to addressing Zn deficiency in sub-Saharan Africa

The cost effectiveness of agronomic biofortification of crops using soil or foliar-applied ZnSO₄.7H₂O was estimated. Where the outcomes of public health interventions are not measured in monetary terms, decision-makers cannot rely on conventional tools of economic evaluation, such as internal rates of return or benefit-cost ratios, to determine whether the 'investment' in an intervention represents a good use of scarce resources. Instead, their relative cost-effectiveness can be assessed by comparing the average cost of saving one DALY across interventions, or against benchmarks.

Table 4 Summary of the effects of zinc (Zn)-enriched fertiliser on Zn and PA concentration in the grain of maize, rice and wheat. 'n' is the number of data points contributing to the meta-analysis in

which individual studies were stratified by crop, cultivar, location and Zn application rate and pooled by application method. Q1 and Q3=first and third quartiles, respectively

Crop	Application via	n	Mean	SD	Min	Q1	Median	Q3	Max
Zn concentration in the grain, % of control									
Maize	Soil	12	128	18	106	114	123	139	173
Rice		30	111	15	84	102	107	119	157
Wheat		158	143	57	51	105	119	174	373
Maize	Foliar	1	130	–	130	–	130	–	130
Rice		34	127	19	107	114	125	140	172
Wheat		38	178	55	112	143	163	203	333
PA concentration in the grain, % of control									
Maize	Soil	3	96	7	7	92	93	104	104
Wheat		26	91	15	65	83	89	98	121
Rice	Foliar	3	86	1	85	85	87	87	87
Wheat		15	97	10	82	89	99	102	121
Yield, % of control									
Maize	Soil	12	110	10	95	100	111	120	125
Rice		15	111	8	102	103	111	117	129
Wheat		47	188	246	90	99	110	129	1,607
Maize	Foliar	1	98	–	98	–	98	–	98
Rice		15	100	7	84	98	102	104	109
Wheat		32	142	173	77	99	102	111	1,071

Scenario one modelled a policy to enrich granular fertilisers currently distributed under government subsidy schemes with Zn. The cost *per* DALY saved ranged from US\$ 624 to 5747 in Burkina Faso and Ghana,

respectively (Table 5; Fig. 2; Supplementary Table 11). Scenario two modelled a policy to enrich subsidised and non-subsidised granular fertilisers used as basal dressings and cost *per* DALY saved ranged from US\$ 977 to

Fig. 1 Concentration of zinc (Zn) in the grains of maize (circles), rice (triangles) and wheat (squares) following Zn application via soil (open) or foliage (filled). Y-axis represents the concentration as a percentage of control. Vertical lines mark Harvest Plus breeding targets for maize and wheat (dashed) and rice (dash-dot)




Table 5 Effect of different zinc (Zn) fertilisation scenarios on dietary Zn and phytate supplies and estimated risk of Zn deficiency in ten example countries in sub-Saharan Africa. Scenario '0' is baseline; 1 and 2 model enrichment of granular fertilisers, either

subsidised or subsidised and non-subsidised; 3 and 4 model application of foliar Zn sprays to 50 and 75 % of target crops. Scenarios that are cost-effective in comparison to WHO (*) or World Bank and WHO (**) benchmarks are highlighted

Country	Scenario	Dietary Zn supply mg capita ⁻¹ d ⁻¹	Dietary phytate supply	Zn deficiency risk %	DALYs lost due to Zn deficiency 100 k population ⁻¹	Programme cost US\$ '000 s year ⁻¹	Cost <i>per</i> DALY saved US\$ year ⁻¹
Burkina Faso	0	11.2	3,617	49.6	1288		
	1	11.2	3,615	49.4	1282	584	624*
	2	–	–	–	–	–	–
	3	11.7	3,606	43.3	1123	1,267	49**
	4	12.0	3,601	40.4	1048	1,900	51**
Ethiopia	0	8.2	1,830	31.6	329		
	1	8.4	1,818	28.9	300	31,983	1,302*
	2	–	–	–	–	–	–
	3	9.5	1,821	18.7	195	11,392	98**
	4	10.2	1,816	14.6	152	17,089	111**
Ghana	0	8.0	1,371	27.1	162		
	1	8.0	1,370	26.7	160	2,791	5,747
	2	8.1	1,366	25.8	155	10,640	5,893
	3	8.4	1,358	22.7	136	2,088	332*
	4	8.6	1,351	20.9	125	3,133	347*
Kenya	0	8.9	1,858	27.0	280		
	1	8.9	1,851	26.2	272	6,021	1,830*
	2	9.2	1,822	23.4	242	29,583	1,932*
	3	9.5	1,851	21.4	222	3,811	162**
	4	9.8	1,847	19.2	199	5,716	172**
Malawi	0	8.9	2,700	54.8	768		
	1	9.2	2,663	49.6	695	15,675	1,431
	2	9.3	2,641	46.6	653	25,115	1,456
	3	9.7	2,696	43.5	610	3,132	132**
	4	10.1	2,694	38.7	542	4,698	138**
Mali	0	11.9	2,795	25.5	495		
	1	–	–	–	–	–	–
	2	12.0	2,787	24.8	482	6,324	3,428
	3	12.4	2,769	21.5	419	2,072	194**
	4	12.7	2,756	19.8	385	3,108	203**
Nigeria	0	8.6	1,751	24.1	406		
	1	8.7	1,749	23.6	397	20,791	1,593
	2	8.7	1,745	23.0	388	45,002	1,613
	3	9.1	1,737	20.0	337	9,790	89**
	4	9.3	1,729	18.2	307	14,685	94**
Senegal	0	8.4	1,820	36.0	471		
	1	8.4	1,816	35.2	461	1,261	964*
	2	8.5	1,810	34.2	447	3,027	977*
	3	9.0	1,786	27.6	362	656	46**
	4	9.3	1,769	24.2	317	984	49**

Table 5 (continued)

Country	Scenario	Dietary Zn supply $\text{mg capita}^{-1} \text{d}^{-1}$	Dietary phytate supply	Zn deficiency risk %	DALYs lost due to Zn deficiency $100 \text{ k population}^{-1}$	Programme cost US\$ '000 s year ⁻¹	Cost <i>per</i> DALY saved US\$ year ⁻¹
Tanzania	0	8.0	2,037	41.1	341		
	1	8.4	2,033	40.4	335	9,630	3,547
	2	8.5	2,028	39.6	328	19,813	3,573
	3	9.0	2,026	33.8	280	7,323	267*
	4	8.8	2,021	30.6	253	10,985	279*
Zambia	0	7.1	2,100	65.7	664		
	1	7.3	2,076	61.3	620	7,222	1,237*
	2	7.4	2,059	58.2	589	12,358	1,244*
	3	7.8	2,097	53.1	537	1,817	108**
	4	8.1	2,095	47.4	480	2,726	112**

5893 in Senegal and Ghana, respectively (Table 5; Fig. 2; Supplementary Table 11). Variation in cost effectiveness between countries was partly a function of the baseline disease burden attributable to Zn deficiency with higher burdens leading to lower costs *per* DALY saved. Foliar application is likely to be a more efficient use of Zn by avoiding fixation of Zn in the soil. Costs *per* DALY saved in scenario three, in which 50 % of cereal production received foliar Zn fertiliser, ranged from US\$ 46 to 332 in Senegal and Ghana, respectively, while in scenario four, in which 75 % of cereal production received foliar Zn fertiliser, the cost ranged from US\$ 49 to 347, also in Senegal and Ghana, respectively (Table 5; Fig. 2; Supplementary Table 11).

The World Health Organization (WHO 2001) and the World Bank (World Bank 1993) have provided benchmarks to assess the cost-effectiveness of health interventions; if the cost of saving a DALY is below the benchmark then it is considered a good investment. The WHO benchmark is calculated in relative terms as 300 % of a country's *per-capita* Gross Domestic Product (GDP), hence in 2009, from US\$ 678 to 5550 *capita*⁻¹ in Malawi and Ghana, respectively (World Bank 2014; Supplementary Table 11). The World Bank benchmark is in absolute terms, taking a value of US\$ 150 *per* DALY in 1990 as a base year which is equivalent to US\$ 246 in 2009 after adjusting for inflation. Thus, even in the poorest country, the World Bank benchmark is lower and harder to meet than that of WHO. According to the WHO benchmark, pursuing Zn enrichment of soil-applied granular fertilisers appears to be cost-effective in Burkina Faso, Ethiopia, Kenya, Senegal

and Zambia, but not other countries, while foliar application of Zn appears to be cost-effective in all countries. According to the World Bank benchmark, the only cost-effective scenarios are foliar application of Zn in Burkina Faso, Ethiopia, Kenya, Malawi, Mali, Nigeria, Senegal and Zambia (Table 5; Fig. 2; Supplementary Table 11).

Discussion

Baseline dietary Zn supplies and deficiency prevalence

Robust estimates of dietary Zn supplies and risk of deficiency underpin the evaluation of approaches to address Zn deficiency. There are potential weaknesses in using FBS food supply data including underestimating food supply as some subsistence production is not captured, or overestimating supply by failing to account for household-level food waste (FAO 2001). These weaknesses have been discussed extensively elsewhere (de Haen et al. 2011; Wessells et al. 2012; Joy et al. 2014). A further source of error may arise from composition data derived from sources that will not capture local variation in elemental composition of crops and there is a lack of spatially-resolved food composition data in sub-Saharan Africa (Joy et al. 2014; Joy et al. 2015). However, in the absence of wide-scale analysis of biomarkers of nutrient status, e.g., blood serum, estimating national dietary supplies of bioavailable Zn remains a valuable method of




Fig. 2 Impact and cost-effectiveness of four zinc (Zn) fertilisation scenarios in ten countries in sub-Saharan Africa: Burkina Faso (BF), Ethiopia (ET), Ghana (GH), Kenya (KE), Malawi (MW), Mali (ML), Nigeria (NG), Senegal (SN), Tanzania (TZ) and Zambia (ZM). Impact is defined as the reduction in disease burden attributable to Zn deficiency and is quantified in disability-adjusted life-years (DALYs; Supplementary Table 10). Cost-effectiveness is quantified in US\$ *per* DALY saved (Supplementary Table 11).

Scenario 1 models enrichment of subsidised granular fertilisers; Scenario 2 models enrichment of subsidised and non-subsidised granular fertilisers; Scenarios 3 and 4 model foliar application of Zn to 50 and 75 % of cereals, respectively. Not all countries are represented in Scenarios 1 and 2 due to lack of data. The text size represents the absolute number of DALYs saved annually (highly dependent on the country's population size): from smallest to largest, <25,000, 25–50,000, 50–75,000 and 75,000+

estimating the prevalence of Zn deficiency at a national level (Gibson et al. 2008).

An alternative to FBSs is to use food consumption data captured in nationally-representative household surveys (Fielder et al. 2008). These data are available for eight of the ten focus countries in this study (Ethiopia, Ghana, Kenya, Malawi, Mali, Nigeria, Tanzania and Zambia). For example, Fielder et al. (2013) estimate mean and median intakes of Zn in Zambia of 5.8 and 4.4 mg *capita*⁻¹ day⁻¹ and a 73.1 % prevalence of inadequate Zn intakes, compared to US Institute of Medicine dietary requirements of Zn. An advantage of household surveys is that they allow sub-national resolution of dietary estimates and shed light on distributional issues. One drawback is that they rely on household member recall, which is subject to misreporting, both intentional and unintentional (Archer et al. 2013; Molteldo et al. 2014). Zinc intakes and status can also be measured directly through analysis of dietary composites or concentration of Zn in blood plasma or serum samples. Through such methods, high

prevalence of Zn deficiency have been reported previously in sub-populations in Burkina Faso (e.g., Müller et al. 2003), Ethiopia (e.g., Abebe et al. 2007; Kassu et al. 2008; Stoecker et al. 2009), Kenya (e.g., Siekmann et al. 2003), Malawi (e.g., Siyame et al. 2013), Nigeria (e.g., Gegios et al. 2010), Tanzania (e.g., Veenemans et al. 2011) and Zambia (e.g., Duggan et al. 2005).

New baseline Zn deficiency estimates correlated well with the IHME estimates of DALYs lost due to Zn deficiency (Spearman's Rank, $r=0.588$, $p=0.018$, $d.f.=9$). This is expected as both the present study and the IHME DALY estimates were derived from the underlying data and methodology developed by Wessells et al. (2012) (Wessells and Brown 2012; Lim et al. 2012). However, there is no correlation between the results of Joy et al. (2014) and Wessells and Brown (2012) which is surprising given the similar underlying methodologies. This may have arisen because the studies used different reference years and food composition tables and prevalence of deficiency in the Wessells and Brown (2012) study are based on estimated dietary

supplies of ‘bioavailable’ Zn. Also, Wessells and Brown (2012) assumed certain milling and fermentation practices of cereals and other crops and adjusted the concentrations of Zn and PA accordingly. For example, estimated dietary PA supplies in Ethiopia are 2802 mg *capita*⁻¹day⁻¹ in the Joy et al. (2014) study, but 1724 mg *capita*⁻¹day⁻¹ in the Wessells and Brown (2012) study in which 59 % of wheat, 90 % of maize, millet and sorghum and 100 % of other cereals (i.e., teff) are assumed to be fermented (Supplementary Table 2).

Zinc deficiency confers increased risk of diarrhoea and is a potential underlying cause of stunting which is defined as having a height-for-age more than two standard deviations below the median of the WHO growth reference (WHO 1995). Although deficiency estimates in the current study, which are based on dietary intakes of Zn and PA, show a general positive relationship with the WHO estimates of childhood stunting prevalence, the relationship is not statistically significant ($P > 0.1$) for both the absolute and log-transformed values. Other potential underlying causes of stunting include caloric deficiency (Stein et al. 2003), mother’s body mass index (Mamiro et al. 2005) access to clean drinking water (Esrey et al. 1988) and sanitation and hygiene practices (Fink et al. 2011; Spears et al. 2013).

Effect of Zn-enriched fertilisers on concentrations of Zn and PA in the grains of maize, rice and wheat

The efficacy of applied Zn in increasing grain Zn concentration depends in part on the crop species and cultivar. Across studies, the effect of soil-applied Zn on grain Zn concentrations was greater in maize and wheat than in rice (Table 4; Fig. 1). This may be due to abiotic factors such as the reducing conditions and high organic matter content typically found in anaerobic flooded paddy soil (Alloway 2008), or biotic factors such as root morphology or root exudates (Widodo et al. 2010). Cereal crops can respond to Zn deficiency stress by releasing compounds capable of chelating soil-bound Zn including low-molecular-weight organic acids and a class of non-protein amino acids known as phytosiderophores (Kochian 1993; Hoffland et al. 2006; Suzuki et al. 2006, 2008; Widodo et al. 2010). Alternatively, the lower efficacy in rice may have been due to the higher soil pH values and lower baseline DTPA-extractable Zn concentrations in maize and wheat trials than rice trials, possibly leading to lower concentrations of Zn in grains from control plots and a

greater response to Zn application (Table 3). In addition, average Zn application rates were greater in maize and wheat trials than rice (Table 3).

The effect of foliar-applied Zn was greater in wheat than in rice (Table 4; Fig. 1). Biotic factors including the ability to remobilize Zn from ageing leaves to the grain may be responsible, while the rate at which remobilization occurs may be dependent on the Zn nutritional status of the plant which will, in turn, be affected by soil properties. Also, a portion of the foliar-applied Zn may run down the stem and reach the rhizosphere where availability to the plant root will depend on soil properties.

Several studies reported significant differences in the Zn concentration of grains and the efficacy of soil or foliar Zn application between cultivars of rice and wheat (Yilmaz et al. 1997; Ekiz et al. 1998; Erdal et al. 2002; Wissuwa et al. 2008; Yang et al. 2011a; Khoshgoftarmansh et al. 2012; Phattarakul et al. 2012; Wei et al. 2012; Ghasemi et al. 2013; Mabesa et al. 2013), suggesting that agronomic and crop breeding biofortification efforts should be aligned. Only one study investigated different cultivars of maize, finding no significant difference between two cultivars (Kanwal et al. 2010).

The majority of trials reviewed here were conducted in Western, Central and Eastern Asia where Zn-deficiency in crops commonly arises in calcareous soils with pH > 7.5 (Table 1; Alloway 2008). In highly-weathered tropical soils, Zn deficiency may be a product of leaching and low total Zn content (Alloway 2008). Only two of the studies reviewed here included trials located in Africa, both of soil-applied Zn, reporting an 18 % increase in the concentration of Zn in the grain of maize on borderline Zn-deficient soils in Zimbabwe (Manzeke et al. 2014) and a 4 % increase in the concentration of Zn in the grain of wheat grown on Zn-deficient soils in Zambia (Zou et al. 2012). Clearly, more studies are required across the varied environmental conditions found in sub-Saharan Africa to verify the estimates of the effects of applied Zn on grain Zn concentration.

Although soil-application of Zn is likely to improve yields of crops grown on Zn-deficient soils, it could inhibit the absorption of other nutrients such as copper, while foliar sprays with high Zn concentration could damage leaf cuticles. Reduced yields could lead to a ‘concentration effect’, where Zn in the leaves or shoot is distributed to fewer or smaller grains. Although this may increase concentrations of Zn in the edible portion, it

would clearly be undesirable. Yield data were reported for 122 treatment plots in total with soil applications appearing to generally improve yields while foliar applications had no obvious effect (Table 4). The yields of just six of the treatment plots were <90 % of the relevant control plot and all of these plots also exhibited increased concentration of Zn in the grain relative to the control plot (Supplementary Table 7). However, removing these data points from the study had minimal impact on the estimated efficacy and cost-effectiveness of the four Zn-fertilisation scenarios: median effects across studies of soil applied Zn would remain unchanged for maize, rice and wheat and median effect of foliar-applied Zn would be unchanged for maize and would be 124 and 160 % for rice and wheat, compared to 125 and 163 %.

Zinc is distributed unevenly across cereal grain fractions, with higher concentrations in the bran and embryo than the endosperm. Thus, milling and processing generally reduce the concentration of Zn in the edible product (Bityutskii et al. 2002; Ozturk et al. 2006; Liang et al. 2008; Cakmak et al. 2010; Joy et al. 2015). Many of the studies included in the meta-analysis only reported Zn concentrations in whole grain, with and without application of Zn. It is possible that greater concentration of Zn in the whole grain with Zn fertilisation is a result of increased concentrations in the bran and embryo and not the endosperm. However, Cakmak et al. (2010) examined Zn concentrations across the different fractions of wheat grain and reported that the greatest relative increase in Zn concentrations with Zn fertilisation is likely to be in the endosperm. In addition, those studies that reported milled or polished grain Zn concentrations generally found positive effects of Zn fertilisation. For example, Wei et al. (2012) found that foliar application increased Zn concentration in the polished grains of three rice cultivars by 18–28 % and Zhang et al. (2012) found significantly greater Zn concentrations in 60–65 % extraction wheat flour (i.e., bran and germ removed) with soil ($P<0.05$) and foliar ($P<0.001$) applications of Zn. Despite these findings, when looking across the studies reviewed here, application of Zn via the soil increased concentrations of Zn in brown rice (median=110, Q1=102, Q3=120 % of control, $n=27$) but not white rice (median=99, Q1=98, Q3=104 % of control, $n=3$) and foliar application increased concentrations more in brown (median=130, Q1=114, Q3=147 % of control, $n=28$) than white (median=117, Q1=112, Q3=122 %

of control, $n=6$) rice. Thus, agronomic biofortification of rice may be less effective at increasing Zn in the diet than assumed and future studies could confirm or allay this concern by reporting data for both whole grain and polished rice.

No studies were identified that reported the effect of Zn application via soil on PA concentration in the grain of rice, nor via foliar spray in maize. Were such data available, it might be that estimates of efficacy and cost-effectiveness of the agronomic biofortification strategies would improve. In addition, this review excluded data reporting the effects of residual soil-applied Zn in subsequent crops. Thus, applied Zn that is not taken up by the crop or permanently ‘fixed’ within mineral phases is potentially available for subsequent crop uptake and some studies show a cumulative increase in grain Zn concentrations in successive seasons following Zn application (Srivastava et al. 2009; Wang et al. 2012; Abid et al. 2013; Manzeke et al. 2014), although positive residual effects are not always found (Lu et al. 2012; Yang et al. 2011a). Therefore, the efficacy of Zn fertilisation might be under-estimated here by excluding residual effects, especially in lower-pH soils. Conversely, meta-analyses are subject to systematic bias due to preferential reporting and publishing of ‘positive’ findings (Dickersin et al. 1992). This may lead to an over-estimate of the efficacy of Zn-enrichment on concentrations of Zn in grains.

Cereals contribute ca. 50 % or more of energy intake across 46 countries in Africa, but root and tuber crops also contribute >30 % of energy supplies in ten countries (Joy et al. 2014). Concentration of Zn in the tuber of potato (*Solanum tuberosum* L.) is generally low (i.e., 10–20 mg kg⁻¹) due to limited translocation of Zn from shoots to tubers via the phloem, although there is significant variation between genotypes and concentrations up to ca. 30 mg kg⁻¹ are achievable with foliar Zn application (White and Broadley 2011; White et al. 2012). Hence, Zn biofortification of potatoes appears to be feasible, in principle. The prospect for supplying Zn via granular fertilisers to cassava and sweet potato (*Ipomoea batatas* L.) is likely to be more limited as these crops are generally grown with few external inputs (Kelly 2006), while a foliar fertilisation programme must consider that the leaves of these crops are consumed in some cultures in sub-Saharan Africa. Leafy vegetables contribute little to dietary energy, but substantially to dietary Zn, intakes due to greater concentrations of Zn in leaves than grains, tubers or fruits

(Broadley et al. 2007). For example, Joy et al. (2015) report concentrations of Zn in edible leaves from Malawi of *ca.* 40–70 mg kg⁻¹ DW. The concentration of Zn in edible leaves is dependent on both environmental factors, such as the concentration of extractable Zn in the soil, and genetic factors which show high heritability in some, but not all, species (Wu et al. 2007, 2008; Broadley et al. 2010); thus there is scope for biofortification of edible leaves through agronomic or breeding approaches.

Effect of Zn fertilisers on Zn deficiency prevalence

Assumptions were made regarding the coverage of fertilisers and hence the proportion of crops that could receive fertiliser enriched with Zn. The proportion of maize, rice, wheat and teff receiving fertiliser was derived from the ratio of national N demand and usage. Demand may be underestimated (thus coverage overestimated) as some crops were not included (e.g., horticultural and oil crops), or may be over-estimated (thus coverage underestimated) as some non-target crops such as millet and sorghum are grown extensively in sub-Saharan Africa with little fertiliser applied (Ahmed et al. 2000). In addition, fertiliser usage data were generally derived from national government statistics (IFDC 2011; 2012a, b, c, d, e, f; 2013b, c) and are likely to vary in accuracy. Fertiliser usage data for 2009 were used due to the availability of IFDC reports. However, usage, and thus the potential reach of a programme to enrich granular fertilisers with Zn, is likely to vary annually depending on prices, farmer purchasing power and government subsidy programmes. For example, estimated N consumption in Zambia in 2009 was 39,400 t based on total fertiliser consumption of 171,000 t in 2007/08 (IFDC 2013c). By 2012/13, estimated fertiliser consumption had increased to 250,000–300,000 t (IFDC 2013c).

A further limitation of the study is that average changes in the composition of maize, rice and wheat (a product of the average effect of Zn-enriched fertilisers on crop composition and coverage of fertiliser usage) were applied to all of the national supply of these food items provided in the Food Balance Sheets. However, a national programme to introduce Zn enrichment via soil or foliar applied fertilisers will only alter the composition of crops produced in-country. Thus, the effect on average Zn concentration may be over-estimated as imported crops are not enriched while some of the

benefits of the fortification may not be captured as exported crops are enriched.

Estimating the cost-effectiveness of an agronomic biofortification approach to address Zn deficiency in sub-Saharan Africa

In the current study, costs of Zn and knapsack sprayers were considered whereas other costs, including for agricultural extension, intra-national distribution of consumables and quality control, were not. This is likely to underestimate the cost of the modelled strategies. Estimating full costs would require detailed study of national agricultural extension services, laboratory capacities etc.

The EAR cut-point approach might underestimate the cost-effectiveness of a population-level fortification programme to alleviate Zn deficiency as only the reduction in deficiency prevalence is considered. In this study, ‘deficient’ status was defined as dietary Zn supply below the mean national EAR. Increasing Zn concentration in staple foods and in the national diet moves a proportion of the population from below to above the EAR cut point. However, those who remain below the EAR may still have derived health benefits from increased Zn intake, (e.g., individuals who move from ‘severe’ to ‘mild’ Zn deficiency), and this benefit is not captured. Moreover, a non-linear relationship between the level of dietary micronutrient deficiencies and the severity of related health outcomes is normally assumed (Stein et al. 2005). Hence, even if an intervention does not completely eliminate a deficiency, it will have a relatively larger impact when alleviating the more severe levels of the deficiency.

Several studies have demonstrated that Zn-enrichment of granular fertilisers can be a cost-effective strategy due to improvements in crop yield (e.g., van Asten et al. 2004; Harris et al. 2007; Cakmak 2009). From the limited yield data available in the studies reviewed here, it appears that soil-applied Zn has a small (*ca.* 10 %) positive impact on yield of maize, rice and wheat while foliar application has minimal effect. The lack of yield response with foliar sprays may be because post-flowering applications were preferred as these late applications have a greater impact on grain Zn concentration with a smaller impact on grain yield (Cakmak et al. 2010; Mabesa et al. 2013). The very high yield responses (i.e., >150 % greater than control) found in a few cases are likely to be due to

severe crop deficiency of Zn and these plots also exhibited great response in grain Zn concentration. Thus, an economic argument for the use of Zn fertilisers due to yield improvements will be highly dependent on the soil characteristics.

The feasibilities of the different scenarios require consideration. Previously, it has been suggested that the yield of crop varieties bred for high-Zn concentration must be maintained or improved if farmer acceptance is to be encouraged (Welch and Graham 2004) and yield improvements have been an important driver of the uptake of Zn-enriched fertilisers in Turkey in areas of Zn-deficient soils (Cakmak 2009). Further studies are required to test whether potential yield improvements due to granular or foliar Zn application are sufficient to drive their uptake among resource-poor smallholder farmers in sub-Saharan Africa. Governments or international donors might be persuaded to subsidise or mandate Zn-enrichment of fertilisers due to the potential public health benefits, possibly implemented through existing fertiliser subsidy schemes. To the authors' knowledge, acceptance of micronutrient sprays by smallholder farmers in sub-Saharan Africa has not been studied and is likely to depend on observable benefits such as yield improvements. In addition, it is questionable whether knapsack sprayers are suitable for foliar application of Zn to maize, which might be *ca.* 2 m in height at tasselling, compared to *ca.* 1 m for mature stands of rice, wheat and teff.

Comparison of agronomic biofortification with other strategies to alleviate Zn deficiency

A fertiliser approach can be compared directly against other Zn interventions. Crop breeding is another strategy to potentially decrease the prevalence and disease burden of Zn deficiency. The HarvestPlus (H+) programme is developing nutrient-rich staple crops through exploitation of existing genotypic variation including in wild relatives, setting target Zn concentrations of 38, 28 and 38 mg kg⁻¹ DW in whole maize grain, polished rice grain and whole wheat grain, respectively (Bouis and Welch 2010; Velu et al. 2014). Stein et al. (2006) estimated that biofortification of crops by breeding for high Zn concentration would be one-to-three orders of magnitude more cost effective than the fertiliser approaches modelled here. However, the potential of new varieties to deliver

greater concentrations of Zn in the grain depends on there being plant-accessible Zn stores in the soil, thus breeding and agronomic biofortification strategies are likely to be complementary.

Fieldier et al. (2013) estimated that fortifying maize meal with a premix containing Zn at large-scale mills in Zambia could save 5657 DALYs annually of which 1757 were due to Zn deficiency, at a cost of US\$ 401 *per* DALY saved. The cost *per* DALY saved is favourable compared to application of Zn via the soil and equivalent or slightly more expensive than via foliar spray, although it should be noted that this is not a direct comparison as the premix also contained iron and vitamin A. Flour fortification during milling currently has limited reach in Zambia as few households purchase maize flour from large, centralised milling factories and those that do are generally wealthier with greater baseline Zn intakes (Fieldier et al. 2013). Thus, while application of Zn to crops via the soil is approximately 10-fold more expensive than via foliar sprays or fortification of flour at centralised mills, it has the potential to reach more households and consequently be more equitable in outcome.

In the studies reviewed here, median Zn concentrations in grain from control plots were 19.0 (Q1=15.4, Q3=22.0, *n*=7) and 15.8 (Q1=9.8, Q3=25.2, *n*=141) mg kg⁻¹ for whole maize and wheat grains, respectively, and 18.8 (Q1=15.2, Q3=23.9, *n*=6) and 18.8 (Q1=13.4, Q3=27.6, *n*=44) mg kg⁻¹ for polished and brown rice, respectively. That there was no difference in median Zn concentration between polished and brown rice samples across studies is surprising and cannot be explained with the available data. If brown and polished rice are considered together and median increases found in reviewed studies are applied, Zn concentrations of 23.4, 16.9 and 22.4 mg kg⁻¹ in maize, rice and wheat appear achievable using soil-applied Zn, and 24.7, 19.8 and 30.6 mg kg⁻¹ using foliar-applied Zn. Even with 100 % coverage of soil or foliar-applied Zn, these concentrations are well below the H+ breeding targets (Fig. 1; White and Broadley 2009; Bouis and Welch 2010). Thus, while agronomic biofortification of staple grains with Zn may be a useful strategy to mitigate inadequate dietary Zn supplies, the elimination of Zn deficiency will require complementary approaches including crop breeding, dietary diversification and possibly fortification during processing. If synergies can be

exploited when pursuing such a combined approach, the cost-effectiveness of these interventions might also improve.

Zinc is just one of a number of micronutrients with widespread risk of deficiency in sub-Saharan Africa. For example, in Malawi, approximately 80 % of the population is at risk of selenium (Se) deficiency due to low concentrations of Se in edible portions of crops grown on low-pH soils (Chilimba et al. 2011; Hurst et al. 2013; Joy et al. 2015). Adding Se to the staple crop maize via subsidised fertiliser could supply adequate amounts of Se in the diets of the *ca.* 1.5 million households who benefit from the Malawi government Fertiliser Input Subsidy Scheme, at a total cost of 250–550 US\$ k year⁻¹ (Chilimba et al. 2012). This is approximately 50-fold cheaper than scenario one in the present study, at US\$ 0.016–0.035 *capita*⁻¹ year⁻¹ compared to US\$ 1.08 *capita*⁻¹ year⁻¹, assuming that costs are spread equally across the national population. However, unlike for Zn, the disease burden of Se deficiency has not yet been quantified so cost *per* DALY saved cannot be estimated using the same frameworks. Agronomic fortification via soil-applied fertiliser requires only 5 g ha⁻¹ year⁻¹ of Se compared to *ca.* 10–25 and 1–4 kg ha⁻¹ year⁻¹ of Zn via soil and foliar application methods, respectively. Thus, although the unit cost of Se is greater than that of Zn, fortification with Zn is more expensive. However, combining multiple elements such as Zn and Se in granular or foliar fertilisers could deliver wider health benefits and improve the cost-effectiveness of agronomic biofortification strategies.

Conclusions

Agronomic biofortification of crops with elements important for human health has been advocated as a public health strategy to address mineral element deficiencies in humans that can have severe consequences for the well-being of individuals and the welfare of affected societies. We systematically reviewed the literature for studies of the impact of Zn fertiliser on Zn and PA concentrations in the grains of maize, rice and wheat. In a simplified meta-analysis, the median effects of soil-applied Zn on the concentration of Zn in the grains of

maize, rice and wheat were 23, 7 and 19 % increases above the control, respectively, while the corresponding figures for foliar applied Zn were 30, 25 and 63 %.

We focused on ten countries in sub-Saharan Africa that currently implement fertiliser subsidy schemes or have strong governmental control over fertiliser imports and therefore have the necessary leverage to implement and enforce Zn enrichment of granular fertiliser. For the nine countries with the necessary data available, enriching subsidised fertiliser with Zn could save a total of 63 k DALYs year⁻¹ lost due to Zn deficiency with cost effectiveness ranging from US\$ 624 to 5747 DALY⁻¹ saved. Enriching subsidised and non-subsidised fertilisers in the eight countries with necessary data could save a total of 83 k DALYs year⁻¹ with cost effectiveness ranging from US\$ 977 to 5893 DALY⁻¹. Foliar sprays may be a more cost-effective approach, saving 375 and 523 k DALYs year⁻¹ for 50 and 75 % coverage of cereals, respectively, at a cost of US\$ 46 to 347 DALY⁻¹ although it is likely that there would be significant administrative costs in implementing such a programme and these costs were not considered here.

Cost-effectiveness of the fertiliser approach varies and, if compared against international cost-effectiveness benchmarks, these results indicate that adoption of a fertiliser approach needs to be assessed on a case-by-case basis to allow decision-makers to optimise the allocation of scarce resources to alternative and complementary public health interventions. Generally, the cost-effectiveness of foliar-applied Zn appears to be equivalent to fortification of staple flours at centralised milling facilities. Soil-applied Zn appears to be more expensive but has the potential advantage of reaching more households. Moreover, synergies might be realised if agronomic (fertilisation) and genetic (breeding) biofortification efforts are combined, potentially improving both impact and cost-effectiveness of these interventions.

Acknowledgments Funding for EJM's Studentship is provided by the University of Nottingham, U.K. and the British Geological Survey (BGS). This study was also supported through a Royal Society-DFID Capacity Building Initiative Network Grant entitled "Strengthening African Capacity in soil geochemistry to inform agricultural and health policies" (AN130007). The study is published with the permission of the Director of BGS. The authors have no conflicts of interest to declare.

References

- Abebe Y, Bogale A, Hambidge KM, Stoecker BJ, Arbide I, Teshome A, Krebs NF, Westcott JE, Bailey KB, Gibson RS (2007) Inadequate intakes of dietary zinc among pregnant women from subsistence households in Sidama, Southern Ethiopia. *Public Health Nutr* 11:379–386. doi:10.1017/S1368980007000389
- Abid M, Ahmed N, Qayyum MF, Shaaban M, Rashid A (2013) Residual and cumulative effect of fertilizer zinc applied in wheat-cotton production system in an irrigated arid soil. *Plant Soil Environ* 59:505–510
- Ahmad W, Watts MJ, Imtiaz M, Ahmed I, Zia MH (2012) Zinc deficiency in soils, crops and humans: a review. *Agrochimica* 56:65–97
- Ahmed MM, Sanders JH, Nell WT (2000) New sorghum and millet cultivar introduction in Sub-Saharan Africa: impacts and research agenda. *Agr Syst* 64:55–65. doi:10.1016/S0308-521X(00)00013-5
- Alloway BJ (2008) Zinc in soils and crop nutrition, 2nd edn. International Zinc Association and International Fertilizer Industry Association, Brussels
- Archer E, Hand GA, Blair SN (2013) Validity of U.S. nutritional surveillance: National Health and Nutrition Examination Survey caloric energy intake data, 1971–2010. *PLoS One* 8:e76632. doi:10.1371/journal.pone.0076632
- Bitvutskii NP, Magnitkiy SV, Korobeynikova LP, Lukina EI, Soloviova AN, Patevitch VG, Lapshina IN, Matveeva GV (2002) Distribution of iron, manganese, and zinc in mature grain and their mobilization during germination and early seedling development in maize. *J Plant Nutr* 25:635–653. doi:10.1081/PLN-120003387
- Bouis HE, Welch RM (2010) Biofortification—a sustainable agricultural strategy for reducing micronutrient malnutrition in the global south. *Crop Sci* 50:S20–S32. doi:10.2135/cropsci2009.09.0531
- Broadley MR, White PJ, Hammond JP, Zelko I, Lux A (2007) Zinc in plants. *New Phytol* 173:677–702. doi:10.1111/j.1469-8137.2007.01996.x
- Broadley MR, Ó Lochlainn S, Hammond JP, Bowen HC, Cakmak I, Eker S, Erdem H, King GJ, White PJ (2010) Shoot zinc (Zn) concentration varies widely within *Brassica oleracea* L. and is affected by soil Zn and phosphorus (P) levels. *J Horticult Sci Biotechnol* 85:375–380
- Brümmer G, Tiller KG, Herms U, Clayton PM (1983) Adsorption-desorption and/or precipitation-dissolution processes of zinc in soils. *Geoderma* 31:337–354. doi:10.1016/0016-7061(83)90045-9
- Cakmak I (2002) Plant nutrition research: priorities to meet human needs for food in sustainable ways. *Plant Soil* 247:3–24. doi:10.1023/a:1021194511492
- Cakmak I (2004) Identification and correction of widespread zinc deficiency in Turkey—a success story (a NATO-Science for Stability Project). *Proceedings of the International Fertiliser Society* 552. International Fertiliser Society, York, UK
- Cakmak I (2008) Enrichment of cereal grains with zinc: agronomic or genetic biofortification? *Plant Soil* 302:1–17. doi:10.1007/s11104-007-9466-3
- Cakmak I (2009) Enrichment of fertilizers with zinc: an excellent investment for humanity and crop production in India. *J Trace Elem Med Biol* 23:281–289. doi:10.1016/j.jtemb.2009.05.002
- Cakmak I, Kalayci M, Ekiz H, Braun HJ, Kiliç Y, Yilmaz A (1999) Zinc deficiency as a practical problem in plant and human nutrition in Turkey: a NATO-science for stability project. *Field Crop Res* 60:175–188. doi:10.1016/S0378-4290(98)00139-7
- Cakmak I, Kalayci M, Kaya Y, Torun AA, Aydin N, Wang Y, Arisoy Z, Erdem H, Yazici A, Gokmen O, Ozturk L, Horst WJ (2010) Biofortification and localization of zinc in wheat grain. *J Agric Food Chem* 58:9092–9102. doi:10.1021/jf101197h
- Central Statistics Agency (CSA 2011) *Agriculture in figures: key findings of the 2008/09–2010/11 agricultural sample surveys for all sectors and seasons. Country summary. Comprehensive Africa Agriculture Development Program (CAADP) Ethiopia Study, Final Report, Volume 1, 2009. Cited in IFDC (2012a)*
- Chilimba ADC, Young SD, Black CR, Rogerson KB, Ander EL, Watts MJ, Lammel J, Broadley MR (2011) Maize grain and soil surveys reveal suboptimal dietary selenium intake is widespread in Malawi. *Sci Rep* 1:72. doi:10.1038/srep00072
- Chilimba ADC, Young SD, Black CR, Meacham MC, Lammel J, Broadley MR (2012) Agronomic biofortification of maize with selenium (Se) in Malawi. *Field Crop Res* 125:118–128. doi:10.1016/j.fcr.2011.08.014
- de Haen H, Klasen S, Qaim M (2011) What do we really know? Metrics for food insecurity and undernutrition. *Food Policy* 36:760–769. doi:10.1016/j.foodpol.2011.08.003
- Dickersin K, Min YI, Meinert CL (1992) Factors influencing publication of research results: follow-up of applications submitted to two institutional review boards. *J Am Med Assoc* 267:374–378. doi:10.1001/jama.267.3.374
- Duggan C, MacLeod WB, Krebs NF, Westcott JL, Fawzi WW, Premji ZG, Mwanakasale V, Simon JL, Yeboah-Antwi K, Hamer DH (2005) Plasma zinc concentrations are depressed during acute phase response in children with falciparum malaria. *J Nutr* 135:802–807
- Eichert T, Fernández V (2012) Uptake and release of elements by leaves and other aerial plant parts, Chapter 4. In: Marschner P (ed) *Marschner's mineral nutrition of higher plants*, 3rd edn. Academic, London. doi:10.1016/B978-0-12-384905-2.00004-2
- Ekiz H, Bağcı SA, Kiral AS, Eker S, Gültekin I, Alkan A, Cakmak I (1998) Effects of zinc fertilization and irrigation on grain yield and zinc concentration of various cereals grown in zinc-deficient calcareous soils. *J Plant Nutr* 21:2245–2256. doi:10.1080/01904169809365558
- Erdal I, Yilmaz A, Taban S, Eker S, Cakmak I (2002) Phytic acid and phosphorus concentrations in seeds of wheat cultivars grown with and without zinc fertilization. *J Plant Nutr* 25: 113–127. doi:10.1081/PLN-100108784
- Erenoglu B, Nikolic M, Romheld V, Cakmak I (2002) Uptake and transport of foliar applied zinc (^{65}Zn) in bread and durum wheat cultivars differing in zinc efficiency. *Plant Soil* 241:251–257. doi:10.1023/A:1016148925918
- Erenoglu EB, Kutman UB, Ceylan Y, Yıldız B, Cakmak I (2011) Improved nitrogen nutrition enhances root uptake, root-to-shoot translocation and remobilization of zinc (^{65}Zn) in wheat. *New Phytol* 189:438–448. doi:10.1111/j.1469-8137.2010.03488.x

- Esrey SA, Habicht J-P, Latham MC, Sisler DG, Casella G (1988) Drinking water source, diarrheal morbidity, and child growth in villages with both traditional and improved water supplies in rural Lesotho, southern Africa. *Am J Public Health* 78: 1451–1455. doi:10.2105/AJPH.78.11.1451
- Fernández V, Eichert T (2009) Uptake of hydrophilic solutes through plant leaves: current state of knowledge and perspectives of foliar fertilization. *CRC Crit Rev Plant Sci* 28:36–68. doi:10.1080/07352680902743069
- Field AP, Gillett R (2010) How to do a meta-analysis. *Br J Math Stat Psychol* 63:665–694. doi:10.1348/000711010X502733
- Fielder JL, Smitz MF, Dupriez O, Friedman J (2008) Household income and expenditure surveys: a tool for accelerating the development of evidence-based fortification programs. *Food Nutr Bull* 29:306–319
- Fielder JL, Lividini K, Kabaghe G, Zulu R, Tehinse J, Bermudez OI, Jallier V, Guyonnet C (2013) Assessing Zambia's industrial fortification options: getting beyond changes in prevalence and cost-effectiveness. *Food Nutr Bull* 34:501–519. doi: 00000034/00000004/art00013
- Fink G, Günther I, Hill K (2011) The effect of water and sanitation on child health: evidence from the demographic and health surveys 1986–2007. *Int J Epidemiol* 40:1196–1204. doi:10.1093/ije/dyr102
- Food and Agriculture Organization of the United Nations (1984) Better farming series. Chapters 22 and 24. Available online: <http://www.fao.org/docrep/006/t0309e/t0309e00.HTM> [accessed September 2014]
- Food and Agriculture Organization of the United Nations (2014a) Food balance sheets. Available online: http://faostat3.fao.org/faostat-gateway/go/to/download/FB/*/E [accessed July 2014]
- Food and Agriculture Organization of the United Nations (2014b) Crop production data. Available online: http://faostat3.fao.org/faostat-gateway/go/to/download/Q/*/E [accessed July 2014]
- Food and Agriculture Organization of the United Nations (2014c) Crop production definitions. Available online: <http://faostat.fao.org/site/384/default.aspx> [accessed July 2014]
- Food and Agriculture Organization of the United Nations (FAO) (2001) Food balance sheets: a handbook. FAO, Rome
- Gegios A, Amthor R, Maziya-Dixon B, Egesi C, Mallowa S, Nungo R, Gichuki S, Mbanaso A, Manary MJ (2010) Children consuming cassava as a staple food are at risk for inadequate zinc, iron, and vitamin A intake. *Plant Foods Hum Nutr* 65:64–70
- Genc Y, Humphries JM, Lyons GH, Graham RD (2005) Exploiting genotypic variation in plant nutrient accumulation to alleviate micronutrient deficiency in populations. *J Trace Elem Med Biol* 18:319–324. doi:10.1016/j.jtemb.2005.02.005
- Ghasemi S, Khoshgoftarmansh AH, Afyuni M, Hadadzadeh H (2013) The effectiveness of foliar applications of synthesized zinc-amino acid chelates in comparison with zinc sulfate to increase yield and grain nutritional quality of wheat. *Europ J Agron* 45:68–74. doi:10.1016/j.eja.2012.10.012
- Gibson RS (2012) Zinc deficiency and human health: etiology, health consequences, and future solutions. *Plant Soil* 361: 291–299. doi:10.1007/s11104-012-1209-4
- Gibson RS, Hotz C, Temple L, Yeudall F, Mtitimuni B, Ferguson E (2000) Dietary strategies to combat deficiencies of iron, zinc, and vitamin A in developing countries: development, implementation, monitoring, and evaluation. *Food Nutr Bull* 21:219–231
- Gibson RS, Hess SY, Hotz C, Brown KH (2008) Indicators of zinc status at the population level: a review of the evidence. *Brit J Nutr* 99:S14–S23. doi:10.1017/S0007114508006818
- Graham RD, Ascher JS, Hynes SC (1992) Selecting zinc-efficient cereal genotypes for soils of low zinc status. *Plant Soil* 146: 241–250. doi:10.1007/BF00012018
- Grice WJ (1990) Tea planter's handbook. Tea Research Foundation of Central Africa, Malawi
- Harris D, Rashid A, Miraj G, Arif M, Shah H (2007) 'On-farm' seed priming with zinc sulphate solution—a cost-effective way to increase the maize yields of resource-poor farmers. *Field Crop Res* 102:119–127. doi:10.1016/j.fcr.2007.03.005
- Harris D, Rashid A, Miraj G, Arif M, Yunas M (2008) 'On-farm' seed priming with zinc in chickpea and wheat in Pakistan. *Plant Soil* 306:3–10. doi:10.1007/s11104-007-9465-4
- Hoagland DR (1948) Lectures on the inorganic nutrition of plants, 2nd edn. Chronica Botanica Company, Waltham
- Hoffland E, Wei C, Wissuwa M (2006) Organic anion exudation by lowland rice (*Oryza sativa* L.) at zinc and phosphorus deficiency. *Plant Soil* 283:155–162. doi:10.1007/s11104-005-3937-1
- Hurst R, Siyame EWP, Young SD, Chilimba ADC, Joy EJM, Black CR, Ander EL, Watts MJ, Chilima B, Gondwe J, Kang'ombe D, Stein AJ, Fairweather-Tait SJ, Gibson RS, Kalimira AA, Broadley MR (2013) Soil-type influences human selenium status and underlies widespread selenium deficiency risks in Malawi. *Sci Rep* 3:1–6. doi:10.1038/srep01425
- Institute of Health Metrics and Evaluation (2014) Global health data exchange, country-level global burden of disease data. Available online: <http://ghdx.healthdata.org/> [accessed July 2014]
- International Fertilizer Development Center (2011) Improving fertilizer markets in West Africa: The fertilizer supply chain in Mali. IFDC, Alabama, U.S.A.. Available online: www.ifdc.org/R-D/Research/Mali_tech_Final11913/ [accessed June 2014]
- International Fertilizer Development Center (2012a) Ethiopia fertilizer assessment. IFDC, Alabama, U.S.A.. Available online: www.ifdc.org/R-D/Research/Ethiopia-Fertilizer-Assessment/ [accessed June 2014]
- International Fertilizer Development Center (2012b) Ghana fertilizer assessment. IFDC, Alabama, U.S.A.. Available online: www.ifdc.org/R-D/Research/Ghana-Fertilizer-Assessment/ [accessed June 2014]
- International Fertilizer Development Center (2012c) Kenya fertilizer assessment. IFDC, Alabama, U.S.A.. Available online: www.ifdc.org/R-D/Research/Kenya-Fertilizer-Assessment/ [accessed June 2014]
- International Fertilizer Development Center (2012d) Improving fertilizer markets in West Africa: The fertilizer supply chain in Nigeria. IFDC, Alabama, U.S.A.. Available online: www.ifdc.org/R-D/Research/Nigeria_tech_Final11913/ [accessed June 2014]
- International Fertilizer Development Center (2012e) Improving fertilizer markets in West Africa: The fertilizer supply chain in Senegal. IFDC, Alabama, U.S.A.. Available online: www.ifdc.org/R-D/Research/Senegal_tech_rev11913.pdf [accessed June 2014]

- International Fertilizer Development Center (2012f) Tanzania fertilizer assessment. IFDC, Alabama, U.S.A.. Available online: www.ifdc.org/R-D/Research/Tanzania-Fertilizer-Assessment/ [accessed June 2014]
- International Fertilizer Development Center (2013a) NEPAD Policy Study: Practices and policy options for the improved design and implementation of fertilizer subsidy programs in sub-Saharan Africa. IFDC, Alabama, U.S.A.. Available online: <http://www.ifdc.org/Documents/NEPAD-fertilizer-study-EN-web/> [accessed June 2014]
- International Fertilizer Development Center (2013b) Malawi fertilizer assessment. IFDC, Alabama, U.S.A.. Available online: www.ifdc.org/R-D/Research/Malawi-Fertilizer-Assessment/ [accessed June 2014]
- International Fertilizer Development Center (2013c) Zambia fertilizer assessment. IFDC, Alabama, U.S.A.. Available online: www.ifdc.org/R-D/Research/Zambia-Fertilizer-Assessment/ [accessed June 2014]
- Jayne TS, Rashid S (2013) Input subsidy programs in sub-Saharan Africa: a synthesis of recent evidence. *Agric Econ* 44:1–16. doi:10.1111/agec.12073
- Johnson SE, Lauren JG, Welch RM, Duxbury JM (2005) A comparison of the effects of micronutrient seed priming and soil fertilization on the mineral nutrition of chickpea (*Cicer arietinum*), lentil (*Lens culinaris*), rice (*Oryza sativa*) and wheat (*Triticum aestivum*) in Nepal. *Exp Agric* 41:427–448. doi:10.1017/S0014479705002851
- Joy EJM, Ander EL, Young SD, Black CR, Watts MJ, Chilimba ADC, Chilima B, Siyame EWP, Kalimbara AA, Hurst R, Fairweather-Tait SJ, Stein AJ, Gibson RS, White PJ, Broadley MR (2014) Dietary mineral supplies in Africa. *Physiol Plant* 151:208–229. doi:10.1111/ppl.12144
- Joy EJM, Broadley MR, Young SD, Black CR, Chilimba ADC, Ander EL, Barlow TS, Watts MJ (2015) Soil type influences crop mineral composition in Malawi. *Sci Total Environ* 505:587–595. doi:10.1016/j.scitotenv.2014.10.038
- Kanwal S, Rahmatullah A, Ranjha AM, Ahmed R (2010) Zinc partitioning in maize grain after soil fertilization with zinc sulphate. *Int J Agric Biol* 12:299–302, doi: 10.13140/2.1.1484.3845
- Kassu A, Yabutani T, Mulu A, Tessema B, Ota F (2008) Serum zinc, copper, selenium, calcium, and magnesium levels in pregnant and non-pregnant women in Gondar, northwest Ethiopia. *Biol Trace Elem Res* 122:97–106. doi:10.1007/s12011-007-8067-6
- Kelly VA (2006) Factors affecting demand for fertilizer in sub-Saharan Africa. Agriculture and rural development discussion paper 23. The World Bank, Washington DC
- Khoshgoftarmansh AH, Sharifi HR, Afuni D, Schulin R (2012) Classification of wheat genotypes by yield and densities of grain zinc and iron using cluster analysis. *J Geochem Explor* 121:49–54. doi:10.1016/j.gexplo.2012.06.002
- Kochian LV (1993) Zinc absorption from hydroponic solution by plant roots. In: Robson AD (ed) Zinc in soils and plants. Kluwer Academic Publishers, Dordrecht, pp 45–58
- Kutman UB, Yildiz B, Ozturk L, Cakmak I (2010) Biofortification of durum wheat with zinc through soil and foliar applications of nitrogen. *Cereal Chem* 87:1–9. doi:10.1094/CCHEM-87-1-0001
- Kutman UB, Yildiz B, Cakmak I (2011) Improved nitrogen status enhances zinc and iron concentrations both in the whole grain and the endosperm fraction of wheat. *J Cereal Sci* 53:118–125. doi:10.1016/j.jcs.2010.10.006
- Liang J, Li Z, Tsuji K, Nakano K, Robert Nout MJ, Hamer RJ (2008) Milling characteristics and distribution of phytic acid and zinc in long-, medium- and short-grain rice. *J Cereal Sci* 48:83–91. doi:10.1016/j.jcs.2007.08.003
- Lim SS, Vos T, Flaxman AD et al (2012) A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the global burden of disease study 2010. *Lancet* 380:2224–2260. doi:10.1016/S0140-6736(12)61766-8
- Lindsay WL, Norvell WA (1978) Development of a DTPA soil test for zinc, iron, manganese, and copper. *Soil Sci Soc Am J* 42:421–428. doi:10.2136/sssaj1978.03615995004200030009x
- Loneragan JF, Grunes DL, Welch RM, Aduayi EA, Tengah A, Lazar VA, Cary EE (1982) Phosphorus accumulation and toxicity in leaves in relation to zinc supply. *Soil Sci Soc Am J* 46:345–352. doi:10.2136/sssaj1982.03615995004600020027x
- Lu XC, Tian XH, Cui J, Zhao AQ, Yang XW, Mai W (2011) Effects of combined phosphorus-zinc fertilization on grain zinc nutritional quality of wheat grown on potentially zinc-deficient calcareous soil. *Soil Sci* 176:684–690. doi:10.1097/SS.0b013e3182331635
- Lu XC, Cui J, Tian XH, Ogunniyi JE, Gale WJ, Zhao AQ (2012) Effects of zinc fertilization on zinc dynamics in potentially zinc-deficient calcareous soil. *Agron J* 104:963–969. doi:10.2134/agronj2011.0417
- Mabesa RL, Impa SM, Grewal D, Johnson-Beebout SE (2013) Contrasting grain-Zn response of biofortification rice (*Oryza sativa* L.) breeding lines to foliar Zn application. *Field Crop Res* 149:223–233. doi:10.1016/j.fcr.2013.05.012
- Makono RCJ, Chanika CM (2008) Density and fertiliser requirement of Catimor coffee in smallholder coffee farmers' fields in Malawi. Horticulture commodity group annual report. Malawi government ministry of agriculture and food security. Available online: www.cabi.org/gara/FullTextPDF/2008/20083323841.pdf [accessed September 2014]
- Mamiro PS, Kolsteren P, Roberfroid D, Tatala S, Opsomer AS, Van Camp JH (2005) Feeding practices and factors contributing to wasting, stunting, and Iron-deficiency anaemia among 3–23 month old children in Kilosa District, rural Tanzania. *J Health Popul Nutr* 23:222–230
- Manzeke GM, Mapfumo P, Mtambanengwe F, Chikowo R, Tendayi T, Cakmak I (2012) Soil fertility management effects on maize productivity and grain zinc content in smallholder farming systems of Zimbabwe. *Plant Soil* 361:57–69. doi:10.1007/s11104-012-1332-2
- Manzeke GM, Mtambanengwe F, Nezomba H, Mapfumo P (2014) Zinc fertilization influence on maize productivity and grain nutritional quality under integrated soil fertility management in Zimbabwe. *Field Crop Res* 166:128–136. doi:10.1016/j.fcr.2014.05.019
- Marschner H (1993) Zinc uptake from soils, Chapter 5. In: Robson AD (ed) Zinc in soils and plants. Kluwer Academic Publishers, Dordrecht
- Martín-Ortiz D, Hernández-Apaolaza L, Gárate A (2009) Efficiency of a NPK fertilizer with adhered zinc lignosulfonate as a zinc source for maize (*Zea mays* L.). *J Agric Food Chem* 57:9071–9078. doi:10.1021/jf9017965

- Miller LV, Krebs NF, Hambidge KM (2007) A mathematical model of zinc absorption in humans as a function of dietary zinc and phytate. *J Nutr* 137:135–141. doi:10.1017/S000711451200195X
- Ministry of Agriculture and Food Security, date unknown. National Fertiliser Strategy. Ministry of Agriculture and Food Security, Lilongwe, Malawi. Available online: http://fsg.afre.msu.edu/mgt/caadp/format_for_national_fertilizer_strategy9.pdf [accessed September 2014]
- Moltedo A, Troubat N, Lokshin M, Sajaia Z (2014) Analyzing food security using household survey data: streamlined analysis with ADePT software. World Bank, Washington, DC
- Müller O, Garenne M, Reitmaier P, van Zweeken AB, Kouyate B, Becher H (2003) Effect of zinc supplementation on growth in West African children: a randomized double-blind placebo-controlled trial in rural Burkina Faso. *Int J Epidemiol* 32:1098–1102. doi:10.1093/ije/dyg190
- Murray CJ (1994) Quantifying the burden of disease: the technical basis for disability-adjusted life years. *Bull World Health Organ* 72:429–445
- Ortiz-Monasterio JI, Palacios-Rojas N, Meng E, Pixley K, Trethowan R, Peña RJ (2007) Enhancing the mineral and vitamin content of wheat and maize through plant breeding. *J Cereal Sci* 46:293–307. doi:10.1016/j.jcs.2007.06.005
- Ozturk L, Yazici MA, Yucel C, Torun A, Cekic C, Bagci A, Ozkan H, Braun H-J, Sayers Z, Cakmak I (2006) Concentration and localization of zinc during seed development and germination in wheat. *Physiol Plant* 128:144–152. doi:10.1111/j.1399-3054.2006.00737.x
- Phattarakul N, Rerkasem B, Li LJ, Wu LH, Zou CQ, Ram H, Sohu VS, Kang BS, Surek H, Kalayci M, Yazici A, Zhang FS, Cakmak I (2012) Biofortification of rice grain with zinc through zinc fertilization in different countries. *Plant Soil* 361:131–141. doi:10.1007/s11104-012-1211-x
- Rengel Z, Batten GD, Crowley DE (1999) Agronomic approaches for improving the micronutrient density in edible portions of field crops. *Field Crop Res* 60:27–40. doi:10.1016/S0378-4290(98)00131-2
- Salgueiro MJ, Zubillaga MB, Lysionek AE, Caro RA, Weill R, Boccio JR (2002) The role of zinc in the growth and development of children. *Nutrition* 18:510–519
- Sandstead HH (2000) Causes of iron and zinc deficiencies and their effects on brain. *J Nutr* 130:S347–S349
- Shivay YS, Kumar D, Prasad R, Ahlawat IPS (2008) Relative yield and zinc uptake by rice from zinc sulphate and zinc oxide coatings onto urea. *Nutr Cycl Agroecosyst* 80:181–188. doi:10.1007/s10705-007-9131-5
- Shrimpton R, Gross R, Darnton-Hill I, Young M (2005) Zinc deficiency: what are the most appropriate interventions? *BMJ* 330:347–349. doi:10.1136/bmj.330.7487.347
- Siekman JH, Allen LH, Bwibo NO, Demment MW, Murphy SP, Neumann CG (2003) Kenyan school children have multiple micronutrient deficiencies, but increased plasma vitamin B-12 is the only detectable micronutrient response to meat or milk supplementation. *J Nutr* 133:S3972–S3980
- Siyame EWP, Hurst R, Wawer AA, Young SD, Broadley MR, Chilimba ADC, Ander EL, Watts MJ, Chilima B, Gondwe J, Kang'ombe D, Kalimbira A, Fairweather-Tait SJ, Bailey KB, Gibson RS (2013) A high prevalence of zinc- but not iron-deficiency among women in rural Malawi: a cross-sectional study. *Int J Vitam Nutr Res* 83:176–187. doi:10.1024/0300-9831/a000158
- Slaton NA, Wilson CE, Ntamatungiro S, Norman RJ, Boothe DL (2001) Evaluation of zinc seed treatments for rice. *Agron J* 93:152–157. doi:10.2134/agronj2001.931152x
- Spears D, Ghosh A, Cumming O (2013) Open defecation and childhood stunting in India: an ecological analysis of new data from 112 districts. *PLoS One* 8:e73784. doi:10.1371/journal.pone.0073784
- Srivastava PC, Singh AP, Kumar S, Ramachandran V, Shrivastava M, D'souza SF (2009) Comparative study of a Zn-enriched post-methanation bio-sludge and Zn sulfate as Zn sources for a rice-wheat crop rotation. *Nutr Cycl Agroecosyst* 85:195–202. doi:10.1007/s10705-009-9258-7
- Stein AJ (2010) Global impacts of human mineral malnutrition. *Plant Soil* 335:133–154. doi:10.1007/s11104-009-0228-2
- Stein AJ (2014) Rethinking the measurement of undernutrition in a broader health context: should we look at possible causes or actual effects? *Global Food Security* 3:193–199. doi:10.1016/j.gfs.2014.09.003
- Stein AD, Barnhart HX, Hickey M, Ramakrishnan U, Schroeder DG, Martorell R (2003) Prospective study of protein-energy supplementation early in life and of growth in the subsequent generation in Guatemala. *Am J Clin Nutr* 78:162–167. doi:10.1016/S0140-6736(08)60205-6
- Stein AJ, Meenakshi JV, Qaim M, Nestel P, Sachdev HPS, Bhutta ZA (2005) Analyzing the health benefits of biofortified staple crops by means of the Disability-Adjusted Life Years approach: a handbook focusing on iron, zinc and vitamin A. HarvestPlus Technical Monograph 4, International Food Policy Research Institute, Washington and International Center for Tropical Agriculture, Cali. Available online: <http://www.harvestplus.org/content/analyzing-health-benefits-biofortified-staple-crops-means-disability-adjusted-life-years-app> [accessed July 2014]
- Stein AJ, Nestel P, Meenakshi JV, Qaim M, Sachdev HPS, Bhutta ZA (2006) Plant breeding to control zinc deficiency in India: how cost-effective is biofortification? *Public Health Nutr* 10:492–501. doi:10.1017/S1368980007223857
- Stoecker BJ, Abebe Y, Hubbs-Tait L, Kennedy TS, Gibson RS, Arbide I, Teshome A, Westcott J, Krebs NF, Hambidge KM (2009) Zinc status and cognitive function of pregnant women in Southern Ethiopia. *Eur J Clin Nutr* 63:916–918. doi:10.1038/ejcn.2008.77
- Suzuki M, Takahashi M, Tsukamoto T, Watanabe S, Matsushashi S, Yazaki J, Kishimoto N, Kikuchi S, Nakanishi H, Mori S, Nishizawa NK (2006) Biosynthesis and secretion of mugineic acid family phytosiderophores in zinc-deficient barley. *Plant J* 48:85–97. doi:10.1111/j.1365-313X.2006.02853.x
- Suzuki M, Tsukamoto T, Inoue H, Watanabe S, Matsushashi S, Takahashi M, Nakanishi H, Mori S, Nishizawa N (2008) Deoxymugineic acid increases Zn translocation in Zn-deficient rice plants. *Plant Mol Biol* 66:609–617. doi:10.1007/s11103-008-9292-x
- Trierweiler JF, Lindsay WL (1969) EDTA-ammonium carbonate soil test for zinc. *Soil Sci Soc Am J* 33:49–54. doi:10.2136/sssaj1969.03615995003300010017x
- Tye AM, Young SD, Crout NMJ, Zhang H, Preston S, Barbosa-Jefferson VL, Davison W, McGrath SP, Paton GI, Kilham K, Resende L (2003) Predicting the activity of Cd²⁺ and Zn²⁺ in

- soil pore water from the radio-labile metal fraction. *Geochim Cosmochim Acta* 67:375–385. doi:10.1016/S0016-7037(02)01138-9
- United Nations Department of Social and Economic Affairs, Population Division (2013) World population prospects: the 2012 revision. United Nations, New York
- Van Asten PJA, Barro SE, Wopereis MCS, Defoer T (2004) Using farmer knowledge to combat low productive spots in rice fields of a Sahelian irrigation scheme. *Land Degrad Dev* 15: 383–396. doi:10.1002/ldr.619
- Veenemans J, Milligan P, Prentice AM, Schouten LRA, Inja N, van der Heijden AC, de Boer LCC, Jansen EJS, Koopmans AE, Enthoven WTM, Kraaijenhagen RJ, Demir AY, Uges DRA, Mbugi EV, Savelkoul HFJ, Verhoef H (2011) Effect of supplementation with zinc and other micronutrients on malaria in Tanzanian children: a randomised trial. *PLoS Med* 8: e1001125. doi:10.1371/journal.pmed.1001125
- Velu G, Ortiz-Monasterio I, Cakmak I, Hao Y, Singh RP (2014) Biofortification strategies to increase grain zinc and iron concentrations in wheat. *J Cereal Sci* 59:365–372. doi:10.1016/j.jcs.2013.09.001
- Wang JW, Mao H, Zhao HB, Huang DL, Wang ZH (2012) Different increases in maize and wheat grain zinc concentrations caused by soil and foliar applications of zinc in Loess Plateau, China. *Field Crop Res* 135:89–96. doi:10.1016/j.fcr.2012.07.010
- Wanzala-Mlobela M, Fuentes P, Mkumbwa S (2013) Practices and policy options for the improved design and implementation of fertilizer subsidy programs in sub-Saharan Africa. NEPAD policy document. IFDC, Alabama
- Wei Y, Shohag MJI, Yang X (2012) Biofortification and bioavailability of rice grain zinc as affected by different forms of foliar zinc fertilization. *PLoS One* 7:e45428. doi:10.1371/journal.pone.0045428
- Welch RM, Graham RD (2004) Breeding for micronutrients in staple food crops from a human nutrition perspective. *J Exp Bot* 55:353–364. doi:10.1093/jxb/erh064
- Wessells KR, Brown KH (2012) Estimating the global prevalence of zinc deficiency: results based on zinc availability in national food supplies and the prevalence of stunting. *PLoS One* 7:e50568. doi:10.1371/journal.pone.0050568
- Wessells KR, Singh GM, Brown KH (2012) Estimating the global prevalence of inadequate zinc intake from national food balance sheets: effects of methodological assumptions. *PLoS One* 7:e50565. doi:10.1371/journal.pone.0050565
- White PJ, Broadley MR (2009) Biofortification of crops with seven mineral elements often lacking in human diets—iron, zinc, copper, calcium, magnesium, selenium and iodine. *New Phytol* 182:49–84. doi:10.1111/j.1469-8137.2008.02738.x
- White PJ, Broadley MR (2011) Physiological limits to zinc biofortification of edible crops. *Front Plant Sci* 2:1–11. doi:10.3389/fpls.2011.00080
- White JG, Zasoski RJ (1999) Mapping soil micronutrients. *Field Crop Res* 60:11–26. doi:10.1016/S0378-4290(98)00130-0
- White PJ, Broadley MR, Hammond JP, Ramsay G, Subramanian NK, Thompson J, Wright G (2012) Bio-fortification of potato tubers using foliar zinc-fertiliser. *J Horticult Sci Biotechnol* 87: 123–129
- Widodo BMR, Rose T, Frei M, Pariaska-Tanaka J, Yoshihashi T, Thomson M, Hammond JP, Aprile A, Close TJ, Ismail AM, Wissuwa M (2010) Response to zinc deficiency of two rice lines with contrasting tolerance is determined by root growth maintenance and organic acid exudation rates, and not by zinc-transporter activity. *New Phytol* 186:400–414. doi:10.1111/j.1469-8137.2009.03177.x
- Wissuwa M, Ismail AM, Graham RD (2008) Rice grain zinc concentrations as affected by genotype, native soil-zinc availability, and zinc fertilization. *Plant Soil* 306:37–48. doi:10.1007/s11104-007-9368-4
- World Bank (2014) National gross domestic product *per capita*. Available online: <http://data.worldbank.org/indicator/NY.GDP.PCAP.CD> [accessed October 2014]
- World Bank (1993) World development report. World Bank, Washington DC
- World Health Organization and Food and Agriculture Organization of the United Nations (2004) Vitamin and mineral requirements in human nutrition, 2nd edn. WHO, FAO, Geneva
- World Health Organization of the United Nations (1995) Expert committee report: physical status: the use and interpretation of anthropometry. Technical report series 854. WHO, Geneva
- World Health Organization of the United Nations (2001) Macroeconomics and health: investing in health for economic development. Report of the commission on macroeconomics and health. WHO, Geneva
- World Health Organization of the United Nations (2009) Global health risks: mortality and burden of disease attributable to selected major risks. WHO, Geneva
- Wu J, Schat H, Sun R, Koornneef M, Wang XW, Aarts MGM (2007) Characterization of natural variation for zinc, iron and manganese accumulation and zinc exposure response in *Brassica rapa* L. *Plant Soil* 291:167–180. doi:10.1007/s11104-006-9184-2
- Wu J, Yuan YX, Zhang XW, Zhao J, Song X, Li Y, Li X, Sun R, Koornneef M, Aarts MGM, Wang XW (2008) Mapping QTLs for mineral accumulation and shoot dry biomass under different Zn nutritional conditions in Chinese cabbage (*Brassica rapa* L. ssp. *pekinensis*). *Plant Soil* 310:25–40. doi:10.1007/s11104-008-9625-1
- Xue YF, Yue SC, Zhang YQ CZL, Chen XP YFC, Cakmak I, McGrath SP, Zhang FS, Zou CQ (2012) Grain and shoot zinc accumulation in winter wheat affected by nitrogen management. *Plant Soil* 361:153–163. doi:10.1007/s11104-012-1510-2
- Yang XW, Tian XH, Gale WJ, Cao YX, Lu XC, Zhao AQ (2011a) Effect of soil and foliar zinc application on zinc concentration and bioavailability in wheat grain grown on potentially zinc-deficient soil. *Cereal Res Commun* 39:535–543
- Yang XW, Tian XH, Lu XC, Gale WJ, Cao YX (2011b) Foliar zinc fertilization improves the zinc nutritional value of wheat (*Triticum aestivum* L.) grain. *Afr J Biotechnol* 10:14778–14785. doi:10.5897/AJB11.780
- Yilmaz A, Ekiz H, Torun B, Gültekin I, Karanlık S, Bağcı SA, Cakmak I (1997) Effect of different zinc application methods on grain yield and zinc concentration in wheat cultivars grown on zinc-deficient calcareous soils. *J Plant Nutr* 20: 461–471. doi:10.1080/01904169709365267
- Zhang YQ, Sun YX, Ye YL, Karim MR, Xue YF, Yan P, Meng QF, Cui ZL, Cakmak I, Zhang FS, Zou CQ (2012) Zinc biofortification of wheat through fertilizer applications in different locations of China. *Field Crop Res* 125:1–7. doi:10.1016/j.fcr.2011.08.003

- Zhao AQ, Xinchun L, Chen Z, Tian X, Yang X (2011) Zinc fertilization methods on zinc absorption and translocation in wheat. *J Agric Sci* 3:28–35. doi:[10.5539/jas.v3n1p28](https://doi.org/10.5539/jas.v3n1p28)
- Zhao AQ, Tian XH, Cao YX, Lu XC, Liu T (2014) Comparison of soil and foliar zinc application for enhancing grain zinc content of wheat when grown on potentially zinc-deficient calcareous soils. *J Sci Food Agric* 94:2016–2022. doi:[10.1002/jsfa.6518](https://doi.org/10.1002/jsfa.6518)
- Zou CQ, Zhang YQ, Rashid A, Ram H, Savasli E, Arisoy RZ, Ortiz-Monasterio I, Simunji S, Wang ZH, Sohu V, Hassan M, Kaya Y, Onder O, Lungu O, Yaqub Mujahid M, Joshi AK, Zelenskiy Y, Zhang FS, Cakmak I (2012) Biofortification of wheat with zinc through zinc fertilization in seven countries. *Plant Soil* 361:119–130. doi:[10.1007/s11104-012-1369-2](https://doi.org/10.1007/s11104-012-1369-2)