




Directional Metric Pseudo Subregularity of Set-valued Mappings: a General Model

Huynh Van Ngai¹ · Nguyen Huu Tron¹ · Nguyen Van Vu¹ · Michel Théra^{2,3} 

Received: 10 October 2018 / Accepted: 3 October 2019 / Published online: 25 November 2019
© Springer Nature B.V. 2019

Abstract

This paper investigates a new general pseudo subregularity model which unifies some important nonlinear (sub)regularity models studied recently in the literature. Some slope and abstract coderivative characterizations are established.

Keywords Abstract subdifferential · Metric regularity · Directional metric regularity · Metric subregularity · Directional Hölder metric subregularity · Directional metric pseudo-subregularity · Coderivative · Slope

Mathematics Subject Classification (2010) 49J52 · 49J53 · 90C30

1 Introduction

Over the past decades, mathematical ideas based on the use of advanced techniques of generalized differentiation have allowed to make significant advances in the study of generalized equations, that is inclusions governed by set-valued mappings. These inclusions/generalized equations cover many important problems in various areas of mathematics and applied sciences such as physics, mechanics, and economics: equations; inequality systems;

Tribute to Professor Alexander Kruger on his sixty-fifth birthday. With recognition for research achievement and friendship

Research of the first author is supported by VIASM.

Research of the second author is funded by Vietnam National Foundation for Science and Technology Development (NAFOSTED) under grant number 101.01-2018.309.

Research of the third author is supported by Vietnam Ministry of Education and Training under grant number B2018-DQN-05

Research of the last author is supported by the Australian Research Council (ARC) grant DP160100854 and benefited from the support of the FMJH Program PGMO and from the support of EDF.

✉ Michel Théra
michel.thera@unilim.fr

Extended author information available on the last page of the article.

variational inequalities; complementarity problems; optimal control, and various other topics. Also, due to their importance, as well as from their theoretical point of view and their broad applicability, variational systems have attracted the interest of many mathematicians.

It is now well known that metric regularity is one of the pillars when studying these generalized equations. This concept has emerged in the last decades, mainly after the contribution of Borwein [4], even if claimed by Ioffe [17, 18], “*the roots of this concept go back to a circle of fundamental regularity ideas of classical analysis embodied in such results as the implicit function theorem, the Banach open mapping theorem, theorems of Lyusternik and Graves, on the one hand, and the Sard theorem and the Thom-Smale transversality theory, on the other*”. Nowadays, this concept is commonly regarded as central for studying the existence and behavior of solutions of nonlinear equations under small perturbations of the data. The crucial role of metric regularity in Optimization and Variational Analysis as well as researcher’s interest in this field can be found in many seminal works including for instance [4, 7, 11, 16–19, 29], and references therein. By the time and demand of use and applications, variants of this property have emerged suitable to practical problems. Weaker/stronger versions: calmness, (strong) (Hölder) metric sub/regularity, semiregularity or equivalent versions: pseudo Lipschitz, linear openness were studied and have proved to have an important role in various applications in Mathematics, especially in Variational Analysis and Optimization [5, 17, 18, 21–23, 29], *etc.*

As recently proposed by Arutyunov [1], Gfrerer [10], Ngai-Théra [14], Ngai-Tron-Théra [12], Ngai-Tron-Tinh [26], a new line of research is to build directional models for these objects. Characterizations of these concepts have been established and successfully applied to study optimality conditions for mathematical programs, for calculating tangent cones, *etc.* This notion of directional regularity is an extension of an earlier notion used by Bonnans and Shapiro [3] to study sensitivity analysis. Later, Ioffe [15] introduced and investigated an extension called relative metric regularity which covers many notions of metric regularity in the literature. In [27] Penot studied this property to establish second-order optimality conditions.

The paper concerns a new type of directional metric regularity. In this article, motivated by the works mentioned above, especially those of (co)authors and by Gfrerer, we build a general new model which formally unifies all (sub)regularity models in the literature; especially it covers both directional Hölder metric subregularity introduced by Ngai, Tron and Tinh and metric pseudo subregularity explored by Gfrerer. This property is given in Definition 3.2. Our aims are to give characterizations of the property. First, we establish a slope characterization and after that we move to a subdifferential/coderivative or a limit critical set characterization. Section 2 introduces the mathematical notation and basic definitions. In Section 3, we present the motivation of this paper and some slope characterizations of directional pseudo regularity. In the final section, we explore how the use of an abstract subdifferential may derive to characterizations of metric directional pseudo subregularity in terms of coderivatives.

2 Preliminaries and Notations

For the convenience of the reader, we include in this section the material concerning set-valued analysis and variational analysis that will be extensively used throughout the sequel. We use the monographs of Mordukhovich [24], Ioffe [19], Rockafellar & Wets [29] and Penot [28] as our desk-copies.

For our purposes, we are going to work in the framework of real Banach spaces. If X is such a space, we denote by $\| \cdot \|$ the associated norm and by $d(x, \Omega)$ the distance from $x \in X$ to the subset Ω of X , that is, $d(x, \Omega) := \inf\{\|x - y\| : y \in \Omega\}$. Given X , we denote the topological dual (continuous dual) by X^* , by $\| \cdot \|^*$ the dual norm of $\| \cdot \|$, by $\mathbb{B}_X = \{x \in X : \|x\| \leq 1\}$ the closed unit ball, by $\mathbb{S}_X = \{x \in X : \|x\| = 1\}$ the unit sphere, by $\mathbb{B}(x, r)$ the closed ball with center x and radius r , respectively.

By a set-valued mapping (also named by some authors multifunction), we mean a mapping T from X into the subsets (possibly empty) of another Banach space Y and we use the notation $T : X \rightrightarrows Y$. The graph of T denoted by $\text{gph } T$ is the set of those points in $X \times Y$ such that $y \in T(x)$, while $T^{-1} : Y \rightrightarrows X$, the inverse of T (always defined), is given by $(x, y) \in \text{gph } T \iff (y, x) \in \text{gph } T^{-1}$. We say that T is closed if its graph is closed with respect to the product topology on $X \times Y$. Given a set $K \subset X$, we use the notation cone K for the conic hull of K , that is for the set of all conic combinations $\sum_{i=1}^n \lambda_i x_i$ of points of K where $\lambda_i \geq 0$ for each index i .

Given an extended real-valued function $f : X \rightarrow \mathbb{R} \cup \{+\infty\}$, we use the notation $\text{cl}f$ to denote the lower semicontinuous envelope of f defined by $\text{cl}f(x) = \liminf_{u \rightarrow x} f(u)$, and $\text{Dom } f$ will refer to the domain of f , that is, the set of those points $x \in X$ such that $f(x)$ is finite. We recall that the convex subdifferential of f at $x \in \text{Dom } f$ is the set

$$\partial f(x) := \{x^* \in X^* : \langle x^*, y - x \rangle \leq f(y) - f(x) \text{ for all } y \in X\},$$

with the convention that $\partial f(x) = \emptyset$, when $f(x) = +\infty$.

For the purpose of this study, we use the concept of slope $|\nabla f|(x)$ of a function f at $x \in \text{Dom } f$. This is a quantity introduced by De Giovanni, Marino & Tosques [6] and defined by

$$|\nabla f|(x) := \begin{cases} 0 & \text{if } x \text{ is a local minimum point of } f \\ \limsup_{y \rightarrow x, y \neq x} \frac{[f(x) - f(y)]_+}{\|x - y\|} & \text{otherwise.} \end{cases}$$

where the notation a_+ means a if $a \geq 0$ and 0 if $a < 0$. For $x \notin \text{Dom } f$, we set $|\nabla f|(x) = +\infty$.

When f is a convex function defined on a Banach space and x is not a minimum point, then according to Ioffe [16, Proposition 3.8] (see also Azé & Corvelec [2, Proposition 3.2])

$$|\nabla f|(x) = \sup_{y \neq x} \frac{f(x) - f(y)}{\|x - y\|} \quad \text{and} \quad |\nabla f|(x) = d(0, \partial f(x)).$$

In the following sections, we make use of the notion of abstract subdifferential operator from X into the subsets of X^* . Such an object, denoted by ∂ , satisfies the following conditions:

- (C1) If $f : X \rightarrow \mathbb{R}$ is a convex function which is continuous around $\bar{x} \in X$ and $\beta : \mathbb{R} \rightarrow \mathbb{R}$ is a continuously differentiable function at $t = f(\bar{x})$, then

$$\partial(\beta \circ f)(\bar{x}) \subseteq \{\beta'(f(\bar{x}))x^* \in X^* : \langle x^*, y - \bar{x} \rangle \leq \beta(f(y)) - \beta(f(\bar{x})) \quad \forall y \in X\};$$

- (C2) $\partial f(x) = \partial g(x)$ if $f(y) = g(y)$ for all y in a neighborhood of x ;
- (C3) Let $f_1 : X \rightarrow \mathbb{R} \cup \{+\infty\}$ be a lower semicontinuous function and $f_2, \dots, f_n : X \rightarrow \mathbb{R}$ be Lipschitz functions. If $f_1 + f_2 + \dots + f_n$ attains a local minimum at x_0 , then for any $\varepsilon > 0$, there exist $x_i \in x_0 + \varepsilon \mathbb{B}_X$, $x_i^* \in \partial f_i(x_i)$, $i = 1, 2, \dots, n$ such that $|f_i(x_i) - f_i(x_0)| < \varepsilon$, and $\|x_1^* + x_2^* + \dots + x_n^*\| < \varepsilon$.

We recall that the indicator function δ_C of a closed set C in X is the function defined by $\delta_C(x) = 0$ when $x \in C$ and $\delta_C(x) = +\infty$, otherwise. Given an abstract subdifferential ∂ , the set $\mathbb{N}(C, x) := \partial\delta_C(x)$ is called the *normal cone* to C at x associated with ∂ .

(C4) $\mathbb{N}(C, x)$ is assumed to be a cone for any closed subset C of X .

Let $F : X \rightrightarrows Y$ and $(\bar{x}, \bar{y}) \in \text{gph } F$. Given an abstract subdifferential ∂ and normal cone associated with ∂ , the set-valued mapping $D^*F(\bar{x}, \bar{y}) : Y^* \rightrightarrows X^*$ defined by

$$D^*F(\bar{x}, \bar{y})(y^*) = \{x^* \in X^* : (x^*, -y^*) \in N(\text{gph } F, (\bar{x}, \bar{y}))\} \text{ for all } y^* \in Y^*,$$

is called the coderivative of F associated with ∂ .

We assume further that ∂ satisfies a separable property in the following sense:

(C5) If f is a separable function defined on $X \times Y$, that is, $f(x, y) := f_1(x) + f_2(y)$, $(x, y) \in X \times Y$, where $f_1 : X \rightarrow \mathbb{R} \cup \{+\infty\}$, $f_2 : Y \rightarrow \mathbb{R} \cup \{+\infty\}$, then

$$\partial f(x, y) = \partial f_1(x) \times \partial f_2(y), \quad \text{for all } (x, y) \in X \times Y.$$

It is well known that the proximal subdifferential in Hilbert spaces, the Fréchet subdifferential in Asplund spaces, the viscosity subdifferentials in smooth spaces as well as the Ioffe and the Clarke-Rockafellar subdifferentials in the setting of general Banach spaces are subdifferentials verifying the conditions (C1)-(C5).

3 Slope Characterizations of Directional Pseudo Subregularity

For the understanding of the paper, we recall the definitions of metric sub-regularity and Hölder metric subregularity. We say that a set-valued mapping $F : X \rightrightarrows Y$ between metric spaces X, Y is *metrically subregular* at a point $(\bar{x}, \bar{y}) \in \text{gph } F$ with constant $\tau > 0$, if there exists a neighbourhood U of \bar{x} such that

$$d(x, F^{-1}(\bar{y})) \leq \tau d(\bar{y}, F(x)) \text{ for all } x \in U. \tag{3.1}$$

If in relation (3.1) we replace $d(\bar{y}, F(x))$ by $d(\bar{y}, F(x))^\gamma$, with $\gamma > 0$, then F is said to be *Hölder metrically subregular* at $(\bar{x}, \bar{y}) \in \text{gph } F$ with modulus τ and order γ . For such a situation, (3.1) becomes

$$d(x, F^{-1}(\bar{y})) \leq \tau d(\bar{y}, F(x))^\gamma \text{ for all } x \in U. \tag{3.2}$$

Throughout the rest of the paper, we assume given Banach spaces X and Y_i , $(i = 1, \dots, m)$, as well as a finite family of set-valued mappings with closed graph $T_i : X \rightrightarrows Y_i$, $(i = 1, \dots, m)$. We note $T := (T_1, \dots, T_m) : X \rightrightarrows Y_1 \times \dots \times Y_m$, the set-valued mapping defined by $T(x) := T_1(x) \times \dots \times T_m(x)$ and we consider $\gamma := (\gamma_1, \dots, \gamma_m) \in \mathbb{R}^m$ such that $\gamma_i \geq 1$ for $i = 1, \dots, m$. To begin with, let us first recall the definition of γ -metric pseudo subregularity (γ -MPSR, for short) w.r.t. a given direction u and order γ .

Definition 3.1 (directional pseudo subregularity,[11]) A set-valued mapping $T = (T_1, \dots, T_m) : X \rightrightarrows Y_1 \times \dots \times Y_m$ is said to be $(\gamma_1, \dots, \gamma_m)$ -metrically pseudo subregular ($\gamma_i \geq 1$) in the direction u at $(\bar{x}, \bar{y}) \in \text{gph } T$ with modulus $\tau > 0$ iff there exist $\varepsilon > 0$ and $\delta > 0$ such that

$$d(x, T^{-1}(\bar{y})) \leq \tau \sum_{i=1}^m \|x - \bar{x}\|^{1-\gamma_i} d(\bar{y}_i, T_i(x)) \tag{3.3}$$

for $x \neq \bar{x}$ in $\mathbb{B}(\bar{x}, \varepsilon) \cap (\bar{x} + \text{cone } \mathbb{B}(u, \delta))$.

As observed by Gfrerer, when $m = 1$, γ -metric pseudo subregularity in the direction $u = 0$ implies Hölder subregularity of order $\frac{1}{\gamma}$. Note also that for $m = 1$, if we choose $T = T_1$ and $\gamma = \gamma_1$, then (3.3) becomes:

$$d(x, T^{-1}(\bar{y}))^\gamma \leq \|x - \bar{x}\|^{\gamma-1} d(x, T^{-1}(\bar{y})) \leq \tau d(\bar{y}, T(x)). \tag{3.4}$$

Hence, if a set-valued mapping $T : X \rightrightarrows Y$ is γ -MPSR at (\bar{x}, \bar{y}) in the direction u , then T is directionally Hölder metrically subregular of order γ at (\bar{x}, \bar{y}) in the direction u as mentioned in [26].

The new idea in this contribution is to consider a general metric pseudo subregularity model called (γ, h) -pseudo subregularity associated with a given function $h := (h_1, \dots, h_m) : X \rightarrow \mathbb{R}_+^m$ and with $\gamma := (\gamma_1, \dots, \gamma_m) \in \mathbb{R}^m$ with $\gamma_i \geq 1$ for $i = 1, \dots, m$. To facilitate ease of reading, we shall introduce some useful real-valued functions corresponding to $T = (T_1, \dots, T_m) : X \rightrightarrows Y_1 \times \dots \times Y_m$ and $\gamma = (\gamma_1, \dots, \gamma_m)$. For each j , we define $\rho_{T_j}(\cdot) = d(\bar{y}_j, T_j(\cdot))$ and set

$$\varphi_{T_j}(x) := \begin{cases} \frac{\rho_{T_j}(x)}{h_j(x)^{\gamma_j-1}} & \text{if } h_j(x) > 0, \\ 0 & \text{otherwise.} \end{cases}$$

Now let us denote by φ_T the sum $\varphi_T = \sum_{j=1}^m \varphi_{T_j}$, by $\psi_T = \text{cl } \varphi_T$ the lower semicontinuous envelope of φ_T and by S the sublevel set $[\psi_T \leq 0]$. Throughout the rest of the document, we make use of the following assumption:

(A) $h_1, \dots, h_m : X \rightarrow \mathbb{R}_+$ are continuous functions and satisfy

$$h_i(x) = 0 \implies \rho_{T_i}(x) = 0. \tag{3.5}$$

Definition 3.2 ((γ, h) -metric pseudo subregularity) Let $T = (T_1, \dots, T_m) : X \rightrightarrows Y_1 \times \dots \times Y_m$ be given and let $(\bar{x}, \bar{y}) \in \text{gph } T$. The mapping T is said to be (γ, h) -metrically pseudo subregular in the direction u at $(\bar{x}, \bar{y}) \in \text{gph } T$ for $h = (h_1, \dots, h_m)$ if there exist $\tau > 0, \varepsilon > 0$ and $\delta > 0$ such that

$$d(x, T^{-1}(\bar{y})) \leq \tau \sum_{1 \leq i \leq m, h_i(x) > 0} \frac{d(\bar{y}_i, T_i(x))}{h_i(x)^{\gamma_i-1}} \tag{3.6}$$

for $x \in \mathbb{B}(\bar{x}, \varepsilon) \cap (\bar{x} + \text{cone } \mathbb{B}(u, \delta))$ with $\sum_{i=1}^m h_i(x) > 0$.

For the direction $u := 0$, we will say briefly that T is (γ, h) -pseudo subregular at (\bar{x}, \bar{y}) (that is, the inequality (3.6) is satisfied for all x near \bar{x}). Let us note that when considering some special mappings h , one recovers the concepts of directional metric subregularity mentioned above. For instance,

- if we choose the functions $h_i(x) = \|x - \bar{x}\|$, one gets metric pseudo subregularity;
- if one considers the case $m = 1, h_1(x) := d(x, T^{-1}(\bar{y}))$ one gets directional Hölder metric subregularity;
- more generally, let be given m subsets $S_i, i = 1, \dots, m$, of $T_i^{-1}(\bar{y})$ containing \bar{x} . In some situations, for suitable sets S_i , when the distances $d(x, S_i)$ can be computed explicitly, the pseudo-subregularity model with respect to $h_i(x) := d(x, S_i), i = 1, \dots, m$. seems to be more convenient than the Hölder metric subregularity one.

Especially, when $\gamma_i = 1 (i = 1, \dots, m)$, one gets the usual metric subregularity.

As mentioned in the introduction, formally this pseudo-subregularity model covers all linear/nonlinear subregularity/error bound models considered in the literature. To see this, let us consider again a multifunction $F : X \rightrightarrows Y$ between two metric spaces X and Y , and for given $(\bar{x}, \bar{y}) \in \text{gph } F$, consider the inclusion

$$S := \{x \in X : \bar{y} \in F(x)\}. \tag{3.7}$$

By means of a suitable residual function $\varphi : X \rightarrow [0, +\infty)$, that is a function such that

$$x \in S \iff \varphi(x) = 0,$$

an usual subregularity model for F at (\bar{x}, \bar{y}) , called φ -subregularity, is defined as the following error bound property:

$$d(x, S) \leq \tau\varphi(x),$$

for all x near \bar{x} , for some $\tau > 0$. Obviously, φ -subregularity can be regarded as pseudo-subregularity by taking

$$h(x) := \frac{d(\bar{y}, F(x))}{\varphi(x)}, \quad x \in X.$$

Let us give further an example for complementarity problems.

Example 3.1 Consider the complementarity system defined by

$$S := \{x \in \mathbb{R}^n : q(x) = 0, \quad f(x) \geq 0, \quad g(x) \geq 0, \quad \langle f(x), g(x) \rangle = 0\}, \tag{3.8}$$

where, $q : \mathbb{R}^n \rightarrow \mathbb{R}^d$, $f := (f_1, \dots, f_m) : \mathbb{R}^n \rightarrow \mathbb{R}^m$; $g := (g_1, \dots, g_m) : \mathbb{R}^n \rightarrow \mathbb{R}^m$.

It is well known that a complementarity system like (3.8) covers, as a particular case, Karush-Kuhn-Tucker (KKT) systems associated to optimization problems, (see, e.g [8, 9, 20]). Given a solution $\bar{x} \in S$, we say that the complementarity system (3.8) satisfies an error bound property at \bar{x} , if there is $\tau > 0$ such that for all x near \bar{x} , one has

$$d(x, S) \leq \tau \left(\|q(x)\| + \sum_{i=1}^m \min\{f_i(x), g_i(x)\} \right). \tag{3.9}$$

This error bound plays a crucial role in the convergence analysis of some algorithms (e.g, the Newton-type methods) for solving system (3.8). It is easy to observe that the error bound property (3.9) is equivalent to the following pseudo-subregularity:

$$d(x, S) \leq \alpha \left(\|q(x)\| + \sum_{i=1}^m (-f_i(x))_+ + \sum_{i=1}^m (-g_i(x))_+ + \sum_{i=1}^m \frac{|f_i(x)g_i(x)|}{h_i(x)} \right), \tag{3.10}$$

where, $(-f_i(x))_+ := \max\{-f_i(x), 0\}$, and $h_i(x) := |\max\{f_i(x), g_i(x)\}|$, $i = 1, \dots, m$. In the error bound relation (3.9), the residual term for both the sign constraints and the complementarity constraint is represented by the function $\sum_{i=1}^m \min\{f_i(x), g_i(x)\}$, while the residual terms in relation (3.10) with respect to the sign constraints and to the complementarity constraint are separably represented. In some situations, this latter representation could be more convenient. In particular, regarding the residual term for the complementarity constraint, the functions $\min\{f_i(x), g_i(x)\}$ ($i = 1, \dots, m$) are generally less regular than the ones $f_i(x), g_i(x)$, e.g, the functions $\min\{f_i(x), g_i(x)\}$ being not necessarily smooth, even if f_i, g_i are smooth.

Throughout the paper, it is convenient to keep in mind the notation used in [14]:

$x \xrightarrow{u} \bar{x}$ is meant to be $x \rightarrow \bar{x}$ if $u = 0$ and $x \rightarrow \bar{x}$ and $\frac{x-\bar{x}}{\|x-\bar{x}\|} \rightarrow \frac{u}{\|u\|}$ otherwise, as well.

The rest of this section will be devoted to establish some characterizations of (γ, h) -metric pseudo subregularity (Definition 3.2). For such a purpose, the next proposition will be useful. First of all, we need the following lemma which gives a relation between the sets S and $T^{-1}(\bar{y})$.

Lemma 3.1 *Let $T : X \rightrightarrows Y$ be a set-valued mapping. Then, one has $S \subset T^{-1}(\bar{y})$. Furthermore, if T has closed graph then $S = T^{-1}(\bar{y})$.*

Proof The proof is straightforward. □

Proposition 3.1 *Suppose that a closed mapping T is not (γ, h) -metrically pseudo subregular in the direction u at $(\bar{x}, \bar{y}) \in \text{gph } T$. Then, for each real sequence $\tau_k \downarrow 0$, of non-negative numbers, there exists a sequence $x_k \xrightarrow{u} \bar{x}$ which satisfies for large integers k the following conditions:*

- i $d(x_k, S) > 0$;
- ii $\psi_T(x_k) \leq \frac{\tau_k}{1-\sqrt{\tau_k}}d(x_k, S)$;
- iii $|\nabla\psi_T|(x_k) \leq \sqrt{\tau_k}$.

Consequently, one has

$$\liminf_{\substack{x \xrightarrow{u} \bar{x}, x \notin S, \\ \psi_T(x)/d(x,S) \rightarrow 0}} \{|\nabla\psi_T|(x)\} = 0. \tag{3.11}$$

Proof Let us note from Lemma 3.1 that $S = T^{-1}(\bar{y})$ since $\text{gph } T$ is closed. According to the definition of metric pseudo subregularity, we can find a sequence $\tilde{x}_k \xrightarrow{u} \bar{x}$ such that

$$\tilde{x}_k \notin S, \quad \tau_k d(\tilde{x}_k, S) > \sum_{\substack{j=1, \dots, m \\ h_j(\tilde{x}_k) > 0}} \frac{d(\bar{y}_j, T_j(\tilde{x}_k))}{h_j(\tilde{x}_k)^{\gamma_j-1}} = \sum_{j=1}^m \varphi_{T_j}(\tilde{x}_k) \geq \psi_T(\tilde{x}_k). \tag{3.12}$$

Hence, the lower semicontinuous (lsc for short) function ψ_T inherits the following property:

$$0 < \psi_T(\tilde{x}_k) < \tau_k d(\tilde{x}_k, S), \quad k = 1, 2, \dots \tag{3.13}$$

Let $\varepsilon_k := \psi_T(\tilde{x}_k) > 0$ and $\lambda_k := \min \left\{ \sqrt{\tau_k} d(\tilde{x}_k, S), \frac{\varepsilon_k}{\tau_k} \right\} > 0$. By the Ekeland variational principle, there exists for each k an element \hat{x}_k such that

- $\|\hat{x}_k - \tilde{x}_k\| \leq \lambda_k$;
- $\psi_T(\tilde{x}_k) - \frac{\varepsilon_k}{\lambda_k} \|\hat{x}_k - \tilde{x}_k\| \geq \psi_T(\hat{x}_k)$;
- the function $f_k : x \in X \mapsto \psi_T(x) + \frac{\varepsilon_k}{\lambda_k} \|x - \hat{x}_k\|$ attains its global minimum at \hat{x}_k .

We are going to establish the following facts:

- i $\hat{x}_k \notin S, k = 1, 2, \dots$;
- ii $\hat{x}_k \xrightarrow{u} \bar{x}$;
- iii $\psi_T(\hat{x}_k) \leq \frac{\tau_k}{1-\sqrt{\tau_k}}d(\hat{x}_k, S)$, for k sufficiently large;
- iv $|\nabla\psi_T|(\hat{x}_k) \leq \sqrt{\tau_k}$, for k sufficiently large.

Assuming that (i) does not hold, one has

$$\tau_k d(\tilde{x}_k, S) \leq \tau_k \|\tilde{x}_k - \hat{x}_k\| \leq \tau_k \lambda_k \leq \varepsilon_k = \psi_T(\tilde{x}_k),$$

in contradiction with (3.13).

To prove (ii), let us note that $\lambda_k = o(\|\tilde{x}_k - \bar{x}\|)$ by virtue of the inequality

$$\lambda_k \leq \sqrt{\tau_k}d(\tilde{x}_k, S) \leq \sqrt{\tau_k}\|\tilde{x}_k - \bar{x}\|.$$

Set $\mu_k := \frac{\|\hat{x}_k - \bar{x}\|}{\|\tilde{x}_k - \bar{x}\|}$. After involving the triangle inequality, one deduces

$$\left(1 - \frac{\|\hat{x}_k - \tilde{x}_k\|}{\|\tilde{x}_k - \bar{x}\|}\right) \|\tilde{x}_k - \bar{x}\| \leq \|\hat{x}_k - \bar{x}\| \leq \left(1 + \frac{\|\hat{x}_k - \tilde{x}_k\|}{\|\tilde{x}_k - \bar{x}\|}\right) \|\tilde{x}_k - \bar{x}\|.$$

Recalling that $\|\hat{x}_k - \tilde{x}_k\| \leq \lambda_k$, we conclude that $\mu_k \rightarrow 1$ as well as $\|\hat{x}_k - \bar{x}\| \rightarrow 0$. If $u \neq 0$, then a few straightforward calculations give us

$$\begin{aligned} \frac{\hat{x}_k - \bar{x}}{\|\hat{x}_k - \bar{x}\|} - \frac{u}{\|u\|} &= \left(\frac{1}{\mu_k}\right) \left(\frac{\tilde{x}_k - \bar{x}}{\|\tilde{x}_k - \bar{x}\|} - \frac{u}{\|u\|}\right) \\ &\quad + \left(\frac{1}{\mu_k}\right) \frac{\hat{x}_k - \tilde{x}_k}{\|\tilde{x}_k - \bar{x}\|} + \left(\frac{1}{\mu_k} - 1\right) \frac{u}{\|u\|}. \end{aligned} \tag{3.14}$$

Using (3.14) it yields $\left\| \frac{\hat{x}_k - \bar{x}}{\|\hat{x}_k - \bar{x}\|} - \frac{u}{\|u\|} \right\| \rightarrow 0$, and therefore, (ii) is proved.

For establishing (iii), we invoke (3.13) and obtain

$$\frac{\psi_T(\hat{x}_k)}{d(\hat{x}_k, S)} \leq \frac{\psi_T(\tilde{x}_k)}{d(\hat{x}_k, S)} \leq \frac{\tau_k d(\tilde{x}_k, S)}{d(\hat{x}_k, S)}.$$

Since

$$|d(\hat{x}_k, S) - d(\tilde{x}_k, S)| \leq \|\hat{x}_k - \tilde{x}_k\| \leq \lambda_k \leq \sqrt{\tau_k}d(\tilde{x}_k, S),$$

for large k we get

$$\frac{\psi_T(\hat{x}_k)}{d(\hat{x}_k, S)} \leq \frac{\tau_k d(\tilde{x}_k, S)}{d(\tilde{x}_k, S) - \sqrt{\tau_k}d(\tilde{x}_k, S)} = \frac{\tau_k}{1 - \sqrt{\tau_k}}.$$

Hence, (iii) follows immediately.

In order to verify (iv), remember that $f_k(\cdot)$ attains a minimum at \hat{x}_k . As a result, the inequality

$$\psi_T(x) + \frac{\varepsilon_k}{\lambda_k} \|x - \hat{x}_k\| \geq \psi_T(\hat{x}_k)$$

holds when x is nearby \hat{x}_k . Equivalently,

$$\frac{\psi_T(\hat{x}_k) - \psi_T(x)}{\|\hat{x}_k - x\|} \leq \frac{\varepsilon_k}{\lambda_k}$$

for all $x \neq \hat{x}_k$ belonging to a neighborhood of \hat{x}_k . In summary, we have when k is large enough

$$|\nabla \psi_T|(\hat{x}_k) \leq \frac{\varepsilon_k}{\lambda_k} = \max \left\{ \frac{\psi_T(\tilde{x}_k)}{\sqrt{\tau_k}d(\tilde{x}_k, S)}, \tau_k \right\} \leq \max \{ \sqrt{\tau_k}, \tau_k \} = \sqrt{\tau_k}. \tag{3.15}$$

Letting $x_k = \hat{x}_k$, the whole proof is established. □

Based on Proposition 3.1, the next theorem offers a sufficient criterion for metric pseudo subregularity.

Theorem 3.1 *Let T, γ, h and ψ_T be defined as above. Suppose that the condition*

$$\liminf_{\substack{x \xrightarrow{u} \bar{x}, x \notin S, \\ \psi_T(x)/d(x, S) \rightarrow 0}} \left\{ |\nabla \psi_T|(x) \right\} > 0 \tag{3.16}$$

is fulfilled. Then, the set-valued mapping T is (γ, h) -metrically pseudo subregular w.r.t. the direction u at $(\bar{x}, \bar{y}) \in \text{gph } T$.

Proof When (3.16) is valid, the set-valued mapping T must be (γ, h) -metrically pseudo subregular as a direct consequence of Proposition 3.1. □

Corollary 3.1 *Under the same assumptions of Theorem 3.1, if now we have*

$$\liminf_{\substack{x \xrightarrow{u} \bar{x}, x \notin S, \\ \psi_T(x)/\|x-\bar{x}\| \rightarrow 0}} \left\{ |\nabla \psi_T|(x) \right\} > 0, \tag{3.17}$$

then, the set-valued mapping T is (γ, h) -metrically pseudo subregular w.r.t. the direction u at $(\bar{x}, \bar{y}) \in \text{gph } T$.

Proof Relation (3.17) implies the one in (3.16). □

Using Theorem 3.1 and taking $h_i(x) = \|x - \bar{x}\|$, one obtains a slope characterization of directional pseudo subregularity of T .

Proposition 3.2 (slope characterization) *If*

$$\liminf_{\substack{x \xrightarrow{u} \bar{x}, x \notin S \\ \|x-\bar{x}\|^{-\gamma} \varphi_{T_i}(x) \rightarrow 0}} \left| \nabla \text{cl} \left(\sum_{i=1}^m \frac{\rho_{T_i}(\cdot)}{\|\cdot - \bar{x}\|^{\gamma_i-1}} \right) \right|(x) > 0, \tag{3.18}$$

then T is γ -MPSR in the direction u at (\bar{x}, \bar{y}) .

Similarly, considering the case $m = 1$, $h_1(x) := d(x, T^{-1}(\bar{y}))$, we derive a characterization of directional Hölder metric subregularity of T .

Proposition 3.3 (slope characterization) *Suppose that the following condition*

$$\liminf_{\substack{x \xrightarrow{u} \bar{x}, x \notin S, \\ \frac{d(\bar{y}, T(x))}{\|x-\bar{x}\|d(x, T^{-1}(\bar{y}))^{\gamma-1}} \rightarrow 0}} \left| \nabla \text{cl} \frac{d(\bar{y}, T(\cdot))}{d(\cdot, T^{-1}(\bar{y}))^{\gamma-1}} \right|(x) > 0 \tag{3.19}$$

holds. Then T is directional Hölder γ -metric subregular in the direction u at (\bar{x}, \bar{y}) .

In many applications, it is sufficient to focus on the case $m = 1$. For such a situation, when applying Theorem 3.1 we have to deal with the slope of the quotient of two functions. The next lemma will be useful to the computation of the slope of such a function.

Lemma 3.2 *Let $f, g : X \rightarrow \mathbb{R} \cup \{+\infty\}$ be lsc extended real-valued functions and let $x \in \text{Dom } f \cap \text{Dom } g$ satisfying $f(x) \geq 0$ and $g(x) > 0$. Suppose in addition that g is continuous at x . Under these assumptions, one has*

$$\left| \nabla \left(\frac{f}{g} \right) \right|(x) \geq \frac{|\nabla f|(x)}{g(x)} - \frac{f(x)}{g(x)^2} |\nabla g|(x). \tag{3.20}$$

Proof Let us denote $\Theta := \frac{f}{g}$. If x is an isolated point of $\text{Dom } \Theta$, then x is a local minimum of both Θ and f , so $|\nabla \Theta|(x) = |\nabla f|(x) = 0$, and the conclusion is trivial. On the contrary,

we fix a sequence of nonnegative reals $(\epsilon_k) \downarrow 0$. Then, there is a sequence $(\delta_k) \downarrow 0$ for which the following property holds true:

$$\|z - x\| \leq \delta_k \implies g(x) - g(z) \leq (|\nabla g|(x) + \epsilon_k)\|x - z\|. \tag{3.21}$$

For each k , we may select $z_k \in \mathbb{B}(x, \delta_k) \setminus \{x\}$ such that

$$f(x) - f(z_k) \geq (|\nabla f|(x) - \epsilon_k)\|x - z_k\|. \tag{3.22}$$

Due to the continuity of g , it is possible to assume $g(z_k) > 0$. We have

$$\begin{aligned} \frac{\Theta(x) - \Theta(z_k)}{\|x - z_k\|} &= \frac{\frac{f(x)}{g(x)} - \frac{f(z_k)}{g(z_k)}}{\|x - z_k\|} \\ &= \frac{1}{g(z_k)} \cdot \frac{f(x) - f(z_k)}{\|x - z_k\|} - \frac{f(x)}{g(x)g(z_k)} \cdot \frac{g(x) - g(z_k)}{\|x - z_k\|} \\ &\geq \frac{1}{g(z_k)} (|\nabla f|(x) - \epsilon_k) - \frac{f(x)}{g(x)g(z_k)} (|\nabla g|(x) + \epsilon_k). \end{aligned}$$

From the last inequality we derive

$$\begin{aligned} \limsup_{k \rightarrow \infty} \frac{\Theta(x) - \Theta(z_k)}{\|x - z_k\|} &\geq \limsup_{k \rightarrow \infty} \left(\frac{1}{g(z_k)} (|\nabla f|(x) - \epsilon_k) - \frac{f(x)}{g(x)g(z_k)} (|\nabla g|(x) + \epsilon_k) \right) \\ &= \frac{1}{g(x)} |\nabla f|(x) - \frac{f(x)}{g(x)^2} |\nabla g|(x). \end{aligned}$$

Since $|\nabla \Theta|(x) = \limsup_{z \rightarrow x} \frac{\Theta(x) - \Theta(z)}{\|x - z\|}$, we obtain

$$|\nabla \Theta|(x) \geq \limsup_{k \rightarrow \infty} \frac{\Theta(x) - \Theta(z_k)}{\|x - z_k\|} \geq \frac{1}{g(x)} |\nabla f|(x) - \frac{f(x)}{g(x)^2} |\nabla g|(x). \quad \square$$

Invoking Lemma 3.2, we obtain Lemma 3.3 used in the sequel for proving Proposition 3.4.

Lemma 3.3 *Let $T : X \rightrightarrows Y$ be a given set-valued mapping and let $(\bar{x}, \bar{y}) \in \text{gph } T$. Suppose that $h : X \rightarrow \mathbb{R}_+$ is locally Lipschitz around \bar{x} and that the subsequent condition is valid as well*

$$\limsup_{x \xrightarrow{u} \bar{x}, x \notin S} \frac{d(x, S)}{h(x)} < +\infty \tag{3.23}$$

for some $u \in X$. Then,

$$\liminf_{\substack{x \xrightarrow{u} \bar{x}, x \notin S, \\ \frac{h(x)^{1-\gamma} \rho_T(x)}{d(x, S)} \rightarrow 0}} \left| \nabla \left(\frac{\text{cl } \rho_T}{h^{\gamma-1}} \right) \right|(x) \geq \liminf_{\substack{x \xrightarrow{u} \bar{x}, x \notin S, \\ h(x)^{-\gamma} \rho_T(x) \rightarrow 0}} \left\{ \frac{|\nabla \text{cl } \rho_T|(x)}{h(x)^{\gamma-1}} \right\}. \tag{3.24}$$

Proof Applying Lemma 3.2 with $f = \text{cl } \rho_T$ and $g = h^{\gamma-1}$ we deduce

$$\left| \nabla \left(\frac{\text{cl } \rho_T}{h^{\gamma-1}} \right) \right|(x) \geq \frac{|\nabla \text{cl } \rho_T|(x)}{h(x)^{\gamma-1}} - \frac{\text{cl } \rho_T(x)}{h(x)^{\gamma-1}} \frac{|\nabla(h^{\gamma-1})|(x)}{h(x)^{\gamma-1}} \tag{3.25}$$

for each $x \notin S$. According to [13], we have

$$|\nabla(h^{\gamma-1})|(x) = (\gamma - 1)h(x)^{\gamma-2}|\nabla h|(x). \tag{3.26}$$

But since h is locally Lipschitz, it holds that $\kappa(x) := |\nabla h|(x)$ is locally bounded around \bar{x} . Let us rewrite (3.25) as follows

$$\begin{aligned} \left| \nabla \left(\frac{\text{cl } \rho_T}{h^{\gamma-1}} \right) \right|(x) &\geq \frac{|\nabla \text{cl } \rho_T|(x)}{h(x)^{\gamma-1}} - (\gamma - 1)\kappa(x) \frac{\text{cl } \rho_T(x)}{[h(x)]^\gamma} \\ &\geq \frac{|\nabla \text{cl } \rho_T|(x)}{h(x)^{\gamma-1}} - (\gamma - 1)\kappa(x) \frac{\rho_T(x)}{h(x)^{\gamma-1}d(x, S)} \cdot \frac{d(x, S)}{h(x)}. \end{aligned} \tag{3.27}$$

Combining (3.23), (3.25) with (3.27) we infer

$$\liminf_{\substack{x \xrightarrow{u} \bar{x}, x \notin S, \\ \frac{h(x)^{1-\gamma} \rho_T(x)}{d(x, S)} \rightarrow 0}} \left| \nabla \left(\frac{\text{cl } \rho_T}{h^{\gamma-1}} \right) \right|(x) \geq \liminf_{\substack{x \xrightarrow{u} \bar{x}, x \notin S, \\ \frac{h(x)^{1-\gamma} \rho_T(x)}{d(x, S)} \rightarrow 0}} \frac{|\nabla \text{cl } \rho_T|(x)}{h(x)^{\gamma-1}}. \tag{3.28}$$

Assume that (x_k) is a sequence in X such that

$$x_k \xrightarrow{u} \bar{x}, x_k \notin S, \frac{h(x_k)^{1-\gamma} \rho_T(x_k)}{d(x_k, S)} \rightarrow 0. \tag{3.29}$$

By virtue of (3.23), we have

$$\limsup_{k \rightarrow \infty} \frac{d(x_k, S)}{h(x_k)} < +\infty,$$

which yields

$$\limsup_{k \rightarrow \infty} \left\{ h(x_k)^{-\gamma} \rho_T(x_k) \right\} = \limsup_{k \rightarrow \infty} \left\{ \frac{h(x_k)^{1-\gamma} \rho_T(x_k)}{d(x_k, S)} \cdot \frac{d(x_k, S)}{h(x_k)} \right\} = 0. \tag{3.30}$$

As a result, we get

$$\liminf_{k \rightarrow \infty} \left\{ \frac{|\nabla \text{cl } \rho_T|(x_k)}{h(x_k)^{\gamma-1}} \right\} \geq \liminf_{\substack{x \xrightarrow{u} \bar{x}, x \notin S, \\ h(x)^{-\gamma} \rho_T(x) \rightarrow 0}} \left\{ \frac{|\nabla \text{cl } \rho_T|(x)}{h(x)^{\gamma-1}} \right\}. \tag{3.31}$$

Combining (3.29) with (3.31), it holds that

$$\liminf_{\substack{x \xrightarrow{u} \bar{x}, x \notin S, \\ \frac{h(x)^{1-\gamma} \rho_T(x)}{d(x, S)} \rightarrow 0}} \frac{|\nabla \text{cl } \rho_T|(x)}{h(x)^{\gamma-1}} \geq \liminf_{\substack{x \xrightarrow{u} \bar{x}, x \notin S, \\ h(x)^{-\gamma} \rho_T(x) \rightarrow 0}} \left\{ \frac{|\nabla \text{cl } \rho_T|(x)}{h(x)^{\gamma-1}} \right\}. \tag{3.32}$$

In summary, (3.28) and (3.32) give us

$$\liminf_{\substack{x \xrightarrow{u} \bar{x}, x \notin S, \\ \frac{h(x)^{1-\gamma} \rho_T(x)}{d(x, S)} \rightarrow 0}} \left| \nabla \left(\frac{\text{cl } \rho_T}{h^{\gamma-1}} \right) \right|(x) \geq \liminf_{\substack{x \xrightarrow{u} \bar{x}, x \notin S, \\ h(x)^{-\gamma} \rho_T(x) \rightarrow 0}} \left\{ \frac{|\nabla \text{cl } \rho_T|(x)}{h(x)^{\gamma-1}} \right\}. \tag{3.33}$$

This completes the proof of Lemma 3.3. □

Based on the previous results, we present a robust version of Theorem 3.1 for the case $m = 1$ which might be more comfortable in practice.

Proposition 3.4 *Let T, \bar{x}, \bar{y} and h satisfy all assumptions of Lemma 3.3. Then, under the following condition*

$$\liminf_{\substack{x \xrightarrow{u} \bar{x}, x \notin S, \\ h(x)^{-\gamma} \rho_T(x) \rightarrow 0}} \frac{|\nabla \text{cl } \rho_T|(x)}{h(x)^{\gamma-1}} > 0, \tag{3.34}$$

the set-valued mapping T is γ -metrically pseudo subregular at \bar{x} for \bar{y} in the direction u .

Proof The proof follows by combining Theorem 3.1 with Lemma 3.3. □

4 Coderivative Characterization of Directional Metric Pseudo Subregularity

Theorem 3.1 provides two sufficient conditions ensuring the validity of directional pseudo subregularity through the slopes corresponding to suitable functions. Gfrerer in his work [11] dealt with such a property using the notion of coderivatives. In order to study the directional metric regularity property, the authors of [25], introduced the notion of limiting critical set around a reference point $(\bar{x}, \bar{y}) \in \text{gph } T$. Following a similar trend, we shall develop an infinitesimal criterion for directional metric pseudo subregularity using the coderivative associated with an abstract subdifferential ∂ . Hereinafter, we assume that the abstract subdifferential ∂ satisfies the following quotient fuzzy rule:

(C₆): Let $f_1, f_2 : X \rightarrow \mathbb{R} \cup \{+\infty\}$ be two locally Lipschitz functions around $x_0 \in X$ with $f_1(x_0) \geq 0, f_2(x_0) > 0$. For any $\varepsilon > 0$ one has

$$\partial \left(\frac{f_1}{f_2} \right) (x_0) \subseteq \bigcup_{x_1, x_2 \in B(x_0, \varepsilon)} \left\{ \frac{f_2(x_0) \partial f_1(x_1) - f_1(x_0) \partial f_2(x_2)}{f_2(x_0)^2} + \varepsilon B_{X^*} \right\}.$$

Note that (C₆) is valid for all usual subdifferentials used in the literature of variational analysis.

Proposition 4.1 *Let $T : X \rightrightarrows Y = Y_1 \times \dots \times Y_m$ be a closed set-valued mapping between two Banach spaces X and Y and $(\bar{x}, \bar{y}) \in \text{gph } T$. Suppose given functions $h_i : X \rightarrow \mathbb{R}_+, (i = 1, \dots, m)$, locally Lipschitz around \bar{x} . Let $u \in X$ and $\gamma \geq 1$ be given for which the following condition is fulfilled*

$$\limsup_{x \xrightarrow{u} \bar{x}, x \notin S} \max_{i=1, \dots, m} \frac{d(x, S)}{h_i(x)} < +\infty. \tag{4.35}$$

If T is not metrically pseudo subregular in the direction u at (\bar{x}, \bar{y}) , then there exist some real sequence $(t_k) \downarrow 0$ together with $(u_k, v_k) \in S_{X \times Y}, u_k^ \in X^*$ and $v_k^* \in Y^*$ such that*

- (a). $\lim_{k \rightarrow \infty} \|u_k\| = 1, \lim_{k \rightarrow \infty} \| \|u\| u_k - u \| = 0;$
- (b). $\lim_{k \rightarrow \infty} \max_{i=1, \dots, m} \{ (t_k)^{1-\gamma_i} \|v_{ki}\| \} = 0;$
- (c). *the vector $(u_k^*, -\tilde{v}_k^*)$ belongs to $N(\text{gph } T, (\bar{x} + t_k u_k, \bar{y} + t_k v_k))$, where $\tilde{v}_k^* \in Y^*$ is given by $\tilde{v}_{ki}^* = h_i(\bar{x} + t_k u_k)^{1-\gamma_i} v_{ki}^*$;*
- (d). $\lim_{k \rightarrow \infty} \left\{ \frac{\sum_{i=1}^m \langle v_{ki}^*, h_i(\bar{x} + t_k u_k)^{1-\gamma_i} v_{ki} \rangle}{\| (h_1(\bar{x} + t_k u_k)^{1-\gamma_1} v_{k1}, \dots, h_m(\bar{x} + t_k u_k)^{1-\gamma_m} v_{km}) \|} \right\} = 1;$
- (e). $\lim_{k \rightarrow \infty} \|u_k^*\| = 0, \lim_{k \rightarrow \infty} \|v_{ki}^*\| = 1.$

Proof For convenience, we assume that the distance on Y is given by

$$\|y\|_Y := \|y_1\|_{Y_1} + \dots + \|y_m\|_{Y_m}.$$

Further, with the aim of simplifying the notation, let us use $\|\cdot\|$ to indicate the norm in any Banach space. By Proposition 3.1, we can find some sequence $x_k \xrightarrow{u} \bar{x}$ such that

- (i) $x_k \notin S$;
- (ii) $\lim_{k \rightarrow \infty} \frac{\psi_T(x_k)}{d(x_k, S)} = 0$;
- (iii) $\lim_{k \rightarrow \infty} \left\{ |\nabla \psi_T|(x_k) \right\} = 0$.

Denoting $\alpha_k := \|x_k - \bar{x}\| > 0$ and assuming $\psi_T(x_k) = \beta_k d(x_k, S)$ where $(\beta_k) \downarrow 0$, for each k , choose some positive parameters σ_k and η_k satisfying $\sigma_k = o(\psi_T(x_k))$ and $\eta_k < \sigma_k$. Making σ_k and η_k smaller if necessary, we may suppose

- $2\eta_k + \sigma_k < \psi_T(x_k)$;
- $\varphi_T(z) > \psi_T(x_k) - \sigma_k$ whenever $\|z - x_k\| \leq \eta_k$.

By (4.35), we have $\min_{j=1, \dots, m} \{h_j(x_k)\} > 0$. Let $\tau_k := |\nabla \psi_T|(x_k)$, then according to the definition, x_k is a local minimum to the function $\psi_T(\cdot) + (\tau_k + \sigma_k)\|\cdot - x_k\|$. Hence, there exists $0 < r_k \leq \min_{j=1, \dots, m} \{h_j(x_k)^{\nu_j} \eta_k\}$ such that

$$\psi_T(x_k) = \min_{\|x - x_k\| \leq 2r_k} \left\{ \psi_T(x) + (\tau_k + \sigma_k)\|x - x_k\| \right\}. \tag{4.36}$$

By the definition of a lsc envelope, we may select some $(\hat{x}_k, \hat{y}_k) \in \text{gph } T$ which fulfills the two properties below:

$$\|\hat{x}_k - x_k\| \leq \sigma_k r_k \quad \text{and} \quad \sum_{j=1}^m \frac{\|\bar{y}_j - \hat{y}_{kj}\|}{h_j(\hat{x}_k)^{\nu_j - 1}} \leq \psi_T(x_k) + \sigma_k r_k. \tag{4.37}$$

Indeed, since $0 < r_k \leq \min_{j=1, \dots, m} \{h_j(x_k)^{\nu_j} \eta_k\}$, h_j are continuous and $\psi_T(x_k) = \text{cl } \varphi_T(x_k)$, we can choose $\hat{x}_k \in X$ such that $\|\hat{x}_k - x_k\| \leq \sigma_k r_k$ and

$$\varphi_T(\hat{x}_k) \leq \psi_T(x_k) + \sigma_k r_k / 2. \tag{4.38}$$

Moreover, there exists $\hat{y}_k \in T_j(\hat{x}_k)$ such that

$$\|\hat{y}_{kj} - \bar{y}_j\| \leq d(\bar{y}_j, T_j(\hat{x}_k)) + \frac{\sigma_k r_k}{2 \sum_{j=1}^m h_j(\hat{x}_k)^{-\nu_j + 1}}.$$

It follows that

$$\begin{aligned} \sum_{j=1}^m \frac{\|\bar{y}_j - \hat{y}_{kj}\|}{h_j(\hat{x}_k)^{\nu_j - 1}} &\leq \sum_{j=1}^m \frac{d(\bar{y}_j, T_j(\hat{x}_k))}{h_j(\hat{x}_k)^{\nu_j - 1}} + \sigma_k r_k \sum_{j=1}^m \frac{1}{h_j(\hat{x}_k)^{\nu_j - 1}} \left(2 \sum_{j=1}^m \frac{1}{h_j(\hat{x}_k)^{\nu_j - 1}} \right)^{-1} \\ &= \sum_{j=1}^m \frac{d(\bar{y}_j, T_j(\hat{x}_k))}{h_j(\hat{x}_k)^{\nu_j - 1}} + \sigma_k r_k / 2 \\ &= \varphi_T(\hat{x}_k) + \sigma_k r_k / 2. \end{aligned}$$

From these estimations and (4.38), one obtains that

$$\sum_{j=1}^m \frac{\|\bar{y}_j - \hat{y}_{kj}\|}{h_j(\hat{x}_k)^{\nu_j-1}} \leq \psi_T(x_k) + \sigma_k r_k/2 + \sigma_k r_k/2 = \psi_T(x_k) + \sigma_k r_k.$$

Using (4.36), we deduce

$$\begin{aligned} \sum_{j=1}^m \frac{\|\bar{y}_j - \hat{y}_{kj}\|}{h_j(\hat{x}_k)^{\nu_j-1}} &\leq \psi_T(x) + (\tau_k + \sigma_k)\|x - x_k\| + \sigma_k r_k \\ &\leq \delta_{\text{gph } T}(x, y) + \sum_{j=1}^m \frac{\|\bar{y}_j - y_j\|}{h_j(x)^{\nu_j-1}} + (\tau_k + \sigma_k)\|x - \hat{x}_k\| \\ &\quad + (\tau_k + \sigma_k)\|x_k - \hat{x}_k\| + \sigma_k r_k. \end{aligned} \tag{4.39}$$

Define an extended real-valued function $f_k : X \times Y \rightarrow \mathbb{R} \cup \{+\infty\}$ by the formula

$$f_k(x, y) := \delta_{\text{gph } T}(x, y) + \sum_{j=1}^m \frac{\|\bar{y}_j - y_j\|}{h_j(x)^{\nu_j-1}} + (\tau_k + \sigma_k)\|x - \hat{x}_k\|. \tag{4.40}$$

Because $\text{gph } T$ is closed, the continuity of h_j implies that f_k is lsc. Observe that $f_k(\hat{x}_k, \hat{y}_k) = \sum_{j=1}^m \frac{\|\bar{y}_j - \hat{y}_{kj}\|}{h_j(\hat{x}_k)^{\nu_j-1}}$ and that

$$f_k(\hat{x}_k, \hat{y}_k) \leq \inf_{\|x - x_k\| \leq 2r_k} f_k(x, y) + \epsilon_k,$$

where $\epsilon_k := (\tau_k + \sigma_k)\|x_k - \hat{x}_k\| + \sigma_k r_k > 0$. Setting $\lambda_k := r_k \sqrt{\sigma_k} > 0$ and applying the Ekeland variational principle, take $(\tilde{x}_k, \tilde{y}_k) \in X \times Y$ such that

- $\|(\hat{x}_k, \hat{y}_k) - (\tilde{x}_k, \tilde{y}_k)\| \leq \lambda_k$;
- $f_k(\hat{x}_k, \hat{y}_k) - \frac{\epsilon_k}{\lambda_k} \|(\hat{x}_k, \hat{y}_k) - (\tilde{x}_k, \tilde{y}_k)\| \geq f_k(\tilde{x}_k, \tilde{y}_k)$;
- $(\tilde{x}_k, \tilde{y}_k)$ is a minimum to the function $(x, y) \in X \times Y \mapsto f_k(x, y) + \frac{\epsilon_k}{\lambda_k} \|(x, y) - (\tilde{x}_k, \tilde{y}_k)\|$ subject to the constraint $\|x - x_k\| \leq 2r_k$.

The last condition along with the properties $(C_1) - (C_6)$ of the subdifferential operator show that, there are some elements $\tilde{x}_k^i, \tilde{x}_{kj}^4 \in X, \tilde{y}_k^i \in Y, \tilde{y}_{kj}^4 \in Y_j$ and also $\tilde{x}_k^{i*}, \tilde{x}_{kj}^{4*} \in X^*, \tilde{y}_k^{i*} \in Y^*, \tilde{y}_{kj}^{4*} \in Y_j^*$ which fulfill the following conditions:

- i $\max \{ \|\tilde{x}_k - \tilde{x}_k^i\|, \|\tilde{x}_k - \tilde{x}_{kj}^4\|, \|\tilde{y}_k - \tilde{y}_k^i\|, \|\tilde{y}_k - \tilde{y}_{kj}^4\| \} \leq \nu_k$, where the parameter ν_k is chosen such that $\nu_k \leq \min \{ \lambda_k, \beta_k \|\bar{y}_1 - \tilde{y}_{k1}\|, \dots, \beta_k \|\bar{y}_m - \tilde{y}_{km}\| \}$;
- ii $(\tilde{x}_k^{1*}, \tilde{y}_k^{1*}) \in \partial_{X \times Y} \delta_{\text{gph } T}(\tilde{x}_k^1, \tilde{y}_k^1) = N(\text{gph } T, (\tilde{x}_k^1, \tilde{y}_k^1))$;
- iii $\tilde{x}_k^{2*} \in \partial_X \|\cdot - \hat{x}_k\|(\tilde{x}_k^2)$;
- iv $(\tilde{x}_k^{3*}, \tilde{y}_k^{3*}) \in \partial_{X \times Y} \|\cdot, \cdot\|(\tilde{x}_k^3, \tilde{y}_k^3)$;
- v $\tilde{y}_{kj}^{4*} \in \partial_{Y_j} \|\bar{y}_j - \cdot\|(\tilde{y}_{kj}^4)$;
- vi $\tilde{x}_{kj}^{4*} \in \partial_X (h_j^{1-\nu_j})(\tilde{x}_{kj}^4) = (1 - \nu_j)h_j(\tilde{x}_{kj}^4)^{-\nu_j} (\partial_X h_j)(\tilde{x}_{kj}^4)$;
- vii $\|\tilde{x}_k^{1*} + (\tau_k + \sigma_k)\tilde{x}_k^{2*} + \frac{\epsilon_k}{\lambda_k} \tilde{x}_k^{3*} + \sum_{j=1}^m \|\bar{y}_j - \tilde{y}_{kj}^4\| \tilde{x}_{kj}^{4*}\| \leq \sigma_k$;
- viii $\|\tilde{y}_k^{1*} + \frac{\epsilon_k}{\lambda_k} \tilde{y}_k^{3*} + h_j(\tilde{x}_{kj}^4)^{1-\nu_j} \tilde{y}_{kj}^{4*}\| \leq \sigma_k$.

We shall establish the conclusion of Proposition 4.1 step-by-step through several auxiliary facts. □

Fact 1 It holds that

$$\lim_{k \rightarrow \infty} \frac{\|\hat{x}_k - x_k\|}{\|x_k - \bar{x}\|} = \lim_{k \rightarrow \infty} \frac{\|\tilde{x}_k^i - x_k\|}{\|x_k - \bar{x}\|} = \lim_{k \rightarrow \infty} \frac{\|\tilde{x}_{kj}^4 - x_k\|}{\|x_k - \bar{x}\|} = 0; \tag{4.41a}$$

$$\lim_{k \rightarrow \infty} \frac{h_j(\hat{x}_k)}{h_j(x_k)} = \lim_{k \rightarrow \infty} \frac{h_j(\tilde{x}_k^i)}{h_j(x_k)} = \lim_{k \rightarrow \infty} \frac{h_j(\tilde{x}_{kj}^4)}{h_j(x_k)} = 1; \tag{4.41b}$$

$$\lim_{k \rightarrow \infty} \left\{ \frac{1}{\psi_T(x_k)} \sum_{j=1}^m \frac{\|\bar{y}_j - \hat{y}_{kj}\|}{h_j(\hat{x}_k)^{\nu_j-1}} \right\} = 1; \tag{4.41c}$$

$$\lim_{k \rightarrow \infty} \left\{ \frac{1}{\psi_T(x_k)} \sum_{j=1}^m \frac{\|\bar{y}_j - \tilde{y}_{kj}^i\|}{h_j(\tilde{x}_k^i)^{\nu_j-1}} \right\} = 1, i = 1, 2, 3; \tag{4.41d}$$

$$\lim_{k \rightarrow \infty} \left\{ \frac{1}{\psi_T(x_k)} \sum_{j=1}^m \frac{\|\bar{y}_j - \tilde{y}_{kj}^4\|}{h_j(\tilde{x}_{kj}^4)^{\nu_j-1}} \right\} = 1. \tag{4.41e}$$

Proof of Fact 1 Firstly, we establish (4.41a). Equality $\lim_{k \rightarrow \infty} \frac{\|\hat{x}_k - x_k\|}{\|x_k - \bar{x}\|} = 0$ is trivial due to the choice of \hat{x}_k . Further, since

$$\max \{ \|\tilde{x}_k^i - \tilde{x}_k\|, \|\tilde{x}_{kj}^4 - \tilde{x}_k\| \} \leq \lambda_k = o(\|x_k - \bar{x}\|),$$

whereas $\|\tilde{x}_k - \hat{x}_k\| \leq \lambda_k = o(\|x_k - \bar{x}\|)$, one gets

$$\max \{ \|\tilde{x}_k^i - x_k\|, \|\tilde{x}_{kj}^4 - x_k\| \} = o(\|x_k - \bar{x}\|),$$

which implies (4.41a).

For (4.41b), let us involve the Lipschitz property of each function h_j . Indeed, for each $j = 1, 2, \dots, m$ there is $L_j > 0$ and $r_j > 0$ for which one has

$$\left. \begin{matrix} \|x - \bar{x}\| \leq r_j, \|x' - \bar{x}\| \leq r_j \\ \|x - x'\| \leq r_j \end{matrix} \right\} \implies |h_j(x) - h_j(x')| \leq L_j \|x - x'\|. \tag{4.42}$$

Thus, when k is large, it holds that

$$|h_j(\hat{x}_k) - h_j(x_k)| \leq L_j \|\hat{x}_k - x_k\| \leq L_j \sigma_k r_k. \tag{4.43}$$

This allows us to write

$$\left| \frac{h_j(\hat{x}_k)}{h_j(x_k)} - 1 \right| \leq L_j \frac{r_k}{h_j(x_k)} \sigma_k. \tag{4.44}$$

By interchanging in turn the role between \hat{x}_k with $\tilde{x}_{kj}, \tilde{x}_k^i$ and repeating the arguments above, we deduce

$$\max \left\{ \left| \frac{h_j(\tilde{x}_{kj}^4)}{h_j(x_k)} - 1 \right|, \left| \frac{h_j(\tilde{x}_k^i)}{h_j(x_k)} - 1 \right| \right\} \leq L_j \frac{r_k}{h_j(x_k)} (\sigma_k + 2\sqrt{\sigma_k}). \tag{4.45}$$

Since $r_k \leq h_j(x_k)^\nu \eta_k$, (4.41b) follows from (4.44) and (4.45).

In the next step, we prove (4.41c). According to the choice of \hat{x}_k , it is possible to derive

$$h_j(\hat{x}_k)^{1-\nu_j} \|\bar{y}_j - \hat{y}_{kj}\| \geq h_j(\hat{x}_k)^{1-\nu_j} d(\bar{y}_j, T_j(\hat{x}_k)) = \varphi_{T_j}(\hat{x}_k).$$

As a result

$$\sum_{j=1}^m \frac{\|\bar{y}_j - \hat{y}_{kj}\|}{h_j(\hat{x}_k)^{\nu_j-1}} \geq \varphi_T(x_k) > \psi_T(x_k) - \sigma_k.$$

Further, we have $\sum_{j=1}^m \frac{\|\tilde{y}_j - \hat{y}_{kj}\|}{h_j(\hat{x}_k)^{\nu_j-1}} < \psi_T(x_k) + \sigma_k r_k$ by the choice of \hat{y}_{kj} . Thus, the limit (4.41c) is a direct consequence of the fact that $\sigma_k = o(\psi_T(x_k))$.

Based on (4.41b), we can derive that the proof of (4.41e) is similar to the one of (4.41d). Thus, it is sufficient to verify (4.41d) only. Indeed, fix an index $i \in \{1, 2, 3\}$. We infer from (4.41b) that

$$\lim_{k \rightarrow \infty} \left\{ \frac{1}{\psi_T(x_k)} \sum_{j=1}^m \frac{\|\tilde{y}_j - \hat{y}_{kj}\|}{h_j(\tilde{x}_k^i)^{\nu_j-1}} \right\} = 1.$$

According to the choice of \tilde{y}_{kj}^i we get

$$\|\tilde{y}_{kj}^i - \hat{y}_{kj}\| \leq \|\tilde{y}_{kj} - \hat{y}_{kj}\| + \|\tilde{y}_{kj}^i - \hat{y}_{kj}\| \leq 2\lambda_k = 2r_k\sqrt{\sigma_k},$$

which yields

$$\begin{aligned} \frac{\|\tilde{y}_{kj}^i - \hat{y}_{kj}\|}{\psi_T(x_k)h_j(\tilde{x}_k^i)^{\nu_j-1}} &\leq 2\sqrt{\sigma_k} \frac{r_k}{\psi_T(x_k)h_j(\tilde{x}_k^i)^{\nu_j-1}} \\ &\leq 2\sqrt{\sigma_k} \frac{h(x_k)^{\nu_j}\eta_k}{\psi_T(x_k)h_j(\tilde{x}_k^i)^{\nu_j-1}}. \end{aligned}$$

Recalling that $\eta_k < \sigma_k = o(\psi_T(x_k))$, we obtain (4.41d) by applying (4.41b). □

Fact 2 We have $\bar{y} \notin T(\tilde{x}_k^1)$.

Let us define

$$t_k := \|(\tilde{x}_k^1 - \bar{x}, \tilde{y}_k^1 - \bar{y})\|, \tag{4.46a}$$

$$u_k := (t_k)^{-1}(\tilde{x}_k^1 - \bar{x}), \tag{4.46b}$$

$$v_{ki} := (t_k)^{-1}(\tilde{y}_{ki}^1 - \bar{y}_i), \quad i = 1, \dots, m. \tag{4.46c}$$

Then the following relations are valid as well

$$\|(u_k, v_k)\| = 1, \tag{4.47a}$$

$$\lim_{k \rightarrow \infty} \|u_k\| = 1, \tag{4.47b}$$

$$\lim_{k \rightarrow \infty} \|u\|u_k - u\| = 0, \tag{4.47c}$$

$$\lim_{k \rightarrow \infty} \{(t_k)^{1-\gamma_i} \|v_{ki}\|\} = 0. \tag{4.47d}$$

Proof of Fact 2 Firstly, by the triangle inequality:

$$\|x_k - \tilde{x}_k^1\| \leq \|x_k - \hat{x}_k\| + \|\hat{x}_k - \tilde{x}_k\| + \|\tilde{x}_k - \tilde{x}_k^1\| \leq \sigma_k r_k + \lambda_k + v_k = o(\eta_k).$$

Hence, for k large $\|x_k - \tilde{x}_k^1\| < \eta_k$. Thus, we infer from the choice of η_k that

$$\varphi_T(\tilde{x}_k^1) > \psi_T(\tilde{x}_k^1) - \sigma_k > \eta_k.$$

This shows that $\bar{y} \notin T(\tilde{x}_k^1)$. Particularly, one has $\tilde{x}_k^1 \neq \bar{x}$, which implies $t_k > 0$. Hence, the elements u_k, v_{ki} are well-defined. The equality (4.47a) is trivial by the definitions (4.46a)–(4.46c). To prove (4.47b), we note that $\|\tilde{x}_k^1 - \bar{x}\| \sim \|x_k - \bar{x}\| = \alpha_k$ according to (4.41a). It follows from the choice of \hat{y}_k that (see (4.37))

$$\|\hat{y}_{ki} - \bar{y}_i\| \leq h_i(\hat{x}_k)^{\nu_i-1}(\psi_T(x_k) + \sigma_k r_k) = o(\alpha_k),$$

which allows for writing

$$\begin{aligned} \|\tilde{y}_{ki}^1 - \bar{y}_i\| &\leq \|\tilde{y}_{ki}^1 - \tilde{y}_{ki}\| + \|\tilde{y}_{ki} - \hat{y}_{ki}\| + \|\hat{y}_{ki} - \bar{y}_i\| \\ &\leq \|\tilde{y}_k^1 - \tilde{y}_k\| + \|\tilde{y}_k - \hat{y}_k\| + \|\hat{y}_{ki} - \bar{y}_i\| \\ &\leq \nu_k + \lambda_k + \|\hat{y}_{ki} - \bar{y}_i\| = o(\alpha_k). \end{aligned}$$

Thus, the limit in (4.47b) is obtained directly from (4.46a) and (4.46b).

With the aim of verifying (4.47c), let us define $\hat{u}_k := \frac{\|u\|}{\|x_k - \bar{x}\|}(x_k - \bar{x}) - u$. For such a notation, we have

$$\begin{aligned} \|u\|u_k - u &= (t_k)^{-1} \|u\|(\tilde{x}_k^1 - \bar{u}) - u \\ &= \|u_k\|(\hat{u}_k + u) - u + \|u\| \|u_k\| \frac{\tilde{x}_k^1 - x_k}{\|x_k - \bar{x}\|} \\ &= \|u_k\|\hat{u}_k + (\|u_k\| - 1)u + \|u\| \|u_k\| \frac{\tilde{x}_k^1 - x_k}{\|x_k - \bar{x}\|}. \end{aligned}$$

Observe that $x_k \xrightarrow{u} \bar{x}$; we get $\|\hat{u}_k\| \rightarrow 0$, and hence, (4.47c) follows from (4.47b) and (4.41a).

Finally, we establish (4.47d). Due to (4.46a), (4.46c) we may write

$$(t_k)^{1-\gamma_i} \|v_{ki}\| = \frac{\|\tilde{y}_k^1 - \bar{y}\|}{\|(\tilde{x}_k^1 - \bar{x}, \tilde{y}_k^1 - \bar{y})\|^{\gamma_i}} \leq \frac{\|\tilde{y}_k^1 - \bar{y}\|}{\|\tilde{x}_k^1 - \bar{x}\|^{\gamma_i}} \stackrel{k \rightarrow \infty}{\sim} \frac{\|\tilde{y}_k^1 - \bar{y}\|}{\|x_k - \bar{x}\|^{\gamma_i}}.$$

Recalling the estimation for $\|\tilde{y}_k^1 - \bar{y}\|$ as above, we find

$$\begin{aligned} \|\tilde{y}_{ki}^1 - \bar{y}_i\| &\leq \nu_k + \lambda_k + \|\hat{y}_{ki} - \bar{y}_i\| \\ &\leq \nu_k + \lambda_k + h_i(\hat{x}_k)^{\gamma_i-1} (\psi_T(x_k) + \sigma_k r_k). \end{aligned}$$

We know that $\limsup_{k \rightarrow \infty} \frac{h_i(x_k)}{\|x_k - \bar{x}\|} < +\infty, \lim_{k \rightarrow \infty} \frac{\psi_T(x_k)}{d(x_k, S)} = 0$. Therefore, (4.41a) and (4.41b)

imply $\limsup_{k \rightarrow \infty} \frac{h_i(\hat{x}_k)^{\gamma_i-1} \psi_T(x_k)}{\|x_k - \bar{x}\|^{\gamma_i}} = 0$. Since $r_k \leq h_i(x_k)^{\gamma_i} \eta_k$ and $\lambda_k = o(r_k), \nu_k = o(r_k)$, we conclude that $\|\tilde{y}_{ki}^1 - \bar{y}_i\| = o(\|x_k - \bar{x}\|^{\gamma_i})$. Thus, (4.47d) is thereby proved. \square

Fact 3 For each $i = 1, 2, \dots, m$ it holds that

$$\lim_{k \rightarrow \infty} \left\{ [h_i(x_k)]^{\gamma_i-1} \|\tilde{y}_{ki}^1\| \right\} = 1. \tag{4.48}$$

Proof of Fact 3 Invoking (viii), there exists $w_{ki}^* \in \mathbb{B}_{Y^*}$ such that

$$\sigma_k w_{ki}^* = \tilde{y}_{ki}^{1*} + \frac{\epsilon_k}{\lambda_k} \tilde{y}_{ki}^{3*} + h(x_{ki}^4)^{1-\gamma_i} \tilde{y}_{ki}^{4*}.$$

By virtue of (iv), the sequence (\tilde{y}_{ki}^{3*}) is bounded in norm, so

$$\lim_{k \rightarrow \infty} \|\tilde{y}_{ki}^{1*} + h(x_{ki}^4)^{1-\gamma_i} \tilde{y}_{ki}^{4*}\| = 0. \tag{4.49}$$

By using (4.41b), it follows that $h_i(\tilde{x}_{ki}^4) > 0$ as k is sufficiently large (because of $h_i(\tilde{x}_{ki}^4) \sim h_i(x_k)$). This implies $\tilde{y}_{ki}^{4*} \neq \bar{y}_i$ unless a finite many of indexes k . Taking into account (v) and

since the function $\|\bar{y}_i - \cdot\|$ is convex continuous on Y_j , we can say that \tilde{y}_{ki}^4 is a minimum of the function $\|\bar{y}_i - \cdot\| - \langle \tilde{y}_{ki}^{4*}, \cdot - \tilde{y}_{ki}^4 \rangle$. Thus,

$$\|\tilde{y}_{ki}^{4*}\| = 1, \quad \|\bar{y}_i - \tilde{y}_{ki}^4\| = -\langle \tilde{y}_{ki}^{4*}, \bar{y}_i - \tilde{y}_{ki}^4 \rangle. \tag{4.50}$$

The latter permits us to obtain

$$\begin{aligned} h(x_{ki}^4)^{1-\gamma_i} - \|\tilde{y}_{ki}^{1*} + h(x_{ki}^4)^{1-\gamma_i} \tilde{y}_{ki}^{4*}\| &\leq \|\tilde{y}_{ki}^{1*}\| \\ &\leq h(x_{ki}^4)^{1-\gamma_i} + \|\tilde{y}_{ki}^{1*} + h(x_{ki}^4)^{1-\gamma_i} \tilde{y}_{ki}^{4*}\| \end{aligned} \tag{4.51}$$

Combining (4.41b), (4.49) with (4.51) we obtain (4.48). □

Fact 4 Let us define with respect to each k the elements below:

$$u_k^* := \tilde{x}_k^{1*}, \tag{4.52a}$$

$$v_{ki}^* := -h_i(\bar{x} + t_k u_k)^{\gamma_i-1} \tilde{y}_{ki}^{1*}, \tag{4.52b}$$

$$\tilde{v}_{ki}^* := h_i(\bar{x} + t_k u_k)^{1-\gamma_i} v_{ki}^*. \tag{4.52c}$$

For such elements, we have

$$\lim_{k \rightarrow \infty} \|v_{ki}^*\| = 1, \tag{4.53a}$$

$$(u_k^*, -\tilde{v}_{k1}^*, \dots, -\tilde{v}_{km}^*) \in N(\text{gph } T, (\bar{x} + t_k u_k, \bar{y} + t_k v_k)), \tag{4.53b}$$

$$\lim_{k \rightarrow \infty} \|u_k^*\| = 0, \tag{4.53c}$$

$$\lim_{k \rightarrow \infty} \frac{\sum_{i=1}^m \langle v_{ki}^*, h_i(\bar{x} + t_k u_k)^{1-\gamma_i} v_{ki} \rangle}{\|(h_1(\bar{x} + t_k u_k)^{1-\gamma_1} v_{k1}, \dots, h_m(\bar{x} + t_k u_k)^{1-\gamma_m} v_{km})\|} = 1. \tag{4.53d}$$

Proof of Fact 4 The relations (4.53a) is derived from (4.52b) and (4.48), while (4.53b) is a consequence of (ii). In order to obtain (4.53c), we invoke (iii), (iv), (vi), (vii), (4.41e) and (4.48). Indeed, thanks to (vii), there is $\hat{u}_k^* \in \mathbb{B}_{X^*}$ such that

$$\sigma_k \hat{u}_k^* = \tilde{x}_k^{1*} + (\tau_k + \sigma_k) \tilde{x}_k^{2*} + \frac{\epsilon_k}{\lambda_k} \tilde{x}_k^{3*} + \sum_{i=1}^m (1 - \gamma_i) \frac{\|\bar{y}_i - \tilde{y}_{ki}^4\|}{h_i(\tilde{x}_{ki}^4)^{\gamma_i}} \tilde{u}_{ki}^*, \tag{4.54}$$

where $\tilde{u}_{ki}^* \in \partial_X h_i(\tilde{x}_{ki}^4)$ satisfies $\tilde{x}_{ki}^{4*} = (1 - \gamma_i) h_i(\tilde{x}_{ki}^4)^{-\gamma_i} \tilde{u}_{ki}^*$. According to (4), (4) and by the Lipschitz property of each function h_i , it is possible to check that

$$\limsup_{k \rightarrow \infty} \left\{ \max \left\{ \|\tilde{x}_k^{2*}\|, \|\tilde{x}_k^{3*}\|, \|\tilde{u}_{k1}^*\|, \dots, \|\tilde{u}_{km}^*\| \right\} \right\} < +\infty. \tag{4.55}$$

Denoting $\gamma^* = \max\{\gamma_1, \dots, \gamma_m\}$, (4.54) yields

$$\begin{aligned} \|\tilde{x}_k^{1*}\| &\leq \sigma_k \|\hat{u}_k^*\| + (\tau_k + \sigma_k) \|\tilde{x}_k^{2*}\| + \frac{\epsilon_k}{\lambda_k} \|\tilde{x}_k^{3*}\| \\ &+ (\gamma^* - 1) \left(\max_{i=1, \dots, m} \frac{1}{h_i(\tilde{x}_{ki}^4)} \right) \left(\max_{i=1, \dots, m} \|\tilde{u}_{ki}^*\| \right) \sum_{i=1}^m \frac{\|\bar{y}_i - \tilde{y}_{ki}^4\|}{h_i(\tilde{x}_{ki}^4)^{\gamma_i-1}}. \end{aligned} \tag{4.56}$$

Using (4.41b) and observing that $\limsup_{k \rightarrow \infty} \frac{d(x_k, S)}{h_i(x_k)} < +\infty$, we deduce

$$\limsup_{k \rightarrow \infty} \frac{d(x_k, S)}{h_i(\tilde{x}_{ki}^4)} < +\infty.$$

As a result, (4.53c) is obtained under the combination of (4.55), (4.56), (4.41e) and the assumption that $\frac{\psi_T(x_k)}{d(x_k, S)} \rightarrow 0$.

With the aim of establishing (4.53d), we define some quantities

$$\begin{cases} a_{ki} := h_i(\bar{x} + t_k u_k)^{1-\gamma_i} \langle v_{ki}^*, v_{ki} \rangle, a_k := \sum_{i=1}^m a_{ki}, \\ b_{ki} := h_i(\bar{x} + t_k u_k)^{1-\gamma_i} \|v_{ki}\|, b_k := \sum_{i=1}^m b_{ki}. \end{cases}$$

Then, we have

$$\begin{cases} -a_{ki} = (t_k)^{-1} \langle \tilde{y}_{ki}^{1*}, \tilde{y}_{ki}^1 - \bar{y}_i \rangle, \\ b_{ki} = (t_k)^{-1} h_i(h_i(\tilde{x}_k^1))^{1-\gamma_i} \|\tilde{y}_{ki}^1 - \bar{y}_i\|. \end{cases} \tag{4.57}$$

Since $\tilde{y}_{ki}^{1*} + \frac{\epsilon_k}{\lambda_k} \tilde{y}_{ki}^{3*} + h_i(\tilde{x}_k^4)^{1-\gamma_i} \tilde{y}_{ki}^{4*} = \sigma_k w_{ki}^*$ (see in the proof of Fact 3), it follows that

$$\begin{aligned} -t_k h_i(\tilde{x}_k^4)^{\gamma_i-1} a_{ki} &= \langle h_i(\tilde{x}_k^4)^{\gamma_i-1} \tilde{y}_{ki}^{1*}, \tilde{y}_{ki}^1 - \bar{y}_i \rangle = \sigma_k h_i(\tilde{x}_k^4)^{\gamma_i-1} \langle \tilde{w}_{ki}^*, \tilde{y}_{ki}^1 - \bar{y}_i \rangle \\ &\quad - \frac{\epsilon_k}{\lambda_k} h_i(\tilde{x}_k^4)^{\gamma_i-1} \langle \tilde{y}_{ki}^{3*}, \tilde{y}_{ki}^1 - \bar{y}_i \rangle - \langle \tilde{y}_{ki}^{4*}, \tilde{y}_{ki}^1 - \bar{y}_i \rangle. \end{aligned}$$

Consequently, we find

$$\begin{aligned} & t_k h_i(\bar{x} + t_k u_k)^{\gamma_i-1} \left\{ b_{ki} - \left[\frac{h_i(\tilde{x}_k^4)}{h_i(\tilde{x}_k^1)} \right]^{\gamma_i-1} a_{ki} \right\} \\ &= \|\bar{y}_i - \tilde{y}_{ki}^1\| \left\{ \sigma_k h_i(\tilde{x}_k^4)^{\gamma_i-1} \frac{\langle \tilde{w}_{ki}^*, \tilde{y}_{ki}^1 - \bar{y}_i \rangle}{\|\bar{y}_i - \tilde{y}_{ki}^1\|} \right\} \\ &\quad - \|\bar{y}_i - \tilde{y}_{ki}^1\| \left\{ \frac{\epsilon_k}{\lambda_k} h_i(\tilde{x}_k^4)^{\gamma_i-1} \frac{\langle \tilde{y}_{ki}^{3*}, \tilde{y}_{ki}^1 - \bar{y}_i \rangle}{\|\bar{y}_i - \tilde{y}_{ki}^1\|} \right\} \\ &\quad + \|\bar{y}_i - \tilde{y}_{ki}^1\| \left\{ 1 - \frac{\langle \tilde{y}_{ki}^{4*}, \tilde{y}_{ki}^1 - \bar{y}_i \rangle}{\|\bar{y}_i - \tilde{y}_{ki}^1\|} \right\}. \end{aligned} \tag{4.58}$$

As in the proof of Fact 3 (see (4.50))

$$\|\tilde{y}_{ki}^{4*}\| = 1, \|\bar{y}_i - \tilde{y}_{ki}^4\| + \langle \tilde{y}_{ki}^{4*}, \bar{y}_i - \tilde{y}_{ki}^4 \rangle = 0,$$

and the latter implies

$$\langle \tilde{y}_{ki}^{4*}, \tilde{y}_{ki}^1 - \bar{y}_i \rangle = \|\bar{y}_i - \tilde{y}_{ki}^4\| + \langle \tilde{y}_{ki}^{4*}, \tilde{y}_{ki}^1 - \tilde{y}_{ki}^4 \rangle. \tag{4.59}$$

Moreover, recall that $v_k \leq \beta_k \|\bar{y}_i - \tilde{y}_{ki}\|$ (cf. (i)). Therefore, from the fact $\max\{\|\tilde{y}_{ki}^1 - \tilde{y}_{ki}\|, \|\tilde{y}_{ki}^4 - \tilde{y}_{ki}\|\} \leq v_k$ (by virtue of (i)) one has

$$\lim_{k \rightarrow \infty} \frac{\|\tilde{y}_{ki}^1 - \tilde{y}_{ki}\|}{\|\bar{y}_i - \tilde{y}_{ki}\|} = \lim_{k \rightarrow \infty} \frac{\|\tilde{y}_{ki}^4 - \tilde{y}_{ki}\|}{\|\bar{y}_i - \tilde{y}_{ki}\|} = 0, \tag{4.60a}$$

$$\lim_{k \rightarrow \infty} \frac{\|\bar{y}_i - \tilde{y}_{ki}^1\|}{\|\bar{y}_i - \tilde{y}_{ki}\|} = \lim_{k \rightarrow \infty} \frac{\|\bar{y}_i - \tilde{y}_{ki}^4\|}{\|\bar{y}_i - \tilde{y}_{ki}\|} = 1. \tag{4.60b}$$

Combining (4.60a) with (4.60b), we obtain

$$\limsup_{k \rightarrow \infty} \left(\frac{|\langle \tilde{y}_{ki}^{4*}, \tilde{y}_{ki}^1 - \tilde{y}_{ki}^4 \rangle|}{\|\bar{y}_i - \tilde{y}_{ki}^1\|} \right) \leq \limsup_{k \rightarrow \infty} \left(\frac{\|\tilde{y}_{ki}^{4*}\| \|\tilde{y}_{ki}^1 - \tilde{y}_{ki}^4\|}{\|\bar{y}_i - \tilde{y}_{ki}^1\|} \right) = 0.$$

Taking into account the estimation

$$\left| 1 - \frac{\langle \tilde{y}_{ki}^{4*}, \tilde{y}_{ki}^1 - \bar{y}_i \rangle}{\|\bar{y}_i - \tilde{y}_{ki}^1\|} \right| \leq \left| 1 - \frac{\|\bar{y}_i - \tilde{y}_{ki}^4\|}{\|\bar{y}_i - \tilde{y}_{ki}\|} \right| + \frac{|\langle \tilde{y}_{ki}^{4*}, \tilde{y}_{ki}^1 - \tilde{y}_{ki}^4 \rangle|}{\|\bar{y}_i - \tilde{y}_{ki}^1\|},$$

we find

$$\lim_{k \rightarrow \infty} \left(1 - \frac{\langle \tilde{y}_{ki}^{4\star}, \tilde{y}_{ki}^1 - \bar{y}_i \rangle}{\|\bar{y}_i - \tilde{y}_{ki}^1\|} \right) = 0.$$

Hence, we can infer from (4.58) that

$$t_k h_i(\bar{x} + t_k u_k)^{\gamma_i - 1} \left\{ b_{ki} - \left[\frac{h_i(\tilde{x}_{ki}^4)}{h_i(\tilde{x}_k^1)} \right]^{\gamma_i - 1} a_{ki} \right\} = o(\|\bar{y}_i - \tilde{y}_{ki}^1\|), \tag{4.61}$$

which implies

$$b_{ki} - \left[\frac{h_i(\tilde{x}_{ki}^4)}{h_i(\tilde{x}_k^1)} \right]^{\gamma_i - 1} a_{ki} = o(b_{ki}). \tag{4.62}$$

Taking the sum over the index i in (4.62), we reach the conclusion

$$b_k - \sum_{i=1}^m \left\{ \left[\frac{h_i(\tilde{x}_{ki}^4)}{h_i(\tilde{x}_k^1)} \right]^{\gamma_i - 1} a_{ki} \right\} = o(b_k). \tag{4.63}$$

Observing that

$$\begin{aligned} \limsup_{k \rightarrow \infty} \frac{|a_{ki}|}{b_{ki}} &= \limsup_{k \rightarrow \infty} \left\{ h_i(\tilde{x}_k^1)^{\gamma_i - 1} \frac{|\langle \tilde{y}_{ki}^{1\star}, \bar{y}_i - \tilde{y}_{ki}^1 \rangle|}{\|\bar{y}_i - \tilde{y}_{ki}^1\|} \right\} \\ &\leq \limsup_{k \rightarrow \infty} \left\{ h_i(\tilde{x}_k^1)^{\gamma_i - 1} \|\tilde{y}_{ki}^{1\star}\| \right\} = 1, \end{aligned} \tag{4.64}$$

we deduce

$$\begin{aligned} b_k - a_k &= b_k - \sum_{i=1}^m \left\{ \left[\frac{h_i(\tilde{x}_{ki}^4)}{h_i(\tilde{x}_k^1)} \right]^{\gamma_i - 1} a_{ki} \right\} \\ &+ \sum_{i=1}^m \left\{ \left\{ \left[\frac{h_i(\tilde{x}_{ki}^4)}{h_i(\tilde{x}_k^1)} \right]^{\gamma_i - 1} - 1 \right\} a_{ki} \right\} = o(b_k). \end{aligned} \tag{4.65}$$

This shows that $\lim_{k \rightarrow \infty} \left(1 - \frac{a_k}{b_k} \right) = 0$, which is equivalent to (4.53d).

Combining Facts 1, 2, 3, 4 we obtain a full proof for Proposition 4.1. □

Definition 4.1 ((γ, h) -limiting critical set) Let T, γ and h as similar as in Definition 3.2 and let $(\bar{x}, \bar{y}) \in \text{gph } T$. For some fixed element $u \in X$, we define the limiting critical set $\text{SCR}_{\gamma, h} T(\bar{x}, \bar{y})(u)$ with respect to T, γ, h and u at the reference point (\bar{x}, \bar{y}) as follows. A pair $(v, u^\star) \in Y \times X^\star$ lies in $\text{SCR}_{\gamma, h} T(\bar{x}, \bar{y})(u)$ if it is possible to find some sequences $(t_k) \downarrow 0, (v_k, u_k^\star) \xrightarrow{Y \times X^\star} (v, u^\star)$ and $(u_k, v_k^\star) \in S_X \times S_{Y^\star}$ which fulfill simultaneously the following conditions:

- i $\lim_{k \rightarrow \infty} \|u_k - u\| = 0$;
- ii the pair (x_k, y_k) with $x_k := \bar{x} + t_k u_k, y_k := \bar{y}_i + (t_k)^{\gamma_i} v_{ki}$ is in $\text{gph } T$ but $\bar{y} \notin T(x_k)$;
- iii $(u_k^\star, -h_1(x_k)^{1-\gamma_1} v_{k1}^\star, \dots, -h_m(x_k)^{1-\gamma_m} v_{km}^\star) \in N(\text{gph } T, (x_k, y_k))$ and one has

$$\lim_{k \rightarrow \infty} \left\{ \frac{\sum_{i=1}^m \langle v_{ki}^\star, h_i(x_k)^{1-\gamma_i} (y_{ki} - \bar{y}_i) \rangle}{\| (h_1(x_k)^{1-\gamma_1} (y_{k1} - \bar{y}_1), \dots, h_m(x_k)^{1-\gamma_m} (y_{km} - \bar{y}_m)) \|} \right\} = 1.$$

Using this new notion, we are now ready to present the infinitesimal characterization for the property of (γ, h) -metric pseudo subregularity. The next theorem is in this sense.

Theorem 4.1 *Let T, γ, h and (\bar{x}, \bar{y}) as in Definition 4.1. Suppose that each function h_i is locally Lipschitz around \bar{x} and that*

$$\limsup_{x \rightarrow \bar{x}, x \notin S} \frac{d(x, S)}{h_i(x)} < +\infty, \quad i = 1, \dots, m. \tag{4.66}$$

If $(0, 0) \notin \text{SCR}_{\gamma, h} T(\bar{x}, \bar{y})(u)$, then the set-valued mapping T is (γ, h) -metrically pseudo subregular in the direction $u \in X$ at (\bar{x}, \bar{y}) .

Proof The proof is almost based on Proposition 4.1. Assume that the norm in $Y^* = Y_1^* \times \dots \times Y_m^*$ coincides with the maximum norm

$$\|y^*\|_{Y^*} = \|(y_1^*, \dots, y_m^*)\|_{Y^*} = \max \{ \|y_1^*\|_{Y_1^*}, \dots, \|y_m^*\|_{Y_m^*} \}.$$

Suppose on the contrary that T is not (γ, h) -metrically pseudo subregular in the direction u at (\bar{x}, \bar{y}) . Let $t_k > 0, (u_k, v_k) \in S_{X \times Y}$, and $(u_k^*, v_k^*) \in X^* \times Y^*$ be the sequences in the conclusion of Proposition 4.1. Let us define

$$x_k := \bar{x} + t_k u_k, \quad y_k := \bar{y} + t_k v_k; \tag{4.67}$$

then it is clear that

$$(u_k^*, -h_1(x_k)^{1-\gamma_1} v_{k1}^*, \dots, -h_m(x_k)^{1-\gamma_m} v_{km}^*) \in N(\text{gph } T, (x_k, y_k)), \tag{4.68a}$$

$$\lim_{k \rightarrow \infty} \|u_k^*\| = 0, \quad \lim_{k \rightarrow \infty} \|v_{ki}^*\| = 1, \tag{4.68b}$$

$$\lim_{k \rightarrow \infty} \|u_k\| = 1, \quad \lim_{k \rightarrow \infty} \| \|u\| u_k - u \| = 0, \tag{4.68c}$$

$$\lim_{k \rightarrow \infty} \{ (t_k)^{1-\gamma_1} \|v_{k1}\| \} = \dots = \lim_{k \rightarrow \infty} \{ (t_k)^{1-\gamma_m} \|v_{km}\| \} = 0, \tag{4.68d}$$

$$\lim_{k \rightarrow \infty} \left\{ \frac{\sum_{i=1}^m (v_{ki}^*, h_i(x_k)^{1-\gamma_i} v_{ki})}{\| (h_1(x_k)^{1-\gamma_1} v_{k1}, \dots, h_m(x_k)^{1-\gamma_m} v_{km}) \|} \right\} = 1. \tag{4.68e}$$

Further, as proved in Fact 2, we also have $\bar{y} \notin T(\bar{x} + t_k u_k) = T(x_k)$. Setting

$$\hat{t}_k := \|u_k\| t_k, \tag{4.69a}$$

$$\hat{u}_k := \|u_k\|^{-1} u_k, \quad \hat{v}_{ki} := (t_k)^{1-\gamma_i} \|u_k\|^{-\gamma_i} v_{ki}, \tag{4.69b}$$

$$\hat{u}_k^* := \|v_k^*\|^{-1} u_k^*, \quad \hat{v}_k^* := \|v_k^*\|^{-1} v_k^*, \tag{4.69c}$$

we obtain

$$x_k = \bar{x} + \hat{t}_k \hat{u}_k, \quad y_{ki} = \bar{y}_i + (\hat{t}_k)^{\gamma_i} \hat{v}_{ki}, \tag{4.70a}$$

$$(\hat{u}_k^*, -h_1(x_k)^{1-\gamma_1} \hat{v}_{k1}^*, \dots, -h_m(x_k)^{1-\gamma_m} \hat{v}_{km}^*) \in N(\text{gph } T, (x_k, y_k)), \tag{4.70b}$$

$$\| \hat{u}_k \| = 1, \quad \| \hat{v}_k^* \| = 1, \tag{4.70c}$$

$$\lim_{k \rightarrow \infty} \| \hat{v}_{ki} \| = 0, \quad \lim_{k \rightarrow \infty} \| \hat{u}_k^* \| = 0. \tag{4.70d}$$

On the other hand, combining (4.69c) with (4.68b) and (4.68e), we find

$$\begin{aligned} & \lim_{k \rightarrow \infty} \left\{ \frac{\sum_{i=1}^m \langle \hat{v}_{ki}^*, h_i(x_k)^{1-\gamma_i} (y_{ki} - \bar{y}_i) \rangle}{\| (h_1(x_k)^{1-\gamma_1} (y_{k1} - \bar{y}_1), \dots, h_m(x_k)^{1-\gamma_m} (y_{km} - \bar{y}_m)) \|} \right\} \\ &= \lim_{k \rightarrow \infty} \left\{ \|v_k^*\|^{-1} \frac{\sum_{i=1}^m \langle v_{ki}^*, h_i(x_k)^{1-\gamma_i} v_{ki} \rangle}{\| (h_1(x_k)^{1-\gamma_1} v_{k1}, \dots, h_m(x_k)^{1-\gamma_m} v_{km}) \|} \right\} \\ &= \lim_{k \rightarrow \infty} \left\{ \left(\max_{i=1, \dots, m} \|v_{ki}^*\| \right)^{-1} \frac{\sum_{i=1}^m \langle v_{ki}^*, h_i(x_k)^{1-\gamma_i} v_{ki} \rangle}{\| (h_1(x_k)^{1-\gamma_1} v_{k1}, \dots, h_m(x_k)^{1-\gamma_m} v_{km}) \|} \right\} \\ &= 1. \end{aligned}$$

In addition, taking into account the representation

$$\|u\| \hat{u}_k - u = \|u_k\|^{-1} (\|u\| u_k - u) + (\|u_k\|^{-1} - 1) u,$$

(4.68c) implies $\lim_{k \rightarrow \infty} \| \|u\| \hat{u}_k - u \| = 0$. In summary, it follows that $(0, 0)$ belongs to the set $\text{SCR}_{\gamma, h} T(\bar{x}, \bar{y})(u)$, which contradicts the assumption of Theorem 4.1. \square

Remark 4.1 By letting $h_i(x) = \|x - \bar{x}\|$, Theorem 4.1 subsumes somewhat studied in [11, Theorem 1]. Taking $m = 1$, $h_1 = d(x, T^{-1}(\bar{y}))$, then Theorem 4.1 recovers the results presented in the works [25, 26].

Note that certain applications concern set-valued mapping having a convex (and closed) graph. For such situations, the counterpart of Theorem 4.1 might be also fulfilled. The next result is in this sense.

Proposition 4.2 (Set-valued mapping with convex graph) *Suppose that the set-valued mapping $T : X \rightrightarrows Y$ has closed convex graph. Fix some given $(\bar{x}, \bar{y}) \in \text{gph } T$ and $u \in X$. Let γ and h be the same as in Theorem 4.1. If T is (γ, h) -metrically pseudo subregular in the direction u at (\bar{x}, \bar{y}) , then one has $(0, 0) \notin \text{SCR}_{\gamma, h} T(\bar{x}, \bar{y})(u)$.*

Proof Replacing T by $\tilde{T}(\cdot) := T(\cdot + \bar{x}) - \bar{y}$ if necessary, we may assume $\bar{x} = 0$ and $\bar{y} = 0$. Let T satisfy all assumptions of Proposition 4.2 but the pair $(0, 0)$ be in the set $\text{SCR}_{\gamma, h} T(0, 0)(u)$. The definition of $\text{SCR}_{\gamma, h} T(0, 0)(u)$ shows that, there exist a real sequence $s_k \downarrow 0$ together with some sequences $(u_k, v_k^*) \in S_X \times S_{Y^*}$ and $(v_k, u_k^*) \xrightarrow{Y \times X^*} (0, 0)$ which fulfill the conditions below:

- $\lim_{k \rightarrow \infty} \| \|u\| u_k - u \| = 0$;
- we have $y_k \in T(x_k)$ but $0 = \bar{y} \notin T(x_k)$, in which $x_k = s_k u_k \in X$ and $y_{ki} = (s_k)^{\gamma_i} v_{ki}$;
- $(u_k^*, -h_1(x_k)^{1-\gamma_1} v_{k1}^*, \dots, -h_m(x_k)^{1-\gamma_m} v_{km}^*)$ is an element of $N(\text{gph } T, (x_k, y_k))$;
- $\lim_{k \rightarrow \infty} \left\{ \frac{\sum_{i=1}^m \langle v_{ki}^*, h_i(x_k)^{1-\gamma_i} y_{ki} \rangle}{\| (h_1(x_k)^{1-\gamma_1} y_{k1}, \dots, h_m(x_k)^{1-\gamma_m} y_{km}) \|} \right\} = 1$.

Let τ, δ and r be positive real parameters such that

$$d(x, T^{-1}(0)) \leq \tau \sum_{i: h_i(x) > 0} h_i(x)^{1-\gamma_i} d(0, T_i(x)) \tag{4.71}$$

whenever $x \in \text{cone } \mathbb{B}(u, \delta)$ with $0 < \|x\| < r$. Since $\lim_{k \rightarrow \infty} \| \|u\|u_k - u \| = 0$, x_k will be in $\text{cone } \mathbb{B}(u, \delta)$ after skipping a few first indexes k . Hence, it is possible to apply (4.71) at $x = x_k$

$$\begin{aligned} d(x_k, T^{-1}(0)) &\leq \tau \sum_{i=1}^m h_i(x_k)^{1-\gamma_i} d(0, T_i(x_k)) \\ &\leq \tau \sum_{i=1}^m h_i(x_k)^{1-\gamma_i} \|y_{ki}\|. \end{aligned} \tag{4.72}$$

Recall that $x_k \notin T^{-1}(0)$. By virtue of (4.72), for some $0 < \sigma_k < 1$ there exists $z_k \in T^{-1}(0)$ which fulfills the inequalities

$$0 < \|x_k - z_k\| \leq \tau(1 + \sigma_k) \sum_{i=1}^m h_i(x_k)^{1-\gamma_i} \|y_{ki}\|. \tag{4.73}$$

Because the set $\text{gph } T$ is convex, we may derive from the choice of u_k^* and v_k^* that

$$\langle u_k^*, z_k - x_k \rangle + \sum_{i=1}^m \langle -h_i(x_k)^{1-\gamma_i} v_{ki}^*, -y_{ki} \rangle \leq 0. \tag{4.74}$$

As a result, we obtain

$$\begin{aligned} &\frac{\sum_{i=1}^m h_i(x_k)^{1-\gamma_i} \|y_{ki}\|}{\|z_k - x_k\|} \frac{\sum_{i=1}^m \langle v_{ki}^*, h_i(x_k)^{1-\gamma_i} y_{ki} \rangle}{\|(h_1(x_k)^{1-\gamma_1} y_{k1}, \dots, h_m(x_k)^{1-\gamma_m} y_{km})\|} \\ &\leq \frac{\langle u_k^*, x_k - z_k \rangle}{\|z_k - x_k\|}. \end{aligned} \tag{4.75}$$

Combining (4.73) with (4.75), we deduce

$$\begin{aligned} &\frac{1}{\tau(1 + \sigma_k)} \frac{\sum_{i=1}^m \langle v_{ki}^*, h_i(x_k)^{1-\gamma_i} y_{ki} \rangle}{\|(h_1(x_k)^{1-\gamma_1} y_{k1}, \dots, h_m(x_k)^{1-\gamma_m} y_{km})\|} \\ &\leq \frac{\langle u_k^*, x_k - z_k \rangle}{\|z_k - x_k\|} \leq \|u_k^*\|. \end{aligned} \tag{4.76}$$

Passing to the limit w.r.t k in both sides of (4.76), we obtain the following estimate

$$\liminf_{k \rightarrow \infty} \left\{ \frac{1}{\tau(1 + \sigma_k)} \frac{\sum_{i=1}^m \langle v_{ki}^*, h_i(x_k)^{1-\gamma_i} y_{ki} \rangle}{\|(h_1(x_k)^{1-\gamma_1} y_{k1}, \dots, h_m(x_k)^{1-\gamma_m} y_{km})\|} \right\} \leq 0. \tag{4.77}$$

Since the left-hand side of (4.77) takes a positive value, we reach a contradiction. This completes our proof. □

As arising from many applications, for instance, in generalized equations, we restrict our consideration to the case $m = 1$ and $T = f + F$ where $f : X \rightarrow Y = Y_1$ is C^1 and $F : X \rightrightarrows Y$ has a closed and convex graph. Under some robust condition imposed on the given data f and the abstract subdifferential ∂ , the next proposition has the advantage of offering a necessary and sufficient condition for (γ, h) -metric pseudo subregularity.

Proposition 4.3 *Let $\gamma = \gamma_1 \in [1, 2)$ and $h = h_1 : X \rightarrow \mathbb{R}$ be the same as in statement of Theorem 4.1. Suppose in addition that the Jacobian map ∇f is Lipschitz around \bar{x} while the coderivative associated with ∂ obeys the sum rule*

$$D^*(f + F)(x, f(x) + z) = \nabla f(x)^* + D^*F(x, z) \tag{4.78}$$

for every $(x, z) \in \text{gph } F$ near $(\bar{x}, \bar{z}) \in \text{gph } F$. Then, $T = f + F$ is (γ, h) -metrically pseudo subregular in the direction u at $(\bar{x}, \bar{y}) \in \text{gph } T$ if and only if

$$(0, 0) \notin \text{SCR}_{\gamma, h} T(\bar{x}, \bar{y})(u).$$

Proof It is sufficient to prove only the necessary part. Without any loss of generality, we may assume $\bar{x} = 0, f(\bar{x}) = \bar{y} = 0$. Suppose T is (γ, h) -metrically pseudo subregular in the direction u at $(0, 0)$. Let κ, δ, ϵ be positive real numbers under which the following estimation

$$d(x, T^{-1}(0)) \leq \tau h(x)^{1-\gamma} d(0, T(x)) \tag{4.79}$$

holds whenever $x \neq 0 \in \text{cone } \mathbb{B}(u, \delta) \cap \epsilon \mathbb{B}$. If $(0, 0) \in \text{SCR}_{\gamma, h} T(0, 0)(u)$, we may select for each k , some elements $s_k > 0, u_k \in \mathbf{S}_X, v_k \in Y, u_k^* \in X^*$ and $v_k^* \in \mathbf{S}_{Y^*}$ such that

- (a). the pair (x_k, y_k) is in $\text{gph } T$ with $x_k = s_k u_k$ and $y_k = f(x_k) + z_k = (s_k)^\gamma v_k$;
- (b). $\lim_{k \rightarrow \infty} \| \|u\| u_k - u \| = \lim_{k \rightarrow \infty} \|v_k\| = \lim_{k \rightarrow \infty} \|u_k^*\| = 0$;
- (c). $(u_k^*, -h(x_k)^{1-\gamma} v_k^*) \in N(\text{gph } T, (x_k, y_k))$;
- (d). $\lim_{k \rightarrow \infty} \left\{ \frac{\langle v_k^*, h(x_k)^{1-\gamma} y_k \rangle}{\|h(x_k)^{1-\gamma} y_k\|} \right\} = 1$.

In view of (c), one has

$$h(x_k)^{\gamma-1} u_k^* \in D^*T(x_k, y_k)(v_k^*).$$

Applying the sum rule formula to $T = f + F$ at (x_k, z_k) , the latter inclusion yields

$$h(x_k)^{\gamma-1} u_k^* - \nabla f(x_k)^* v_k^* \in D^*F(x_k, z_k)(v_k^*). \tag{4.80}$$

According to the definition of a coderivative, and noticing that $\text{gph } F$ is closed and convex, we deduce

$$\langle h(x_k)^{\gamma-1} u_k^* - \nabla f(x_k)^* v_k^*, x - x_k \rangle + \langle -v_k^*, z - z_k \rangle \leq 0 \tag{4.81}$$

when $(x, z) \in \text{gph } F$. Taking (b) into account, we may apply (4.79) with $x = x_k$ and get

$$d(x_k, T^{-1}(0)) \leq \tau h(x_k)^{1-\gamma} d(0, T(x_k)) \leq \tau h(x_k)^{1-\gamma} \|y_k\|. \tag{4.82}$$

Recall that $S = T^{-1}(0)$. Let $\sigma_k > 0$ be such that $\limsup_{k \rightarrow \infty} \frac{\sigma_k}{d(x_k, S)} = 0$ and let $\hat{x}_k \in S \setminus \{x_k\}$ satisfy

$$\|x_k - \hat{x}_k\| \leq d(x_k, S) + \min \left\{ \sigma_k, \frac{1}{k^2} h(x_k)^{1-\gamma} \|y_k\| \right\}. \tag{4.83}$$

Substituting $x = \hat{x}_k$ and $z = -f(\hat{x}_k)$ in (4.81), we get

$$\langle h(x_k)^{\gamma-1} u_k^* - \nabla f(x_k)^* v_k^*, \hat{x}_k - x_k \rangle + \langle -v_k^*, -f(\hat{x}_k) - z_k \rangle \leq 0. \tag{4.84}$$

Consequently, after replacing z_k by $y_k - f(x_k)$, (4.84) reads

$$\begin{aligned} & \langle h(x_k)^{\gamma-1} u_k^*, \hat{x}_k - x_k \rangle + \langle v_k^*, y_k \rangle \\ & \leq \langle v_k^*, -f(\hat{x}_k) + f(x_k) + \nabla f(x_k)(\hat{x}_k - x_k) \rangle. \end{aligned} \tag{4.85}$$

Combining inequalities (4.82), (4.83) and (4.85), yields

$$\begin{aligned}
 & h(x_k)^{\gamma-1} \frac{\langle u_k^*, \hat{x}_k - x_k \rangle}{\|\hat{x}_k - x_k\|} + h(x_k)^{\gamma-1} \left(\tau + \frac{1}{k^2} \right)^{-1} \frac{\langle v_k^*, y_k \rangle}{\|y_k\|} \\
 & \leq \frac{1}{\|\hat{x}_k - x_k\|} \langle v_k^*, -f(\hat{x}_k) + f(x_k) + \nabla f(x_k)(\hat{x}_k - x_k) \rangle. \tag{4.86}
 \end{aligned}$$

Setting $\hat{u}_k := \hat{x}_k - x_k$ and applying the Taylor expansion to f gives:

$$f(\hat{x}_k) = f(x_k) + \nabla f(x_k)(\hat{u}_k) + \int_0^1 [\nabla f(x_k + t\hat{u}_k) - \nabla f(x_k)](\hat{u}_k) dt.$$

From the Lipschitz continuity of the Jacobian ∇f , we have

$$\limsup_{k \rightarrow \infty} \frac{\sup_{t \in [0,1]} \{ \|\nabla f(x_k + t\hat{u}_k) - \nabla f(x_k)\| \}}{\|\hat{u}_k\|} < +\infty,$$

and therefore, we obtain

$$\limsup_{k \rightarrow \infty} \frac{\|f(\tilde{x}_k) - f(x_k) - \nabla f(x_k)(\hat{u}_k)\|}{\|\hat{u}_k\|^2} < +\infty. \tag{4.87}$$

This leads to the estimation

$$\begin{aligned}
 & \frac{\langle u_k^*, \hat{x}_k - x_k \rangle}{\|\hat{x}_k - x_k\|} + \left(\tau + \frac{1}{k^2} \right)^{-1} \frac{\langle v_k^*, y_k \rangle}{\|y_k\|} \\
 & \leq \theta_k h(x_k)^{1-\gamma} \|\hat{x}_k - x_k\|, \tag{4.88}
 \end{aligned}$$

in which the real sequence (θ_k) is bounded. Due to the choice of \hat{x}_k , it is possible to write

$$d(x_k, S) \leq \|\hat{x}_k - x_k\| \leq d(x_k, S) + \sigma_k,$$

which permits to deduce that

$$\limsup_{k \rightarrow \infty} \frac{\|\hat{x}_k - x_k\|}{h(x_k)} = \limsup_{k \rightarrow \infty} \left(\frac{\|\hat{x}_k - x_k\|}{d(x_k, S)} \cdot \frac{d(x_k, S)}{h(x_k)} \right) < +\infty. \tag{4.89}$$

Taking into account that $\|u_k^*\| \rightarrow 0$, and passing to the limit as $k \rightarrow \infty$ in (4.88), we obtain that

$$\liminf_{k \rightarrow \infty} \left\{ \left(\tau + \frac{1}{k^2} \right)^{-1} \frac{\langle v_k^*, y_k \rangle}{\|y_k\|} \right\} \leq 0.$$

However, the latter relation obviously contradicts (d). Thus, the proof is complete. □

Remark 4.2 The conclusion of the preceding proposition is still valid for $\gamma = 2$ if we choose $h(x) = \|x - \bar{x}\|$. Indeed, following the proof above, we have

$$\|y_k\| = (s_k)^\gamma \|v_k\| = \|x_k\|^\gamma \|v_k\| = h(x_k)^\gamma \|v_k\|. \tag{4.90}$$

Reminding (4.82), this yields $d(x_k, S) = o(h(x_k))$, and hence, from the choice of \hat{x}_k , this implies

$$\|\hat{x}_k - x_k\| = o(h(x_k)). \tag{4.91}$$

Combining this relation with (4.88), we reach a contradiction as in the previous proof.

Acknowledgments We gratefully acknowledge the referees for their constructive comments and suggestions.


References

1. Arutyunov, A.V., Avakov, E.R., Izmailov, A.F.: Directional regularity and metric regularity. *SIAM J. Optim.* **18**(3), 810–833 (2007). <https://doi.org/10.1137/060651616>
2. Azé, D., Corvellec, J.N.: On the sensitivity analysis of Hoffman constants for systems of linear inequalities. *SIAM J. Optim.* **12**(4), 913–927 (2002)
3. Bonnans, J.F., Shapiro, A.: *Perturbation Analysis of Optimization Problems*. Springer, New York (2000)
4. Borwein, J.M.: Stability and regular points of inequality systems. *J. Optim. Theory Appl.* **48**(1), 9–52 (1986)
5. Cibulka, R., Fabian, M., Kruger, A.: On semiregularity of mappings. *J. Math. Anal. Appl.* **473**, 811–836 (2019)
6. De Giorgi, E., Marino, A., Tosques, M.: Evolution problems in metric spaces and steepest descent curves. *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur.* (8) **68**(3), 180–187 (1980). In Italian. English translation: Ennio De Giorgi, *Selected Papers*, Springer, Berlin 2006, 527–533
7. Dontchev, A.L., Rockafellar, R.T.: *Implicit Functions and Solution Mappings. A View from Variational Analysis*, 2 Edn. Springer Series in Operations Research and Financial Engineering. Springer, New York (2014)
8. Facchinei, F., Fischer, A., Herrich, M.: An Ip-newton method: nonsmooth equations, kkt systems, and nonisolated solutions. *Math. Program.* **146**(1–2, Ser. A), 1–36 (2014)
9. Facchinei, F., P.J.S.: Finite-dimensional variational inequalities and complementarity problems. In: Vol. II. Springer Series in Operations Research. Springer, New York (2003)
10. Gfrerer, H.: On directional metric subregularity and second-order optimality conditions for a class of nonsmooth mathematical programs. *SIAM J. Optim.* **23**(1), 632–665 (2013). <https://doi.org/10.1137/120891216>
11. Gfrerer, H.: On metric pseudo-(sub)regularity of multifunctions and optimality conditions for degenerated mathematical programs. *Set-Valued Var. Anal.* **22**(1), 79–115 (2014). <https://doi.org/10.1007/s11228-013-0266-z>
12. Huynh, V.N., Nguyen, H.T., Théra, M.: Directional Hölder metric regularity. *J. Optim. Theory Appl.* **171**(3), 785–819 (2016)
13. Huynh, V.N., Théra, M.: Error bounds in metric spaces and application to the perturbation stability of metric regularity. *SIAM J. Optim.* **19**(1), 1–20 (2008). <https://doi.org/10.1137/060675721>
14. Huynh, V.N., Théra, M.: Directional metric regularity of multifunctions. *Math. Oper. Res.* **40**(4), 969–991 (2015). <https://doi.org/10.1287/moor.2014.0705>
15. Ioffe, A.D.: On regularity concepts in variational analysis. *J. Fixed Point Theory Appl.* **8**(2), 339–363 (2010). <https://doi.org/10.1007/s11784-010-0021-0>
16. Ioffe, A.D.: Convexity and variational analysis. In: Bailey, D.H., Bauschke, H.H., Borwein, P., Garvan, F., Théra, F., D. V.J., Wolkowicz, H. (eds.) *Computational and Analytical Mathematics*, Springer Proc. Math. Stat., vol. 50, pp. 411–444. Springer, New York (2013). <https://doi.org/10.1007/978-1-4614-7621-4-19>
17. Ioffe, A.D.: Metric regularity – a survey. Part I. Theory. *J. Aust. Math. Soc.* **101**(2), 188–243 (2016). <https://doi.org/10.1017/S1446788715000701>
18. Ioffe, A.D.: Metric regularity – a survey. Part II. Applications. *J. Aust. Math. Soc.* **101**(3), 376–417 (2016). <https://doi.org/10.1017/S1446788715000695>
19. Ioffe, A.D.: *Variational analysis of regular mappings. Theory and applications*. Springer monographs in mathematics springer (2017)
20. Izmailov, A.F., Solodov, M.V.: Newton-type methods for optimization and variational problems. In: Springer Series in Operations Research and Financial Engineering, Springer
21. Kruger, A.Y.: Error bounds and Hölder metric subregularity. *Set-Valued Var. Anal.* **23**(4), 705–736 (2015)
22. Kruger, A.Y.: Error bounds and metric subregularity. *Optimization* **64**(1), 49–79 (2015)
23. Kruger, Alexander.Y.: Nonlinear metric subregularity. *J. Optim. Theory Appl.* **171**(3), 820–855 (2016). <https://doi.org/10.1007/s10957-015-0807-8>
24. Mordukhovich, B.S.: *Variational Analysis and Generalized Differentiation. I: Basic Theory* Grundlehren Der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 330. Springer, Berlin (2006)
25. Ngai, H.V., Tinh, P.N.: Metric subregularity of multifunctions: First and second order infinitesimal characterizations. *Math. Oper. Res.* **40**(3), 703–724 (2015). <https://doi.org/10.1287/moor.2014.0691>
26. Ngai, H.V., Tron, N.H., Tinh, P.N.: Directional hölder metric subregularity and application to tangent cones. *J. Convex Anal.* **24**(2), 417–457 (2017)

27. Penot, J.P.: Metric regularity, openness and Lipschitzian behavior of multifunctions. *Nonlinear Anal.* **13**(6), 629–643 (1989). [https://doi.org/10.1016/0362-546X\(89\)90083-7](https://doi.org/10.1016/0362-546X(89)90083-7)
28. Penot, J.P.: *Calculus without Derivatives Graduate Texts in Mathematics*, vol. 266. Springer, New York (2013)
29. Rockafellar, R.T., Wets, R.J.B.: *Variational Analysis*. Springer, Berlin (1998)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Affiliations

Huynh Van Ngai¹ · Nguyen Huu Tron¹ · Nguyen Van Vu¹ · Michel Théra^{2,3} 

Huynh Van Ngai
ngaivn@yahoo.com

Nguyen Huu Tron
nguyenhutrong@qnu.edu.vn

Nguyen Van Vu
nguyenvanvu@qnu.edu.vn

¹ Department of Mathematics, Quy Nhon University, 170 An Duong Vuong, Quy Nhon, Binh Dinh Vietnam

² XLIM UMR-CNRS 7252, Université de Limoges, Limoges, France

³ Centre for Informatics and Applied Optimisation (CIAO), Federation University Australia, Mount Helen, Australia