

A geo-logical solution to the lottery paradox, with applications to conditional logic

Hanti Lin · Kevin T. Kelly

Received: 8 August 2011 / Accepted: 8 August 2011 / Published online: 6 September 2011
© Springer Science+Business Media B.V. 2011

Abstract We defend a set of acceptance rules that avoids the lottery paradox, that is closed under classical entailment, and that accepts uncertain propositions without ad hoc restrictions. We show that the rules we recommend provide a semantics that validates exactly Adams' conditional logic and are exactly the rules that preserve a natural, logical structure over probabilistic credal states that we call *probalogic*. To motivate probalogic, we first expand classical logic to *geo-logic*, which fills the entire unit cube, and then we project the upper surfaces of the geo-logical cube onto the plane of probabilistic credal states by means of standard, linear perspective, which may be interpreted as an extension of the classical principle of indifference. Finally, we apply the geometrical/logical methods developed in the paper to prove a series of trivialization theorems against question-invariance as a constraint on acceptance rules and against rational monotonicity as an axiom of conditional logic in situations of uncertainty.

Keywords Lottery paradox · Uncertain acceptance · Ramsey test · Conditional logic · Belief revision · Framing effects

1 The lottery paradox

If Bayesians are right, one's credal state should be a probability measure p over propositions, where probabilities represent degrees of belief. It seems that one also *accepts* propositions in light of p . Acceptance of proposition A is sometimes portrayed as a

H. Lin · K. T. Kelly (✉)
Department of Philosophy, Carnegie Mellon University, Pittsburgh, PA 15213, USA
e-mail: kk3n@andrew.cmu.edu

H. Lin
e-mail: hantil@andrew.cmu.edu

momentous inference making A certain, in the sense that one would bet one's life against nothing that A is true (e.g., Levi 1967). But that extreme standard would eliminate almost all ordinary examples of accepted propositions. We therefore entertain a more modest view of acceptance, according to which the set of propositions accepted in light of p should, in some sense, aptly capture some characteristics of p to others or, in everyday cognition, to ourselves. That view is non-inferential in the sense that p is not conditioned on the propositions accepted, but it is inferential in another sense—the accepted propositions may serve as premises in arguments whose conclusions are also accepted in the same, weak sense.

It seems that high probability short of full certainty suffices for acceptance, a view now referred to as the *Lockean thesis*. But the Lockean rule licenses acceptance of inconsistent sets of propositions, however high the threshold $r < 1$ is set. For there exists a fair lottery with more than $1/(1-r)$ tickets. It is accepted that some ticket wins, since that proposition carries probability 1. But for each ticket, it is also accepted that the ticket loses, since that proposition has probability greater than r . So an inconsistent set of propositions is accepted. That is Kyburg's (1961) *lottery paradox*.

To elude the paradox, one must abandon either the full Lockean thesis or classical consistency. Kyburg pursued the second course by rejecting the classical inference rule that A, B jointly imply $A \wedge B$, so that the collection of propositions of form "ticket i does not win" does not entail "no ticket wins". Most responses side with classical logic and constrain the Lockean thesis in some manner to avoid contradictions. For example, Jeffrey (1970) recommended that the entire practice of acceptance be abandoned in favor of reporting probabilities. Levi (1967) rejected the idea that acceptance can be based on probability alone, since utilities should also be consulted. Or one may impose as a necessary condition that accepted propositions be certain (van Fraassen 1995; Arló-Costa and Parikh 2005). Or one may restrict the Lockean thesis to cases in which no logical contradiction happens to result (Pollock 1995; Ryan 1996; Douven 2002).

Our approach is different. Instead of restricting the Lockean thesis, we *revise* it. In particular, we defend an *unrestricted* rule of acceptance that is *contradiction-free* and yet capable of accepting *uncertain* propositions—even propositions of fairly low probability. Like the Lockean rule, the proposed rule has a parameter that controls its strictness. When the parameter is tuned toward 1, the proposed rule is almost indistinguishable from the classical logical closure of the Lockean rule; but as the parameter drops toward 0, the proposed rule's geometry shifts steadily away from that of the Lockean rule so as to avert the lottery paradox.

The rule we recommend was invented by Levi (1996, p. 286), who saw no justification for it except as a loose approximation to an alternative rule he took to be justified by decision-theoretic means (1967, 1969).¹ We provide two justifications of the rule. The first is that it *preserves logical structure*, in the sense that it accepts

¹ Levi writes: "I do not know how to derive it from a view of the cognitive aims of inquiry [i.e. seeking more information and avoiding error] that seems attractive" (1996, p. 286). We rediscovered the rule as a consequence of our work on Ockham's razor. The problem was to extend the Ockham efficiency theorem (Kelly 2008) from methods that choose theories to methods that update probabilistic degrees of belief on theories. That required a concept of retraction of credal states, expounded in Kelly (2011). We thank Teddy Seidenfeld for bringing the prior publication of the rule to our attention.

stronger propositions in stronger probabilistic credal states. The crux is to order probabilistic credal states according to relative logical strength, as Boolean algebra does for propositions. We do so in two steps. First, we start with a sigma algebra of propositions (closed under negation and countable disjunction) and then *extend* that sigma algebra to cover the entire unit cube by introducing a new connective \neg_d interpreted as negation to *degree* d , so that $\neg_0\phi$ is equivalent to ϕ and $\neg_1\phi$ is equivalent to $\neg\phi$. The resulting logical structure is called *geologic* (Sect. 4). Next, we *view* geologic through the *picture plane* of possible credal states to obtain a logical structure over credal states that we call *probalogic* (Sects. 5 and 6). Then it is natural to require that every acceptance rule preserves probalogical structure when it maps probabilistic credal states to standard, Boolean propositions.

The requirement that acceptance rules preserve probalogical structure has an appealing consequence for the theory of acceptance: we show that the rules we recommend are exactly the rules that preserve probalogical structure (Sect. 7). In contrast, no plausible logical structure on probability measures is preserved by the Lockean rule or its variants (Sect. 8).

Our second justification of the proposed acceptance rules concerns the logic of conditionals and defeasible reasoning. Frank P. Ramsey proposed an influential, epistemic condition for acceptance of conditional statements, now commonly referred to as the *Ramsey test*:

If two people are arguing ‘If A , then B ?’ and are both in doubt as to A , they are adding A hypothetically to their stock of knowledge and arguing on that basis about B ; so that in a sense ‘If A , B ’ and ‘If A , $\neg B$ ’ are contradictories. We can say that they are fixing their degrees of belief in B given A . (Ramsey 1929, footnote 1)²

Suppose that an agent is in a probabilistic credal state p and adopts an acceptance rule. We propose the following interpretation of the Ramsey test: the agent accepts the (flat) conditional ‘if A then B ’ when, by the acceptance rule she adopts, she would accept B in the credal state $p(\cdot|A)$ that results from p by conditioning on A . Thus, conditional acceptance is reduced to Bayesian conditioning and acceptance of non-conditional propositions. This natural semantics allows one to characterize the axioms of conditional logic in terms of their geometrical constraints on acceptance rules, in much the same way that axioms of modal logic are standardly characterized in terms of constraints on accessibility relations among worlds. Accordingly, for each of the axioms in Adams’ (1975) conditional logic, we solve for its geometrical constraint on acceptance rules (Sect. 9). These constraints are shown to be satisfied by the rules that preserve probalogical structure, so the probalogic-preserving rules validate Adams’ logic with respect to the Ramsey test (Sect. 10). Conversely, Adams’ logic is shown to be *complete* with respect to the Ramsey test when acceptance follows probalogic-preserving rules (Sect. 12). The result is a new probabilistic semantics: it defines validity simply as preservation of acceptance, which improves upon Adams’ (1975) ϵ – δ semantics; and it allows for accepting propositions of low probabilities,

² We take the liberty of substituting A , B for p , q in Ramsey’s text.

which improves upon Pearl's (1989) infinitesimal semantics. Thus, the recommended acceptance rules are vindicated both by probalogic and by conditional logic.

One might hope for validating a stronger logic of flat conditionals than Adams', e.g. system **R** (Lehmann and Magidor 1992) or, stronger still, the AGM axioms for belief revision (Harper 1975; Alchourrón et al. 1985). We close the door on that hope with a new trivialization theorem (Sect. 11). In light of that result, we propose that Adams' conditional logic reflects Bayesian ideals better than AGM belief revision does.

Finally, the acceptance rules we recommend are sensitive to framing effects determined by an underlying question. One might hope that the advantages of the proposed rules could be obtained without question-dependence. Again, we close the door on that hope with a series of trivialization theorems (Sects. 13 and 14), employing the geometrical techniques described above. We conclude that, all things considered, the advantages of the recommended acceptance rules within questions justify their dependence on questions.

2 The geometry of the lottery paradox

Let $\mathcal{E} = \{E_i : i \in I\}$ be a countable collection of mutually exclusive and jointly exhaustive propositions over some underlying set of possibilities. Let κ (either ω or some finite n) denote the cardinality of \mathcal{E} . We think of \mathcal{E} as a *question* in context whose *potential answers* are the various E_i . Let \mathcal{A} be the least collection of propositions containing \mathcal{E} that is closed under negation and countable disjunction and conjunction, and let \mathcal{P} denote the set of all (countably additive) probability measures defined on \mathcal{A} . We think of \mathcal{P} as the space of *probabilistic credal states* over answers to question \mathcal{E} . Occasionally we write \mathcal{P}_κ or \mathcal{E}_κ to emphasize the cardinality of \mathcal{E} .

We assume that acceptance rules produce sets of propositions that are closed under classical entailment so that, without loss of generality, each acceptance rule may be viewed as a map $\alpha : \mathcal{P} \rightarrow \mathcal{A}$, where proposition $\alpha(p)$ is understood as the strongest proposition accepted in light of probability measure p . Then proposition A is *accepted* by rule α at credal state p , written $p \Vdash_\alpha A$, if and only if $\alpha(p)$ entails A . The *acceptance zone* of A under α is defined as the set of all credal states at which A is accepted by α .

For example, the *Lockean* acceptance rule with threshold set to r in the unit interval is just the mapping:

$$\lambda_r(p) = \bigwedge \{A \in \mathcal{A} : p(A) \geq r\}. \quad (1)$$

Each probability measure p in \mathcal{P} can be represented with respect to \mathcal{E} as the κ -dimensional vector $(p(E_i) : i \in I)$ with components in the unit interval summing to one. In the context of question \mathcal{E} , we identify p with its vector, so that the i th component p_i equals $p(E_i)$. In the case of three answers, \mathcal{P} corresponds to the set of all such 3-vectors, which is the equilateral triangle in \mathbb{R}^3 whose corners have Cartesian coordinates $e_1 = (1, 0, 0)$, $e_2 = (0, 1, 0)$ and $e_3 = (0, 0, 1)$ (Fig. 1). To avoid ambiguity,




Fig. 1 The space \mathcal{P}_3 of probabilistic credal states




Fig. 2 Acceptance zone for $E_2 \vee E_3$ under λ_r

we let $(e_i)_j$ pick out the j th component of e_i . Reformulate the Lockean rule (1) as follows³:

$$\lambda_r(p) = \bigwedge \{ \neg E_i : p(\neg E_i) \geq r \text{ and } i \in I \}; \tag{2}$$

$$= \bigwedge \{ \neg E_i : p_i \leq 1 - r \text{ and } i \in I \}. \tag{3}$$

By this formulation, the acceptance zone of $\neg E_1$ under λ_r with respect to question \mathcal{E}_3 is depicted in Fig. 2. The Lockean rule is *geometrical*—its acceptance zone for $\neg E_1$

³ This is equivalent to the original formulation because, first, every proposition A is equivalent to the conjunction of all propositions of form $\neg E_i$ that are entailed by A and, second, propositions of form $\neg E_i$ that are entailed by A are at least as probable as A .




Fig. 3 Acceptance zones under λ_r

has a definite, trapezoidal *shape* that results from truncating the triangular space \mathcal{P}_3 parallel to one side. As threshold r is dropped, the trapezoid becomes thicker. The acceptance zones of $\neg E_2$ and $\neg E_3$ are included in Fig. 3a. By closure under entailment, proposition E_1 is accepted exactly when both $\neg E_2$ and $\neg E_3$ are accepted, so the corner, diamond-shaped zones license acceptance of potential answers to \mathcal{E} . When $r \leq 2/3$, the propositions $\neg E_1, \neg E_2, \neg E_3$ are all accepted at the probability measures contained in the small, dark, central triangle (Fig. 3b). But that set of propositions is inconsistent so, by closure under entailment, the dark, central triangle is the acceptance zone of the inconsistent proposition \perp . That is just the lottery paradox for thresholds $r \leq 2/3$ (interpret E_i as the proposition “ticket i wins”).

Geometrically, the lottery paradox arises because the Lockean rule’s acceptance zones for the various propositions $\neg E_i$ crash clumsily into one another as the probability threshold r decreases. It is easy to design alternative acceptance zones that *bend* progressively as they approach the center of the triangle so that they eventually meet without overlapping like the leaves of a camera shutter (Fig. 4). The proposed acceptance zones are almost indistinguishable from those of the Lockean rule when r is close to 1. As r approaches 0, the bending becomes more pronounced and the lottery paradox is avoided.




Fig. 4 Progressively bent zones that avert collision

A special, symmetric case of the proposed rule, which we call the *symmetric camera shutter rule*, modifies the Lockean rule as follows. Test whether answer E_i to \mathcal{E} should be rejected at credal state p by considering, not probability p_i itself, but the probability ratio:

$$\sigma(p)_i = \frac{p_i}{\max_j p_j},$$

resulting in the modified rule:

$$v_r(p) = \bigwedge \{ \neg E_i : \sigma(p)_i \leq 1 - r \text{ and } i \in I \}. \tag{4}$$

The symmetric camera shutter rule is algebraically the same as the Lockean rule (3) except that probability is divided by the probability distribution’s mode. Say that acceptance rule α is *everywhere consistent* if and only if $p \not\llcorner_\alpha \perp$ for each p in \mathcal{P} , and say that α is *non-skeptical* if and only if for each E_i in \mathcal{E} there exists p in \mathcal{P} such that $p(E_i) < 1$ and $p \Vdash_\alpha E_i$. Then:

Proposition 1 *Let \mathcal{E} contains at least two answers. The symmetric camera shutter rule v_r is everywhere consistent and non-skeptical, for each r such that $0 < r < 1$.*

Proof For everywhere consistency, note that since $\sum_i p_i = 1$, so there exists $i \in I$ such that $p_i = \max_j p_j$. Then, since $r > 0$,

$$\sigma(p)_i = 1 \not\leq 1 - r$$

so $p \not\llcorner_{v_r} \neg E_i$, by formula (4). It follows that $p \not\llcorner_{v_r} \perp$. For non-skepticism, let E_i be an arbitrary answer, and it suffices to show that E_i is accepted by v_r at some credal state p such that $p_i < 1$. Let $p_i = 1/(2 - r)$. Since \mathcal{E} contains at least two answers, choose j in I distinct from i and let $p_j = (1 - r)/(2 - r)$. Since a probability distribution is normalized, $p_k = 0$ for all $k \neq i, j$. Note that p_i is the mode of p , since $r > 0$. So for each $k \neq i$:

$$\begin{aligned} \sigma(p)_k &\leq 1 - r, \\ \sigma(p)_i &= 1 \not\leq 1 - r, \end{aligned}$$

since $r > 0$. Hence $p \Vdash_{v_r} E_i$, by formula (4), with $p_i = 1/(2 - r) < 1$, since $r < 1$. □

On the other hand:

Proposition 2 *Suppose that \mathcal{E} is countably infinite. The Lockean rule λ_r is either skeptical or somewhere inconsistent, for each r such that $0 \leq r \leq 1$.*

Proof If the Lockean rule is not skeptical, then $r < 1$, and thus there exists p in \mathcal{P}_ω such that $p_i \leq 1 - r$, for each $i \in I$. So by formula (3), $\lambda_r(p) = \perp$, and hence λ_r is somewhere inconsistent. □

3 Respect for logic

The range of acceptance rule $\alpha : \mathcal{P} \rightarrow \mathcal{A}$ has a natural, Boolean logical structure:

$$(\mathcal{A}, \leq, \vee, \wedge, \perp, \top),$$

where the partial order \leq corresponds to classical entailment or relative strength of propositions and \vee and \wedge are the least upper bound and the greatest lower bound with respect to \leq , which correspond to the usual propositional operations of disjunction and conjunction.⁴ If there were also a motivated logical structure

$$(\mathcal{P}, \leq, \vee, \wedge, \perp, \top)$$

on the space of probabilistic credal states, in which \leq is intended, again, to reflect relative strength, then an obvious constraint on acceptance rules would be to *preserve* logical structure in the sense that:

$$p \leq q \Rightarrow \alpha(p) \leq \alpha(q); \quad (5)$$

$$\alpha(p \vee q) = \alpha(p) \vee \alpha(q); \quad (6)$$

$$\alpha(p \wedge q) = \alpha(p) \wedge \alpha(q); \quad (7)$$

$$\alpha(e_i) = E_i; \quad (8)$$

$$\alpha(\top) = \top; \quad (9)$$

$$\alpha(\perp) = \perp. \quad (10)$$

Any plausible logical structure over \mathcal{P} should also satisfy the following constraint:

$$\text{the unit vectors } e_i, \text{ for } i \in I, \text{ are exactly the strongest credal states in } \mathcal{P}. \quad (11)$$

Then we already have the following assurance against inconsistency:

Proposition 3 (no lottery paradox) *Suppose that acceptance rule α and relative strength \leq over \mathcal{P} satisfy conditions (5), (8), and (11). Then α is everywhere consistent.*

Proof Suppose for reductio that for some credal state p , $\alpha(p) = \perp$. Then, by condition (11), there exists a strongest state e_i such that $e_i \leq p$. So $\alpha(e_i) \leq \alpha(p)$, by (5). Then by (8), we have that $E_i = \alpha(e_i) \leq \alpha(p) = \perp$. So $E_i \leq \perp$, which is false in the Boolean logical structure of \mathcal{A} . \square

Therefore, the lottery paradox witnesses the failure of the Lockean rule to preserve logical structure. But the lottery paradox is only the most glaring consequence of the Lockean rule's disrespect for logical structure. It is plausible to suppose that with respect to question \mathcal{E} , if one's credal state p accords non-maximal probability to answer

⁴ In algebraic logic, $A \leq B$ means that A is at least as strong as B .




Fig. 5 Deeper trouble for the Lockean rule

E_i , compared to all the alternative answers to \mathcal{E} , then upon receiving the information that refutes exactly E_i , the posterior credal state $p(\cdot | \neg E_i)$ is at least as strong as p ; namely:

$$p_i \neq \max_j p_j \implies p(\cdot | \neg E_i) \leq p. \tag{12}$$

But then the Lockean rule again fails to preserve relative strength, i.e., it violates condition (5). Recall from Fig. 3 that a consistent Lockean rule’s acceptance zone for E_2 is a diamond. The diamond has the wrong shape—its sides meet at an angle that is too acute. For consider a credal state p very close to the inner apex of the diamond, as depicted in Fig. 5. Let $q = p(\cdot | \neg E_3)$. By condition (12), we have that $q \leq p$. But point q lies on the side of the triangle opposite e_3 because $q_3 = 0$, and q lies on the ray from e_3 that passes through p because $q_1/q_2 = p_1/p_2$. So $\lambda(q) = \neg E_3 \not\leq E_2 = \lambda(p)$. Therefore, $q \leq p$ but $\lambda(q) \not\leq \lambda(p)$, which violates (5). That is another counterintuitive way to fail to preserve logical order even when the lottery paradox does not arise.

The preceding argument illustrates a further point: intuitions about relative strength of credal states are tied to conditioning. The boundaries of acceptance zones determined by the Lockean rule do not follow the geometrical rays that correspond to the trajectories of probabilistic credal states under conditioning. For that reason, the Lockean rule is a bad choice for trying to explicate the acceptance of conditionals in terms of conditional probabilities. Specifically, consider the following interpretation of the Ramsey test. Let A, B be arbitrary propositions in \mathcal{A} . Let the flat conditional with antecedent A and consequent B be expressed by $A \Rightarrow B$. (The arrow notation ‘ \Rightarrow ’ does not denote a binary operation on propositions; it is simply a suggestive way to refer to the ordered pair (A, B) . We propose that the Ramsey test be explicated by the following acceptance condition for flat conditionals⁵:

$$p \Vdash_\alpha A \Rightarrow B \iff p(\cdot | A) \Vdash_\alpha B \text{ or } p(A) = 0. \tag{13}$$

So when the antecedent has nonzero probability, this semantic rule says that flat conditional $A \Rightarrow B$ is accepted at credal state p if and only if, when one “adds A

⁵ If $p(A) \neq 0$, $p(\cdot | A)$ is defined to be $p(\cdot \wedge A) / p(A)$; otherwise it is undefined.

hypothetically to one's stock of knowledge" and thereby hypothetically conditions p on A to obtain posterior credal state $p(\cdot|A)$, one accepts the consequent B in the posterior state. Consider again the consistent Lockean rule λ and credal state p in Fig. 5. Then, as is evident from the picture, we have:

$$\begin{aligned} p &\Vdash_{\lambda} E_2, \\ p &\not\Vdash_{\lambda} \neg E_3 \Rightarrow E_2. \end{aligned}$$

Note that $\neg E_3$ is entailed by E_2 , so Lockean rule λ instructs one to retract her acceptance of E_2 when a logical consequence of E_2 is learned or supposed. That contradicts intuitions about scientific method and violates almost all logics of conditionals that have interested logicians, including Adams (1975) conditional logic.⁶ On the other hand, to preserve acceptance under logically entailed information, it suffices to require conditions (5) and (12). For by (12), credal state q would have been at least as strong as credal state p and hence, by (5), any proposition accepted in p remains accepted in q , e.g., E_2 .

The angles formed by the sides of the acceptance zones are crucial to the preservation of logical structure. The acceptance rules we recommend—the camera shutter rules—do have acceptance zones with the correct angles at their corners and, therefore, do not encounter any of the preceding logical difficulties. We will show that the camera shutter rules preserve a very natural logical structure on state space \mathcal{P} and, therefore, yield a soundness and completeness theorem for Adams' conditional logic that is simpler and more natural than Adams' original version (1975).

4 Geologic

Consider classical, infinitary propositional logic, which allows for countable disjunction and conjunction.⁷ Start with propositional constants \perp , \top and propositional variables $V_{\kappa} = \{E_i : i \in I\}$, where the countable index set I has cardinality κ . Let $\bigvee_j \phi_j$ and $\bigwedge_j \phi_j$ be countable disjunction and conjunction, respectively. Let language L_{κ} be the least set containing the propositional constants in V_{κ} that is closed under negation, countable disjunction, and countable conjunction. We interpret the propositional variables to be mutually exclusive and exhaustive. Under that restriction, each truth assignment is a κ -dimensional basis vector e_i . Let \mathcal{B}_{κ} denote the set of all such vectors. The valuation function for classical logic is definable as follows. In the base case:

$$v_{e_i}(E_j) = e_i \cdot e_j; \quad v_{e_i}(\top) = 1; \quad v_{e_i}(\perp) = 0,$$

⁶ Specifically, the principle that acceptance be preserved under logically implied information can be shown to be equivalent to Cautious Monotonicity, given two other axioms in Adams' conditional logic (a.k.a system P): Right Weakening, And. That system will be discussed in detail later.

⁷ For classic studies concerning completeness infinitary logic, cf. Karp (1964) and Barwise (1969). Our applications make no reference to completeness or to proof systems for infinitary logic.

where \cdot denotes the vector *inner product* $x \cdot y = \sum_{i \in I} x_i y_i$. In the inductive case:

$$v_{e_i}(\neg\phi) = 1 - v_{e_i}(\phi); \quad v_{e_i}\left(\bigvee_j \phi_j\right) = \max_j(v_{e_i}(\phi_j));$$

$$v_{e_i}\left(\bigwedge_j \phi_j\right) = \min_j(v_{e_i}(\phi_j)).$$

Logical entailment is definable in terms of valuation as follows:

$$\phi \models \psi \iff v_{e_i}(\phi) \leq v_{e_i}(\psi), \quad \text{for all } i \in I.$$

Let the *proposition* $\llbracket \phi \rrbracket_\kappa$ expressed by ϕ in language L_κ denote the set of all assignments in \mathcal{B}_κ in which ϕ evaluates to 1. Each proposition $\llbracket \phi \rrbracket_\kappa$ is represented uniquely by its *valuation vector*:

$$v_\kappa(\phi) = (v_{e_i}(\phi) : i \in I),$$

which belongs to 2^κ . Define the following relations and operations over 2^κ :

$$u \leq v \iff u_i \leq v_i, \quad \text{for all } i \text{ in } I; \tag{14}$$

$$(\neg v)_i = 1 - v_i; \tag{15}$$

$$\left(\bigvee_j v^j\right)_i = \max_j v_i^j; \tag{16}$$

$$\left(\bigwedge_j v^j\right)_i = \min_j v_i^j. \tag{17}$$

Then the structure of classical, infinitary logic is captured⁸ by the mathematical structure:

$$\mathcal{L}_\kappa = \left(2^\kappa, \leq, \bigvee, \bigwedge, 1, 0\right).$$

Figure 6a illustrates \mathcal{L}_3 , which bears a suggestive resemblance to the unit cube $[0, 1]^3$ (Fig. 6b), but it is really just a string-and-bead figure whose strings happen to be sized and stretched to outline a cube. However, one can *extend* classical propositional logic on L_κ to a *fuzzy language* L_κ^* that generates fuzzy propositions covering the entire κ -dimensional unit cube $[0, 1]^\kappa$.⁹ A *fuzzy proposition* is just a *fuzzy subset* (Zadeh

⁸ I.e., the Lindenbaum-Tarski algebra of language L_κ is isomorphic to \mathcal{L}_κ .

⁹ The idea may sound similar to multi-valued logic, but it is quite different. In multi-valued logic, (discrete) logical formulas in L_κ are interpreted over an expanded, continuous space of assignments (Hajek 1998; Novak et al. 2000)—such logics generate a discrete, weakening of classical logic, rather than a continuous, conservative extension of classical logic.




Fig. 6 Bead-and-string logic versus geologic

1965) of \mathcal{B}_κ , which is representable by a *fuzzy characteristic function* from \mathcal{B}_κ to $[0, 1]$ and, hence, by a *fuzzy valuation vector* v in $[0, 1]^\kappa$. Formula (14) represents the fuzzy subset relation and formulas (15) through (17) correspond to fuzzy complement, intersection, and union over fuzzy propositions.

Here is one natural way to extend classical logic over L_κ to cover the κ -dimensional unit cube. For each real number d in the unit interval, let the *partial negation* $\neg_d \phi$ be understood as the negation of ϕ to *degree* d , interpreted as follows:

$$v_{e_i}(\neg_d \phi) = d v_{e_i}(\neg\phi) + (1 - d)v_{e_i}(\phi).$$

In particular, $\neg_0\phi$ is equivalent to ϕ , whereas $\neg_1\phi$ is equivalent to $\neg\phi$. Between these extremes, $\neg_{1/2}\phi$ hovers semantically midway between ϕ and $\neg\phi$. Let L_κ^* be the result of expanding language L_κ with \neg_d . Otherwise, the preceding definitions of valuation function v_{e_i} and valuation vector $(v_{e_i}(\phi) : i \in I)$ remain unaltered.¹⁰ Partial negation never generates values outside of the unit interval, so all valuation vectors for L_κ^* are in the unit cube $[0, 1]^\kappa$. Conversely, every vector v in $[0, 1]^\kappa$ is the valuation vector of some formula in L_κ^* , namely:

$$\bigvee_{i \in I} (\neg_{1-v_i} E_i \wedge E_i).$$

So the propositions expressible by the fuzzy language \mathcal{L}_κ^* correspond to the vectors in the κ -dimensional unit cube $[0, 1]^\kappa$. Therefore, we refer to the logic just defined as *geologic*.

Formulas (14)–(17) still make sense for fuzzy valuation functions (because they correspond to the standard definitions of the fuzzy set theoretic operations). Therefore,

¹⁰ More directly, one can simply introduce a new unary connective $a\phi$ called *scalar multiple* interpreted by $v_{e_i}(a\phi) = av_{e_i}(\phi)$. But we found it harder to motivate usage of such a connective.




Fig. 7 Geological operations

the *structure* of geologic is:

$$\mathcal{L}_\kappa^* = ([0, 1]^\kappa, \leq, \vee, \wedge, 1, 0).$$

Since the valuation definition for geologic is exactly the same as for classical logic over the fragment L_κ , it follows that \mathcal{L}_κ^* restricted to L_κ is just \mathcal{L}_κ —in other words, geologic is a conservative extension of classical, infinitary logic.

Since the operations in \mathcal{L}_κ^* correspond to fuzzy set theoretical operations on propositions, it is immediate that the geological operations satisfy associativity, commutativity, distributivity, and the De Morgan rules (Zadeh 1965). Excluded middle and disjunctive syllogism, on the other hand, can fail spectacularly for propositions in the unit cube’s interior. For example, let c denote the center $(\frac{1}{2}, \dots, \frac{1}{2}, \dots)$ of the unit cube. Then:

$$\begin{aligned} \neg c &= c; \\ c \vee \neg c &= c; \\ (e_1 \vee c) \wedge \neg c &= c. \end{aligned}$$

In spite of that, we think of geologic as the natural *extension* of classical logic to fuzzy propositions. Associativity, commutativity, distributivity, and the De Morgan rules are all motivated by symmetries of the unit cube. Excluded middle is not motivated by symmetry—it is a mere artifact of an impoverished syntax. Furthermore, unlike modal logic, which is also a conservative extension of classical propositional logic, geologic arises from the addition of an *extensional* negation.

Filling the interior of the Boolean algebra to make it a genuine cube provides an explanatory, geometrical perspective on classical logic. For example, given points v and u in the unit cube, find the smallest parallelepiped solid $S(v, u)$ containing v and u




Fig. 8 Geometry of the De Morgan rules

whose sides are parallel to the sides of the cube. Then the uppermost vertex of $S(v, u)$ is $v \vee u$ and the lowermost vertex of $S(v, u)$ is $v \wedge u$ (Fig. 7a). The parallelepiped $S(v, u)$ is like a sub-crystal within the cube, which is another reason for thinking of geologic as *geological*.¹¹ The geometry of full geological negation is just reflection through the center c of the cube, which is a natural generalization of Boolean complementation. To construct the partial negation $\neg_d v$ of v , first reflect v through c to obtain the full negation $\neg v$. Now draw a straight line segment between v and $\neg v$. Then $\neg_d v$ is the point that lies proportion d of the way from v to $\neg v$ along the line segment (Fig. 7b). Consider the classical De Morgan rules. Since full negation involves projection through the center c of the cube, think of c as the aperture of a pinhole camera. It is a familiar fact that projection through an aperture inverts the image. But the disjunction $v \vee u$ is the top vertex of the parallelepiped spanning v and u . Projecting the parallelepiped through the aperture inverts it and turns the top vertex into the bottom vertex—the conjunction of the projection of v with the projection of u (Fig. 8).

5 Logic from a probabilistic perspective

For our purposes, the point of geologic is that it affords a unified perspective on logic and probability.¹² The set \mathcal{P}_3 of possible credal states is a horizontal, triangular plane through the unit cube of geological propositions (Fig. 6b). Thus, credal state space

¹¹ Note that the same geometrical relationships would hold even if the unit cube were stretched along its various axes to form a prism. We will return to that theme in the last section of the paper.

¹² Due to the truth-functionality of conjunction in fuzzy logic, the fuzzy logic community tends to view fuzzy logic in isolation from probability theory, rather than as a tool for understanding probability theory, as we propose.

\mathcal{P}_3 has a natural embedding *within* geologic. That embedding generalizes to each countable cardinality κ .¹³

Valuation and probability assignment can both be viewed as inner products within the geological cube:

$$\begin{aligned} v_{e_i}(u) &= e_i \cdot u, \\ p(b) &= p \cdot b, \end{aligned}$$

where u is a vector in $[0, 1]^\kappa$ corresponding to an arbitrary, geological proposition, v_{e_i} is a valuation function corresponding to a classical assignment, b is a Boolean valuation vector in 2^κ corresponding to a classical proposition, and p is a probability measure/vector in \mathcal{P}_κ .

Say that probability measure p is *uniform* with respect to \mathcal{E} if and only if p assigns only zero or a fixed value to the answers in \mathcal{E} . The *support* of p is the disjunction of all elements of \mathcal{E} that p assigns non-zero probability to (recall that \mathcal{E} is countable). The *classical principle of indifference* is a mapping σ that associates each uniform probability distribution p with its support. For example, σ associates $(\frac{1}{2}, \frac{1}{2}, 0)$ with the classical proposition $\sigma(\frac{1}{2}, \frac{1}{2}, 0) = (1, 1, 0)$. Construct a ray from \perp through uniform distribution p and then $\sigma(p)$ is the (classical) proposition on the upper surface of the unit cube that the ray points to (Fig. 9a). Algebraically, $\sigma(p)$ is the (unique) scalar multiple of p in the unit cube that has at least one component equal to 1, which amounts to the formula encountered earlier in the definition of the symmetric camera shutter acceptance rules:

$$\sigma(p)_i = \frac{p_i}{\max_j p_j}, \quad \text{for } i \in I.$$

Say that geological proposition u is *fully satisfiable* if and only if there exists e_i such that $v_{e_i}(u) = 1$, i.e. u has a component equal to 1. So $\sigma(p)$ is the (unique), fully satisfiable, geological valuation vector that is proportional to p . In classical logic, the mapping $\sigma(p)$ is defined only for uniform p , but it is defined for *all* p in geologic, since the (continuous) upper surface of the geological cube covers the entire triangle of probability measures (Fig. 9b). Now *every* probability measure p has a unique, geological proposition $\sigma(p)$ that stands to p in much the same way that the support of p stands to uniform p .

Mapping σ has a heuristic interpretation. Think of the unit cube as a room with tiled walls (Fig. 10). Imagine that there is a digital camera embedded in the baseboard of the room at corner \perp . Think of the triangle \mathcal{P}_3 as the picture plane corresponding to the 2-dimensional image received by the camera. Then the inverse σ^{-1} of σ is the classical perspective rendering of the room's interior on the picture plane. The perspective is extreme because the camera is literally embedded in the lower corner

¹³ Perhaps the destiny of geologic is to liberalize all of the type restrictions on the preceding two equations, so that the entire unit cube does triple-duty as generalized propositions, generalized inconsistent or indeterministic assignments, and generalized non-normalized degrees of belief. We leave that solid philosophy unexplored in this paper.




Fig. 9 Indifference as projection




Fig. 10 A literal, probabilistic perspective on logic

of the room, so the floor and adjacent walls are tangent to the camera’s view and are rendered as the boundaries of the triangular picture.

The point of the preceding detour through geologic is that the picture plane is the space \mathcal{P}_3 of probability measures on \mathcal{A}_3 and the walls and ceiling of the office are the fully satisfiable propositions in geologic. So Fig. 10 literally illustrates geologic from a *probabilistic perspective*. That perspective sheds new light on the lottery paradox and its associated conundrums. In particular, note the similarity between the acceptance zones of the proposed, paradox-avoiding rule (Fig. 4) and the projected coordinate lines of the unit cube (Fig. 10b). The boundaries of the former always follow the latter.

6 Probalogic

We understand “logic” in the broad, pragmatic sense that logic is wherever logical structure is. If the logical structure pertains to relative strength of credal states, then there is a logic of such states, even though the states in question are not necessarily propositional and the logical relations among them are not plausibly interpreted as arguments. And if the structure happens to be relative to pragmatic factors such as a question that elevates the significance of certain propositions as relevant answers, then logic, itself, is pragmatic—we do not insist that logic must in some sense be prior to or independent of such considerations. Our view accords with an ancient tradition according to which logic is a tool or organon for *inquiry*, which typically begins with some question and ends with an answer thereto. In this section, we introduce a logic of probabilistic credal states in the broad, pragmatic sense just outlined.

When credence is modeled as qualitative belief in a proposition, it is straightforward to judge the relative strength of credal states in terms of the classical, logical strength of the propositions believed:

$$B\phi \leq B\psi \iff \phi \leq \psi.$$

We propose, in a similar spirit, that probabilistic credal states inherit their logical strength from their unique, geological images:

$$p \leq q \iff \sigma(p) \leq \sigma(q) \tag{18}$$

$$\iff \sigma(p)_i \leq \sigma(q)_i \text{ for all } i \in I. \tag{19}$$

Disjunction \vee and conjunction \wedge are standardly defined, respectively, as the least upper bound and the greatest lower bound with respect to \leq . We call the resulting logical structure on probability measures *probalogic*:

$$(\mathcal{P}, \leq, \vee, \wedge).$$

Probalogic is just geologic from a probabilistic perspective.

Consider arbitrary credal state p in \mathcal{P}_3 . Which credal states are probalogically at least as weak as p ? First, project p up to geological proposition $\sigma(p)$ on the upper surface of the geological cube. The geological consequences of $\sigma(p)$ consist of the parallelepiped containing \top whose sides are parallel to the sides of the unit cube and whose bottom-most corner is $\sigma(p)$ (Fig. 11). Since $\sigma(p)$ is incident to an upper surface of the cube, the parallelepiped is, in this case, a rectangle lying entirely in one upper face of the unit cube (or, in degenerate cases, entirely within an upper edge of the unit cube). The credal states probalogically weaker than p are contained within the linear perspective projection of that rectangle onto the picture plane \mathcal{P}_3 . Note that, according to the usual rules of linear perspective, parallel sides of the rectangles meet at *vanishing points*, which correspond to the corners of \mathcal{P}_3 that are not closest to p . Similarly, the geological propositions in the range of σ that are geologically at least as strong as $\sigma(p)$ are in the rectangle with sides parallel to the sides of the unit cube that has $\sigma(p)$ as its upper corner and the nearest unit vector e_i to $\sigma(p)$ as its lower




Fig. 11 Probalogical strength

corner. So the inverse image of that rectangle under σ is the set of the credal states that are probalogically at least as strong as $\sigma(p)$. We call the partial order \leq so defined *relative probalogical strength*.

Probalogical disjunction, conjunction, and negation can be defined similarly, as the projections of the corresponding, geological disjunction:

$$p \vee q = \sigma^{-1}(\sigma(p) \vee \sigma(q)); \tag{20}$$

$$p \wedge q = \sigma^{-1}(\sigma(p) \wedge \sigma(q)); \tag{21}$$

$$\neg p = \sigma^{-1}(\neg\sigma(p)). \tag{22}$$

Since the geological disjunction of two propositions on the upper surface of the unit cube is also on the upper surface of the unit cube, \mathcal{P}_3 is closed under probalogical disjunction. Geometrically, these logical operations can be constructed as perspective renderings of the corresponding geological operations on the cube (Figs. 12, 13, and 14). Probalogical constants and operations are not necessarily defined. In finite questions, \top denotes the uniform distribution, but in countably infinite questions there is no such distribution. There is no interpretation of \perp . Letting $\perp = \top$ is obviously unappealing, but any choice of \perp that is off-center is equally implausible. Geological negation is closed over the lower edges of the upper faces of the unit cube, but is not closed elsewhere over the upper faces of the unit cube, so probalogical negation is defined only over the perimeter of \mathcal{P} . Furthermore, if $\sigma(p)$ and $\sigma(q)$ are on different upper faces of the unit cube, then the conjunction $\sigma(p) \wedge \sigma(q)$ lies below the upper faces of the unit cube, so $p \wedge q = \sigma^{-1}(\sigma(p) \wedge \sigma(q))$ is undefined in that case.

Although we will not pursue the idea in this paper, there is a way to expand \mathcal{P} to a space over which probalogical conjunction and disjunction are closed. Some assumptions are so certain that one does not even conceive of their falsity—e.g., that a particle cannot have two distinct momenta at the same time but can have a definite




Fig. 12 Probological conjunction and disjunction within a face




Fig. 13 Probological conjunction and disjunction across faces




Fig. 14 Negation around the perimeter




Fig. 15 σ extended to measures normalized to a value ≤ 1

momentum and position at the same time. But when experience gets strange, we may come to doubt our basic assumptions without having thought yet of any concrete alternatives. In such cases, a natural response is to transfer probability mass to a non-descript “catchall hypothesis” absent from the original algebra \mathcal{A}_3 . Within \mathcal{A}_3 , the resulting credal state appears to be normalized to a value less than 1. Accordingly, let \mathcal{P}^* denote the set of all additive measures p on \mathcal{A} such that $0 \leq p(\top) \leq 1$. Then the problem of closure under negation and conjunction is solved by plausibly extending σ to a bijection between \mathcal{P}^* and the entire unit cube as follows (Fig. 15):

$$\sigma^*(p)_i = p(\top) \cdot \frac{p_i}{\max_j p_j}, \text{ for } i \in I.$$

So Eqs. 18–22, with σ replaced by σ^* , induce a probalogical structure on \mathcal{P}^* that is closed under the probalogical operations of conjunction, disjunction, and negation.

7 Acceptance that respects probalogic

A *probalogical* acceptance rule ν is an acceptance rule that preserves probalogical structure in the sense of morphism conditions (5)–(8).¹⁴ As described in the preceding section, condition (7) is understood to hold only when $p \wedge q$ is defined over \mathcal{P} .

Recall the camera-shutter-like acceptance rules introduced above as one geometrical strategy for solving the lottery paradox. The rules can be stated a bit more generally, by allowing the threshold r and the strictness of the inequality to vary with i . Say that acceptance rule ν is a *camera shutter rule* for \mathcal{E} if and only if there exist thresholds $\{r_i : i \in I\}$ in the unit interval and inequalities $\{\triangleleft_i : i \in I\}$ that are either \leq or $<$, such that for each p in \mathcal{P} and $i \in I$:

¹⁴ Note that (5) is redundant, for it is derivable from (6).

1. $v(p) = \bigwedge \{ \neg E_i : \sigma(p)_i \triangleleft_i 1 - r_i \text{ and } i \in I \}$;
2. if $\triangleleft_i = \leq$ then $r_i > 0$;
3. if $\triangleleft_i = <$ then $r_i < 1$.

Note that 0 is omitted in the second condition to make it possible to not accept $\neg E_i$, and 1 is omitted in the third condition to make it possible to accept $\neg E_i$ —else morphism condition (8) would be violated trivially. The main result of this section is that, over countable dimensions, the camera shutter rules are precisely the rules that preserve probalogic.

Theorem 1 (representation of probalogical rules) *Suppose that \mathcal{E} is countable. Then an arbitrary acceptance rule is probalogical if and only if it is a camera shutter rule.*

The proof proceeds by a series of lemmas. Let p, q be in \mathcal{P} . Define:

$$q \leq_i p \iff \sigma(q)_i \leq \sigma(p)_i.$$

Lemma 1 *Suppose that $q \leq_i p$. Then $p = (p \vee e_i) \wedge (p \vee q)$.*

Proof See Fig. 16. By the definition of probalogic in terms of geologic, it suffices to show that

$$\sigma(p) = \sigma((p \vee e_i) \wedge (p \vee q)).$$

By geologic, the j th component of the right hand side expands to:

$$\min(\max(\sigma(p)_j, \sigma(e_i)_j), \max(\sigma(p)_j, \sigma(q)_j)).$$

Since $(e_i)_i = 1$, it follows that $\max(\sigma(p)_i, \sigma(e_i)_i) = 1$. Since $\sigma(q)_i \leq \sigma(p)_i$, it follows that $\max(\sigma(p)_i, \sigma(q)_i) = \sigma(p)_i$. So $\sigma((p \vee e_i) \wedge (p \vee q))_i = \sigma(p)_i$. Now let E_j be in \mathcal{E} for $j \neq i$. Then $(e_i)_j = 0$, so $\max(\sigma(p)_j, \sigma(e_i)_j) = \sigma(p)_j$. In general, $\min(x, \max(x, y)) = x$, so we have as well that $\sigma((p \vee e_i) \wedge (p \vee q))_j = \sigma(p)_j$. \square

Lemma 2 *Let v satisfy morphism conditions (6), (7), and (8). Let $i \in I$. Then:*

$$p \Vdash_v \neg E_i \text{ and } q \leq_i p \implies q \Vdash_v \neg E_i.$$




Fig. 16 Proof of Lemma 1

Proof Suppose that $p \Vdash \neg E_i$ and that $q \leq_i p$. Since $p \Vdash \neg E_i$, it follows that $p_i < \max_k p_k$. For otherwise, $e_i \leq p$, so by morphism condition (5), $e_i \Vdash \neg E_i$, contrary to morphism condition (8). Since it is also the case that $q \leq_i p$, Lemma 1 yields that $p = (p \vee e_i) \wedge (p \vee q)$. Suppose for reductio that $v(q)$ is logically compatible with E_i . Then by morphism condition (6), $v(p \vee q)$ is compatible with E_i . By morphism condition (8), $v(e_i)$ is compatible with E_i . So again by morphism condition (6), $v(p \vee e_i)$ is compatible with E_i . So $v(p) = v((p \vee e_i) \wedge (p \vee q))$ is compatible with E_i , by morphism condition (7) and by the fact that E_i is an atom in algebra \mathcal{A} . But $p \Vdash_v \neg E_i$. Contradiction. Hence, $q \Vdash_v \neg E_i$. \square

Proof of Theorem 1 For the only if side, let $i \in I$. Define:

$$1 - r_i = \sup\{\sigma(p)_i : p \in \mathcal{P} \text{ and } p \Vdash_v \neg E_i\}.$$

Suppose that $\sigma(p)_i < 1 - r_i$. Then $p \Vdash_v \neg E_i$, by Lemma 2. Suppose that $\sigma(p)_i > 1 - r_i$. Then $p \not\Vdash_v \neg E_i$, by the definition of $1 - r_i$. Finally, suppose that $\sigma(p)_i = \sigma(q)_i = 1 - r_i$. Consider the case in which there exists r in \mathcal{P} such that $\sigma(r) = 1 - r_i$ and $r \Vdash_v \neg E_i$. Then $p \Vdash_v \neg E$ and $q \Vdash_v \neg E$, by Lemma 2. In the alternative case, it is immediate that $p \not\Vdash_v \neg E$ and $q \not\Vdash_v \neg E$. Thus, $p \Vdash_v \neg E_i$ if and only if $q \Vdash_v \neg E_i$. Set $\triangleleft_i = \leq$ in the former case and set $\triangleleft_i = <$ in the latter case. In the former case, suppose for reductio that $r_i = 0$. Then $v(e_i) \Vdash \neg E_i$, contradicting morphism condition (8), so $r_i > 0$, as required. In the latter case, suppose for reductio that $r_i = 1$. Then $v(e_i) \not\Vdash E_i$, contradicting morphism condition (8), so $r_i > 0$, as required. For the if side of the theorem, suppose that v is a camera shutter rule for countable \mathcal{E} . For morphism condition (5), suppose that $p \leq q$. Then $\sigma(p)_i \leq \sigma(q)_i$, for each $i \in I$. Then $q \Vdash_v \neg E_i$ implies $p \Vdash_v \neg E_i$, so $v(p) \leq v(q)$. For morphism condition (6), let $v(p) = A$ and $v(q) = B$, so:

$$A = \bigwedge \{\neg E_i : \sigma(p)_i \triangleleft_i 1 - r_i\};$$

$$B = \bigwedge \{\neg E_i : \sigma(q)_i \triangleleft_i 1 - r_i\}.$$

Let $\mathcal{D} = \{\neg E_i : A \leq E_i \text{ and } B \leq E_i\}$ and note that $A \vee B = \bigwedge \mathcal{D}$. Suppose that $\neg E_i$ is in \mathcal{D} . Then $\sigma(p)_i \triangleleft_i 1 - r_i$ and $\sigma(q)_i \triangleleft_i 1 - r_i$. Hence, $\max(\sigma(p)_i, \sigma(q)_i) \triangleleft_i 1 - r_i$. Thus, $v(p \wedge q) \leq \neg E_i$. Suppose that $\neg E_i$ is not in \mathcal{D} . Then either $\sigma(p)_i \not\triangleleft_i 1 - r_i$ or $\sigma(q)_i \not\triangleleft_i 1 - r_i$, so $\max(\sigma(p)_i, \sigma(q)_i) \not\triangleleft_i 1 - r_i$ and, thus, $v(p \wedge q) \not\leq \neg E_i$. Hence, $\neg E_i$ is in \mathcal{D} if and only if $v(p \wedge q) \leq \neg E_i$. Therefore, $v(p \vee q) = \bigwedge \mathcal{D} = A \vee B$. The dual argument works for morphism condition (7). \square

Recall that the conditions (5)–(7) omit preservation of negation and of the infinitary versions of disjunction and conjunction. There are good reasons to drop those conditions.

Proposition 4 *In finite dimensions, no probabilological acceptance rule preserves infinite conjunction and disjunction.*

Proof Consider probabilological acceptance rule v for question $\{E_i : i \in I\}$. By morphism condition (8), $v(e_1) = E_1$ and $v(e_2) = E_2$. Let L be the straight line connecting

e_1 with e_2 . Note that no uniform distribution with infinite support is encountered along this line, so it is continuous. So by morphism condition (5), there is a boundary point b such that $q \Vdash_\nu E_1$, for all q closer to e_1 than b , and $q \not\Vdash_\nu E_1$, for all q farther from e_1 than b . Let m be the mid-point of L . Consider the case in which p is between m and e_1 . Consider the case in which $b \Vdash_\nu E_1$. Let $\{p_i : i \in \mathbb{N}\}$ be a discrete sequence of points in line segment $\overline{e_1 b}$ that converges to b and let $\{q_i : i \in \mathbb{N}\}$ be a discrete sequence of points in line segment $\overline{m b}$ that converges to b . Then:

$$\bigvee_i p_i = b = \bigwedge_i q_i.$$

Suppose that $b \not\Vdash_\nu E_1$. Then $\nu(\bigvee_i p_i) \neq \bigvee_i \nu(p_i)$. Alternatively, suppose that $b \Vdash_\nu E_1$. Then $\nu(\bigwedge_i q_i) \neq \bigwedge_i \nu(q_i)$. □

Proposition 5 *In finite dimensions, no probalogical acceptance rule also preserves probalogical negation.*

Proof Let $p = (\frac{2}{3}, \frac{1}{3}, 0)$. Assume, for reductio, that acceptance rule ν is probalogical and preserves probalogical negation as well. So by Proposition 1, ν is a camera shutter rule. Suppose that ν rejects E_2 in p . So $\sigma(e_2) = \frac{1}{2} \triangleleft_2 1 - r_2$. Note, in Fig. 14, that $\neg p = (0, \frac{1}{3}, \frac{2}{3})$. So by preservation of negation, ν does not reject E_2 at $\neg p$. Thus: $\sigma(e_2) = \frac{1}{2} \not\triangleleft_2 1 - r_2$, which is a contradiction. The case in which ν does not reject E_2 in p is similar. The argument generalizes to arbitrary, finite dimensions. □

On the other hand, setting each $r_i = \frac{1}{2}$ almost preserves negation, in the sense that negation is preserved at all points on the perimeter of the triangle *except* at the six probability assignments with range $\{0, \frac{1}{3}, \frac{2}{3}\}$. But even so, no other setting for the r_i other than $\frac{1}{2}$ has that property, so the demands imposed by negation preservation are unreasonably strict.

8 Acceptance that does not respect probalogic

The acceptance rules we recommend, the camera shutter rules, are exactly the rules that preserve probalogical structure. Alternative acceptance rules proposed by Kyburg (1961) and by Pollock (1995) fail to preserve probalogical structure—actually, they fail to preserve any plausible logical structure.

Each Kyburgian acceptance rule χ_r is a Lockean rule without closure under conjunction:

$$\chi_r = \{A \in \mathcal{A} : p(A) \geq r\}.$$

Let question \mathcal{E} be ternary and set $r = \frac{2}{3}$. In Fig. 17, the set $\chi_{\frac{2}{3}}(c)$ of propositions accepted at the center $c = (\frac{1}{3}, \frac{1}{3}, \frac{1}{3})$ is indicated by a solid line and the set $\chi_{\frac{2}{3}}(e_3)$ is indicated by a dashed line. Rule $\chi_{\frac{2}{3}}$ does not preserve logical order in any plausible




Fig. 17 Kyburgian acceptance rule

sense, for corner e_3 is at least as strong as center c , but $\chi_{\frac{2}{3}}^2(e_3)$ is, intuitively, not at least as strong as $\chi_{\frac{2}{3}}^2(c)$ due to the retraction of $e_1 \vee e_2$.

There is, therefore, a hidden dilemma in Kyburg’s thesis that one should give up closure of accepted propositions under conjunction. On the one hand, if only \top is accepted at the uniform measure c , then there is no lottery paradox and, hence, there is no motivation for failing to close the accepted propositions under conjunction. On the other hand, if some proposition other than \top is accepted at c —say, a disjunction D that is incompatible with E_i —then, using the same argument as above, when one jumps from the center c to the stronger state e_i , one must accept E_i (which has probability one) and retract D (which has probability zero) and thus one must fail to expand the set of accepted propositions. In contrast, all camera shutter rules preserve probabilistic logic.

Pollock (1995), Ryan (1996), and Douven (2002) all propose what we will call *Pollockian* variants of the Lockean acceptance rule. The basic idea is to *restrict* the Lockean rule to cases in which it produces no paradox. The idea is illustrated, for ternary \mathcal{E} , in Fig. 18. The basic difference between Pollockian and Lockean rules in 3-dimensions is that the former return \top whenever the latter return \perp (compare to Fig. 3). The choice of \top as a substitute for \perp is natural enough, on grounds of symmetry, but due to the *shape* of Pollockian acceptance zones, there still exists no single logical structure that all Pollockian rules preserve.

Proposition 6 *Suppose that \mathcal{E} is ternary. Let \leq be an arbitrary partial order on \mathcal{P} whose binary least upper bound operation \vee is totally defined. Then there exists at least one Pollockian acceptance rule that is not a structure preserving map from $(\mathcal{P}, \leq, \vee)$ to $(\mathcal{A}, \leq, \vee)$.*

Proof Suppose the contrary for reductio. Let π_r be a Pollockian rule. When $r > \frac{2}{3}$, as in Fig. 18a, the rule π_r accepts $E_1 \vee E_2$ at $p = (\frac{1}{2}, \frac{1}{2}, 0)$ and $E_2 \vee E_3$ at $q = (0, \frac{2}{3}, \frac{1}{3})$, respectively, whose disjunction is $E_1 \vee E_2 \vee E_3 = \top$. So, to preserve disjunction, $p \vee q$ must lie within the white triangle, where \top is accepted. If we let r approach $\frac{2}{3}$ from




Fig. 18 Pollockian acceptance rules

above, as in Fig. 18b, the white triangle converges to the center point $c = (\frac{1}{3}, \frac{1}{3}, \frac{1}{3})$, so $p \Upsilon q = c$. Now consider the case in which $r < \frac{2}{3}$ (Fig. 18c). By preservation of disjunction, we have:

$$\begin{aligned}
 \top &= \pi_r(c) \\
 &= \pi_r(p \Upsilon q) \\
 &= \pi_r(p) \vee \pi_r(q) \\
 &= (E_1 \vee E_2) \vee E_2 \\
 &= E_1 \vee E_2.
 \end{aligned}$$

Hence $\top = E_1 \vee E_2$, a contradiction. □

A dilemma for Pollockian theorists is that, on the one hand, symmetry precludes accepting anything other than \top at the center point c , but that implies that there is no logical structure on \mathcal{P} that all Pollockian rules preserve. In contrast, all camera shutter rules preserve probalagic.

9 The geometry of conditional logic

As illustrated in Fig. 5, acceptance zones with the wrong shape can invalidate plausible principles of nonmonotonic reasoning. In fact, each axiom of the logic for flat conditionals corresponds to a definite, geometrical constraint on acceptance zones. The correspondences are established in this section and are used below to demonstrate that each probalogical rule validates a plausible set of axioms for conditional logic due to Adams (1975).

The acceptance condition of a conditional is defined by (13) as an explication of Ramsey test:

$$p \Vdash_{\alpha} A \Rightarrow B \iff p(\cdot|A) \Vdash_{\alpha} B \text{ or } p(A) = 0.$$

The set of axioms known as Adams' conditional logic (Adams 1975) or system P have been widely recognized as central to conditional and nonmonotonic reasoning

(Kraus et al. 1990). They state closure properties for a set of accepted conditionals. Here we rewrite them as closure properties for the set of conditionals accepted at a fixed credal state p under a fixed acceptance rule α (where the horizontal line means material implication):

$$\text{(Reflexivity)} \frac{}{p \Vdash_{\alpha} A \Rightarrow A}$$

$$\text{(Left Equivalence)} \frac{p \Vdash_{\alpha} A \Rightarrow B}{p \Vdash_{\alpha} C \Rightarrow B} \text{ if } A \text{ is classically equivalent to } C.$$

$$\text{(Right Weakening)} \frac{p \Vdash_{\alpha} A \Rightarrow B}{p \Vdash_{\alpha} A \Rightarrow C} \text{ if } B \text{ classically entails } C.$$

$$\text{(And)} \frac{p \Vdash_{\alpha} A \Rightarrow B \quad p \Vdash_{\alpha} A \Rightarrow C}{p \Vdash_{\alpha} A \Rightarrow (B \wedge C)}$$

$$\text{(Or)} \frac{p \Vdash_{\alpha} A \Rightarrow C \quad p \Vdash_{\alpha} B \Rightarrow C}{p \Vdash_{\alpha} (A \vee B) \Rightarrow C}$$

$$\text{(Cautious Monotonicity)} \frac{p \Vdash_{\alpha} A \Rightarrow B \quad p \Vdash_{\alpha} A \Rightarrow C}{p \Vdash_{\alpha} (A \wedge B) \Rightarrow C}$$

Say that acceptance rule α *validates* an axiom for conditional logic if and only if, for each credal states p , α together with p satisfies that axiom. Say that α *validates* a set of axioms if and only if α validates each axiom in that set. Some axioms in Adams' logic are validated trivially.

Proposition 7 *Each acceptance rule validates And Left Equivalence, and Right Weakening.*

Proof Immediate from the modeling assumption: with respect to each credal state and each acceptance rule, there is a strongest accepted proposition that entails all the other accepted propositions. \square

Proposition 8 *Let α be an acceptance rule. Then, α validates Reflexivity if and only if α accepts every certain proposition in the following sense: $p \Vdash_{\alpha} A$ for each credal state p in \mathcal{P} and each proposition A in \mathcal{A} such that $p(A) = 1$.*

Proof For the only if side, suppose that α validates Reflexivity. Suppose further that $p(A) = 1$. Then we have that $p \Vdash_{\alpha} A \Rightarrow A$ (by Reflexivity), and thus that $p(\cdot|A) \Vdash_{\alpha} A$ (note that $p(\cdot|A)$ exists), and hence that $p \Vdash_{\alpha} A$ (because $p = p(\cdot|A)$). So α accepts every certain proposition. For the converse, let α be an acceptance rule and p a credal state. Either credal state $p(\cdot|A)$ is undefined, and thus we have that $p \Vdash_{\alpha} A \Rightarrow A$ by default. Or $p(\cdot|A)$ is defined, and thus $p(A|A) = 1$ and then $p(\cdot|A) \Vdash_{\alpha} A$ (by acceptance of every certain proposition) and hence we have that $p \Vdash_{\alpha} A \Rightarrow A$ (by definition). \square

Axioms Cautious Monotonicity and Or impose substantial geometrical constraints on acceptance rules. Let A be a proposition in \mathcal{A} . Let $\mathcal{P}|A$ denote the set of all p in \mathcal{P} such that $p(A) = 1$, which we will call the *facet* of simplex \mathcal{P} for proposition A . The *line segment* with endpoints p, q in simplex \mathcal{P} is defined by convex combination:¹⁵

$$\overline{pq} = \{ap + (1 - a)q : a \in [0, 1]\}.$$

Say that q is a *projection* of p from facet $\mathcal{P}|\neg A$ onto facet $\mathcal{P}|A$ if and only if (i) there exists a line segment L through p with endpoint q in $\mathcal{P}|A$ and the other endpoint in facet $\mathcal{P}|\neg A$ and (ii) p is not in the complementary facet $\mathcal{P}|\neg A$. Projection is equivalent to Bayesian conditioning:

Lemma 3 *Credal state q is a projection of p from facet $\mathcal{P}|\neg A$ onto facet $\mathcal{P}|A$ if and only if $p(\cdot|A)$ is defined and $q = p(\cdot|A)$.*

Proof This lemma is trivially true when p is in $\mathcal{P}|A$ or in $\mathcal{P}|\neg A$, so suppose that p is neither in $\mathcal{P}|A$ nor in $\mathcal{P}|\neg A$ and, thus, that both $p(\cdot|A)$ and $p(\cdot|\neg A)$ are defined. For the if side, consider line segment $L = \overline{p(\cdot|A) p(\cdot|\neg A)}$, whose endpoints are in $\mathcal{P}|A$ and $\mathcal{P}|\neg A$, respectively. Note that p lies on L , since for each B in \mathcal{A} ,

$$p(B) = p(B|A)p(A) + p(B|\neg A)p(\neg A) = a p(B|A) + (1 - a) p(B|\neg A),$$

where $a = p(A)$. Therefore, $p(\cdot|A)$ is a projection of p from $\mathcal{P}|\neg A$ onto $\mathcal{P}|A$. For the only if side, suppose that q is a projection of p from facet $\mathcal{P}|\neg A$ onto facet $\mathcal{P}|A$. So q is in $\mathcal{P}|A$ and there exists credal state r in $\mathcal{P}|\neg A$ such that line segment \overline{qr} contains p . Then, p lies in the interior of \overline{qr} , since p is neither in $\mathcal{P}|A$ nor in $\mathcal{P}|\neg A$. So there exists a in the open interval $(0, 1)$ such that $p = aq + (1 - a)r$. Then it suffices to show that $q = p(\cdot|A)$. Consider the case in which $E_i \not\leq A$. Then $E_i \leq \neg A$. Since q is in facet $\mathcal{P}|A$, we have that $q(E_i) = 0 = p(E_i|A)$. Now consider the case in which $E_i \leq A$. Then since r is in facet $\mathcal{P}|\neg A$, we have that $r(E_i) = 0$, so $p(E_i) = aq(E_i)$. Similarly, we have that $q(A) = 1$ and $r(A) = 0$, so $p(A) = a \cdot 1 + 0 = a$. Hence, $q(E_i) = p(E_i)/a = p(E_i)/p(A) = p(E_i)p(A|E_i)/p(A) = p(E_i|A)$. So $q(\cdot)$ agrees with $p(\cdot|A)$ for all E_i in \mathcal{E} and, thus, for all B in \mathcal{A} , as required. \square

Proposition 9 (geometry of Cautious Monotonicity) *Let α be an acceptance rule. Then, α validates Cautious Monotonicity if and only if the following condition holds: for each credal state p and for each proposition A , if α accepts A at p , then α accepts A at the projection of p on the facet $\mathcal{P}|B$, for each logical consequence B of A (as long as the projection exists). In light of Lemma 3, the condition may be restated as:*

$$p \Vdash_\alpha A, \quad A \leq B, \quad \text{and} \quad p(\cdot|B) \text{ is defined} \implies p(\cdot|B) \Vdash_\alpha A. \quad (23)$$

Proof The proof of the only if side involves unpacking the definitions and checking that the projection condition (23) is simply an instance of Cautious Monotonicity. For the if side, assume that the projection condition (23) holds. Suppose that $p \Vdash_\alpha A \implies B$

¹⁵ Addition is defined as vector addition; multiplication is defined as scalar multiplication.

and $p \Vdash_{\alpha} A \Rightarrow C$. It suffices to show that $p \Vdash_{\alpha} (A \wedge B) \Rightarrow C$. If $p(\cdot|A \wedge B)$ is undefined, then by default $p \Vdash_{\alpha} (A \wedge B) \Rightarrow C$. So suppose that $p(\cdot|A \wedge B)$ is defined and, thus, $p(\cdot|A)$ is defined. Then argue as follows:

$$\begin{aligned}
 & p \Vdash_{\alpha} A \Rightarrow B, \quad p \Vdash_{\alpha} A \Rightarrow C \\
 \implies & q \Vdash_{\alpha} B, \quad q \Vdash_{\alpha} C && \text{letting } q = p(\cdot|A), \\
 \implies & q \Vdash_{\alpha} B \wedge C, \quad (B \wedge C) \leq B \\
 \implies & q(\cdot|B) \Vdash_{\alpha} B \wedge C && \text{by condition (23) and the existence of } q(\cdot|B) \\
 & && \text{which equals } p(\cdot|A \wedge B), \\
 \implies & p(\cdot|A \wedge B) \Vdash_{\alpha} B \wedge C && \text{since } p(\cdot|A \wedge B) = q(\cdot|B), \\
 \implies & p(\cdot|A \wedge B) \Vdash_{\alpha} C \\
 \implies & p \Vdash_{\alpha} (A \wedge B) \Rightarrow C. && \square
 \end{aligned}$$

Proposition 10 (geometry of Or) *Let α be an acceptance rule that validates Reflexivity. Then, α validates Or if and only if the following condition holds: for each line segment L connecting two complementary facets $\mathcal{P}|B$ and $\mathcal{P}|\neg B$, and for each proposition A in \mathcal{A} , if α accepts A at both endpoints of L , then α accepts A at each point on L ; in light of Lemma 3, the condition may be restated as:*

$$p(\cdot|B) \Vdash_{\alpha} A, \quad p(\cdot|\neg B) \Vdash_{\alpha} A \implies p \Vdash_{\alpha} A. \tag{24}$$

Proof For the only if side, argue as follows:

$$\begin{aligned}
 & p(\cdot|B) \Vdash_{\alpha} A, \quad p(\cdot|\neg B) \Vdash_{\alpha} B \\
 \implies & p \Vdash_{\alpha} B \Rightarrow A, \quad p \Vdash_{\alpha} \neg B \Rightarrow A \\
 \implies & p \Vdash_{\alpha} (B \vee \neg B) \Rightarrow A && \text{by axiom Or,} \\
 \implies & p(\cdot|B \vee \neg B) \Vdash_{\alpha} A \\
 \implies & p \Vdash_{\alpha} A.
 \end{aligned}$$

For the converse, suppose that $p \Vdash_{\alpha} A \Rightarrow C$ and $p \Vdash_{\alpha} B \Rightarrow C$. It suffices to show that $p \Vdash_{\alpha} (A \vee B) \Rightarrow C$. If both $p(\cdot|A)$ and $p(\cdot|B)$ are undefined, then $p(\cdot|A \vee B)$ is undefined and thus we have that $p \Vdash_{\alpha} (A \vee B) \Rightarrow C$ by default. If one is defined and the other is undefined—say, $p(\cdot|A)$ is defined and $p(\cdot|B)$ is undefined—then $p(B) = 0$ and thus $p(\cdot|A \vee B) = p(\cdot|A)$ is defined, so:

$$\begin{aligned}
 & p \Vdash_{\alpha} A \Rightarrow C \\
 \implies & p(\cdot|A) \Vdash_{\alpha} C \\
 \implies & p(\cdot|A \vee B) \Vdash_{\alpha} C && \text{by } p(\cdot|A \vee B) = p(\cdot|A), \\
 \implies & p \Vdash_{\alpha} (A \vee B) \Rightarrow C.
 \end{aligned}$$

Last, suppose that both $p(\cdot|A)$ and $p(\cdot|B)$ are defined. So $p(\cdot|A \vee B)$ is defined. Then argue for Or as follows:

$$\begin{aligned}
 & p \Vdash_{\alpha} A \Rightarrow C, \quad p \Vdash_{\alpha} B \Rightarrow C \\
 \implies & p(\cdot|A) \Vdash_{\alpha} C, \quad p(\cdot|B) \Vdash_{\alpha} C \\
 \implies & q(\cdot|A) \Vdash_{\alpha} C, \quad q(\cdot|B) \Vdash_{\alpha} C && \text{letting } q = p(\cdot|A \vee B), \\
 & && \text{so } q(\cdot|A) = p(\cdot|A) \text{ and } q(\cdot|B) = p(\cdot|B),
 \end{aligned}$$

$\implies q \Vdash_{\alpha} C \vee \neg A, q \Vdash_{\alpha} C \vee \neg B$ (*) see the explanation below,
 $\implies q \Vdash_{\alpha} C \vee \neg(A \vee B)$ by classical entailment,
 $\implies q \Vdash_{\alpha} C \vee \neg(A \vee B),$ since $q(A \vee B) = 1$ and Proposition 8 applies,
 $q \Vdash_{\alpha} A \vee B$
 $\implies q \Vdash_{\alpha} C$ by classical entailment,
 $\implies p(\cdot|A \vee B) \Vdash_{\alpha} C$
 $\implies p \Vdash_{\alpha} (A \vee B) \Rightarrow C.$

It only remains to establish step (*). By the symmetric roles of A and B , it suffices to show that $q(\cdot|A) \Vdash_{\alpha} C$ implies that $q \Vdash_{\alpha} C \vee \neg A$. If $q(\cdot|\neg A)$ is undefined, then $q(A) = 1 - q(\neg A) = 1 - 0 = 1$ and thus $q = q(\cdot|A) \Vdash_{\alpha} C \leq C \vee \neg A$, so $q \Vdash_{\alpha} C \vee \neg A$. If $q(\cdot|\neg A)$ is defined, then we have both that $q(\cdot|A) \Vdash_{\alpha} C$ (by supposition) and that $q(\cdot|\neg A) \Vdash_{\alpha} \neg A$ (by Reflexivity and Proposition 8). So we have both that $q(\cdot|A) \Vdash_{\alpha} C \vee \neg A$ and that $q(\cdot|\neg A) \Vdash_{\alpha} C \vee \neg A$ (by classical entailment). Hence $q \Vdash_{\alpha} C \vee \neg A$, by the convexity condition (24). \square

10 The geometry of system P

In this section we examine the geometric constraints imposed by the entire system P. It is an easy corollary of the geometrical characterizations in the preceding section that:

Theorem 2 (Lin 2011) *Each probalogical rule validates system P.*

Proof sketch When $|\mathcal{E}| = 3$, one can verify by a picture that probalogical rules satisfy the geometric conditions given in Propositions 7–10. The routine verification can be easily generalized to a proof for all countable dimensional cases. \square

We now proceed to establish a partial converse to Theorem 2. Recall that acceptance zones for answers have the following form under probalogical rules:¹⁶

$$\begin{aligned}
 p \Vdash_{\nu} E_i &\iff E_i = \nu(p) \\
 &\iff E_i = \bigwedge \{ \neg E_j : \sigma(p)_j \triangleleft_j 1 - r_j \} \\
 &\iff \forall j \neq i, \sigma(p)_j \triangleleft_j 1 - r_j \\
 &\iff \forall j \neq i, \frac{p_j}{\max_k p_k} \triangleleft_j 1 - r_j \\
 &\iff \forall j \neq i, \frac{p_j}{p_i} \triangleleft_j 1 - r_j.
 \end{aligned}$$

Namely, answer E_i is accepted if and only if each rival has a sufficiently low odds to E_i . To allow for more generalized rules entertained below (Sect. 15), we relax the conditions that the rejection threshold $1 - r_j$ is in the unit interval and that it is constant for all i . Accordingly, say that the acceptance zone of answer E_i under α is a *blunt*

¹⁶ The last step follows because the condition $p_j / \max_k p_k \triangleleft_j 1 - r_j$ implies that $p_j / \max_k p_k < 1$, for each $j \neq i$. Therefore, $\max_k p_k = p_i$.




Fig. 19 Acceptance zone of E_2

diamond (Fig. 19a) if and only if it takes the following form: there exist thresholds $\{t_{ij} : j \in I \setminus \{i\}\}$ in interval $[0, \infty]$ and inequalities $\{\triangleleft_{ij} : j \in I \setminus \{i\}\}$ that are either \leq or $<$, such that for each $p \in \mathcal{P}$:

1. $p \Vdash_{\alpha} E_i \iff \forall j \neq i, \frac{p_j}{p_i} \triangleleft_{ij} t_{ij}$;
2. if $\triangleleft_{ij} = \leq$ then $t_{ij} < \infty$;
3. if $\triangleleft_{ij} = <$ then $t_{ij} > 0$.

Say that acceptance rule α is *corner-monotone* if and only if (i) $\alpha(e_i) = E_i$ for each $i \in I$, and (ii) for each $p \in \mathcal{P}$ such that $\alpha(p) = E_i$, we have that $\alpha(q) = E_i$ for all q in line segment $\overline{p e_i}$. Corner-monotonicity is a very natural constraint on acceptance rules and it is satisfied by all the rules we have discussed. Our partial converse to Theorem 2 is as follows.

Theorem 3 (blunt diamond, Lin 2011) *Let α be an acceptance rule. If α is everywhere consistent, satisfies corner-monotonicity, and validates system \mathbf{P} , then for each answer E_i to question \mathcal{E} , the acceptance zone of E_i under α is a blunt diamond.*

Proof sketch Here we present a geometric argument for case $|\mathcal{E}| = 3$, which can be easily generalized to each countable dimension. Solve for the acceptance zone of E_2 under α , as depicted in Fig. 19b. By corner-monotonicity, the credal states $\overline{e_2 e_1}$ where α accepts E_2 form a continuous, unbroken line segment with e_2 as an endpoint, which is depicted as the heavy, grey line segment lying on $\overline{e_2 e_1}$. The same is true for side $\overline{e_2 e_3}$.¹⁷ Connect the endpoints of the grey line segments to the opposite corners by straight lines, which enclose the grey blunt diamond at the corner e_2 .

Argue as follows that $p \Vdash_{\alpha} E_2$, for each point p in the blunt diamond. Consider the projection p' of p to the facet $\mathcal{P}(E_2 \vee E_3)$. Note that p' is in the heavy, grey line segment along side $\overline{e_2 e_3}$. On line segment $e_1 p'$, acceptance rule α accepts E_1 at one endpoint (e_1) and accepts E_2 at the other endpoint (p'), so α accepts $E_1 \vee E_2$ at both

¹⁷ There is an issue whether the line segments are open or closed at the endpoints distinct from e_2 , which would give rise to a possible mixture of strict and weak inequalities, as stated in the theorem. That detail is handled in the formal proof in Lin (2011), but is ignored here.

endpoints. Then, by Proposition 10, we have that $p \Vdash_{\alpha} E_1 \vee E_2$. By applying the same argument to the projection of p to the facet for proposition $E_1 \vee E_3$, we have that $p \Vdash_{\alpha} E_3 \vee E_2$. Then $p \Vdash_{\alpha} E_2$, since E_2 is entailed by $E_1 \vee E_2$ plus $E_2 \vee E_3$.

Argue as follows that $q \not\Vdash_{\alpha} E_2$, for each point q outside of the blunt diamond. Since q lies outside of the blunt diamond, there exists at least one answer E_i other than E_2 such that the projection q' of q to the facet $\mathcal{P}(E_2 \vee E_i)$ does not touch the grey line segment along side $\overline{e_2 e_i}$. Suppose for reductio that $q \Vdash_{\alpha} E_2$. Then, by applying Proposition 9 to the projection q' of q , we have that $q' \Vdash_{\alpha} E_2$. But $q' \not\Vdash_{\alpha} E_2$, for q' lies outside of the grey line segment—contradiction. \square

11 AGM geometry is trivial

A popular, stronger system for the logic of flat conditionals, **R**, is obtained from **P** by adding the following axiom (Lehmann and Magidor 1992):

$$\text{(Rational Monotonicity)} \frac{p \not\Vdash_{\alpha} A \Rightarrow \neg B \quad p \Vdash_{\alpha} A \Rightarrow C}{p \Vdash_{\alpha} (A \wedge B) \Rightarrow C}$$

Recall the *probabilistic* Ramsey test assumed in the preceding sections of this paper:

$$p \Vdash_{\alpha} A \Rightarrow B \iff \alpha(p(\cdot|A)) \leq B \text{ or } p(A) = 0.$$

Given this test, validation of system **R** trivializes uncertain acceptance in the sense defined as follows. Say that acceptance rule α is *skeptical* if and only if there is some answer to \mathcal{E} that is accepted by α over no open subset of \mathcal{P} .¹⁸ Say that α is *opinionated* if and only if for each disjunction D of at least two distinct answers to question \mathcal{E} , there is no open subset of \mathcal{P} over which α accepts D as strongest. Finally, **B** is *trivial* if and only if α is either skeptical or opinionated.

Theorem 4 (skepticism or opinionation) *Let question \mathcal{E} has cardinality ≥ 3 . Suppose that acceptance rule α is everywhere consistent, corner-monotone, and validates system **R**. Then α is trivial.*

Since the probabilistic Ramsey test is based on probabilistic conditioning, acceptance rules must respect the geometry of conditioning in order to validate axioms of nonmonotonic reasoning. What Theorem 4 says is that these geometrical constraints become hopelessly severe when one adds rational monotonicity to system **P**. Of course, the situation is quite different if one drops probabilistic conditioning from the Ramsey test.¹⁹ A *conditional* acceptance rule is a mapping $\beta : \mathcal{P} \times \mathcal{A} \rightarrow \mathcal{A}$, where $\beta(p|A) = B$ is interpreted as saying that B is the strongest proposition accepted in

¹⁸ This definition of skepticism is a bit stricter than the one introduced earlier, since non-skeptical methods must accept each answer over some neighborhood of credal states in which it is uncertain, rather than at a single such credal state.

¹⁹ The approach that follows is due to Hannes Leitgeb, who presented his unpublished results at the Opening Celebration of the Center for Formal Epistemology at Carnegie Mellon University in the Summer of




Fig. 20 Two paths

p in light of new information A . Then one can state a new, *non-probabilistic* Ramsey test directly in terms of conditional acceptance:

$$p \Vdash_{\beta} A \Rightarrow B \iff \beta(p|A) \leq B. \tag{25}$$

Such a conditional acceptance rule is an abstract concept that can be filled out in various different ways. For example, say that conditional acceptance rule β is *Bayesian* if and only if there exists a (non-conditional) acceptance rule α such that:

$$\beta(p|A) = \begin{cases} \alpha(p(\cdot|A)) & \text{if } p(A) > 0; \\ \perp & \text{otherwise.} \end{cases} \tag{26}$$

When β is Bayesian, the new information A is used to condition the credal state p to obtain $p(\cdot|A)$ and then some new propositional belief state S' is accepted in light of $p(\cdot|A)$ (the upper path in Fig. 20). If β is Bayesian, then the non-probabilistic Ramsey test for β is equivalent to the probabilistic Ramsey test for α , so Theorem 4 still applies to β . But β need not be Bayesian. For example, β may sidestep Bayesian conditioning entirely by using α to accept a propositional belief state $S = \alpha(p)$ in p and by subsequently applying a propositional belief revision operator $*_p$ (that may depend on p) to convert $\alpha(p)$ into a new propositional belief state $S' = \alpha(p) *_p A$ (the lower path in Fig. 20):

$$\beta(p|A) = \alpha(p) *_p A. \tag{27}$$

In that case, the validation of system **R** depends entirely on the propositional revision operator $*_p$ —probabilistic conditioning and α are both irrelevant, so the geometrical proof of Theorem 4 is also sidestepped. To validate Rational Monotonicity, it is sufficient to require that each $*_p$ be an *AGM belief revision operator* (Harper 1975;

Footnote 19 continued
 2010. The discussion in this section is based on detailed slides he presented at that meeting and on personal communication with him at that time.

Alchourròn et al. 1985), thanks to the translation between nonmonotonic logic and belief revision due to Makinson and Gärdenfors (1991).²⁰

The escape route just described does not really vindicate or explain Rational Monotonicity from a Bayesian perspective, since Bayesian conditioning is bypassed and Rational Monotonicity is simply imposed on the propositional belief revision operator $*_p$. On the other hand, it is an immediate corollary of Theorem 2 that the Bayesian rules of form:

$$\beta(p|A) = v(p(\cdot|A)) \tag{28}$$

all validate system **P** with respect to the non-probabilistic Ramsey test. We propose, therefore, that system **P** reflects Bayesian ideals better than system **R**.

The proof of Theorem 4 proceeds by establishing a slightly stronger result. The following two properties of an acceptance rule are derivable from validation of system **R**: Say that α satisfies *preservation* if and only if, for each credal state p and each proposition E , if (new information) E is consistent with (the prior belief state) $\alpha(p)$, then (the posterior belief state) $\alpha(p(\cdot|E))$ entails $\alpha(p) \wedge E$. Say that α satisfies *inclusion* if and only if, for each credal state p and each proposition E , $\alpha(p(\cdot|E))$ is entailed by $\alpha(p) \wedge E$.²¹ So, to prove Theorem 4, it suffices to prove the following theorem:

Theorem 5 *Let question \mathcal{E} has cardinality ≥ 3 . Suppose that acceptance rule α is everywhere consistent, corner-monotone, and satisfies preservation and inclusion. Then α is trivial.*

The proof of Theorem 5 proceeds by a sequence of lemmas and occupies the balance of this section. Suppose that rule α is everywhere consistent, corner-monotone, and satisfies preservation and inclusion. Suppose further that α is not skeptical. It suffices to show that α is opinionated. Let E_i, E_j be distinct answers to \mathcal{E} . Choose an arbitrary, third answer E_m to \mathcal{E} (since \mathcal{E} is assumed to have at least three answers). Let $\Delta e_i e_j e_m$ denote the two dimensional facet $\mathcal{P}|(E_i \vee E_j \vee E_m)$ (Fig. 21a). Let $\overline{e_i e_m}$ denote the one-dimensional facet $\mathcal{P}|(E_i \vee E_m)$, and similarly for $\overline{e_i e_j}$ and $\overline{e_j e_m}$. Let L_{im} be the set of the credal states on line segment $\overline{e_i e_m}$ at which E_i is accepted by α as strongest; namely:

$$L_{im} = \{p \in \overline{e_i e_m} : \alpha(p) = E_i\}.$$

Lemma 4 *L_{im} is a connected line segment of nonzero length that contains e_i but does not contain e_m .*

²⁰ It is not necessary, though, because to validate system **R** one does not have to require that $*_p$ satisfies the consistency axiom in AGM—but all the other axioms have to be satisfied. We thank David Etlin for pointing this out.

²¹ Inclusion is derivable from system **P** alone. Preservation is derivable from a special case Rational Monotonicity alone in which A is replaced by the tautology \top . They are so named because of their correspondence to the AGM axioms K^*3 and K^*4 .




Fig. 21 Why system R is trivial

Proof By non-skepticism, there exists open subset O of \mathcal{P} over which α accepts E_i as strongest. Let O' be the image $\{o|_{E_i \vee E_m} : o \in O\}$ of O under conditioning on $E_i \vee E_m$. Since O is open, O' is an open subset of $\overline{e_i e_m}$. Note that the conditioning proposition $E_i \vee E_m$ is consistent with the prior belief state E_i , so preservation applies. Since α satisfies preservation, α accepts old belief E_i over O' . It follows that α accepts E_i as strongest over O' , because α is consistent and the only proposition strictly stronger than E_i in the algebra is the inconsistent proposition \perp . So open set O' is included in L_{im} , and thus L_{im} has nonzero length. Then, since α is corner-monotone, L_{im} is a connected line segment that contains e_i . It remains to show that L_{im} does not contain e_m . Suppose for reductio that L_{im} contains e_m . Then L_{im} must be so large that it is identical to $\overline{e_i e_m}$, by corner-monotonicity. By the same argument for showing that there is an open subset O' of $\overline{e_i e_m}$ over which α accepts E_i , we have that there is an open subset O'' of $\overline{e_i e_m}$ over which α accepts E_m . So α accepts both E_m and E_i over O'' , and hence by closure under conjunction, α accepts their conjunction, which is an inconsistent proposition. So α is not consistent—contradiction. \square

Let a be the endpoint of L_{im} that is closest to e_m ; namely, probability measure a is such that:

$$a \in \overline{e_i e_m},$$

$$a(E_m) = \sup\{p(E_m) : p \in L_{im}\}.$$

By the lemma we just proved, point a lies in the interior of side $\overline{e_i e_m}$. Applying the above argument for pair (i, m) to pair (j, m) , we have that the set L_{jm} , defined by

$$L_{jm} = \{p \in \overline{e_j e_m} : \alpha(p) = E_j\},$$

is a connected line segment of nonzero length that contains e_j but does not contain e_m , with endpoint b that lies in the interior of side $\overline{e_j e_m}$. Since both points a, b lie in the interiors of their respective sides, we have the following constructions. Let A be the line that connects a to e_j , B be the line that connects b to e_i , and C be the line that connects e_m through the intersection d of A and B , to point c on side $\overline{e_i e_j}$.

Lemma 5 α accepts E_i as strongest over the interior of Δade_i .

Proof Consider an arbitrary point p in the interior of Δade_i (Fig. 21b). Argue as follows that α accepts $E_i \vee E_j$ at p . Take p as a prior state and consider $\neg E_j$ as the conditioning information. Note that credal state $p(\cdot|\neg E_j)$ falls inside L_{im} , so α accepts E_i as strongest at the posterior credal state $p(\cdot|\neg E_j)$. Then, since α satisfies inclusion, we have that:

$$\alpha(p) \wedge \neg E_j \models E_i$$

(namely the posterior belief state E_i is entailed by the conjunction of the prior belief state and the conditioning information). Then, by the consistency of α and the mutual exclusion among the answers, we have only three possibilities for $\alpha(p)$:

$$\alpha(p) \text{ is either } E_i, \text{ or } E_j, \text{ or } E_i \vee E_j.$$

Rule out the last two alternatives as follows. Suppose for reductio that the prior belief state $\alpha(p)$ is E_j or $E_i \vee E_j$. Consider $\neg E_i$ as the conditioning information, which is consistent with the prior belief state and thus makes preservation applicable. Then, since α satisfies preservation, the posterior belief state $\alpha(p(\cdot|\neg E_i))$ must entail $\alpha(p) \wedge \neg E_i$ (i.e. the conjunction of the prior belief state and the information). But the latter proposition $\alpha(p) \wedge \neg E_i$ equals E_j , by the reductio hypothesis. So $\alpha(p(\cdot|\neg E_i)) = E_j$, by the consistency of α . Hence $p(\cdot|\neg E_i)$ lies on line segment L_{jm} by the construction of L_{jm} —but that is impossible (Fig. 21b). Ruling out the last two alternatives for $\alpha(p)$, we conclude that $\alpha(p) = E_i$. \square

Lemma 6 α accepts E_i as strongest over the interior of $\overline{e_i c}$.

Proof Let p be an arbitrary interior point of Δade_i . So $\alpha(p) = E_i$. Consider proposition $E_i \vee E_j$ as the conditioning information. Then, since α satisfies preservation, the posterior belief state $\alpha(p(\cdot|E_i \vee E_j))$ entails $\alpha(p) \wedge (E_i \vee E_j)$ (i.e. the conjunction of the prior belief state and the information), which equals E_i . Then, by consistency, the posterior belief state is determined:

$$\alpha(p(\cdot|E_i \vee E_j)) = E_i.$$

Let q be an arbitrary point in the interior of $\overline{e_i c}$. Then q can be expressed as $q = p(\cdot|E_i \vee E_j)$ for some point p in the interior of Δade_i (Fig. 21c). So, by the formula we just proved, $\alpha(q) = \alpha(p(\cdot|E_i \vee E_j)) = E_i$, as required. \square

Lemma 7 There is no open subset of $\overline{e_i e_j}$ over which α accepts $E_i \vee E_j$ as strongest.

Proof We have established in the last lemma that α accepts E_i as strongest over the interior of $\overline{e_i c}$. By the same argument, α accepts E_j as strongest over the interior of $\overline{e_j c}$ (Fig. 21c). So if α accepts disjunction $E_i \vee E_j$ as strongest somewhere on $\overline{e_i e_j}$, α does so at some of the three points: e_i , e_j , and c . (We can rule out the first two alternatives; but the for the sake of the lemma, this result suffices.) \square

Since the choice of E_i and E_j is arbitrary, the last lemma generalizes to the following:

Lemma 8 *For each pair of distinct answers E_i, E_j to \mathcal{E} , there is no open subset of $\overline{e_i e_j}$ over which α accepts $E_i \vee E_j$ as strongest.*

The last lemma establishes opinionation for all edges of the simplex. The next step is to extend opinionation to the whole simplex.

Lemma 9 *α is opinionated.*

Proof Suppose for reductio that some disjunction $E_i \vee E_j \vee X$ of at least two distinct answers is accepted by α as strongest over some open subset O of \mathcal{P} . Take $E_i \vee E_j \vee X$ as the prior belief state at each point in O and consider $E_i \vee E_j$ as the conditioning information. So the image O' of O under conditioning on $E_i \vee E_j$ is an open subset of 1-dimensional space $\overline{e_i e_j}$. Let p' be an arbitrary point in O' . Since α satisfies inclusion, posterior belief state $\alpha(p')$ is entailed by $(E_i \vee E_j \vee X) \wedge (E_i \vee E_j)$ (i.e. the conjunction of the prior state and the new information), which equals $E_i \vee E_j$. But $\alpha(p')$ also entails $E_i \vee E_j$, for otherwise the process of conditioning p' on $\neg E_j$ to obtain e_i would violate the fact that α satisfies inclusion and accepts E_i at e_i . So $\alpha(p') = E_i \vee E_j$. Hence α accepts $E_i \vee E_j$ as strongest over open subset O' of $\overline{e_i e_j}$, which contradicts the last lemma. \square

Proof of Theorem 5 Since the last lemma states that α is opinionated, we are done. \square

Proof of Theorem 4 Immediate from Theorem 5. \square

12 A new probabilistic semantics for flat conditionals

Axiom system \mathbf{P} is characteristic of Adams' logic of flat conditionals, so it is not surprising that the probalogical rules yield a new probabilistic semantics for which Adams' logic is sound. In fact, Adams' logic is both sound and complete for the new semantics.

Let \mathcal{L} be a set of sentences that contains atomic propositional letters and is closed under conjunction, disjunction, and negation. Let \Rightarrow be a sentential connective standing for "if ... then ...". The language for the logic of flat conditionals, written $\mathcal{L}_{\Rightarrow}$, is the set of all sentences $\phi \Rightarrow \psi$ with $\phi, \psi \in \mathcal{L}$. Adams (1975) logic of flat conditionals for language $\mathcal{L}_{\Rightarrow}$ is just the system \mathbf{P} that we have stated, except that now it is construed as a system of rules of inference (with the symbol " $p \Vdash_{\alpha}$ " deleted). Say that γ is derivable from Γ in Adams' logic of flat conditionals, written $\Gamma \vdash_{\text{Adams}} \gamma$, if and only if γ is derivable from Γ in a finite number of steps using the rules of inference in system \mathbf{P} .

A probabilistic model of acceptance for language $\mathcal{L}_{\Rightarrow}$ is a triple:

$$M = (\alpha, p, \llbracket \cdot \rrbracket),$$

where $\alpha : \mathcal{P} \rightarrow \mathcal{A}$ is an acceptance rule, p is a probability measure in the domain \mathcal{P} of α , and $\llbracket \cdot \rrbracket$ is a classical interpretation of \mathcal{L} to the codomain \mathcal{A} of α . When

$M = (\alpha, p, \llbracket \cdot \rrbracket)$, say that α is the *underlying* acceptance rule of M . Let $\phi \Rightarrow \psi$ be a flat conditional in $\mathcal{L}_{\Rightarrow}$. *Acceptance* of flat conditional $\phi \Rightarrow \psi$ in model $M = (\alpha, p, \llbracket \cdot \rrbracket)$, written $M \Vdash \phi \Rightarrow \psi$, is defined by the probabilistic Ramsey test:

$$\begin{aligned} M \Vdash \phi \Rightarrow \psi &\iff p \Vdash_{\alpha} \llbracket \phi \rrbracket \Rightarrow \llbracket \psi \rrbracket, \\ &\iff p(\cdot | \llbracket \phi \rrbracket) \Vdash_{\alpha} \llbracket \psi \rrbracket \text{ or } p(\llbracket \phi \rrbracket) = 0. \end{aligned}$$

Let Γ be a set of flat conditionals in $\mathcal{L}_{\Rightarrow}$. *Acceptance* of Γ in model M is defined by: $M \Vdash \Gamma$ if and only if $M \Vdash \gamma$ for all $\gamma \in \Gamma$. Validity is defined straightforwardly, as preservation of acceptance. Let \mathcal{C} be a class of acceptance rules. Say that \mathcal{C} *validates* the inference from Γ to γ , written $\Gamma \Vdash_{\mathcal{C}} \gamma$, if and only if for each probabilistic model M whose underlying acceptance rule is in \mathcal{C} , if $M \Vdash \Gamma$, then $M \Vdash \gamma$.

The proposed probabilistic semantics has the following attractive properties: (i) it is based on the probabilistic Ramsey test for accepting conditionals; (ii) it defines validity simply as preservation of acceptance, which improves upon Adams’ (1975) ϵ – δ semantics; and (iii) it allows for accepting propositions of probabilities significantly less than 1, which improves upon Pearl’s (1989) infinitesimal semantics. To establish the soundness and completeness result for Adams’ logic of flat conditionals, it suffices to assume that the underlying acceptance rule is probalogical, or equivalently, a camera shutter rule:

Theorem 6 (soundness and completeness, Lin 2011) *Let \mathcal{N} be the class of the camera shutter rules. Then, for each finite sentence set Γ and each sentence γ in the language $\mathcal{L}_{\Rightarrow}$ of flat conditionals, $\Gamma \Vdash_{\text{Adams}} \gamma$ if and only if $\Gamma \Vdash_{\mathcal{N}} \gamma$.*

13 Question-invariance

To this point, we have considered acceptance only within a fixed question \mathcal{E} . But one can and should consider the behavior of acceptance rules *across* questions. Let Ω denote some infinite collection of possibilities. A *question* $\mathcal{E} = \{E_i : i \in I\}$ is a countable partition of Ω such that each answer/cell E_i is infinite—the requirement of infinite answers rules out the artificial possibility of a maximally informative question whose answers cannot be strengthened. Let $\mathcal{A}_{\mathcal{E}}$ denote the least collection of propositions containing \mathcal{E} and closed under negation and countable disjunction and conjunction. Let \mathbb{E} denote the set of all such questions over Ω , and let \mathbb{P} denote the set of all countably additive probability measures p such that p is defined on $\mathcal{A}_{\mathcal{E}}$ for some question \mathcal{E} in \mathbb{E} . If p is in \mathbb{P} , let \mathcal{A}_p denote the domain of p and let \mathcal{E}_p denote the (unique) question that generates \mathcal{A}_p . A (*cross-question*) *acceptance rule* is a map β defined on \mathbb{P} such that β always maps p to a proposition in \mathcal{A}_p . Then the rules discussed earlier in this paper can be defined explicitly across questions as follows, where I_p is the index set of the question over which p is defined:

$$\begin{aligned} \lambda_r(p) &= \bigwedge \{ \neg E_i : p_i \leq 1 - r \text{ and } i \in I_p \}; \\ \lambda(p) &= \lambda_{s(p)}(p), \quad \text{where } s(p) = 1 - \frac{1}{2|I_p|}; \\ \nu_r(p) &= \bigwedge \{ \neg E : \sigma(p)_i <_i 1 - r_i \text{ and } i \in I_p \}; \\ \pi_r(p) &= \begin{cases} \top & \text{if } \lambda_r(p) = \perp; \\ \lambda_r(p) & \text{otherwise.} \end{cases} \end{aligned}$$

Rule λ_r is the Lockean rule with a fixed threshold across all questions in \mathbb{E} . Rule ν_r is the probalogical rule. Rule λ is the ad hoc Lockean rule whose threshold is adjusted to avoid lottery paradoxes in finite questions. Rule π_r is the Pollockian rule that substitutes \top for \perp whenever the latter is produced by λ_r .

Say that cross-question acceptance rule β is *question-invariant* if and only if:

$$p(A) = q(A) \implies (p \Vdash_\beta A \iff q \Vdash_\beta A),$$

for each p, q in \mathbb{P} and for each A that is in both \mathcal{A}_p and \mathcal{A}_q . Question-invariance is appealing. First, question-invariance makes it easier to compute whether to accept A in light of $p(A)$, since all of the detailed structure of p aside from $p(A)$ can be ignored. Second, question-invariance allows for the accumulation of accepted propositions as one’s question is refined by new concepts and theories. Third, question-invariance allows individual scientists pursuing distinct questions to *pool* their accepted conclusions. Probalogical rules, however, are not even remotely question-invariant. For example, in a four ticket lottery, the probalogical rule $\nu_{2/3}$ licenses acceptance of “ticket 1 will lose” when the question is “will ticket 1 lose or not?”, but not when the question is “which ticket will win?”. That makes one wonder whether the question-dependence of probalogical rules is a design defect that could have been avoided. We now proceed to demonstrate that *no* question-invariant rule has the three crucial virtues of the probalogical rules: consistency, logical closure, and non-skeptical acceptance of uncertain propositions.

Here is the first sign of trouble. Say that acceptance rule β is *non-skeptical* about answer E in question \mathcal{E} if and only if β accepts E at some probability measure p defined on $\mathcal{A}_\mathcal{E}$ such that $p(E) < 1$. Say that acceptance rule β is *gullible* about E in \mathcal{E} if and only if β accepts E at some p defined on $\mathcal{A}_\mathcal{E}$ such that $p(E) = 0$. Then:

Proposition 11 *Suppose that β is question-invariant. If β is non-skeptical about answer E in ternary question \mathcal{E} , then β is gullible about E in \mathcal{E} .*

Proof Consider the equilateral triangle $\Delta q u v$ depicted in Fig. 22a. Note that p lies on a line parallel to $\overline{e_2 e_3}$ extending the base $\overline{u v}$ of the triangle $\Delta q u v$ and q is at the apex. Suppose that $p \Vdash_\beta E_1$. Then $u, v \Vdash_\beta E_1$, by question-invariance. So $u \Vdash_\beta \neg E_2$ and $v \Vdash_\beta \neg E_3$. Then by question-invariance again, $q \Vdash_\beta \neg E_2$ and $q \Vdash_\beta \neg E_3$. So $q \Vdash_\beta \neg E_2 \wedge \neg E_3 = E_1$. Therefore, if β accepts E_1 at p , then β also accepts E_1 at q . Now we can chain such triangles all the way to the bottom of \mathcal{P}_3 to obtain s such that $s \Vdash E_1$ and $s(E_1) = 0$. Note that if $p(E_1) < 1$, there is room in \mathcal{P}_3 for such a chain. □




Fig. 22 Triangles preserve acceptance

It gets worse. Say that β is *dogmatic* about answer E in question \mathcal{E} if and only if β accepts E at each probability measure defined on $\mathcal{A}_{\mathcal{E}}$.

Proposition 12 *Suppose that β is question-invariant. If β is non-skeptical about answer E in ternary question \mathcal{E} , then β is dogmatic about E in \mathcal{E} .*

Proof Consider the situation depicted in Fig. 22b, in which question-invariant rule β accepts E_1 at s , with $s(E_1) = 0$, and let q be an arbitrary credal state in \mathcal{P}_3 . Then there exists an equilateral triangle with s on its base and with q at its apex, so β also accepts E_1 at the arbitrarily chosen state q . \square

Here is the *coup de grâce*. Say that β is *everywhere inconsistent* if and only if $\beta(p) = \perp$, for all p in \mathbb{P} . Nothing could be more useless than an acceptance rule that accepts the contradiction in every possible credal state and every possible question.

Theorem 7 *Suppose that β is question-invariant. If β is non-skeptical about at least two distinct answers in some ternary question, then β is everywhere inconsistent.*

Proof Suppose that β is non-skeptical about at least two distinct answers E_i, E_j in ternary question \mathcal{E} . Then, by Proposition 12, β accepts $E_i \wedge E_j$ and, thus, \perp at every state in question \mathcal{E} . But \perp has the same probability, namely 0, at every state in every question. So, by question-invariance, \perp is accepted at every state in every question. \square

It follows from the preceding propositions that *none* of the rules listed above is question-invariant. That fact is obvious for probalogical rules and the ad hoc rules, all of which base acceptance explicitly on the underlying question. However, even the logically closed Lockean rule with fixed threshold is question-dependent whenever the threshold is strictly between 0 and 1—for then the rule is neither skeptical nor everywhere inconsistent (at threshold 0 it is everywhere inconsistent and at threshold 1 it is skeptical). If closure under conjunction is dropped, the Lockean rule with a fixed threshold is question-invariant and is non-skeptical, but is also consistent, so it escapes Theorem 7 (recall that set-valued rules are not covered by that proposition).

We are inclined to view Theorem 7 as a *reductio* argument against question-invariance. That conclusion fits naturally with a minimalist, pragmatic interpretation of accepted proposition A as a more or less apt *proxy* for one's underlying credal state p , rather than as new “information” that alters p (e.g., by conditioning p on A). Question-invariance would be *nice*, but it is not rationally mandated under the minimalist conception of acceptance, and its price in terms of logical virtues *within* questions is too high.

14 Refinement-monotonicity

Invariance across all questions is a strong requirement. In this section, we consider the consequences of requiring invariance only over questions that refine or coarsen the given question. Say that \mathcal{E} *refines* \mathcal{F} (or that \mathcal{F} *coarsens* \mathcal{E}) if and only if each answer to \mathcal{E} entails some answer to \mathcal{F} . When \mathcal{E} refines \mathcal{F} , write $\mathcal{E} \leq \mathcal{F}$. By extension, say that p *refines* q (written $p \leq q$) when q is the restriction of p to \mathcal{A}_q , which implies that $\mathcal{E}_p \leq \mathcal{E}_q$. Say that cross-question acceptance rule β is *refinement-invariant* if and only if:

$$p \leq q \implies (p \Vdash_{\beta} A \iff q \Vdash_{\beta} A),$$

for each p, q in \mathbb{P} and for each proposition A in \mathcal{A}_q . However:

Proposition 13 *Refinement-invariance is equivalent to question-invariance.*

Proof Suppose that refinement-invariance holds and that $p(A) = q(A)$. Let $r = (p(A), 1 - p(A))$ over question $\{A, \neg A\}$. Then $p \leq r \leq q$. By refinement-invariance, it follows that $p \Vdash_{\beta} A \iff q \Vdash_{\beta} A$. The converse is immediate.

Refinement-invariance demands that acceptance be preserved under both refinement and coarsening. Since questions tend to become more precise as inquiry proceeds, perhaps it suffices merely to preserve acceptance under refinement. Accordingly, say that β is *refinement-monotone* if and only if:

$$p \leq q \implies \beta(p) \leq \beta(q),$$

for all p, q in \mathbb{P} . Refinement-monotonicity suffices for the accumulation of accepted conclusions as the question is refined and for the pooling of propositions accepted across diverse questions. With respect to the latter, let p, q, r be in \mathbb{P} . Say that r is a *conjunction* of p, q if and only if r is a maximally coarse common refinement of p, q . Then say that β *preserves conjunction* if and only if $\beta(r) \leq \beta(p) \wedge \beta(q)$, for each p and q in \mathbb{P} and for each conjunction r of p and q . Then it is easy to show that:

Proposition 14 *Conjunction-preservation is equivalent to refinement-monotonicity.*

Alas, probalogical rules also violate refinement-monotonicity—as witnessed by the simple lottery example in the preceding section of this paper. Again, the failure is not a defect but an ineluctable consequence of the logical virtues of probalogical rules.




Fig. 23 Acceptance snakes up the triangle

Theorem 8 *Suppose that β is refinement-monotone, validates system \mathbf{P} in each question, and is non-skeptical about both answers in some binary question. Then there exists a facet of at least two dimensions over which β accepts \perp everywhere.*

The alternative rules listed above also violate refinement-monotonicity, even though they all fail to validate system \mathbf{P} . Choosing a probalogical rule at least yields the net advantage of validating \mathbf{P} .

The proof of Theorem 8 proceeds by a sequence of lemmas that rely heavily on the geometrical characterizations of the axioms of \mathbf{P} established in Sect. 9. Consider the binary question $\{E_0, F_0\}$, whose space of credal states is depicted in Fig. 23a as the line next to the triangle. Assume that β is non-skeptical about answers E_0 and F_0 , so that β accepts E_0 at p_0 and F_0 at q . Since E_0 is infinite, split E_0 into infinite answers F_1 and E_1 to produce the refined, ternary question $\{F_0, F_1, E_1\}$ (Fig. 23b). Suppose that β is refinement-monotone. Then proposition F_0 is accepted throughout the line segment L depicted in Fig. 23b, which is defined to be the set of all credal states that refine q . Similarly, proposition $E_0 = E_1 \vee F_1$ is accepted throughout the line segment M , which is the set of all credal states that refine p_0 . Let line segment N connect the right endpoint of L in Fig. 23b to the opposite corner e_1 , intersecting M at credal state u ; then project u to the (one-dimensional) facet for proposition $E_1 \vee F_0$ to obtain credal state p_1 . The following lemma concerns p_1 .

Lemma 10 *Suppose that β is refinement-monotone and validates system \mathbf{P} . Then $p_1 \Vdash_{\beta} E_1$.*

Proof Proposition E_1 is accepted by β at e_1 , by the geometry of Reflexivity (Proposition 8); and F_0 is accepted at each point on L , by construction. So the disjunction $E_1 \vee F_0$ is accepted by β at both endpoints of N . Then, since u lies on N , we have that $u \Vdash_{\beta} E_1 \vee F_0$, by the geometry of Or (Proposition 10). We have noted that $u \Vdash_{\beta} E_1 \vee F_1$. So $u \Vdash_{\beta} E_1$, because $E_1 = (E_1 \vee F_0) \wedge (E_1 \vee F_1)$. Then, since p_1 is the projection of u onto the facet for a logical consequence of E_1 , the geometry of Cautious Monotonicity (Proposition 9) yields that $p_1 \Vdash_{\beta} E_1$, as required. \square

The result is that E_1 is accepted by β with a lower probability than E_0 . Split E_1 into two infinite, exclusive propositions E_2 and F_2 and, thus, obtain the finer, quaternary question $\{F_0, F_1, F_2, E_2\}$. Restrict attention to the two-dimensional, triangular facet

for proposition $F_0 \vee F_2 \vee E_2$, as depicted in Fig. 23c. Construct credal state p_2 as we did for p_1 , and argue similarly that E_2 is accepted at p_2 , with an even lower probability. This construction can be repeated until we obtain a refined, finite question $\{F_0, F_1, \dots, F_n, E_n\}$ such that E_n is accepted at p_n with low probability (Fig. 23d)—so low that p_n is far away from corner e_n and lies on or above the line L . Therefore:

Lemma 11 *Continuing from the preceding lemma, $p_n \Vdash_\beta E_n$.*

Then inconsistency arises:

Lemma 12 *Continuing from the preceding lemma, let line segment $\overline{p_n f_n}$ intersect L at v . Then $v \Vdash_\beta \perp$.*

Proof Proposition E_n is accepted by β at p_n , by construction; and F_0 is accepted at f_n , by the geometry of Reflexivity (Proposition 8). So the disjunction $E_n \vee F_n$ is accepted by β at both endpoints of line segment $\overline{p_n f_n}$. Then, by the geometry of Or (Proposition 10), $v \Vdash_\beta E_n \vee F_n$. But $v \Vdash_\beta F_0$, because v lies on L and thus refines q . Since $\perp = F_0 \wedge (E_n \vee F_n)$, we have that $v \Vdash_\beta \perp$, as required. \square

Here is the *coup de grace*, of which Theorem 8 is an immediate corollary.

Lemma 13 *Continuing from the preceding lemma, let \mathcal{P}_{n+2} be the set of probability measures defined on $\mathcal{A}_{\mathcal{E}_{n+2}}$, where \mathcal{E}_{n+2} is the question $\{F_0, \dots, F_n, E_n\}$. Then β accepts \perp at each credal state p in facet $\mathcal{P}_{n+2}|(F_0 \vee F_n \vee E_n)$.*

Proof Let Δ denote the two-dimensional facet $\mathcal{P}_{n+2}|(F_0 \vee F_n \vee E_n)$. Suppose that v lies in the interior, but not the sides, of Δ . Since \perp is accepted at v , we have that \perp is accepted at the three corners f_0, f_n, e_n of Δ , by projecting v to the three corners and by the geometry of Cautious Monotonicity (Proposition 9). Then, since each side of Δ has endpoints that are corners, we have that \perp is accepted on the three sides of Δ , by the geometry of Or (Proposition 10). Then, since each point on Δ is on a line segment with endpoints on the sides of Δ , we have that \perp is accepted at each credal state on Δ , as required. When v is not in the interior of Δ , v lies on side $\overline{e_n f_0}$ of Δ and, thus, cannot be projected to the opposite corner f_n . But in that case we can apply the geometry of Or (Proposition 10) to line segment $v f_n$ to show that F_n is accepted at every credal state on $v f_n$. Similarly, $F_0 \vee E_n$ is accepted at every credal state on $\overline{w e_n}$, where w is defined to be the intersection of line L and $\overline{f_0 f_n}$. So \perp is accepted at the intersection of $v f_n$ and $\overline{w e_n}$, which is in the interior of Δ —the second case is thus reduced to the first case. \square

15 Probalogic generalized

We close with a natural generalization of the probalogical framework. The uniform probability measure over \mathcal{E} is the center of the simplex \mathcal{P} and serves as the probalogically weakest credal state in \mathcal{P} in the presentation to this point. But, as Levi (1967, 1969) has emphasized, the answers to question \mathcal{E} typically have different *contents* (e.g., “quantum mechanics is true” has a great deal of content but “quantum mechanics is false” has very little). Therefore, a credal state that assigns less probability to an




Fig. 24 Deformation of geologic and corresponding deformation of probalogue

answer that has more content could sensibly be understood as weaker than a uniform state that accords the same probability to all answers. In that case, probalogue should be relative not only to question \mathcal{E} , but to an assignment of contents to the answers to \mathcal{E} . The result is a family of probalogics sensitive both to question \mathcal{E} and to the relative contents of the answers to \mathcal{E} .

We approach the issue as follows. If the answers E_i differ in content, it is natural to weight answers by *weakness* and to think of the neutral credal state as the center of mass of the answers. As a result, the weakest credal state is biased toward answers of low content. In particular, the center of \mathcal{P} is stronger than a state closer to a very weak answer. Recall that probalogue is just the geological cube in perspective. The sides of the cube have equal length. To represent differences in content, deform the cube into an oblong box whose side lengths are inversely proportional to the strengths of the corresponding answers (Fig. 24). Just like the cube, the oblong box may be viewed as a *generalized geological semantics* (recall that geological structure does not uniquely determine the metric). Project the generalized geologic from the box to the triangular credal state space, just as before, to induce a *generalized probalogue* on it. Then the credal states stronger than p are those in the grey region of Fig. 24d. Disjunction and conjunction are defined as before.




Fig. 25 Generalized probabilogical acceptance rule

The weakest proposition in the generalized geologic is (m_1, m_2, m_3) (i.e. the vertex of the box that is most distant from the origin), so its rectilinear projection w to the triangle is the weakest credal state in the corresponding probabilogic. Projection preserves ratios between the rectangular coordinates, so we have: $w = (\frac{m_1}{M}, \frac{m_2}{M}, \frac{m_3}{M})$, where $M = \sum_{i \in I} m_i$. The coordinates of w uniquely determine the generalized probabilogic that has w as the weakest state. Intuitively, the result is like viewing a phone booth, rather than a cubical office, from the origin (Fig. 24d).²² Acceptance rules are still defined as maps that preserve probabilogical structure and they look like Fig. 25. Although the generalized probabilogical acceptance rules appear “oblique”, the boundaries of acceptance zones still follow rays from the corners—so they still validate exactly Adams’ conditional logic. Algebraically, the generalized rules take the following form:

$$\alpha(p) = \bigwedge \left\{ \neg E_i : \frac{p(E_i)/m_i}{\max_j p(E_j)/m_j} \triangleleft_i 1 - r_i \text{ and } i \in I \right\}.$$

The acceptance rules introduced in Levi (1996, p. 286) are the same, except that we allow different thresholds r_i for different answers E_i while Levi does not. As we mentioned at the outset, Levi sees no justification for these rules, relative to his momentous understanding of acceptance as an explicit decision to condition one’s credal state on the accepted proposition and, therefore, to bet one’s life on it against nothing. Our own justification for the rules, grounded in a weaker conception of acceptance as apt description of one’s credal state relative to a question, is again, that they preserve naturally defined logical structures over credal states relative to a question and that they validate exactly Adams’ logic of conditionals.

Acknowledgments The authors are indebted to David Etlin for his extremely detailed and expert commentary on a draft of this paper at the 2011 Formal Epistemology Workshop. We are indebted to Teddy Seidenfeld for informing us that Levi already proposed the camera shutter rule and for providing the citation. We are indebted to Hannes Leitgeb for sharing his alternative approach based on conditional acceptance rules with us. We are also indebted to Clark Glymour, Horacio Arlo-Costa, and Greg Wheeler for useful comments. This work was supported generously by the National Science Foundation under grant 0750681.

²² In terms of projective geometry, the geological transformation is a non-rotational, non-reflective linear transformation and, thus, the induced probabilogical transformation is a projective transformation that fixes all the corners.

Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.

References

- Adams, E. W. (1975). *The logic of conditionals*. Dordrecht: D. Reidel.
- Alchourrón, C. E., Gärdenfors, P., & Makinson, D. (1985). On the logic of theory change: Partial meet contraction and revision functions. *Journal of Symbolic Logic*, *50*, 510–530.
- Arló-Costa, H., & Parikh, R. (2005). Conditional probability and defeasible inference. *Journal of Philosophical Logic*, *34*, 97–119.
- Barwise, K. (1969). Infinitary logic and admissible sets. *Journal of Symbolic Logic*, *34*, 226–252.
- Douven, I. (2002). A new solution to the paradoxes of rational acceptability. *British Journal for the Philosophy of Science*, *53*, 391–410.
- Hajek, P. (1998). *Metamathematics of fuzzy logic*. Dordrecht: Kluwer.
- Harper, W. (1975). Rational belief change, popper functions and counterfactuals. *Synthese*, *30*(1–2), 221–262.
- Jeffrey, R. C. (1970). Dracula meets Wolfman: Acceptance vs. partial belief. In M. Swain (Ed.), *Induction, acceptance, and rational belief*. D. Reidel.
- Karp, C. (1964). *Languages with expressions of infinite length*. Dordrecht: North Holland.
- Kelly, K. (2008). Ockham's razor, truth, and information. In J. van Benthem & P. Adriaans (Eds.), *Handbook of the philosophy of information* (pp. 321–360). Dordrecht: Elsevier.
- Kelly, K. (2011). Ockham's razor, truth, and probability. In P. Bandyopadhyay & M. Forster (Eds.), *Handbook on the philosophy of statistics* (pp. 983–1024). Dordrecht: Elsevier.
- Kraus, S., Lehmann, D., & Magidor, M. (1990). Nonmonotonic reasoning, preferential models and cumulative logics. *Artificial Intelligence*, *44*, 167–207.
- Kyburg, H. (1961). *Probability and the logic of rational belief*. Middletown: Wesleyan University Press.
- Lehmann, D., & Magidor, M. (1992). What does a conditional base entail? *Artificial Intelligence*, *55*, 1–60.
- Leitgeb, H. (2010). *Reducing belief simpliciter to degrees of belief*. Presentation of his unpublished results at the opening celebration of the Center for Formal Epistemology at Carnegie Mellon University in the Summer of 2010 (unpublished results).
- Levi, I. (1967). *Gambling with truth: An essay on induction and the aims of science*. New York: Harper & Row (2nd ed., Cambridge, MA: The MIT Press, 1973).
- Levi, I. (1969). Information and inference. *Synthese*, *19*, 369–391.
- Levi, I. (1996). *For the sake of the argument: Ramsey test conditionals, inductive inference and non-monotonic reasoning*. Cambridge: Cambridge University Press.
- Lin, H. (2011). *A new theory of acceptance that solves the lottery paradox and provides a simplified probabilistic semantics for Adams' logic of conditionals*. Master's thesis, Carnegie Mellon University, Pittsburgh, PA.
- Makinson, D., & Gärdenfors, P. (1991). Relations between the logic of theory change and nonmonotonic logic. In A. Fuhrmann & M. Morreau (Eds.), *The logic of theory change* (pp. 183–205). Springer-Verlag Lecture notes in computer science 465. Berlin: Springer.
- Novak, V., Perfilieva, I., & Mockor, J. (2000). *Mathematical principles of fuzzy logic*. Dordrecht: Kluwer.
- Pearl, J. (1989). Probabilistic semantics for nonmonotonic reasoning: A survey. In *Proceedings of the first international conference on principles of knowledge representation and reasoning (KR '89)* (pp. 505–516). (Reprinted in G. Shafer & J. Pearl (Eds.), *Readings in uncertain reasoning* (pp. 699–710). San Francisco: Morgan Kaufmann).
- Pollock, J. (1995). *Cognitive carpentry*. Cambridge, MA: MIT Press.
- Ramsey, F. P. (1929). General propositions and causality. In H. A. Mellor (Ed.), *Philosophical papers*. Cambridge: Cambridge University Press; 1990.
- Ryan, S. (1996). The epistemic virtues of consistency. *Synthese*, *109*, 121–141.
- van Fraassen, B. (1995). Fine-grained opinion, probability and the logic of full belief. *Journal of Philosophical Logic*, *24*, 349–377.
- Zadeh, L. (1965). Fuzzy sets. *Information and Control*, *8*, 338–353.