

## New copper(II) 2-(alkylamino)tropolonates

Marie C. Barret · Purvi H. Bhatia ·  
Gabriele Kociok-Köhn · Kieran C. Molloy

Received: 25 March 2014 / Accepted: 15 April 2014 / Published online: 8 May 2014  
© Springer International Publishing Switzerland 2014


**Abstract** The copper aminotropolones  $\text{Cu}[\text{ON}(\text{R}')\text{C}_7\text{H}_4\text{R}-4]_2$  [ $\text{R} = \text{H}$ ,  $\text{R}' = \text{Me}$  (**13**),  $\text{Et}$  (**14**),  $n\text{-Pr}$  (**15**),  $n\text{-Bu}$  (**16**),  $\text{Bz}$  (**17**),  $\text{MenOCH}_2\text{CH}_2$  (**20**);  $\text{R} = i\text{-Pr}$ ,  $\text{R}' = \text{Me}$  (**18**),  $n\text{-Pr}$  (**19**),  $\text{MenOCH}_2\text{CH}_2$  (**21**)] have been prepared from the corresponding aminotropolones  $\text{HN}(\text{R}')\text{OC}_7\text{H}_4\text{R}-4$  (**1–7**) by reacting with copper(II) acetate in aqueous ethanol. **20**, **21** contain the flavourant, menthol, as part of the ligand. The structures of **5** ( $\text{R} = \text{H}$ ,  $\text{R}' = \text{Bz}$ ), a hydrogen-bonded dimer, **14** and **20**, both incorporating square-planar, four-coordinate copper centres, have been determined by X-ray crystallography. The antibacterial activities of complexes **13**, **17**, **20** and **21** have been assayed against *Staphylococcus wagneri*, an in vitro model of plaque inhibition effects, and found to be more active than a commercial toothpaste formulation, but less active than the *O,O*-chelated copper(II) complex of ethylmaltol.

### Introduction

Hinokitiol, a naturally occurring  $\alpha$ -hydroxyketone which can be extracted from wood (Thujapalicata, Taiwan Hinoki) continues to attract interest as a ligand due to its attractive biological properties (antimicrobial [1, 2], insecticidal [3], cytotoxicity on tumour cells [4]) which make it a favourable chelating agent for metals which might offer biological synergism. As a result, numerous metal derivatives of both hinokitiol (I,  $\text{R} = i\text{-Pr}$ ) [5], and its parent, tropolone (I,  $\text{R} = \text{H}$ ) [6–11], have been prepared and analysed. More

recently, tropolone has been elaborated to form mesogenic complexes when coordinated to metals [12, 13]. Our own work in this area stems from an interest in copper, zinc and tin complexes which can be incorporated into dental formulations [14], including complexes of this triad with ligands such as maltol and ethylmaltol as well as I [15–17]. In this regard, the ligand must be orally acceptable, be able to deliver the metal in the correct oxidation state and concentration (i.e. soluble complexes) and have reasonable long-term stability.

As part of this general interest, we have become interested in the related 2-(alkylamino)tropolone ligand system (II), which offers a complimentary  $\kappa^2\text{-O,N}$  chelation mode to the  $\kappa^2\text{-O,O}$  mode inherent in I. Relatively few reports have appeared concerning the chelation of metals such as  $\text{Cu}(\text{II})$  [18–20] and  $\text{Zn}(\text{II})$  [21] with II, to which we now add our own findings relating to copper. In particular, as well as broadening the scope of simple  $\text{ML}_2$  complexes, we report on the incorporation of a flavouring component, menthol, onto the ligand periphery. As one of the drawbacks of these complexes for oral delivery is their metallic taste even at low concentrations, this approach may go some way to mitigating this problem.



M. C. Barret · P. H. Bhatia · G. Kociok-Köhn ·  
K. C. Molloy (✉)  
Department of Chemistry, University of Bath, Claverton Down,  
Bath BA2 7AY, UK  
e-mail: kcm68win@outlook.com

### Experimental

Starting materials: tropolone was obtained commercially from Avocado, and hinokitiol was provided by Unilever

Research; both were used as received without further purification. Reactions were carried out in air unless otherwise specified. Elemental analyses were performed on a Carlo-Erba Strumentazione E. A. model 1106 micro-analyser, and the temperature of the furnace was set to 500 °C. The results were duplicated, and the mean of the duplicated measurements was the final result. All  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra were recorded on a Bruker Avance (300 MHz) Fourier transform spectrometer. All spectra were recorded in *d*-chloroform and are referenced to residual solvent peaks. Peak positions were recorded in  $\delta$  ppm with abbreviations s, d, t and m denoting singlet, doublet, triplet and multiplet, respectively. All coupling constants (*J*) are quoted in Hertz. Infrared spectra were recorded as a thin oil film sandwiched between NaCl plates in the frequency range 4,000–400  $\text{cm}^{-1}$  on a Nicolet 510P FT-IR spectrometer. All absorptions are quoted in  $\text{cm}^{-1}$ .

#### Synthesis of 2-*p*-toluenesulfonyltropolone

Following a previously reported procedure [22], tropolone (5.00 g, 41.0 mmol) and *p*-toluenesulfonyl chloride (7.81 g, 41.0 mmol) were mixed together in dry pyridine (20 mL). After 30 min, the mixture was filtered and the product obtained was then washed with cold water and dried in vacuo, yielding the product as a white solid (11.18 g, 98 %). Analysis: Found [calc. for  $\text{C}_{14}\text{H}_{12}\text{O}_4\text{S}$ ]: C 61.0 (60.9) %; H 4.4 (4.3) %. M. p.: 156–158 °C (lit: 156.5–157.5 °C) [23].

#### Synthesis of 2-*p*-toluenesulfonylhinokitiol

The method described above [22] was applied to hinokitiol (5.00 g, 30.5 mmol). Hinokitiol and *p*-toluenesulfonyl chloride (5.81 g, 30.5 mmol) were mixed in dry pyridine (20 mL); the product was then washed with cold water and dried in vacuo, to obtain a pale yellow solid (9.31 g, 96 %). Analysis: Found (calc. for  $\text{C}_{17}\text{H}_{18}\text{O}_4\text{S}$ ): C 64.6 (64.2) %; H 6.1 (5.7) %.

The 2-alkylaminotropones and 2-alkylaminohinokitiols were prepared following literature procedures; [23, 24] the methodology described by Rasika Dias et al. [23] was used for (1), (2), (6), (7), and the one from Pietra et al. [24] was repeated for (3)–(5) and (8)–(10).

#### Synthesis of 2-methylaminotropone (1)

2-Methylaminotropone was prepared using a solution of 2-*p*-toluenesulfonyltropolone (2.00 g, 7.24 mmol) in dichloromethane (30 mL) and aqueous methylamine (40 %) (10 mL). The mixture was stirred overnight at room temperature, then the two layers were separated, and the organic layer was washed with water, dried with

magnesium sulphate, and concentrated under vacuum to obtain a yellow solid (0.76 g, 78 %). Analysis: Found (calc. for  $\text{C}_8\text{H}_9\text{NO}$ ): C 61.0 (60.9); H 4.4 (4.3); N 4.1 (4.0) %.  $^1\text{H}$  NMR [ $\delta$ (ppm),  $\text{CDCl}_3$ ]: 2.95 [3H, s,  $\text{NCH}_3$ ]; 6.50–7.19 [5H, m,  $\text{C}_7\text{H}_5$ ].  $^{13}\text{C}$  NMR [ $\delta$ (ppm),  $\text{CDCl}_3$ ]: 28.5 [ $\text{CH}_3$ ]; 107.3, 121.0, 127.3, 135.3, 136.3 [CH]; 155.5 [ $\text{C}_2$ ]; 175.6 [ $\text{C}_3$ ]. IR: 3,293  $\nu$ (N–H); 1,629, 1,590  $\nu$ (C=O); 1,510  $\nu$ (C=C).

#### 2-Ethylaminotropone (2)

The methodology described above was employed with aqueous ethylamine (70 %) (10 mL) and 2-*p*-toluenesulfonyltropolone (2.00 g, 7.24 mmol) in dichloromethane (30 mL). An orange oil was obtained (0.77 g, 71 %). Analysis: Found (calc. for  $\text{C}_9\text{H}_{11}\text{NO}$ ): C 71.9 (72.5); H 7.4 (7.4); N 9.3 (9.4) %.  $^1\text{H}$  NMR [ $\delta$ (ppm),  $\text{CDCl}_3$ ]: 1.31 [3H, t,  $\text{NCH}_2\text{CH}_3$ ]; 3.28 [2H, q,  $\text{NCH}_2\text{CH}_3$ ]; 6.49–7.21 [5H, m,  $\text{C}_7\text{H}_5$ ].  $^{13}\text{C}$  NMR [ $\delta$ (ppm),  $\text{CDCl}_3$ ]: 13.4 [ $\text{CH}_3$ ]; 37.2 [ $\text{CH}_2$ ]; 108.4, 121.7, 128.0, 136.1, 136.9 [CH]; 155.2 [ $\text{C}_2$ ]; 176.2 [ $\text{C}_3$ ]. IR data g.: 3,301  $\nu$ (N–H); 1,630, 1,594  $\nu$ (C=O); 1,515  $\nu$ (C=C).

#### 2-*n*-Propylaminotropone (3)

To a solution of 2-*p*-toluenesulfonyltropolone (2.00 g, 7.24 mmol) in dried DMSO (100 mL) was added *n*-propylamine (10 mL), and the solution was stirred overnight. The mixture was poured into water and extracted with ether; the ether layer was dried with magnesium sulphate; and the solvent removed in vacuo. A brown oil was obtained (0.81 g, 68 %). Analysis: Found [calc. for  $\text{C}_{10}\text{H}_{13}\text{NO}$ ]: C 72.7 (73.6); H 8.0 (8.0); N 8.65 (8.58) %.  $^1\text{H}$  NMR [ $\delta$ (ppm),  $\text{CDCl}_3$ ]: 0.94 [3H, t,  $\text{NCH}_2\text{CH}_2\text{CH}_3$ ]; 1.64 [2H, m,  $\text{NCH}_2\text{CH}_2\text{CH}_3$ ]; 3.18 [2H, m,  $\text{NCH}_2\text{CH}_2\text{CH}_3$ ]; 6.49–7.21 [5H, m,  $\text{C}_7\text{H}_5$ ].  $^{13}\text{C}$  NMR [ $\delta$ (ppm),  $\text{CDCl}_3$ ]: 12.1 [ $\text{CH}_2\text{CH}_2\text{CH}_3$ ]; 22.2 [ $\text{CH}_2\text{CH}_2\text{CH}_3$ ]; 44.9 [ $\text{CH}_2\text{CH}_2\text{CH}_3$ ]; 108.8, 122.1, 128.4, 136.5, 137.3 [CH]; 155.7 [ $\text{C}_2$ ]; 176.5 [ $\text{C}_3$ ]. IR: 3,291  $\nu$ (N–H); 1,637, 1,595  $\nu$ (C=O); 1,512  $\nu$ (C=C).

#### 2-*n*-Butylaminotropone (4)

Following the procedure with *n*-butylamine (10 mL) and 2-*p*-toluenesulfonyltropolone (2.00 g, 7.24 mmol), 4 was obtained (0.89 g, 69 %) as a brown-orange oil. Analysis: Found [calc. for  $\text{C}_{11}\text{H}_{15}\text{NO}$ ]: C 72.6 (74.6); H 8.6 (8.5); N 7.9 (7.9) %.  $^1\text{H}$  NMR [ $\delta$ (ppm),  $\text{CDCl}_3$ ]: 0.94 [3H, t,  $\text{NCH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ ]; 1.40 [2H, m,  $\text{NCH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ ]; 1.63 [2H, m,  $\text{NCH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ ]; 3.22 [2H, t,  $\text{NCH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ ]; 6.49–7.21 [5H, m,  $\text{C}_7\text{H}_5$ ].  $^{13}\text{C}$  NMR [ $\delta$ (ppm),  $\text{CDCl}_3$ ]: 13.5 [ $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ ]; 20.0 [ $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ ]; 30.2 [ $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ ]; 42.3

[CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>]; 108.4, 121.7, 127.9, 136.1, 136.9 [CH]; 155.4 [C<sub>2</sub>]; 176.2 [C<sub>3</sub>]. IR: 3,307  $\nu$ (N–H); 1,636, 1,593  $\nu$ (C=O); 1,516  $\nu$ (C=C).

### 2-Benzylaminotropone (5)

Following the same method as for (3), a yellow solid was obtained using benzylamine (10 mL) and 2-*p*-toluenesulfonyltropone (2.00 g, 7.24 mmol). Yield 1.34 g, 88 %. Recrystallisation from chloroform/cyclohexane gave yellow crystals suitable for X-ray analysis. Analysis: Found [calc. for C<sub>14</sub>H<sub>13</sub>NO]: C 79.5 (79.6); H 6.3 (6.2); N 6.7 (6.6) %. <sup>1</sup>H NMR [ $\delta$ (ppm), CDCl<sub>3</sub>]: 4.44 [2H, s, CH<sub>2</sub>]; 6.43–7.52 [10H, m, C<sub>6</sub>H<sub>5</sub> and C<sub>7</sub>H<sub>5</sub>]. <sup>13</sup>C NMR [ $\delta$ (ppm), CDCl<sub>3</sub>]: 46.5 [CH<sub>2</sub>]; 108.4, 122.2, 127.5, 128.0, 130.1, 131.4, 136.1, 136.9 [CH]; 156.0 [C<sub>2</sub>]; 175.8 [C<sub>3</sub>]. IR: 3,304  $\nu$ (N–H); 1,632, 1,592  $\nu$ (C=O); 1,512  $\nu$ (C=C).

### 2-Methylamino-4-*i*-propyltropone (6)

The method described earlier [23] for (1) was repeated for (6) using 2-*p*-toluenesulfonylhinokitiol (2.00 g, 6.28 mmol) and the corresponding amine in excess in solution in dichloromethane (30 mL). Aqueous methylamine (40 %) (10 mL) was used for (6) (0.57 g, 51 %), aqueous ethylamine (70 %) (10 mL) for (7) (0.92 g, 76 %). Analysis for (6): Found [calc. for C<sub>11</sub>H<sub>15</sub>NO]: C 72.4 (74.6); H 8.3 (8.5); N 7.6 (7.9) %. IR: 3,286  $\nu$ (N–H); 1,633, 1,595  $\nu$ (C=O); 1,514  $\nu$ (C=C).

### 2-*n*-Propylamino-4-*i*-propyltropone (7)

Following the procedure for (3), (7) was obtained using 2-*p*-toluenesulfonylhinokitiol (2.00 g, 6.28 mmol) and <sup>n</sup>propylamine (10 mL). Yields 0.51 g, 39 %. Analysis for (8): Found [calc. for C<sub>13</sub>H<sub>19</sub>NO]: C 74.0 (76.1); H 9.5 (9.3); N 6.2 (6.8). IR: 3,303  $\nu$ (N–H); 1,632, 1,592  $\nu$ (C=O); 1,511  $\nu$ (C=C).

### Synthesis of menthoxyacetic acid (8)

Following a previously reported procedure [25], menthol (15.6 g, 100 mmol) was dissolved in THF (ca. 50 mL) under nitrogen. To this solution was added lithium ribbon (0.80 g, 115 mmol) cut into small pieces, and this mixture was refluxed under nitrogen for 4 h. This solution was then filtered to remove the unreacted lithium, and dry monochloroacetic acid (4.25 g, 45.1 mmol) in anhydrous THF (12.5 mL) was added dropwise to the filtrate. This mixture was gently refluxed for 20 h. After this time, water (ca. 30 mL) was added. The THF and menthol were fractionally distilled from the yellow suspension, which was then cooled in an ice bath and filtered to obtain the solid lithium

salt as a white precipitate. This was dissolved in dilute HCl and extracted using diethyl ether. The resulting ethereal solution was washed with water, dried over magnesium sulphate, filtered, and concentrated under rotary evaporation to produce a yellow oil (2.25 g; 10 %). <sup>1</sup>H NMR (CDCl<sub>3</sub>): 8.10 [H, br s, OH]; 4.13 [1H, d, J<sub>a-b</sub> = 15.0 Hz, C<sup>2</sup>H<sup>a</sup>]; 4.02 [1H, d, J<sub>b-a</sub> = 15.0 Hz, C<sup>2</sup>H<sup>b</sup>]; 3.13 [1H, ddd, J<sub>3-4ax</sub> = J<sub>3-8</sub> = 9.10, J<sub>3-4eq</sub> = 3.00 Hz, C<sup>3</sup>H]; 1.10–2.30 [2H, m, C<sup>4</sup>H<sub>2</sub>; 1H, m, C<sup>5</sup>H; 2H, m, C<sup>6</sup>H<sub>2</sub>; 2H, m, C<sup>7</sup>H<sub>2</sub>; 1H, m, C<sup>8</sup>H; 1H, m, C<sup>9</sup>H]; 0.83, 0.75 [3H, d, J<sub>10-9</sub>/J<sub>11-9</sub> = 6.30, 8.40 Hz, C<sup>10</sup>H<sub>3</sub>/C<sup>11</sup>H<sub>3</sub>]; 0.85 [3H, d, J<sub>12-5</sub> = 8.10 Hz, C<sup>12</sup>H<sub>3</sub>]. <sup>13</sup>C NMR (CDCl<sub>3</sub>): 176 [C<sup>1</sup>]; 66.3 [C<sup>2</sup>]; 80.8 [C<sup>3</sup>]; 40.2 [C<sup>4</sup>]; 31.9 [C<sup>5</sup>]; 34.7 [C<sup>6</sup>]; 22.6 [C<sup>7</sup>]; 48.3 [C<sup>8</sup>]; 26.0 [C<sup>9</sup>]; 21.4, 16.4 [C<sup>10</sup>/C<sup>11</sup>]; 23.5 [C<sup>12</sup>]. IR: 1,720  $\nu$ (C=O); 1,275  $\nu$ (C–O); 3,120  $\nu$ (O–H).

### Synthesis of 2-menthoxyacetamide (9) [26]

Menthoxyacetyl chloride was produced by dissolving menthoxyacetic acid (8) (10.0 g, 46.7 mmol) in thionyl chloride (16.7 mL, 229 mmol) and warming at 50 °C for 3 h. The excess thionyl chloride was removed by warming in a water bath under reduced pressure, and the acid chloride (7.30 g; 67 %) was used directly. Ammonia (19.2 mL) was added directly to the acid chloride, (7.30 g, 31.3 mmol) and stirred at 0 °C for 24 h. The final product was washed with water, extracted with diethyl ether (2 × 20 mL), dried over anhydrous magnesium sulphate, filtered, and dried in vacuo yielding a white solid. Recrystallisation from petroleum-ether 60–80 °C gave white crystals (4.58 g; 46 %; m. p. (°C): 92–94 (Lit: 92–94 [26]). Analysis for (22): Found [calc. for C<sub>12</sub>H<sub>23</sub>O<sub>2</sub>N]: C 66.4 (67.7); H 10.7 (10.7); N 6.2 (6.5) %. <sup>1</sup>H NMR (CDCl<sub>3</sub>): 5.80, 6.59 [2H, br s, NH<sub>2</sub>]; 3.98 [1H, d, J<sub>a-b</sub> = 15.0 Hz, C<sup>2</sup>H<sup>a</sup>]; 3.78 [1H, d, J<sub>b-a</sub> = 15.6 Hz, C<sup>2</sup>H<sup>b</sup>]; 3.10 [1H, ddd, J<sub>3-4ax</sub> = J<sub>3-8</sub> = 11.4, J<sub>3-4eq</sub> = 4.50 Hz, C<sup>3</sup>H]; 1.20–2.20 [2H, m, C<sup>4</sup>H<sub>2</sub>; 1H, m, C<sup>5</sup>H; 2H, m, C<sup>6</sup>H<sub>2</sub>; 2H, m, C<sup>7</sup>H<sub>2</sub>; 1H, m, C<sup>8</sup>H; 1H, m, C<sup>9</sup>H]; 0.84, 0.72 [3H, d, J<sub>10-9</sub>/J<sub>11-9</sub> = 3.90, 6.60 Hz, C<sup>10</sup>H<sub>3</sub>/C<sup>11</sup>H<sub>3</sub>]; 0.87 [3H, d, J<sub>12-5</sub> = 3.30 Hz, C<sup>12</sup>H<sub>3</sub>]. <sup>13</sup>C NMR (CDCl<sub>3</sub>): 174 [C<sup>1</sup>]; 68.1 [C<sup>2</sup>]; 80.8 [C<sup>3</sup>]; 40.5 [C<sup>4</sup>]; 31.8 [C<sup>5</sup>]; 34.8 [C<sup>6</sup>]; 22.6 [C<sup>7</sup>]; 48.4 [C<sup>8</sup>]; 26.4 [C<sup>9</sup>]; 21.3, 16.6 [C<sup>10</sup>/C<sup>11</sup>]; 23.6 [C<sup>12</sup>]. IR: 3,189  $\nu$ (N–H); 1,640  $\nu$ (C=O).

### Synthesis of 2-menthoxyethylamine (10) [26]

In a three-necked round-bottomed flask were placed 2-menthoxyacetamide (9) (4.01 g, 18.9 mmol) and anhydrous THF (30 mL). Lithium aluminium hydride (2.15 g, 56.7 mmol) dissolved in anhydrous THF (20 mL) was added dropwise using a syringe under the flow of nitrogen to the 2-menthoxyacetamide. During the addition, the reaction mixture gave off bubbles and became grey. The

contents of the flask were stirred vigorously and allowed to reflux for 8 h, then cooled to room temperature, and stirred at this temperature overnight. The mixture was cooled to 0 °C by application of an ice bath, and water (5 mL) was added dropwise. A white precipitate was produced, and bubbles of gas were given off. The precipitate was removed by filtration and washed with diethyl ether. The filtrate was extracted with diethyl ether (2 × 25 mL) and 15 % (w/v) sodium hydroxide (5 mL). The organic layers were combined, dried over anhydrous magnesium sulphate, filtered, and concentrated to dryness by rotary evaporation yielding a colourless liquid (2.87 g; 78 %). Analysis for (**23**): Found (calc. for C<sub>12</sub>H<sub>25</sub>ON): C 70.1 (72.4); H 12.2 (12.7); N 5.2 (5.4) %. <sup>1</sup>H NMR (CDCl<sub>3</sub>): 2.75 [2H, t, J<sub>NH-1</sub> = 3.00 Hz, NH<sub>2</sub>]; 3.65 [2H, m, C<sup>1</sup>H<sub>2</sub>]; 3.25 [2H, m, C<sup>2</sup>H<sub>2</sub>]; 2.97 [1H, ddd, J<sub>3-4ax</sub> = J<sub>3-8</sub> = 10.5, J<sub>3-4eq</sub> = 4.50 Hz, C<sup>3</sup>H]; 1.15–2.20 [2H, m, C<sup>4</sup>H<sub>2</sub>; 1H, m, C<sup>5</sup>H; 2H, C<sup>6</sup>H<sub>2</sub>; 2H, m, C<sup>7</sup>H<sub>2</sub>; 1H, m, C<sup>8</sup>H; 1H, m, C<sup>9</sup>H]; 0.83, 0.71 [3H, d, J<sub>10-9</sub>/J<sub>11-9</sub> = 6.60, 7.20 Hz, C<sup>10</sup>H<sub>3</sub>/C<sup>11</sup>H<sub>3</sub>]; 0.86 [3H, d, J<sub>12-5</sub> = 6.00 Hz, C<sup>12</sup>H<sub>3</sub>]. <sup>13</sup>C NMR (CDCl<sub>3</sub>): 39.5 [C<sup>1</sup>]; 69.6 [C<sup>2</sup>]; 78.3 [C<sup>3</sup>]; 41.3 [C<sup>4</sup>]; 30.5 [C<sup>5</sup>]; 33.6 [C<sup>6</sup>]; 21.4 [C<sup>7</sup>]; 47.3 [C<sup>8</sup>]; 26.7 [C<sup>9</sup>]; 19.9, 15.2 [C<sup>10</sup>/C<sup>11</sup>]; 22.3 [C<sup>12</sup>]. IR: 3,205 ν(N–H).

Synthesis of 2-menthoxyethylamino-cyclohept-2,4,6-trien-1-one.H<sub>2</sub>O (**11**)

2-Menthoxyethylamino-cyclohept-2,4,6-trien-1-one.H<sub>2</sub>O (**11**) was prepared by reacting 2-menthoxyethylamine (**10**) (1.00 g, 5.03 mmol), two equivalents of triethylamine (0.93 mL, 6.67 mmol) and 2-*p*-toluenesulfonyltropolone (0.92 g, 3.33 mmol) in DMSO (50 mL). This mixture was stirred for 1 week at room temperature. After completion of the reaction, the reaction mixture was dissolved in water and the two layers were separated using diethyl ether. The organic layer was washed with water, dried over magnesium sulphate, filtered, and rotary evaporated to obtain a brown oil. Concentrated HCl was added dropwise to this mixture until the pH reached 1. This was extracted with diethyl ether and washed with water (2 × 50 mL), dried over magnesium sulphate, filtered, and rotary evaporated to yield a brown oil (0.96 g; 60 %). Analysis for (**11**): Found [calc. C<sub>19</sub>H<sub>31</sub>O<sub>3</sub>N]: C 71.2 (71.0); H 9.9 (9.7); N 4.4 (4.4) %. <sup>1</sup>H NMR (CDCl<sub>3</sub>): 7.20–6.38 [5H, m, C<sup>3</sup>H, C<sup>4</sup>H, C<sup>5</sup>H, C<sup>6</sup>H, C<sup>7</sup>H]; 3.55 [1H, t, NH]; 3.32–3.90 [4H, m, C<sup>8</sup>H<sub>2</sub>, C<sup>9</sup>H<sub>2</sub>]; 3.01 [1H, ddd, J<sub>10-11ax</sub> = J<sub>10-15</sub> = 10.5, J<sub>10-11eq</sub> = 4.50 Hz, C<sup>10</sup>H]; 1.15–2.20 [2H, m, C<sup>11</sup>H<sub>2</sub>; 1H, m, C<sup>12</sup>H; 2H, m, C<sup>13</sup>H<sub>2</sub>; 2H, m, C<sup>14</sup>H<sub>2</sub>; 1H, m, C<sup>15</sup>H; 1H, m, C<sup>16</sup>H]; 0.82, 0.70 [3H, d, J<sub>17-16</sub>/J<sub>18-16</sub> = 6.60, 7.50 Hz, C<sup>17</sup>H<sub>3</sub>/C<sup>18</sup>H<sub>3</sub>]; 0.85 [3H, d, J<sub>19-12</sub> = 6.30 Hz, C<sup>19</sup>H<sub>3</sub>]. <sup>13</sup>C NMR (CDCl<sub>3</sub>): 177 [C<sup>1</sup>]; 156 [C<sup>2</sup>]; 123 [C<sup>3</sup>]; 137 [C<sup>4</sup>]; 129 [C<sup>5</sup>]; 137 [C<sup>6</sup>]; 108 [C<sup>7</sup>]; 66.3 [C<sup>8</sup>]; 40.8 [C<sup>9</sup>]; 80.3 [C<sup>10</sup>]; 43.5 [C<sup>11</sup>]; 31.9 [C<sup>12</sup>]; 34.8 [C<sup>13</sup>]; 21.8 [C<sup>14</sup>]; 48.6 [C<sup>15</sup>];

26.1 [C<sup>16</sup>]; 21.3, 16.6 [C<sup>17</sup>/C<sup>18</sup>]; 23.7 [C<sup>19</sup>]. IR: 3,300 ν(N–H) and ν(O–H); 1,631, 1,595 ν(C=O); 1,519 ν(C=C).

Synthesis of 2-menthoxyethylamino-4-isopropylcyclohept-2,4,6-trien-1-one.H<sub>2</sub>O (**12**)

2-Menthoxyethylamino-4-isopropylcyclohept-2,4,6-trien-1-one.H<sub>2</sub>O (**12**) was prepared in a similar manner to **11** by reacting 2-menthoxyethylamine (**10**) (2.00 g, 10.1 mmol), two equivalents of triethylamine (1.86 mL, 13.3 mmol) and 2-*p*-toluenesulfonylhinokitiol (2.12 g, 6.67 mmol) in DMSO (50 mL) to yield the product as a brown oil (2.54 g; 70 %). Analysis for (**12**): Found [calc. C<sub>22</sub>H<sub>37</sub>O<sub>3</sub>N]: C 73.1 (72.7); H 10.1 (10.2); N 3.2 (3.8) %. <sup>1</sup>H NMR (CDCl<sub>3</sub>): 7.45–6.28 [4H, m, C<sup>3</sup>H, C<sup>5</sup>H, C<sup>6</sup>H, C<sup>7</sup>H]; 3.45 [1H, t, NH]; 3.30–3.95 [4H, m, C<sup>8</sup>H<sub>2</sub>, C<sup>9</sup>H<sub>2</sub>]; 3.01 [1H, ddd, J<sub>10-11ax</sub> = J<sub>10-15</sub> = 10.5, J<sub>10-11eq</sub> = 4.50 Hz, C<sup>10</sup>H]; 1.20–2.20 [2H, m, C<sup>11</sup>H<sub>2</sub>; 1H, m, C<sup>12</sup>H; 2H, m, C<sup>13</sup>H<sub>2</sub>; 2H, m, C<sup>14</sup>H<sub>2</sub>; 1H, m, C<sup>15</sup>H; 1H, m, C<sup>16</sup>H]; 0.83, 0.70 [3H, d, J<sub>17-16</sub>/J<sub>18-16</sub> = 6.00, 7.20 Hz, C<sup>17</sup>H<sub>3</sub>/C<sup>18</sup>H<sub>3</sub>]; 0.81 [3H, d, = 9.00 Hz]; 0.83 [3H, d, J<sub>19-12</sub> = 5.90 Hz, C<sup>19</sup>H<sub>3</sub>]; 2.80 [1H, tt, C<sup>20</sup>H]; 1.15 [3H, d, C<sup>21</sup>H<sub>3</sub>/C<sup>22</sup>H<sub>3</sub>]. <sup>13</sup>C NMR (CDCl<sub>3</sub>): 174 [C<sup>1</sup>]; 158 [C<sup>2</sup>]; 122 [C<sup>3</sup>]; 135 [C<sup>4</sup>]; 127 [C<sup>5</sup>]; 137 [C<sup>6</sup>]; 107 [C<sup>7</sup>]; 68.5 [C<sup>8</sup>]; 37.8 [C<sup>9</sup>]; 78.6 [C<sup>10</sup>]; 39.5 [C<sup>11</sup>]; 30.5 [C<sup>12</sup>]; 33.5 [C<sup>13</sup>]; 22.6 [C<sup>14</sup>]; 47.2 [C<sup>15</sup>]; 24.7 [C<sup>16</sup>]; 19.9, 15.5 [C<sup>17</sup>/C<sup>18</sup>]; 22.9 [C<sup>19</sup>]; 39.8 [C<sup>20</sup>]; 24.6, 24.5 [C<sup>21</sup>/C<sup>22</sup>]. IR: 3,295 ν(N–H); 1,635, 1,593 ν(C=O); 1,513 ν(C=C).

### Synthesis of copper compounds

The method described by us previously [17] was adapted to prepare (**13**)–(**21**). A solution of the appropriate ligand in water/ethanol mix (1:1) was added to a well-stirred solution of copper(II) acetate in water/ethanol mix (1:1). After stirring for 2 h, the solution was refluxed and stirred for another 2 h and then left at room temperature.

#### Cu(2-methylaminotroponate)<sub>2</sub> (**13**)

A dark green solid was obtained using (**1**) (0.81 g, 5.99 mmol) and copper acetate (0.60 g, 2.99 mmol). Yield 99 % (0.98 g). Recrystallisation was from ethanol. Analysis for (**13**): Found [calc. for C<sub>16</sub>H<sub>16</sub>N<sub>2</sub>O<sub>2</sub>Cu]: C 57.9 (57.9); H 4.86 (4.82); N 8.40 (8.44) %. IR: 1,596, 1,569 ν(C=O); 1,518 ν(C=C).

#### Cu(2-ethylaminotroponate)<sub>2</sub> (**14**)

Following the same procedure using (**2**) (0.40 g, 2.68 mmol) and copper acetate (0.27 g, 1.34 mmol), (**14**) was obtained. Recrystallisation from chloroform yielded green crystals (0.46 g, 95 %) suitable for X-ray analysis.

Analysis for (14): Found [calc. for  $C_{18}H_{20}N_2O_2Cu$ ]: C 60.1 (60.0); H 5.7 (5.6); N 7.7 (7.8) %. IR: 1,595, 1,540  $\nu(C=O)$ ; 1,513  $\nu(C=C)$ .

#### Cu(2-*n*-propylaminotroponate)<sub>2</sub> (15)

Using the above methodology with (3) (0.55 g, 2.56 mmol) added to copper acetate (0.26 g, 1.28 mmol). Following this, a green solid was obtained (0.55 g, 86 %). Analysis for (15): Found [calc. for  $C_{20}H_{24}N_2O_2Cu$ ]: C 63.1 (63.5); H 4.8 (4.9); N 5.8 (5.7) %. IR: 1,597, 1,570  $\nu(C=O)$ ; 1,516  $\nu(C=C)$ .

#### Cu(2-benzylaminotroponate)<sub>2</sub> (16)

Using the above methodology with (4) (0.29 g, 1.37 mmol) and copper acetate (0.14 g, 0.68 mmol). A green solid was obtained (0.32 g, 95 %). Analysis for (16): Found [calc. for  $C_{28}H_{24}N_2O_2Cu$ ]: C 68.9 (69.5); H 5.1 (5.0); N 5.8 (5.8) %. IR: 1,593, 1,570  $\nu(C=O)$ ; 1,513  $\nu(C=C)$ .

#### Cu(2-*n*-butylaminotroponate)<sub>2</sub> (17)

A green solid was obtained using (5) (0.46 g, 2.59 mmol) and copper acetate (0.26 g, 1.29 mmol). Yield 0.46 g, 86 %. Analysis for (17): Found [calc. for  $C_{22}H_{28}N_2O_2Cu$ ]: C 63.2 (63.5); H 6.8 (6.7); N 6.8 (6.7) %. IR: 1,597, 1,569  $\nu(C=O)$ ; 1,515  $\nu(C=C)$ .

#### Cu(2-methylamino-4-*i*-propyltroponate)<sub>2</sub> (18)

A green solid was obtained (0.21 g, 42 %) from (6) (0.50 g, 2.82 mmol) added to copper acetate (0.28 g, 1.41 mmol). Analysis for (18): Found [calc. for  $C_{22}H_{28}N_2O_2Cu$ ]: C 63.6 (63.5); H 6.7 (6.7); N 6.5 (6.7) %. IR: 1,606, 1,557  $\nu(C=O)$ ; 1,512  $\nu(C=C)$ .

#### Cu(2-*n*-propylamino-4-*i*-propyltroponate)<sub>2</sub> (19)

A green solid was obtained using (7) (0.50 g, 2.44 mmol) and copper acetate (0.24 g, 1.22 mmol). Yield 0.36 g, 62 %. Analysis for (19): Found [calc. for  $C_{26}H_{36}N_2O_2Cu$ ]: C 66.9 (66.2); H 8.1 (7.6); N 5.7 (5.9) %. IR: 1,591, 1,578  $\nu(C=O)$ ; 1,512  $\nu(C=C)$ .

#### Synthesis of Cu(2-menthoxyethylaminotroponate)<sub>2</sub> (20)

Using the methodology described earlier for 13–19 [17], (11) (0.36 g, 1.12 mmol) was reacted with copper(II) acetate (0.12 g, 0.60 mmol) to produce a dark green precipitate on standing (0.21 g, 27 %). Recrystallisation from hot acetone gave pale green crystals suitable for crystallographic analysis. Analysis for (20): Found [calc.

$C_{38}H_{56}O_4N_2Cu$ ]: C 68.1 (68.3); H 8.4 (8.4); N 4.1 (4.2) %. IR: 1,592, 1,572  $\nu(C=O)$ ; 1,509  $\nu(C=C)$ .

#### Synthesis of Cu(2-menthoxyethylamino-4-*i*-propyltroponate)<sub>2</sub>·H<sub>2</sub>O (21)

In a manner similar to that described for (20), (12) (1.00 g, 2.75 mmol) was reacted with copper(II) acetate (0.29 g, 1.45 mmol). A dark green precipitate was produced on standing. Crystals were obtained from hot toluene, but were too small for crystallographic analysis. Analysis for (21): Found [calc.  $C_{44}H_{70}O_5N_2Cu$ ]: C 68.4 (68.6); H 9.0 (9.1); N 3.4 (3.6) %. IR: 3,200  $\nu(O-H)$ ; 1,595, 1,570  $\nu(C=O)$ ; 1,511  $\nu(C=C)$ .

### X-ray crystallography

Experimental details relating to the single-crystal X-ray crystallographic studies are summarised in Table 1. For all structures, data were collected on a Nonius Kappa CCD diffractometer at either 150(2) (5, 20) or 170(2) K (14) using

**Table 1** Crystallographic data for 5, 14 and 20

Compound reference	5	14	20
Chemical formula	$C_{14}H_{13}NO$	$C_{18}H_{20}CuN_2O_2$	$C_{38}H_{56}CuN_2O_4$
Formula mass	211.25	359.90	668.38
Crystal system	Monoclinic	Monoclinic	Monoclinic
<i>a</i> /Å	5.5780(1)	9.9188(5)	17.9475(9)
<i>b</i> /Å	9.3640(2)	5.1220(1)	5.2632(2)
<i>c</i> /Å	20.8230(3)	17.0174(6)	19.2602(9)
$\beta$ /°	95.0710(9)	115.424(3)	101.034(2)
$V/\text{Å}^3$	1,083.38(3)	780.83(5)	1,785.71(14)
Temperature/K	150(2)	170(2)	150(2)
Space group	$P2_1/c$	$P2_1/c$	$P2_1$
Z	4	2	2
No. of reflections measured	14,213	5,203	17,173
No. of independent reflections	3,152	2,260	5,971
$R_{\text{int}}$	0.0393	0.0306	0.0961
Final $R_1$ values ( $I > 2\sigma(I)$ )	0.0534	0.0283	0.0575
Final $wR(F^2)$ values ( $I > 2\sigma(I)$ )	0.1296	0.0795	0.1308
Final $R_1$ values (all data)	0.0630	0.0302	0.0736
Final $wR(F^2)$ values (all data)	0.1342	0.0814	0.1414
Goodness of fit on $F^2$	1.122	1.077	1.060
Flack parameter			−0.014(13)
CCDC	990207	990208	990209

Mo- $K_{\alpha}$  radiation ( $\lambda = 0.71073 \text{ \AA}$ ). Structure solution followed by full-matrix least squares refinement was performed using the WinGX-1.70 suite of programmes [27], incorporating SHELXS and SHELXL within the SHELX-2013 suite of software [28]. Corrections for absorption (multi-scan) were made in all cases. Specific details: **14**: the asymmetric is one-half of the molecule, and the remainder generated by an inversion centre located at the metal.

### Antibacterial activity


*Staphylococcus waneri* was cultured in BHI broth (100 ml, Oxoid, UK) for 24 h at 37 °C (20 % CO<sub>2</sub>), transferred into centrifuge tubes and spun at 3,500 rpm for 7 min in a Mistral 1,000 centrifuge. The supernatant was decanted, and remaining pellet re-suspended in phosphate-buffered saline (PBS) (5 ml). This process was repeated twice, and

the culture finally re-suspended in PBS (2 ml). The optical density at 610 nm (OD<sub>610</sub>) was then adjusted to 1.0 using PBS.


A known weight of compound was dissolved in DMSO to make a 10 % solution of the active formulation. 200  $\mu$ L of this was then added to the bacterial suspension (190  $\mu$ L) and left for 1 min. After decanting, the culture was rinsed 3 times with water (200  $\mu$ L) and BHI broth (200  $\mu$ L) was added followed by mineral oil (80  $\mu$ L), and the resultant incubated at 37 °C in a Dynatec plate reader. The absorbance at 610 nm was monitored every fifteen minutes for 18 h, and the time taken for culture regrowth to return OD<sub>610</sub> to 0.5 noted.

### Results and discussion

A range of 2-alkylaminotropone ligands (**1–7**) were synthesised from a straightforward reaction between tosylated tropolone



**Scheme 1** Protocols for the synthesis of 2-alkylaminotropone ligands




**Scheme 2** Synthetic sequence for the formation of menthol-flavoured 2-alkylaminotropone ligands

and the appropriate amine (Scheme 1). For those amines that are commercially available as aqueous solutions ( $\text{RNH}_2$ ,  $\text{R} = \text{Me}$ ,  $\text{Et}$ ), the reactions were carried out in  $\text{CH}_2\text{Cl}_2$ , as this facilitates separation of the product into the organic layer. For the remaining amines ( $\text{R} = \text{C}_3\text{H}_7$ ,  $\text{C}_4\text{H}_9$ ,  $\text{C}_6\text{H}_5\text{CH}_2$ ), the reactions were carried out in DMSO. In the cases of **6** and **7**, tosylated hinokitiol was prepared as a mixture of isomers (a, b) in the ratio 9:1 [29], which leads to a mixture of isomeric 2-alkylamino-4-*i*-propyltropone products; no attempt to separate these isomers was made, and the mixture was used without further purification. However, this complication manifested itself in the purification of the alkylaminotropone ligands, which proved more challenging than for **1–5**. In the cases of **6** and **7**, while the ligand identity was confirmed by NMR, purification was effected by the formation of the copper complexes **18** and **19**. Related 2-alkylamino-4-*i*-propyltropone where  $\text{R} = \text{Et}$ , *n*-Bu, Bz were impure as prepared, and while the purity of the resulting copper complexes was enhanced, satisfactory microanalytical data could not be obtained.

In addition to these unfunctionalised substituents, we have also prepared analogues containing the flavourant, menthol (Scheme 2). This follows a sequence in which menthyoxyacetic acid (**8**) is converted to the acetyl chloride and then the amide (**9**) using aqueous ammonia. **9** was then reduced to the amine (**10**), a colourless oil, with excess  $\text{LiAlH}_4$  [26] and which was in turn used in the protocols of Scheme 1 to afford the flavourant-functionalised 2-alkylaminotropones (**11**) and (**12**).

NMR data for **1–7** are unexceptional, but show the presence of the amine  $\text{R}'$  along with resonances for the tropone ring in the correct ratios, while data for **11**, **12** confirm the linking of the 2-menthoxyethylamine with the tropolone. Both these compounds, brown oils, were isolated as monohydrates in analytically pure form after extraction from the reaction mixture; data for the major isomer (note: Scheme 1) are given in the Experimental.


**1–7**, **11**, **12** all show  $\nu(\text{NH})$  and  $\nu(\text{C}=\text{O})$  in the ranges 3,293–3,307 and 1,629–1,637  $\text{cm}^{-1}$ , respectively, in their IR spectra. The structure of **5** has also been determined (Fig. 1) and is that of a  $\text{N}-\text{H}\cdots\text{O}=\text{C}$  hydrogen-bonded dimer, similar to analogues with  $\text{R}' = 2,6\text{-}i\text{-Pr}_2\text{C}_6\text{H}_3$  [30] and  $4\text{-F-C}_6\text{H}_4$  [31],



**Fig. 1** The asymmetric unit of **5** showing the labelling scheme used in the text; thermal ellipsoids are at the 50 % level. Selected geometric data:  $\text{O}(1)-\text{C}(1)$  1.2573(18),  $\text{N}(1)-\text{C}(2)$  1.3457(18),  $\text{N}(1)-\text{C}(8)$  1.449(2),  $\text{C}(1)-\text{C}(7)$  1.432(2),  $\text{C}(1)-\text{C}(2)$  1.485(2),  $\text{C}(2)-\text{C}(3)$  1.3982(19),  $\text{C}(3)-\text{C}(4)$  1.398(2),  $\text{C}(4)-\text{C}(5)$  1.375(2),  $\text{C}(5)-\text{C}(6)$  1.398(2),  $\text{C}(6)-\text{C}(7)$  1.376(2) Å,  $\text{C}(2)-\text{N}(1)-\text{C}(8)$  125.09(12),  $\text{C}(2)-\text{N}(1)-\text{H}(1)$  115.2(13),  $\text{C}(8)-\text{N}(1)-\text{H}(1)$  119.3(13)°. Hydrogen bond:  $\text{H}(1)-\text{O}(1)$  2.9601(16),  $\text{N}(1)\cdots\text{O}(1)$  2.19(2) Å,  $\angle\text{N}(1)-\text{H}(1)\cdots\text{O}(1)$  145.8(17)°. Symmetry operation:  $1-x, -y, -z$


though differing from the monomeric  $\text{R}' = t\text{-Bu}$  [32], the latter structure presumably dictated by packing of the non-planar  $\text{R}'$ . Like the other reported dimers, the  $\text{C}=\text{O}$  [1.2573(18) Å] and  $\text{C}(2)-\text{N}(1)$  [1.3457(11) Å] in **5** suggest some lengthening with respect to the monomeric  $\text{R}' = t\text{-Bu}$  derivative [1.242(4), 1.348(4) Å, respectively] as anticipated [32], though the large errors in these data preclude definitive comment. The C–C bonds around the ring between  $\text{C}(2)-\text{C}(7)$  [1.375(2)–1.398(2) Å] evidence delocalisation, while the two C–C(1) bonds are notably longer [ $\text{C}(1)-\text{C}(2)$  1.485(2),  $\text{C}(1)-\text{C}(7)$  1.432(2) Å]. The nitrogen atom shows a degree of planarity via its bond angles [115.2(13)°–125.09(12)°] implying delocalisation of the lone electron pair.

Copper derivatives (**13–21**) of the aminotropones described above were prepared by reaction of the above



$\text{R} = \text{H}$ ,  $\text{R}' = \text{Me}$  (**13**),  $\text{Et}$  (**14**), *n*-Pr (**15**), *n*-Bu (**16**), Bz (**17**),  $\text{MenOCH}_2\text{CH}_2$  (**20**)

$\text{R} = i\text{-Pr}$ ,  $\text{R}' = \text{Me}$  (**18**), *n*-Pr (**19**),  $\text{MenOCH}_2\text{CH}_2$  (**21**)





**Fig. 2** The asymmetric unit of **14** showing the labelling scheme used in the text; thermal ellipsoids are at the 50 % level. Selected geometric data: Cu–O 1.9322(9), Cu–N 1.9366(10), O–C(3) 1.2945(15), NC(1) 1.4623(15), NC(4) 1.3250(15); O–Cu–O' 180, O–Cu–N 82.35(4), NCu–N' 180, O–Cu–N' 97.65(4)°. Symmetry operation:  $1 - x, 1 - y, 1 - z$

ligands with Cu(II) acetate in aqueous ethanol:

The products are green solids, in which the IR  $\nu(\text{C}=\text{O})$  stretch has shifted 30–70  $\text{cm}^{-1}$  to lower wavenumber, indicating  $\text{C}=\text{O} \rightarrow \text{Cu}$  coordination. Furthermore,  $\nu(\text{NH})$  is absent in all these complexes, which, in the case of **21**, a combination of microanalysis and a broad  $\nu(\text{OH})$  observed in the IR at 3,200  $\text{cm}^{-1}$  suggest is a monohydrate.

The structures of both **14** (Fig. 2) and **20** (Fig. 3) have been determined. The structures of both compounds adopt four-coordinate square-planar geometry as seen in the only other simple analogous copper compound reported to date ( $\text{R}' = \text{Me}$ ) [18], though examples with  $\text{R}' = \text{Et}$  along with further functionalisation of the seven-membered ring [ $5\text{-C}_5\text{H}_4\text{FeC}_5\text{H}_5$ ] also adopt this arrangement [19]. In the case of **14**, the molecule is centrosymmetric about an inversion centre at the metal, while **20** lacks this symmetry by virtue of the flexible pendant group on nitrogen. The Cu–O and

**Fig. 3** The asymmetric unit of **20** showing the labelling scheme used in the text; thermal ellipsoids are at the 50 % level. Selected geometric data: Cu–O(1) 1.931(5), Cu–O(3) 1.923(5), Cu–N(1) 1.926(6), Cu–N(2) 1.929(6), O(1)–C(1) 1.304(9), O(3)–C(21) 1.297(8), N(1)–C(7) 1.317(8), N(2)–C(27) 1.340(9); O(1)–Cu–O(3) 179.7(2), O(1)–Cu–N(1) 82.3(2), O(1)–Cu–N(2) 96.8(2), O(3)–Cu–N(1) 98.0(2), O(3)–Cu–N(2) 82.9(2), N(1)–Cu–N(2) 178.6(3)°



**Fig. 4** Antibacterial activity of new copper 2-alkylaminotroponate complexes **13**, **17**, **20** and **21** along with bis(ethylmaltolato) copper(II) [16] and a commercial toothpaste (Signal) for comparison. The plot shows the time taken for a culture of *Staphylococcus waneri* to regrow to half its original size after treatment with the agent

Cu–N bond lengths (Figure captions) are similar to those in **13** reported by others [18]. All three copper aminotropones show a general lengthening of the C=O by ca. 0.04 Å on complexation, while the NC(ring) bond contracts by ca. 0.02 Å, though in **20** these two C–N bonds are tending towards distinction [1.319(8), 1.340(9) Å], again reflecting the lower symmetry of the species; the nitrogen remains essentially planar in all cases. In the lattice, both **14** and **20** adopt a “slipped stack” arrangement with a  $\pi$ – $\pi$  separation of ca. 4.2 Å.

The antibacterial activity of **13**, **17**, **20** and **21**, along with data for the bis(ethylmaltol)copper complex [16] for comparison, is shown in Fig. 4. This single species biofilm assay (against *Staphylococcus waneri*) is an in vitro model of plaque inhibition effects. The results obtained from this test reflect the antibacterial activity of the complexes and are plotted as graphs of absorbance at 610 nm against time, which shows the delay in biofilm regrowth after treatment. From this, the time taken to return to an  $\text{OD}_{610}$  of 0.5 is



noted; the longer the time for the bacteria to regrow after treatment, the more effective the agent is deemed to be. Figure 4 shows that all four 2-alkylaminotropones are more effective than the formulation in a commercial toothpaste, though none are as active as the copper derivative of ethylmaltol [16], a related but *O,O*-chelating ligand. Interestingly, the presence of the flavourant does not reduce the efficacy of the compounds; in fact, the activity is marginally enhanced for these two species over the other copper 2-alkylaminotropones.

### Supplementary information

CCDC 990207–990209 contains the supplementary crystallographic data for this paper (Table 1). These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).

### References

1. Arima Y, Nakai Y, Hayakawa R, Nishino T (2003) *J Antimicrob Chemother* 51:113
2. Morita Y, Matsumura E, Okabe T, Shibata M, Sugiura M, Ohe T, Tsujibo H, Ishida N, Inamori Y (2003) *Biol Pharm Bull* 26:1487
3. Inamori Y, Sakagami Y, Morita Y, Shibata M, Sugiura M, Kumeda Y, Okabe T, Tsujibo H, Ishida N (2000) *Biol Pharm Bull* 23:995
4. Inamori Y, Tsujibo H, Ohishi H, Ishii F, Mizugaki M, Aso H, Ishida N (1993) *Biol Pharm Bull* 16:521
5. Nomiya K, Onodera K, Tsukagoshi K, Shimada K, Yoshizawa A, Tada-aki I, Sugie A, Tsuruta S, Sato R, Kasuga C (2009) *Inorg Chim Acta* 362:43
6. Irving RJ, Post ML, Povey DC (1973) *J Chem Soc, Dalton Trans* 2:697
7. Berg J-E, Pilotti A-M, Söderholm A-C, Karlsson B (1978) *Acta Crystallog B* 34:3071
8. Griffith WP, Pumphrey CA, Skapski AC (1987) *Polyhedron* 6:891
9. Avdeef A, Costamanga JA, Fackler JPJ (1974) *Inorg Chem* 13:1854
10. Davis AR, Einstein FWB (1974) *Inorg Chem* 13:1880
11. Lockhart TP, Davidson F (1987) *Organometallics* 6:2471
12. Mori A, Yamamoto S-I, Takemoto M, Ujiie S (2002) *Chem Lett* 2:186
13. Mori A, Mori R, Takemoto M, Yamamoto S-I, Kuribayashi D, Uno K, Kubo K, Ujiie S (2005) *J Mater Chem* 15:3005
14. Creeth J, Molloy KC, Wright P (2000) Oral care compositions. Patent WO 00/16736
15. Barret MC, Mahon MF, Molloy KC, Wright P (2000) *Main Group Met Chem* 23:663
16. Barret MC, Mahon MF, Molloy KC, Steed JW, Wright P (2001) *Inorg Chem* 40:4384
17. Barret MC, Mahon MF, Molloy KC, Wright P, Creeth JE (2002) *Polyhedron* 21:1761
18. Steyl G, Muller TJ, Roodt A (2010) *Acta Crystallog Sect E Struct Rep Online* 66:m1508
19. Miyake Y, Watanabe S, Aono S, Nishinaga T, Miyazaki A, Enok T, Miyasaka H, Otani H, Iyoda M (2008) *Chem Commun* 44:6167
20. Nishinaga T, Aono T, Iomura E, Watanabe S, Miyake Y, Miyazaki A, Enoki T, Miyasaka H, Otani H, Iyoda M (2010) *Dalton Trans* 39:2293
21. Meyer N, Lohnwitz K, Zulys A, Roesky PW, Dochnahl M, Blechert S (2006) *Organometallics* 25:3730
22. Reddington MV (1998) *J Chem Soc Perkin Trans II* 143
23. Rasika-Dias HV, Jin W (1996) *Inorg Chem* 35:6546
24. Biggi G, Del Cima F, Pietra F (1973) *J Am Chem Soc* 95:7101
25. Cochran TG, Huitric AC (1971) *J Org Chem* 36:3046
26. Minardi G, Schenone P (1968) *Farmaco Ed Sci* 23:1181
27. Farrugia LJ (1999) *J Appl Crystallogr* 32:837
28. Sheldrick G (2008) *Acta Cryst A* 64:112
29. Yanagisawa T, Kosakai K, Yasunami M, Takase K (1990) *Chem Pharm Bull* 38:3355
30. Hicks FA, Jenkins JC, Brookhart M (2003) *Organometallics* 22:3533
31. Steyl G (2007) *Acta Crystallogr Sect E: Struct Rep Online* 63:o4353
32. Siwatch RK, Kundu S, Kumar D, Nagendran S (2011) *Organometallics* 30:1998